Tegenaria medicinalis Hentz is near *Coelotes atropos*, possibly the same species.

Micrommata canadensis Hentz is represented by the common *Ocyale mirabilis*.

Pylarus bicolor Hentz belongs to the genus *Ariadne*, and is very near *A. insidiatrix*.

J. H. Emerton.

Leipzig, Dec. 16, 1875.

Arctic Lepidoptera in the White Mountains.

Mr. Roland Thaxter sends me a male specimen, taken by Mr. B. P. Mann, above the tree-line, on Mount Washington, which seems to be the first indication of the probable occurrence of *Laria Rossii* Curtis on the mountains, and still further to illustrate the relations between the Arctic and Mount Washington faunæ. The specimen is a male, smaller than Curtis' figure, with the dark band on the secondaries so broadened and diffused as to leave but a discal yellowish patch. The markings on the fore wings seem quite similar, and, in the absence of a larger series of specimens, the more closely scaled and rather darker Mount Washington specimen should not be held to indicate a distinct species. It appears that arctic species are subject, as a rule, to unusual variation. My specimens of *Pachnobia carnea* from Labrador strongly vary in depth of color and distinctness of marking; a specimen, undoubtedly of this species, sent me by Mr. Morrison, from Mount Washington, is very pale. We shall probably find, when more material is collected together, that we have, besides *Anarta melanopa*, *Agrotis islandica*, and *Pachnobia carnea*, also the Bombycid *Laria Rossii* to add to our list of Arctic Lepidoptera found in the Mountains of New Hampshire.

A. R. Grote.

Chrysalis with attached Larval Head.

Mr. Roland Thaxter brought me, Sept. 27th, 1875, the living chrysalis of *Euphœades Troilus*, with the head of the caterpillar still remaining upon it. The head is split, as ordinarily when cast for the last time, and the two hemispheres hug the

base of the right ocellar prominence of the chrysalis, the left ocellar prominence being greatly aborted and entering the heart of the left cephalic hemisphere; the right hemisphere is nothing but a pellicle, connected only with the opposite half, but the left seems to be closely soldered to the insect, although the edges are free almost or quite throughout; in addition, the pellicle and hardened dorsal shield of the first segment of the larva are present, directly behind the left hemisphere; the frontal triangle and all the mouth parts of the larval head remain naturally attached to the left hemisphere; there is nothing else peculiar to the creature excepting its total lack of maxillary and antennal cases, the empty trough for the reception of the former being particularly conspicuous, while that of the latter is closed on the distal half, or, on the right side, even more than that. This makes it all the more probable that the larval head has still an organic connection with the insect. The chrysalis was raised in confinement.

Samuel H. Scudder.

BIBLIOGRAPHICAL RECORD.

Authors and Societies are requested to forward their works to the Editor at the earliest date possible. We ask our readers to inform us of the publication especially of those works which are not generally consulted by entomologists.

B. Pickman Mann.

(Continued from page 128.)

The Cinc. Quart. Journ. Sci., vol. i (continued), contains Nos. 337-339, and vol. ii contains Nos. 340 to 344.

* 337. V. T. CHAMBERS. Prof. Frey, of Zürich, and some American Teneina. p. 193-211.

Criticism of Frey & Boll's * Nordamerikanische Tineen, Stett. Entom. Zeit., 1873, p. 201-224; discussion of the species therein mentioned and described. [See Rec., No. 339.]

* 338. CYRUS THOMAS, Ph.D. From the Introduction to the Synopsis of the Acrididæ of North America. p. 253-269.

Reprint from Thomas' * Syn. Acrid. N. A. (1873), p. 9-25; external and internal structure and terminology of Acrididæ; oviposition; stridulation.

* 339. V. T. CHAMBERS. Notes and Errata on a former paper, on Prof. Frey, and some American Teneina. p. 338-340.

Addition to the article recorded above, Rec., No. 337.