Additions to the List of Newton Noctuidæ.

The following names are to be added to those given in the list on p. 34-39 of the present volume of *PSYCHE*; the abbreviations are the same as in that list.

<i>Diphtera fallax</i> . R. l. Jl.	<i>Glaea viatica</i> . S. Oct. 19.
<i>Apatela Radcliffei</i> . R.; raised. Jn.	" <i>inulta</i> . Several s. Sp.
Lrv. Prunus, Sp.	" <i>deleta</i> . (See below.)
<i>Agrotis badicollis</i> . R. l. Au.	<i>Scopelosoma Graefiana</i> . Several s.
" <i>fennica</i> . Several s. Sp.	Sp. Oct. Nov.
" <i>albipennis</i> . Two s. l. Sp.	<i>Lithophane semiusta</i> . R. s. Sp. 28,
" <i>incivis</i> . R.; three s. Sp.	Oct. 18.
" <i>trabalis</i> . Cocoon found under bark of <i>Pinus strobus</i> in April.	" <i>Georgii</i> . One s. Sp.
Imago appeared in June.	" <i>laticinerea</i> . C. s. Sp.
<i>Hadena impuls</i> a. R. l. Jl.	Oct.
" <i>apamiformis</i> . R. l. Jl. Au.	" <i>tepida</i> . R. s. Sp.
" <i>vultuosa</i> . C. l. Au.	" <i>querquera</i> . R. s. Sp.
" <i>leucoscelis</i> . R. s. Au.	" <i>Baileyi</i> . (Grote, Can.
<i>Dryobota stigmata</i> . R. s. Sp.	Entom., ix, 86). Several s. Sp. Oct. Nov.
<i>Laphygma frugiperda</i> . C. s. Sp.	" <i>viridipallens</i> ¹ (Grote,
<i>Prodenia flavimelia</i> . R. s. Sp.	Can. Entom., ix, 215).
" <i>lineatella</i> . R. l. Jl. [<i>P.</i>	R. s. Sp. Oct.
<i>ornithogalli</i> is to be stricken from the previous list.]	<i>Plusia formosa</i> . One l. Jl.
<i>Gortyna limpida</i> . R. l. Sp.	" <i>verruca</i> . One s. Au.
<i>Arzama densa</i> . R. l. Au.	<i>Galgula hepara</i> . One s. Nov.
<i>Nonagria</i> sp. One l. Au.	<i>Catocala neogama</i> . R. s. q. Au.
<i>Doryodes acutaria</i> . R. l. Au.	" <i>piatrix</i> . R. q. s. Au. Sp.
<i>Platysenta atriciliata</i> . R. l. Jn. Jl.	<i>Homoptera Woodii</i> (Grote, Can.
<i>Parastichtis minuscula</i> . R. l. s. Au.	Entom., ix, 88-89).
Sp.	R. l. Jl.
	<i>Spargoloma 6-punctata</i> . R. l. Jl.

The new species mentioned in the above list is described as follows, by Mr. Grote:—

Glaea deleta n. sp. ♂. Belongs, with *G. tremula*, to the section which I have called *Epiglaea*, in which the thorax has a sharp scale-ridge. Resembles *G. viatica* at first sight, but belongs with *G. pastillicans* to a different section of the genus. Expanse, 46 mm. Of the same color as *G. viatica*, but smaller, and with the transverse posterior line punctate, and the subterminal also broken into spots. Stigmata pale ringed, well sized, concolorous; reniform stained at base. The markings are faint. Hind wings fuscous above; with discal spot and line beneath. Antennæ white at base. Abdomen with ochery anal hairs. (*A. R. Grote.*)

Rare, at sugar, in September. Several (poor) specimens were taken at sugar in October.

Noctuae have been abundant here this fall. Besides the additional species of *Lithophane* mentioned above, I have taken all those given in my former list. The larvæ of *Lithophane disposita* and *L. Bethunei* were found last June at the roots of apple and willow trees.

R. Thaxter.