spicuous upon the margin of the posterior wings upon the upperside, only those just at the anal angle and the two between the median nervules reappear upon the underside. Of these the two at the anal angle are obscure, while those between the median nervules are large and very distinct. They are all crowned with red lines and have their black centres irrorated with shining green scales.

Expanse of wings 28 mm. Type in the collection of the author.

I have assigned this species in a gen-

eral way to *Lycaena* Fabr., feeling unable to refer it to any of the recent subdivisions of the genus instituted by the Indian lepidopterists, without such an examination of the neururation as I do not feel justified in making with only one specimen at my disposal. It is very different in appearance from any other species of the group known to me, and recalls *Pseudodipsas cephenes* Hew., so far as the coloration of the upperside of the secondaries is concerned.

ON AN IMPORTANT CHARACTER, HITHERTO LITTLE NOTICED, IN THE FAMILY BUPRESTIDAE.

BY FREDERICK BLANCHARD.

In Comstock's Introduction to Entomology, part i, the many excellencies of which and its fresh treatment of the subject lead us to hope for the early appearance of the succeeding parts, on page 18, there is given a figure of the underside of *Euchroma gigantea* Linn. in which is shown the antecoxal piece of the metasternum, separated by a short transverse suture, the ends arcuately bent towards and reaching the hind coxae. This appears to be the first distinct reference to this peculiarity of the Buprestidae so far as I can learn. Deyrolle, in his Buprestides de la Malasie, plate 4, figs. 3 and 4 exhibits the same thing, but in fig. 25, illustrating the underside of a species of *Pachyscelis*, the suture is not indicated. There is, however, no reference to the antecoxal piece in the text.

From the examination of a considerable number of genera in this family during the last few years, both native and foreign, it appears that this structure is always present and is of much greater importance in limiting the family than the connate first and second ventrals chiefly depended upon heretofore.

Although the members of the family Buprestidae are usually quite easily recognized, some early errors would have been avoided had this character been observed or appreciated; in our own fauna, notably in the case of the genus *Schizopus* Lec., for the reception of which a distinct family, the Schizopodidae, was created although afterwards suppressed.

It need hardly be said that the existence of an antecoxal piece, seen elsewhere among Coleoptera, so far as I know, only in the Adephaga, where the importance of its existence and of its modifications have been so skilfully demonstrated by Dr. Horn, does not necessarily imply any relationship of the Buprestidae to that series, but adds another, and a most impressive one, to the many known instances of the repetition of characters or structures in widely different families, which have been noticed by writers, and especially by Dr. Horn.

HARRISIMEMNA TRISIGNATA—I found two of these grotesque larvae on *Spiraea tomentosa*, Sept. 22nd, Northborough, Mass. The only food-plant given by Mr. Edwards is "*Syringa*." The larva bored into bits of rotten wood, and "backed out" with the chips. These chips were rolled into neat pellets of almost uniform size, very round,

and evidently kept in shape by some gummy secretion applied by the larva. Each pellet was about one-sixteenth inch in diameter and all could be rolled about without crumbling. When the burrow was finished the larva "backed in" and closed the opening with a thin, transparent, parchment-like door. It took over twenty-four hours to make the burrow and seal it. *C. G. Soule.*

MISCELLANEOUS NOTES.—Mr. C. P. Gillette, formerly entomologist at the Experiment Station at Ames, Iowa, has removed to Colorado to take a similar position at Fort Collins.

A continuation of Mr. J. H. Emerton's New England spiders appears in the last part of the Transactions of the Connecticut Academy. It concerns the Drassidae (33 sp.), Agalenidae (11 sp.), and Dysderidae (2 sp.), with six plates crowded with excellent figures; more than half the species are regarded as new.

Baron Osten Sacken has suggested in the February number of the Entomologists' monthly magazine a new classification of the families of nemocerous Diptera.

A timely and convenient catalogue of the described transformations of North American Coleoptera by W. Beutenmuller will be found in the January number of the journal of the New York microscopical society.

L'ABEILLE, the journal of entomology founded by the late abbé Marseul, and by him carried through twenty-six volumes, is to be continued by the Entomological Society of Paris, to which he bequeathed it. Mr. L. Bedel has been chosen editor, and the twenty-seventh volume is now in press. The journal, as formerly, will be devoted mainly to Old World Coleoptera, and the frequency of its issue will depend upon subscriptions and sales.

PROCEEDINGS OF SOCIETIES.

CAMBRIDGE ENTOMOLOGICAL CLUB.

8 June, 1888.—The 139th meeting was held at 156 Brattle St. Mr. J. H. Emerton was chosen chairman.

Mr. C. W. Woodworth exhibited speci-

mens of a species of *Typhlocyba* found on the rose, to which they are very injurious. He also showed a specimen of a species of Jassidae found on the apple. No jassids have before been found on that tree.

Mr. S. H. Scudder remarked on the number of subsegments in the larvae of butterflies. Mr. Scudder has applied this to the position of *Libythea*, which has been variously placed in the Pierinae, Nymphalidae, and Erycininae.

Mr. Scudder stated that most of the changes in the larvae of the Lepidoptera, such as moults, etc., occur before ten o'clock in the morning.

Mr. S. H. Scudder mentioned the finding of the larva of *Oeneis semidea* which had formed a very slight silken cocoon for pupating. It is now a pupa.

9 November, 1888.—The 140th meeting of the Club was held at 156 Brattle St. Mr. J. H. Emerton was chosen chairman.

Mr. J. H. Emerton exhibited some drawings of spiders made by himself and Mr. Edwin Sheppard for Dr. H. C. McCook of Philadelphia.

Mr. S. H. Scudder showed a collection of butterflies brought together to illustrate mimicry in this country. Considerable discussion of the subject of protective mimicry followed.

Prof. C. H. Fernald spoke of the work of the State experiment stations, established under the new act of Congress appropriating a sum of money to each state for the purpose, and especially of the Hatch Experiment Station of the Amherst Agricultural College, of which station he is entomologist.

Mr. S. H. Scudder read a paper for Miss Caroline G. Soule on a mode of preserving pupae of Sphingidae through the winter.

Mr. H. Hinkley spoke of forcing the early emergence of Sphingidae by keeping the pupae at room temperature.

Mr. S. H. Scudder described how Mr. S. L. Elliot kept his pupae through the winter, and mentioned the opportunity offered by a cold storage company in Boston for wintering hibernating insects.

Mr. H. Hinkley described a double cocoon of *Attacus promethea*.