Stage IV. Head rounded bilobed, full, pale wood brown, faintly mottled with red brown, the upper tubercles black; width 1.4 mm. Body rather slender, smooth, pale greenish brown, faintly, finely, longitudinally lined with whitish, the tubercles rather broadly black marked. A subventral blackish shade line posteriorly. The pale lines are numerous, edged finely with dark brown, crinkly and a little mottled, some of the edgings broader and blacker than the others.

Stage V. Head whitish, mottled, dotted in patches with gray, tubercles and sutures of clypeus black; width 1.7 mm. Body greenish white, slightly tinged with brown, a little lined and marbled with brownish and gray, as before; tubercles black, setae pale. Shields concolorous with the body. Variation in color as in the next stage.

Stage VI. Head rounded, full, slightly bilobed, the clypeus small, somewhat depressed; width 2.2 mm. Whitish green with gray dottings over the lobes, tubercles represented by black spots. Body cylindrical, smooth, uniform, colored as before. The bred larvae were rather brightly colored. Ground color pale, nearly all whitish green, the black dashes heavy subdorsally intersegmentally; orange shading on the cervical shield, and in the subdorsal pale line, which is the only distinct line. The orange shadings are above the black dashes, which in turn border the pale line below. Anal plate orange shaded; tubercles represented by black spots. Another collected example was uniformly brownish, heavily mottled, only the subdorsal line, and a few streaks showing the whitish green ground color.

Food plant oak. The larvae are colored like the bark, and probably rest upon it. A single brood in the year, the winter passed as pupa. Larvae from Brookhaven, Long Island, N. Y. Eggs, June 12th, mature larvae not till late September or October; the development very slow.

The descriptions of the larvae of *Therina* are in some confusion. In the 5th report of

the U. S. entomological commission are four descriptions, two of larvae on oak, and two on pine and spruce. I have shown above that the pupa described (p. 777) as "*seminudaria*" should be credited to *pellucidaria*, and probably the pupa described on p. 841 as "*ferviduria*" belongs to the same, judging from the food plant. On p. 186 "*ferviduria*" is described, possibly correctly; but more probably it is *fiscellaria* Gn., as the moths "reared from the live oak in Florida by Dr. Riley" (*i. e.* Mr. Koebele) are of that species, and Abbot's locality is southern. Following this is a description of "*endrophiaria*," but obviously incorrect, as it differs totally from Goodell's correct one, which is referred to, without comment on the marked discrepancy. The description is taken from the books of the Department of Agriculture (no. 3904), and the bred moth before me proves it to refer to *fiscellaria* Gn.

The larvae of *pellucidaria*, *athasiaria*, *fiscellaria* (and *ferviduria* also?) are alike at maturity, within the normal range of variation, so that they cannot be distinguished with certainty. As to the earlier stages, I am not yet fully informed; but hope to be able to bring out the characters before this series of papers is closed.

THE NAME *LEONIA*.—In *Psyche*, Oct., 1899, p. 416, I discussed *Leonia*, and its ally *Hornia*, without remembering that the name *Leonia* was proposed by Gray many years ago (1840) for a genus of Mollusca. There seems to be no alternative but to change the name of the meloid *Leonia*, so I will propose to substitute *Leonidia*, n. n. It is perhaps a matter of taste whether it is regarded as a distinct genus, or subgenus of *Hornia*.

T. D. A. Cockrell.

CORRECTION OF AN ERROR.—In *Psyche*, vol. vii, p. 252 (June, 1895, I described an interesting Tineid larva, feeding on *Rubus*, as "*Butalis basilaris* Zell," depending upon the supposed accuracy of the determination made for me by Prof. C. H. Fernald. The

name should be changed to *Schreckensteinia festaliella* Hübn. It will then not be an abnormality in the genus, the larva agreeing with the habits and appearance of the European form. (See Walsingham, Pter. Cal. and. Or., p. 1, 1880, and Meyrick, Handb. Brit. Lep., p. 690, 1895). I have not

seen a previous record of the occurrence of this species in the East. Of the other species of *Schreckensteinia*, *felicella* Wals. is said to live on *Orthocarpus* and *erythriella* Clem. in the racemes of *Rhus*.

Harrison G. Dyar.

A NEW VOLUME OF PSYCHE

begins in January, 1900, and continues through three years. The subscription price (payable in advance) is \$5.00 per volume, or \$2.00 per year, postpaid. Numbers are issued on the first day of each month. Libraries and individuals generally ordering through subscription agencies (*which only take annual subscriptions*) will please notice that it is cheaper to subscribe for the entire volume at once directly of us.—Any early volume can be had for \$5.00, unbound. Address Psyche, Cambridge, Mass.

Vols. 1-8, Complete, Unbound	=	=	=	=	=	\$37.00.
Vols. 1-8, and Subscription to Volume 9	=	=	=	=	=	\$41.00.
Vol. 8 contains about 450 pp. and 8 plates, besides other illustrations.						

The Butterflies of the Eastern United States and Canada.

With special reference to New England. By SAMUEL H. SCUDDER.

Illustrated with 96 plates of Butterflies, Caterpillars, Chrysalids, etc. (of which 41 are colored) which include about 2,000 Figures besides Maps and portraits. 1958 Pages of Text.

Vol. 1. Introduction; Nymphalidae.

Vol. 2. Remaining Families of Butterflies.

Vol. 3. Appendix, Plates and Index.

The set, 3 vols., royal 8vo. half levant, \$75.00 net.

HOUGHTON, MIFFLIN & CO., 4 Park St., Boston, Mass.

A. SMITH & SONS, 269 PEARL STREET, New York.

MANUFACTURERS AND IMPORTERS OF

GOODS FOR ENTOMOLOGISTS,

Klaeger and Carlsbad Insect Pins, Setting Boards, Folding Nets, Locality and Special Labels, Forceps, Sheet Cork, Etc. Other articles are being added, Send for List.