LIFE HISTORIES OF NORTH AMERICAN GEOMETRIDAE.—LIV.

BY HARRISON G. DYAR, WASHINGTON, D. C.

Hyperitis trianguliferata Packard. The life history of this species has been described by Professor G. H. French (Can. ent., xviii, 105, 1886) from Californian larvae, with four stages. I present my own descriptions, however, as I have noted some points not mentioned by him and the larva has not been otherwise referred to.

Egg. Elliptical, rounded, one diameter less, no flattening nor depression; truncation marked, oblique. Longitudinal ribs low, obscure, straight, few, about six visible on the broad side, joined by cross-striae, forming cells three times as wide as long; ends reticulate. Pale yellow; size $1 \times .7 \times .6$ mm.

Stage I. Head rounded, testaceous, a little darker on the vertex, eye black, mouth brown. Body moderate, not greatly elongate, rapidly looping or held rigid with a thread. Whitish, a broad dorsal band of smoky olivaceous and a still fainter ventral one, scarcely more than a shade. Feet pale; tubercles very small, brownish; setae moderate, glandular tipped; no shields.

Stage II. Head rounded, the lobes full, sordid whitish with black dottings on the lobes and around the mouth; width 6 mm. Body moderate, greenish, shaded with blackish olivaceous dorsally, with subdorsal and lateral lines indicated by white segmentary dashes. a white stigmatal line with black and gray spots, tubercle iv largest on joints 7 and 8. Venter orange tinted, obscurely lined and marked by blackish blotches at the tubercles. Tubercles roundedly elevated; setae obscure; feet pale with dark lines; a geminate ventral whitish band.

Stage III. Head whitish with pale brown dots forming bands curved transversely, antennae prominent, pale, ocelli black; width 1.1 mm. Body moderate, the segments enlarged centrally by the conical tubercles, especially joint 7. Pale brown; subdorsal line broken into white dashes, the longest one on joint 5, edged with black shading; sides shaded; substigmatal band broad, yellowish white, illy defined but distinct; a paler subventral line; centers of segments subventrally and venter medially black shaded; tubercles conical, black, ii prominent, especially on joint 7. Feet pale; setae very short; anal shield and leg plates pale.

Stage IV. Head round, cuneiform, the lobes full above, held flatly; light and brown mottled on the sides, pale over the face and vertex; width 1.7 mm. Body cylindrical, joints 2 and 3 depressed, their feet appressed, those of joint 3 erected; straight, stick-like, joint 7 humped dorsally, tubercles ii forming high cones on joints 7 and 9, less so on joints 6 and 8, small on joints 5, 11 and 12. Whitish green, washed with brown of light and dark; an oblique lateral streak on joints 3 and 4, the fronts and backs of the high tubercles, irregular marks and blotches on the sides, especially of joints 7 to 9 and a partly continuous subventral band. Ordinary markings lost. Feet of joint 10 brown behind, pale before; thoracic feet bright red shaded. Faint whitish washed longitudinal lines.

The larvae fed on currant (*Ribes sanguineum*); bred from eggs obtained at Kaslo, British Columbia.