LIFE HISTORIES OF NORTH AMERICAN GEOMETRIDAE.—LVIII.

BY HARRISON G. DYAR, WASHINGTON, D. C.

Mesoleuca vasiliata Guenée.

Egg. Elliptical, rounded, normal, the depression marked; all rounded, no flat areas; evenly reticulate with broad low lines, hexagonal, finely punctate shagreened all over. Pale waxy yellow; size .9 x .6 x .5 mm.

Stage I. Head rounded, whitish testaceous, eye black, mouth brown. Body thin and slender but not rapid in motion, becoming rigid erect when disturbed; white, the ends (joints 2-4, 10-13) yellowish tinged in certain lights from their retraction. No marks; no shields; tubercles concolorous, obscure; setae stiff, pale, rather conspicuous.

Stage II. Head rounded bilobed, the apex of clypeus depressed, shining light yellow, ocelli black; width .5 mm. Body slender, rather elongate, normal, coarsely annulate. Setae rather coarse, white; no shields; tubercles invisible; no marks, food green.

Stage III. Head round, bilobed, free, lobes full, shining ochraceous luteous, clypeus depressed, ocelli black; width .8 mm. Body moderate, green from the food, unmarked, the folded incisures whitish; tubercles a little elevated, round, shining, colorless; feet pale; no shields; setae short, pale; subventral fold prominent.

Stage IV. Head rounded, shining, caramel luteous, setae brown, ocelli black; width 1.1 mm. Body cylindrical, slender, the central segments well elongated. Translucent green, incisures narrowly folded, yellowish; tubercles small, round, white, elevated, distinct. The dorsal vessel shows darker and the tracheal line white by transparency. No marks; no shields; thoracic feet brown. About joint 9 a broad stigmatal band begins and runs to the anal feet. It is also slightly indicated on thorax. Setae moderate.

Stage V. Head long, bilobed, green, shagreened, the lobes full, pale greenish, the epistoma, and antennae whitish with a short white band before the eyes; width 1.7 mm. Body moderate, elongate, cylindrical, the segments wrinkly annulate, anal feet large, projected outwardly. Green, anal feet and dorsum of joint 13 shaded in vinous; tubercles elevated, rounded, white like small buttons. Very faint longitudinal white washed lines; dorsal vessel dark green; setae moderately long, distinct. Later the head acquired a red flush over the lobes.

Food plant thimbleberry (*Rubus nutkanus*); the hatched larvae ate this plant and a wild larva was found on it. Larvae from Kaslo, British Columbia.