fourth and one and three-fourths their diameters, respectively. Distal enlargement subcylindric, with a length one-fourth greater than its diameter; terminal segment, distal enlargement somewhat produced, irregularly fusiform. Palpi; first segment short, with a length one-half greater than its diameter, the second nearly twice the length of the first, the third a little longer than the second and the fourth three-fourths longer than the third, somewhat dilated. Mesonotum slaty gray, the grayish submedian lines sparsely haired. Scutellum and postscutellum dark gray. Abdomen thickly haired, dark brown. Wings hyaline, costa light straw. Halteres yellowish basally, fuscous apically. Coxæ and femora mostly fuscous yellowish; tibiæ and tarsi mostly dark brown; claws moderately stout, strongly curved, the pulvilli shorter than the claws. Genitalia; dorsal plate deeply and triangularly divided, the lobes narrowly triangular; ventral plate broad, broadly and roundly emarginate. Type: Cecid. 1422.

A VERNAL BEE (*COLLETES INÆQUALIS* SAY).

BY JOHN H. LOVELL,
Waldoboro, Maine.

In Mr. Frost's interesting note on *Tricrania sanguinipennis* Say, which was published in *PSYCHE*, December, 1912, the name of the bee near whose burrows the beetles were found is given (through an inadvertence on my part) as *Colletes compactus* Cr., when it should be *C. inæqualis* Say. The two species, while bearing a general superficial resemblance, may be distinguished by the difference in the sculpturing of the enclosure on the metathorax or prepodeum; and by the fact that the former is an autumnal species and the latter a vernal species, some three months intervening between the disappearance of *C. inæqualis* and the appearance of *C. compactus*.

I have this season received both sexes from Mr. Frost, to whom I am indebted for the following notes. The bees were abundant, and had constructed numerous burrows near the base of a sloping embankment, about three feet high, sparsely covered with grass. There were also nests where the ground was level and free from vegetation. As the burrows appeared to be unlined and the soil was sandy it was impossible to follow the tunnels to a depth of more than three or four inches. This species of *Colletes* in New England is on the wing for about two months, and is most commonly taken on the aments of the willows. Mr. Frost captured

his first specimens on March 23, and on the 30th observed many visits to the flowers of *Salix*. On March 31 a female was seen looking out from the mouth of nearly every burrow, probably waiting for a warmer temperature, for on April 6, when there was a severe snow squall, no bees were seen flying about the burrows; and a few which had ventured to fly to the willows were numbed by the cold.

Swenk states that he examined a female from Durham, N. H., taken Oct. 5, 1899, and has also seen a few other autumnal specimens of *C. inaequalis*. He regards them as individuals appearing prematurely, which normally would not have come out until the following spring.¹

Colletes compactus flies in New England from about the first of September to the middle of October. I most commonly find it on the inflorescence of the goldenrod, but have one specimen taken on the flowers of *Aster puniceus*.

THE LIFE HISTORY AND HABITS OF *SPALANGIA MUSCIDARUM* RICHARDSON, A PARASITE OF THE STABLE FLY.²

BY HARRY PINKUS.³

Bureau of Entomology, United States Department of Agriculture.

During the summer of 1912, while assisting in the study of the life history of the stable fly (*Stomoxys calcitrans* Linn.), at least two species of parasites were found breeding in large numbers in the puparia of this fly. These parasites belong to the family Pteromalidæ. One has been determined by Mr. C. H. Richardson as *Spalangia muscidarum* Richardson; the others have not been definitely identified. *S. muscidarum* was found most abundant while the other species appeared in smaller numbers. They appear to have similar breeding habits, although little has been done on the undetermined species.

During the investigation of the stable fly a lot of oat straw, placed in a pan and kept moist, was placed in the laboratory yard

¹ ("Specific Characters in the Bee Genus *Colletes*," Contributions from the Dept. of Ent., Univ. Neb., No. 1, p. 32.)

² Published by permission of the Chief of the Bureau of Entomology.

³ Deceased August 27, 1913, at Dallas, Texas.