EXPLANATION OF PLATE I.

In all figures only a portion of the terga (which are symmetrical) has been shown, since the missing portions are exactly like those figured.

Fig. 1. Terga and wing bases of a Plecopteron.

Fig. 2. Terga and wing bases of the Coleopteron *Photuris*.

Fig. 3. Terga and wing bases of *Embia major*.

Fig. 4. Terga and wing bases of the Dermapteron *Echinosoma*.

ON THE OCCURRENCE OF A MERMIS EPIDEMIC
AMONGST GRASSHOPPERS.¹

By R. W. GLASER and A. M. WILCOX.

While engaged in some investigations on grasshoppers, near Dummerston Station, southern Vermont, this past summer (1917), our attention was attracted to a high mortality amongst these insects (*Melanoplus atlantis* and *M. bivittatus*). The two species, especially *M. atlantis*, are extremely bad pests in this region of the country, attacking corn, wheat, oats and clover to such an extent that during certain summers the farmers become nearly frantic. Therefore, the high mortality amongst the grasshoppers, which appeared during the latter part of August and the early part of September, was exceedingly gratifying.

We soon discovered that this mortality was due to a species of Nematode belonging, as we supposed at the time, probably to the family Mermithidæ. Subsequently (Sept. 20, 25 and Oct. 6), we sent large shipments of these worms to Dr. N. A. Cobb, of Washington, D. C., for identification. Dr. Cobb was able to give us only a provisional identification on account of the utter absence of males in all of our shipments. We made collections of parasitized grasshoppers from a large variety of fields and as stated, sent a large number of specimens, but curiously enough no males were found. Dr. Cobb in a letter said: "Nothing I have learned would preclude your specimens from belonging to the same species as that referred to by Leidy under the name of *Mermis ferruginea*, which

¹ Contribution from the Entomological Laboratory of the Bussey Institution in coöperation with the U. S. Bureau of Entomology. Bussey Institution, No. 146.

he says was common in *Locusta carolina* near Philadelphia; but there can be no certainty about the matter until males of the present species are obtained and a comparison made with Leidy's material, which may or may not be in existence." Dr. Cobb further stated that after the nematodes leave the grasshoppers, they make their way into the soil and that their further history is obscure.

In Vermont the nematodes parasitized both *M. atlantis* and *M. bivittatus*. The worms seem to leave the bodies of the grasshoppers when these insects are maturing. We had not the opportunity to observe grasshoppers in the early stages of parasitism, but in August and September dissection of a large number of the insects showed that the worms were located within the body cavity. Later in the season, when the worms are about to emerge, the grasshoppers fall over on one side, kick for a time and then die. In the meantime, the worms gradually bore their way through the body wall and reach the exterior after which they make their way slowly into the earth. Usually only one worm parasitizes a grasshopper, but by dissection we have often found two or three and in one case we found forty. Needless to say, that when an insect contains so many worms the abdomen is considerably swollen.

The length of these female worms varied from two to eight inches. It is extraordinary that with hundreds of hoppers dying everywhere, we were unable to find any males.

A great many nematodes, at one stage of their life cycle, seek water on leaving their hosts and there mature, or wait until another host presents himself. We placed about two dozen of our worms in a bowl of water in which they seemed to flourish for about two weeks. However, the localities where the hoppers, and consequently the worms abounded were free from streams, ponds or marshes of any kind. The Connecticut River flows through a valley at a distance of about one-half mile so it seemed unlikely that the worms would travel so far. In all probability, we thought, the worms make their way into the soil on leaving the insects and this we found true. We placed recently dead parasitized hoppers in boxes containing earth. In about three days the boxes were examined and the worms were found coiled up at a depth of about one foot. Often a number would be coiled up together in one

spot. An examination of the soil in the fields revealed quantities of the worms below the surface at a distance of six inches to one foot. November the 7th and 8th, long after the grasshopper season, the ground now cold was again broken and the nematodes were found coiled up at about the same distance below the surface. Undoubtedly they hibernate in these positions.

During the highest mortality we made a series of dissections in order to determine the per cent. of parasitism. On a place called the Halladay Farm, we obtained the following astonishing high figures by the dissection of *M. bivittatus*.

Sept.	8.	100 ♀ ♀	dissected and worms found in	59%.
"	10.	80 ♀ ♀	" " " " "	76%.
"	12.	50 ♀ ♀	" " " " "	62%.
"	8.	40 ♂ ♂	" " " " "	12%.
"	10.	60 ♂ ♂	" " " " "	21%.
"	12.	100 ♂ ♂	" " " " "	12%.

On a place called the Tarbox Farm, we dissected about equal numbers of *M. bivittatus* and *M. atlantis* and obtained the following:

Sept.	8.	100 ♀ ♀	dissected and worms found in	22%.
"	10.	75 ♀ ♀	" " " " "	25%.
"	12.	60 ♀ ♀	" " " " "	25%.
"	8.	90 ♂ ♂	" " " " "	3%.
"	10.	80 ♂ ♂	" " " " "	5%.
"	12.	100 ♂ ♂	" " " " "	2%.

In both series of dissections it will be noticed that the percentage of parasitism in females is much higher than in males. Since the life-history of the worm is still so obscure we are at present unable to offer any explanation for this fact.

How the grasshoppers become infected is unknown. Since the nematodes are so large when they leave the grasshoppers in order to burrow into the soil, we are under the impression that grasshoppers are the secondary hosts. It is difficult to imagine what animal might constitute the primary host. Perhaps some other insect may furnish the clue to this interesting question.

Next summer we hope to extend our observations and attempt to gain a more complete insight into the life-history of this *Mermis* parasite. Some parasites fluctuate so numerically from one season

to another, that an entirely different condition may, of course, present itself in 1918. We mean that the worms may not be so plentiful for some reason and if this should prove true, it will be difficult to obtain very much information.

From our observations this summer (1917) we firmly believe that the nematodes accomplished an immense reduction in the number of grasshoppers near Dummerston Station, Vermont. This worm, if its life-history is investigated, might offer possibilities for introduction into regions where it does not occur and where grasshoppers are a pest. For this reason, and because we were unable to find any records of such a high degree of parasitism, we thought it best to present these preliminary observations.

THE PULSATILE VESSELS IN THE LEGS OF APHIDIDÆ.

BY CHAS. H. RICHARDSON,

College of Physicians and Surgeons, Columbia University, New York City.

When one of the light-colored aphids, like *Myzus persicæ* Sulz., is mounted alive on a depression slide, a rapid beating motion can be detected with the low power of the microscope in the tibia of each leg just below its juncture with the femur. These centers of activity mark the position of the pulsatile vessels.

The structure of these minute and delicate organs in aphids is difficult to determine, but serial sections through the tibia show that they are undoubtedly tubular. In the large aquatic Hemiptera, where they were first studied, the structure is more easily seen. Berlese¹ describes them as tubular organs crossed obliquely with numerous muscle bands and continuous with a non-pulsating part on either side.

The function of these organs is clearly one of blood propulsion. Locy,² who studied them in the aquatic Hemiptera, was able to discern the direction of the blood currents in the immediate vicinity of the pulsatile vessels, one current moving inward, the other outward. In *Myzus persicæ*, upon which most of my observations

¹ Gli Insetti, Milano, 1909, p. 764, fig. 953.

² American Naturalist, Vol. 18, pp. 13-19, 1884 (1 pl.).