DEMETRIDA (COLEOPTERA: CARABIDAE)
IN THE MOLUCCAS*

BY P. J. DARLINGTON, JR.
Museum of Comparative Zoology

Demetrída is a genus of the carabid tribe Lebiini recognizable by usually rather slender form (Figs. 1, 2); moderate size (5.5 - 12.0 mm in New Guinea); brown, bicolored, or brilliant metallic coloration; and technical characters given by me in 1968. Although a few species occur in New Zealand and New Caledonia, the genus is dominant primarily in Australia and New Guinea. The principal habitats occupied on these two adjacent land masses are surprisingly different. In Australia, adults (I do not know the larvae) of most species live on the trunks of eucalypts and other trees; in New Guinea, most live in understory foliage of rain forest. (In New Guinea, *Demetrída* therefore occupies the niche that *Calleida* occupies in some other parts of the world.) Some New Guinean *Demetrída* occur in the lowlands, but more, including most of the brilliant ones, are found at moderate or sometimes high altitudes in the mountains. The New Guinean members of the genus seem to have radiated independently of the Australian forms. Their radiation is in some ways compared to that of the brilliant birds of paradise that live in the same montane rain forests.

When I revised the New Guinean species (1968), I stated that the genus reached the Moluccas but did not give details. I now record the two species known from these islands. They set the western limit of the genus in the Indo-Australian Archipelago.

Demetrída latangula Darlington (Fig. 1)

Darlington 1968, pp. 147, 158.

Occurrence in the Moluccas: 1 ♂, "Amboina, F. Muir" (Bishop Museum).

This species is widely distributed at low altitudes in New Guinea, although not common there. The Moluccan individual matches New Guinean ones well.

***Demetrída moluccensis* n. sp. (Fig. 2)**

Description. With characters of genus (Darlington 1968); form as in Fig. 2, rather convex; dark reddish brown, appendages somewhat paler; not pubescent, moderately shining, reticulate micro-

*Manuscript received by the editor September 1, 1968

Fig. 1. *Demetrida latangula* DarlingtonFig. 2. *Demetrida moluccensis* n. sp.

sculpture indistinct on head and pronotum, light and imperfect on elytra. *Head* 1.10 and 1.03 width prothorax; eyes normal, prominent, genae short and oblique; front nearly flat, slightly irregularly impressed between eyes especially at sides. *Prothorax* quadrate; width/length 1.03 and 1.05; base/apex 1.17 and 1.15; base/head 0.83 and 0.83; margins narrow, each with seta-bearing puncture at middle but none at base; disc with middle line well impressed, transverse impressions very weak, baso-lateral impressions weak, they and marginal areas punctate; disc otherwise virtually impunctate but faintly transversely stringulose. *Elytra*: width elytra/prothorax 2.05 and 1.98; apices with outer angles well defined, right or slightly obtuse, then broadly emarginate to long spines *c.* opposite 3rd intervals, then emarginate to obtuse sutural angles; striae well impressed, scarcely punctulate; intervals somewhat convex, 3rd 2-punctate. *Inner wings* fully developed. *Lower surface* not distinctly punctate. *Legs*: claws each with *c.* 6 long teeth. *Secondary sexual characters*: ♂ front tarsi narrowly biseriately squamulose (as usual in genus); ♂ middle tibiae nearly straight, briefly tuberculate-serrate (*c.* 3 tubercles) distally on inner edge; ♂ with 2♀ apparently 3 seta-

bearing punctures each side last ventral segment. *Measurements*: length 8.6-8.8 (including spines); width 2.9 mm.

Types. Holotype ♂ (British Mus.) and 1 (probably ♀) paratype (MCZ Type No. 31,672) both from Laboean, Batjan Is., Moluccas, Feb., Mar. (no year given) (Doherty).

Notes. In my key to the species of *Demetrída* of New Guinea (1968, pp. 146ff) this runs to couplet 35 but fits neither species there named, having the prothorax more quadrate than in *forma* Darlington and the ♂ middle tibiae less bent in at apex than in *recta* Darlington and tuberculate-serrate (not so in *recta*).

The specimens here described were seen by Liebke, who labeled them "Genus? spec.?"

REFERENCE

DARLINGTON, P. J., JR.

1968. The carabid beetles of New Guinea. Part III. Harpalinae (continued): Perigonini to Pseudomorphini. Bull. Mus. Comp. Zool. 137 (1): 1-253 (*Demetrída*, pages 140-183).