OCCUPANCY BY *HYLAEUS* OF SUBTERRANEAN HALICTID NESTS (HYMENOPTERA: APOIDEA)¹

BY EDWARD M. BARROWS

Department of Entomology, University of Kansas
Lawrence, Kansas 66045

Hylaeus (Colletidae) is well known to nest in various preformed cavities. It is most commonly reported in hollow stems (Stoekert, 1933; Malyshev, 1936; Michener, 1965; Stephen et al., 1969) but also may occupy cynipid wasp galls (Stoekert, 1933; Laroca, 1971) and dead wood (Perkins, 1899; Michener, 1965). However, nest sites are not limited to plant materials. Smith (1855) reported cells in hollow pieces of "flint stone"; Ferton (1932), in earthworm burrows; and Perkins (1899), in ground. The present paper concerns nests of *H. bisinuatus*² in burrows of a halictine bee in an earthen bank; this is the first record of *Hylaeus* occupancy of nests of another hymenopteran.

H. bisinuatus has also been found in North America nesting in soda-straw traps and nail holes (R. R. Snelling, pers. comm.) and in Europe in hollow briar (*Rubus*) stems (Stoekert, 1933). Thus this species evidently has behavior which is flexible enough to enable it to nest in a variety of preformed holes in a wide geographical area.

In early July to early August, 1974, six nests of *H. bisinuatus* were found among those of *Lasioglossum* (*Dialictus*) *zephyrum* (Halictidae: Halictinae) in a nesting aggregation in vertical banks of the Kansas River, near Eudora, Douglas County, Kansas. Although the *Lasioglossum* nest aggregation is enormous, extending for about 1 km up and down the river, all the *Hylaeus* nests were found in about 10 square meters where the bank faces northwest.

The burrows occupied by *H. bisinuatus* were obviously former nests of *L. zephyrum*. In two nests vacant cells of *L. zephyrum* were near those used by *H. bisinuatus*; abandoned cells of *L. zephyrum* probably would have been found in all nests by following main burrows farther into the soil. Cells of *Hylaeus* were constructed in cells of the *Lasioglossum* still having the shape and wax-like lining

¹Contribution No. 1580 from the Department of Entomology, University of Kansas, Lawrence, Kansas.

²*H. bisinuatus* (Förster, 1871), described from Europe, is a senior synonym of *H. stevensi* (Crawford, 1913), described from North America, where it is probably adventive (R. R. Snelling, pers. comm.).

Manuscript received by the editor, April 5, 1975

characteristic of the halictine. The longitudinal axes of the burrows and cells of the nests were horizontal as is typical for nests of *L. zephyrum* in vertical banks (Batra, 1966). Burrows and nest entrances were similar in size to those of *L. zephyrum*. Thus there is no evidence that the *Hylaeus* did any nest excavation; it appears to have merely used available nests for its own purposes. Furthermore there is no evidence that the *Hylaeus* entered occupied *Lasioglossum* nests, although this possibility exists.

All six *Hylaeus* nests were discovered after their entrances had been closed, each with a sheet of a cellophane-like material flush with the surface of the soil bank. The sheets had striations radiating from their centers and small openings along their margins. A female of *H. bisinuatus* was discovered licking across a nest entrance already closed by a "cellophane" sheet. Evidently she was finishing the closure. She revolved around the entrance 3 times in 18 minutes while she licked 22.5 times per minute ($n = 2 \text{ min}$) across the sheet; during this time she took occasional rests of several seconds each. During each lick the tip of her proboscis moved from her midventral area to a point in front of her head.

For identification she was captured before she finished the closure and two days later (on 5 July) her nest was examined (Fig. 1). All three of the cells in her nest were provisioned with a soft yellow material, most likely a mixture of pollen and nectar as is used by

Fig. 1. Top view of a nest of *Hylaeus bisinuatus* made in a nest of *Lasioglossum zephyrum*; B, main burrow; C, *Hylaeus* cell inside *Lasioglossum* cell; M, cellophane-like membrane secreted by the *Hylaeus*.

other *Hylaeus*, and each contained a larva. One larva became a prepupa on 6 July, a pupa on 7 July, and an adult female on 14 July; the other larvae were small and died.

Nests had from 1 to 3 *Hylaeus* cells ($\bar{x} = 1.7$, $n = 6$ nests), each in a different *Lasioglossum* cell (Fig. 2). Both *Hylaeus* cells which were measured were 5 mm long and 3.5 mm in maximum diameter and at the ends of *Lasioglossum* cells. The walls of the cells had old wax-like lining deposited by *L. zephyrum* (Batra, 1966). *H. bisinuatus* laid down two other cell linings to form its cells: (1) a very thin "cellophane" layer visible under a microscope (30 $\times$) only when light strikes it at a certain angle and (2) a slightly thicker "cellophane" layer over the first one. This thicker layer also covered walls of lateral burrows and parts of main burrows. A sheet of what appeared to be this same second layer closed off each cell and with the unaided eye the halictine cells appeared to be constricted where such a sheet was placed. Complete removal of the cellophane-like material, however, shows the unmodified form of the halictine cell.

H. bisinuatus used cells of *L. zephyrum* that were 0.8 to 4.5 cm ($\bar{x} = 3.3$ cm, $n = 6$ nests) from the nest entrances. Since *L. zephyrum* constructs cells this close to the surface of a soil bank only in spring or early summer, the *Hylaeus* must have used early-season *Lasioglossum* cells made in the year of this study or possibly in the previous year.

ACKNOWLEDGMENTS

I thank Mr. Roy R. Snelling of the Los Angeles County Museum of Natural History for identifying *Hylaeus bisinuatus*. He and Drs. Charles D. Michener and William J. Bell and Ms. Marjorie R. Chabot made important suggestions regarding this note. It is a byproduct of National Science Grant GB-38502 (W. J. Bell and C. D. Michener, coprincipal investigators).

LITERATURE CITED

BATRA, S. W. T.

1966. The life cycle and behavior of the primitively social bee *Lasioglossum zephyrum* (Halictidae). Univ. Kansas Sci. Bull. 46: 359-423.

FERTON, C.

1923. La vie des abeilles et des guêpes, oeuvres choisies, groupées et annotées par Étienne Rabaud et François Picard. Étienne Chiron, ed., 40 Rue de Seine, Paris. 376 pp.

- LAROCA, S.
1971. Notas sôbre a biologia de *Hylaeus cecidonastes* Moure (Hymenoptera, Apoidea). Rev. Brasil. Biol. 32: 285-290.
- MALYSHEV, S. I.
1936. The nesting habits of solitary bees (a comparative study). Eos 11: 201-309.
- MICHENER, C. D.
1965. A classification of the bees of the Australian and South Pacific Regions. Bull. Amer. Mus. Nat. Hist. 130: 1-362.
- PERKINS, R. C. L. in PERKINS, R. C. L. AND A. FOREL.
1899. Fauna Hawaiiensis or the zoology of the Sandwich (Hawaiian) Isles. Vol. 1, Part 1. Hymenoptera, Aculeata. Cambridge Univ. Press. 122 pp.
- SMITH, F.
1855. Catalogue of British Hymenoptera: Part 1 Apidae — bees. London. 252 pp.
- STEPHEN, W. P., G. E. BOHART, AND P. E. TORCHIO.
1969. The biology and external morphology of bees with a synopsis of the genera of northwestern America. Agric. Exp. Station, Oregon St. Univ., Corvallis. 140 pp.
- STOEKHERT, F. K.
1933. Die Bienen Frankens (Hym. Apid.) Beiheft Deuts. Entomol. Zeit. 1-294.