Research Article

Microarray-Based Comparisons of Ion Channel Expression Patterns: Human Keratinocytes to Reprogrammed hiPSCs to Differentiated Neuronal and Cardiac Progeny

Leonhard Linta,¹ Marianne Stockmann,¹ Qiong Lin,² André Lechel,³ Christian Proepper,¹ Tobias M. Boeckers,¹ Alexander Kleger,³ and Stefan Liebau¹

¹ Institute for Anatomy Cell Biology, Ulm University, Albert-Einstein Allee 11, 89081 Ulm, Germany

² Institute for Biomedical Engineering, Department of Cell Biology, RWTH Aachen, Pauwelsstrasse 30, 52074 Aachen, Germany

³ Department of Internal Medicine I, Ulm University, Albert-Einstein Allee 11, 89081 Ulm, Germany

Correspondence should be addressed to Alexander Kleger; alexander.kleger@uni-ulm.de and Stefan Liebau; stefan.liebau@uni-ulm.de

Received 31 January 2013; Accepted 6 March 2013

Academic Editor: Michael Levin

Copyright © 2013 Leonhard Linta et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Ion channels are involved in a large variety of cellular processes including stem cell differentiation. Numerous families of ion channels are present in the organism which can be distinguished by means of, for example, ion selectivity, gating mechanism, composition, or cell biological function. To characterize the distinct expression of this group of ion channels we have compared the mRNA expression levels of ion channel genes between human keratinocyte-derived induced pluripotent stem cells (hiPSCs) and their somatic cell source, keratinocytes from plucked human hair. This comparison revealed that 26% of the analyzed probes showed an upregulation of ion channels in hiPSCs while just 6% were downregulated. Additionally, iPSCs express a much higher number of ion channels compared to keratinocytes. Further, to narrow down specificity of ion channel expression in iPS cells we compared their expression patterns with differentiated progeny, namely, neurons and cardiomyocytes derived from iPS cells. To conclude, hiPSCs exhibit a very considerable and diverse ion channel expression pattern. Their detailed analysis could give an insight into their contribution to many cellular processes and even disease mechanisms.

1. Introduction

Ion channels are comprised of a large variety of differing families of pore proteins. Initially, ion channels were mostly known for their role in the nervous system where they play a crucial role in the signal transmission over neurites and synapses. But in fact they are involved in numerous other cellular processes including cell size regulation, muscle contractions, immune system activation [1], or hormone release [2]. Distinct ion channels are furthermore recognized to be of high importance for excitable cells of the heart: cardiomyocytes of the working myocardium as well as cells of the cardiac conduction system. In the heart, specific ion channels are responsible, for example, for the regulated generation of action potentials and for cardiac muscle contraction strength and time [3]. Additionally, ion channels play an important

role in several differentiation and maturation processes [4–6]. The presented study aims to take a closer look at ion channel expression in human-induced pluripotent stem cells (hiPSCs) to give a start point for further analyses of their distinct roles at an early developmental cell state and differentiation processes.

hiPSCs are generated from somatic cells by timed overexpression of specific transcription factors and strongly resemble pluripotent embryonic stem cells [7, 8]. Pluripotency is amongst others defined by the ability to differentiate into cells of all three germ layers and unlimited symmetrical cell division. This cell system is widely utilized for studies investigating developmental processes or disease mechanisms [9, 10]. Although it has become clear that differentiation processes influence ion channel expression [11], the distinct role of ion channels during these processes is so far only poorly

understood. Nevertheless, it became evident that certain ion channels play a pivotal role in stem cell biology, including cell fate determination, cell cycle regulation, or cytoskeletal reorganization [6, 12–15].

The possibilities of iPSCs include the generation of individual or patient-specific-pluripotent cells, which can be subsequently differentiated into the affected cell types. This is already utilized to study pathomechanisms in a variety of tissues and cell types [9, 16–18]. The so-called channelopathies which are based on mutations in ion channels are the cause of developmental disorders and are the subject of various studies [19, 20]. To elucidate the role of ion channels in cell differentiation, maturation or their role in pathomechanisms a well-founded knowledge of ion channel distribution in pluripotent cells, representing one of the earliest stages of development, is indispensable. In that respect, we compared the expression levels of several ion channels in human keratinocytes with their reprogrammed progeny, hiPSCs. Keratinocytes derived from plucked human hair [21, 22] represent one of the most promising cell sources for the generation of the investigated hiPSCs [23]. We have elucidated ion channel regulation for various channel families and their subtypes. Additionally, we compared the expression levels of ion channel families and subtypes, found to be regulated during reprogramming, with iPSC-differentiated progeny, namely, neurons and cardiomyocytes. These comparisons could be a start point to evaluate the contribution and function of different ion channels, for example, for self-renewal and differentiation processes in stem cells.

2. Material and Methods

2.1. Ethical Statement and Donor Information. After informed consent was given by the donors (ethical agreement by the University of Ulm, number 88/12) hair with intact hair roots was gathered by plucking from the scalp after disinfection. We used hair from healthy volunteers (age between 24 to 45 and both male and female gender).

2.2. hiPSC Generation and Cell Culture. Keratinocytes were obtained from plucked human scalp hair as already described [22]. Keratinocytes were propagated in EpiLife medium with HKGS supplement (both Invitrogen, Carlsbad, CA, USA). hiPSCs were generated from keratinocytes by lentiviral transduction of four reprogramming factors (Oct4, Sox2, Klf4, and cMyc) as described earlier [23]. After the reprogramming on rat embryonic fibroblasts they were maintained feeder-free on Matrigel (BD Biosciences, Franklin Lakes, NJ, USA) coated dishes in mTeSR1 medium (Stemcell Technologies, Vancouver, CA, USA).

2.3. Gene Expression Microarrays. Gene expression microarrays were performed for 6 keratinocyte samples and 9 hiPSC samples with the Agilent Whole Human Genome Microarray Kit (4x44k microarray kit G4112F, Agilent Technologies, Santa Clara, CA, USA). 500 ng of total RNA was used to produce Cy3-CTP-labeled cRNA with the Agilent Low RNA Input Liner Amplification Kit. The cRNA

was purified and 1,65 µg per array was hybridized for 17 h at 65°C and 10 r.p.m. Afterwards, the arrays were washed with Agilent Gene Expression Wash Buffers one and two and finally with acetonitrile for 1 min. The slides were scanned using Scan Control 7.0 software with a resolution of 5 µm. Scan data was extracted with the Feature Extraction 9.1 software. Expression levels were background adjusted and quantile normalized with the GeneSpring GX 12 software. Differential expression between keratinocytes and hiPSCs was analyzed using student's *t*-test. A fold change > 2 and a *P* value < 0.05 was considered significant and highlighted bold (upregulation) or italic (downregulation) in the results table. For comparisons of iPSCs with iPSC-derived neurons published data from GSE34879 (GSM856936, GSM856937, GSM856915, GSM856916) and for cardiomyocytes GSE17579 (GSM438022, GSM438026, GSM438034, GSM438021, GSM438032, GSM438036) were used (both from NCBI Gene Expression Omnibus, <http://www.ncbi.nlm.nih.gov/geo/>). Fold change is shown if > 2.

3. Results

3.1. Differential Expression of Ion Channels in hiPSCs Compared to Keratinocytes. We first compared the expression of various ion channel families from keratinocytes to hiPSCs and from hiPSCs to cardiomyocytes and neurons, respectively. From the 387 probes (Table 1) binding in ion channel genes from parental keratinocytes to hiPSCs, 101 (26%) showed a significant increase in expression (fold change > 2, *P* < 0.05; labeled in bold) while 23 (6%) showed a significant decrease (fold change > -2, *P* < 0.05; labeled in italic). In a second step differentially regulated ion channels from hiPSCs to hiPSC-derived neurons were investigated and we found 29 ion channel transcripts to be upregulated (fold change > 2; labeled in bold) while 6 showed a significant decrease (fold change > -2; labeled in italic). For cardiomyocytes, mRNA levels of only 7 ion channel members were upregulated (fold change > 2; labeled in bold) and 10 mRNA levels showed a significant decrease (fold change > -2; labeled in italic).

3.2. Voltage-Gated Calcium Channels. Voltage-gated calcium channels are crucially involved in the Ca²⁺-influx thereby playing an important role in calcium signaling of virtually all cells. High-voltage-gated calcium channels include the neural N-type channel, the poorly defined brain-specific R-type channel, the closely related P/Q-type channel, and the dihydropyridine-sensitive L-type channels responsible for excitation-contraction coupling of skeletal, smooth, and cardiac muscles as well as for hormone secretion in endocrine cells (reviewed in [24]). While mainly permeable for calcium ions they also show a low permeability for sodium ions. Upon depolarization of the cell they mediate a calcium influx into the cell. The channels consist of the main α subunit as well as regulatory β , $\alpha 2/\delta$, and γ subunits. The α_1 subunit forms the ion conducting pore while the associated subunits have several functions including modulation of gating [25]. CACNA1A mutations are for example involved in ataxia [26]. We have analyzed their expression with 42 probes.

TABLE 1: The expression levels for ion channel genes were compared in hiPSCs (9 samples) and keratinocytes (6 samples) via microarray. Gene names, accession numbers, and gene symbols are shown together with the corresponding fold change and *P* values. Significant changes (defined as fold change > 2, *P* < 0.05) are labeled in bold for upregulation or italic for downregulation. Accession numbers refer to the NCBI or Ensembl (numbers starting with ENST) databases. Analyses of hiPSC-derived neurons or hiPSC-derived cardiomyocytes were performed using published data sets from GSE34879 (neurons) and GSE17579 (cardiomyocytes). These data sets already contain both iPSCs and differentiated progeny. Fold changes > 2 (marked bold) and < -2 (marked italic) are shown in the respective columns.

(a) Voltage-gated calcium channels						
Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Calcium channel, voltage-dependent, P/Q type, alpha 1A subunit, transcript variant 1	NM_000068	CACNA1A	2.26E - 01	-1.32		
Calcium channel, voltage-dependent, N type, alpha 1B subunit	NM_000718	CACNA1B	7.27E - 01	-1.10		
Calcium channel, voltage-dependent, N type, alpha 1B subunit [Source: HGNC Symbol; Acc: 1389]	ENST00000277550	CACNA1B	7.15E - 01	-1.14		
Calcium channel, voltage-dependent, N type, alpha 1B subunit	NM_000718	CACNA1B	6.32E - 09	6.82		
Calcium channel, voltage-dependent, N type, alpha 1B subunit [Source: HGNC Symbol; Acc: 1389]	ENST00000277551	CACNA1B	3.32E - 02	-1.57		
Calcium channel, voltage-dependent, L type, alpha 1C subunit, transcript variant 18	NM_000719	CACNA1C	6.77E - 02	-1.37		
Calcium channel, voltage-dependent, L type, alpha 1C subunit, transcript variant 18	NM_000719	CACNA1C	5.72E - 03	2.37		2.94
Calcium channel, voltage-dependent, L type, alpha 1D subunit, transcript variant 1	NM_000720	CACNA1D	3.83E - 01	1.58		
Calcium channel, voltage-dependent, R type, alpha 1E subunit, transcript variant 3	NM_000721	CACNA1E	3.10 E - 02	-4.48	2.08	
Calcium channel, voltage-dependent, R type, alpha 1E subunit [Source: HGNC Symbol; Acc: 1392]	ENST00000524607	CACNA1E	8.83E - 01	1.05		
Calcium channel, voltage-dependent, L type, alpha 1F subunit	NM_005183	CACNA1F	8.92E - 01	1.02		
Calcium channel, voltage-dependent, T type, alpha 1G subunit, transcript variant 15	NM_198397	CACNA1G	6.42E - 01	-1.12	4.60	
Calcium channel, voltage-dependent, T type, alpha 1G subunit, transcript variant 1	NM_018896	CACNA1G	5.37E - 07	6.96		
Calcium channel, voltage-dependent, T type, alpha 1H subunit, transcript variant 1	NM_021098	CACNA1H	7.38E - 10	6.78		
Calcium channel, voltage-dependent, T type, alpha 1I subunit, transcript variant 1	NM_021096	CACNA1I	2.22E - 11	37.25		
Calcium channel, voltage-dependent, T type, alpha 1I subunit, transcript variant 1	NM_021096	CACNA1I	7.56E - 02	-1.56		
Calcium channel, voltage-dependent, L type, alpha 1S subunit	NM_000069	CACNA1S	5.22E - 02	-1.97		
Calcium channel, voltage-dependent, alpha 2/delta subunit 1	NM_000722	CACNA2D1	1.38E - 09	9.65		
Calcium channel, voltage-dependent, alpha 2/delta subunit 2, transcript variant 1	NM_001005505	CACNA2D2	3.30E - 10	62.50	2.02	
Calcium channel, voltage-dependent, alpha 2/delta subunit 3	NM_018398	CACNA2D3	5.14E - 05	5.42	2.55	

(a) Continued.

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Calcium channel, voltage-dependent, alpha 2/delta subunit 3	AF516696	CACNA2D3	4.68E – 11	33.67		
Calcium channel, voltage-dependent, alpha 2/delta subunit 4	NM_172364	CACNA2D4	2.95E – 01	–1.41		
Calcium channel, voltage-dependent, beta 1 subunit, transcript variant 3	NM_199248	CACNB1	4.31E – 01	–1.14		
Calcium channel, voltage-dependent, beta 1 subunit, transcript variant 1	NM_000723	CACNB1	8.02E – 01	–1.02		
Calcium channel, voltage-dependent, beta 1 subunit, transcript variant 1	NM_000723	CACNB1	6.22E – 01	1.04		
cDNA FLJ45229 fis, clone BRCAN2020972,	AK128769	CACNB2	9.78E – 01	1.02		
Calcium channel, voltage-dependent, beta 2 subunit, transcript variant 1	NM_000724	CACNB2	4.71E – 02	3.45		
Calcium channel, voltage-dependent, beta 3 subunit, transcript variant 1	NM_000725	CACNB3	3.55E – 02	–1.93	9.56	
Calcium channel, voltage-dependent, beta 3 subunit, transcript variant 1	NM_000725	CACNB3	6.61E – 01	1.09		
Calcium channel, voltage-dependent, beta 4 subunit, transcript variant 1	NM_001005747	CACNB4	1.33E – 03	4.43		
Calcium channel, voltage-dependent, gamma subunit 1	NM_000727	CACNG1	9.46E – 01	1.02		
Calcium channel, voltage-dependent, gamma subunit 2	NM_006078	CACNG2	4.19E – 01	–1.29		
Calcium channel, voltage-dependent, gamma subunit 2	NM_006078	CACNG2	7.91E – 01	1.09		
Calcium channel, voltage-dependent, gamma subunit 3	NM_006539	CACNG3	2.21E – 01	–1.95		
Calcium channel, voltage-dependent, gamma subunit 4	NM_014405	CACNG4	4.93E – 01	1.14		
Calcium channel, voltage-dependent, gamma subunit 5	NM_145811	CACNG5	7.49E – 01	–1.13		
Calcium channel, voltage-dependent, gamma subunit 5 [Source: HGNC Symbol; Acc: 1409]	ENST00000169565	CACNG5	2.19E – 01	–1.56		
Calcium channel, voltage-dependent, gamma subunit 6, transcript variant 1	NM_145814	CACNG6	5.56E – 11	46.40		
Calcium channel, voltage-dependent, gamma subunit 7	NM_031896	CACNG7	3.54E – 13	1439.36	3.37	
Calcium channel, voltage-dependent, gamma subunit 7	NM_031896	CACNG7	1.82E – 01	1.26		
Calcium channel, voltage-dependent, gamma subunit 8	NM_031895	CACNG8	2.01E – 01	1.43		
Calcium channel, voltage-dependent, gamma subunit 8	NM_031895	CACNG8	1.18E – 09	9.70		

(b) Sperm-associated cation channels

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Cation channel, sperm associated 1	NM_053054	CATSPER1	3.26E – 01	1.51		
Cation channel, sperm associated 2, transcript variant 2	NM_172095	CATSPER2	1.32 E–04	–2.78		

(b) Continued.

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Cation channel, sperm associated 2, transcript variant 4	NM_172097	CATSPER2	$3.12E - 02$	-1.78		
Cation channel, sperm associated 3	NM_178019	CATSPER3	$1.50E - 03$	1.93		
Cation channel, sperm associated 4	NM_198137	CATSPER4	$2.51E - 01$	1.48		
Cation channel, sperm-associated, beta	NM_024764	CATSPERB	$4.38E - 01$	-1.25		
Cation channel, sperm-associated, gamma	NM_021185	CATSPERG	$3.56E - 01$	1.34		

(c) Nicotinic acetylcholine receptors

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Cholinergic receptor, nicotinic, alpha 1 (muscle), transcript variant 1	NM_001039523	CHRNA1	$8.06E - 01$	-1.10		
Cholinergic receptor, nicotinic, alpha 1 (muscle), (cDNA clone IMAGE: 4124038), with apparent retained intron	BC006314	CHRNA1	$4.64E - 01$	1.47		
Cholinergic receptor, nicotinic, alpha 2 (neuronal)	NM_000742	CHRNA2	$6.05E - 01$	-1.30		
Cholinergic receptor, nicotinic, alpha 3, transcript variant 1	NM_000743	CHRNA3	$1.84E - 10$	34.22		
Cholinergic receptor, nicotinic, alpha 4	NM_000744	CHRNA4	$6.79E - 01$	1.15		
Cholinergic receptor, nicotinic, alpha 4 (neuronal), exon 1	X89741	CHRNA4	$2.79E - 02$	-2.53		
Cholinergic receptor, nicotinic, alpha 5	NM_000745	CHRNA5	$4.03E - 05$	3.41		
Cholinergic receptor, nicotinic, alpha 6, transcript variant 1	NM_004198	CHRNA6	$2.23E - 01$	-1.27	2.21	
Cholinergic receptor, nicotinic, alpha 7, transcript variant 1	NM_000746	CHRNA7	$4.61E - 04$	1.86		
Cholinergic receptor, nicotinic, alpha 7, transcript variant 2	NM_001190455	CHRNA7	$1.56E - 11$	13.24		
Cholinergic receptor, nicotinic, alpha 9	NM_017581	CHRNA9	$6.30E - 06$	6.15		
Cholinergic receptor, nicotinic, alpha 10	NM_020402	CHRNA10	$1.57E - 02$	1.57		
Cholinergic receptor, nicotinic, beta 1 (muscle)	NM_000747	CHRNA1	$8.07E - 04$	-2.47		
Cholinergic receptor, nicotinic, beta 1 (muscle)	NM_000747	CHRNA1	$8.87E - 03$	-1.92		
Cholinergic receptor, nicotinic, beta 2 (neuronal)	NM_000748	CHRNA2	$6.03E - 04$	-3.01		
Cholinergic receptor, nicotinic, beta 2 (neuronal) [Source: HGNC Symbol; Acc: 1962]	ENST00000368476	CHRNA2	$2.98E - 09$	32.05		
Cholinergic receptor, nicotinic, beta 3	NM_000749	CHRNA3	$5.91E - 01$	1.12		
Cholinergic receptor, nicotinic, beta 4	NM_000750	CHRNA4	$1.76E - 02$	4.20		
Cholinergic receptor, nicotinic, delta	NM_000751	CHRNA5	$9.84E - 01$	1.01		
Cholinergic receptor, nicotinic, epsilon	NM_000080	CHRNA6	$3.66E - 01$	-1.20		
Cholinergic receptor, nicotinic, gamma	NM_005199	CHRNA7	$6.70E - 01$	1.14		

(d) Cyclic nucleotide-gated channels

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Cyclic nucleotide-gated channel alpha 1, transcript variant 2	NM_000087	CNGA1	$7.14E - 01$	1.15		

(d) Continued.

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Cyclic nucleotide-gated channel alpha 1, transcript variant 2	NM_000087	CNGA1	1.18E – 02	2.00		
Cyclic nucleotide-gated channel alpha 3, transcript variant 1	NM_001298	CNGA3	2.00E – 01	1.21		
Cyclic nucleotide-gated channel alpha 4	NM_001037329	CNGA4	2.78E – 01	1.82		
Cyclic nucleotide-gated channel beta 1, transcript variant 1	NM_001297	CNGB1	9.22 E–03	–2.05		
Cyclic nucleotide-gated channel beta 1, transcript variant 1	NM_001297	CNGB1	6.36E – 01	–1.13		
Cyclic nucleotide-gated channel beta 3	NM_019098	CNGB3	2.93E – 01	1.78		

(e) GABA receptors

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Gamma-aminobutyric acid (GABA) A receptor, alpha 1, transcript variant 1	NM_000806	GABRA1	1.26E – 01	1.41		
Gamma-aminobutyric acid (GABA) A receptor, alpha 1, transcript variant 1	NM_000806	GABRA1	8.87E – 01	1.05		
Gamma-aminobutyric acid (GABA) A receptor, alpha 2, transcript variant 1	NM_000807	GABRA2	9.70E – 02	1.70		
Gamma-aminobutyric acid (GABA) A receptor, alpha 2, transcript variant 1	NM_000807	GABRA2	4.49E – 01	1.48		
Gamma-aminobutyric acid (GABA) A receptor, alpha 3	NM_000808	GABRA3	2.10E – 01	1.43		
Gamma-aminobutyric acid (GABA) A receptor, alpha 4, transcript variant 1	NM_000809	GABRA4	3.76E – 01	1.43		
Gamma-aminobutyric acid (GABA) A receptor, alpha 4, transcript variant 1	NM_000809	GABRA4	8.30E – 01	1.05		
Gamma-aminobutyric acid (GABA) A receptor, alpha 5, transcript variant 1	NM_000810	GABRA5	1.09E – 08	14.41		
Gamma-aminobutyric acid (GABA) A receptor, alpha 6	NM_000811	GABRA6	1.69E – 01	1.23		
Gamma-aminobutyric acid (GABA) A receptor, beta 1	NM_000812	GABRB1	1.14E – 01	1.43		
Gamma-aminobutyric acid (GABA) A receptor, beta 2, transcript variant 2	NM_000813	GABRB2	1.27E – 02	1.94		
Gamma-aminobutyric acid (GABA) A receptor, beta 3, transcript variant 1	NM_000814	GABRB3	6.70E – 10	85.93		–4.49
Gamma-aminobutyric acid (GABA) A receptor, beta 3, transcript variant 1	NM_000814	GABRB3	1.15E – 09	115.69		
Gamma-aminobutyric acid (GABA) A receptor, delta	NM_000815	GABRD	6.26E – 02	3.02		
Gamma-aminobutyric acid (GABA) A receptor, epsilon	NM_004961	GABRE	5.83 E–05	–5.47		
Gamma-aminobutyric acid (GABA) A receptor, Gamma 1	NM_173536	GABRG1	1.78E – 01	1.24		
Gamma-aminobutyric acid (GABA) A receptor, Gamma 1	NM_173536	GABRG1	1.70E – 02	2.19		
Gamma-aminobutyric acid (GABA) A receptor, Gamma 2, transcript variant 1	NM_198904	GABRG2	6.02E – 01	1.14		

(e) Continued.

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Gamma-aminobutyric acid (GABA) A receptor, Gamma 2, transcript variant 2	NM.000816	GABRG2	$5.28E - 02$	1.72		
Gamma-aminobutyric acid (GABA) A receptor, Gamma 2, transcript variant 2	NM.000816	GABRG2	$9.47E - 09$	8.41		
Gamma-aminobutyric acid (GABA) A receptor, Gamma 3	NM.033223	GABRG3	$1.99E - 01$	2.00		
Gamma-aminobutyric acid (GABA) A receptor, Gamma 3	NM.033223	GABRG3	$1.04E - 01$	1.61		
Gamma-aminobutyric acid (GABA) A receptor, pi	NM.014211	GABRP	$6.62E - 01$	-1.12		17.25
Gamma-aminobutyric acid (GABA) receptor, theta	NM.018558	GABRQ	$1.60E - 01$	1.56		
Gamma-aminobutyric acid (GABA) receptor, theta	NM.018558	GABRQ	$4.76E - 01$	-1.27		
Gamma-aminobutyric acid (GABA) receptor, theta [Source: HGNC Symbol; Acc: 14454]	ENST00000370306	GABRQ	$8.22E - 09$	52.17		
Gamma-aminobutyric acid (GABA) receptor, rho 1	NM.002042	GABRR1	$8.94E - 01$	-1.04		
Gamma-aminobutyric acid (GABA) receptor, rho 2	NM.002043	GABRR2	$2.83E - 01$	1.54		
Gamma-aminobutyric acid (GABA) receptor, rho 3	NM.001105580	GABRR3	$3.05E - 01$	-1.34		

(f) Glycine receptors

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Glycine receptor, alpha 1, transcript variant 2	NM.000171	GLRA1	$5.47E - 01$	-1.29		
Glycine receptor, alpha 2, transcript variant 1	NM.002063	GLRA2	$1.17E - 01$	1.42	11.41	
Glycine receptor, alpha 3, transcript variant 1	NM.006529	GLRA3	$5.16E - 01$	-1.26		
Glycine receptor, alpha 3 [Source: HGNC Symbol; Acc: 4328]	ENST00000274093	GLRA3	$8.33E - 02$	1.64		
Glycine receptor, alpha 3, transcript variant 1	NM.006529	GLRA3	$1.46E - 01$	1.41		
Glycine receptor, alpha 4, transcript variant 1	NM.001024452	GLRA4	$7.75E - 01$	1.17		
Glycine receptor, beta, transcript variant 1	NM.000824	GLRB	$6.38E - 02$	-1.69	6.61	

(g) Ionotropic glutamate receptors

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Glutamate receptor, ionotropic, AMPA 1, transcript variant 1	NM.000827	GRIA1	$7.65E - 01$	-1.20	2.06	
Glutamate receptor, ionotropic, AMPA 2, transcript variant 1	NM.000826	GRIA2	$2.87E - 01$	-1.87	21.26	
Glutamate receptor, ionotropic, AMPA 2, transcript variant 1	NM.000826	GRIA2	$2.72E - 01$	1.33		
Glutamate receptor, ionotropic, AMPA 2, transcript variant 1	NM.000826	GRIA2	$6.54E - 01$	1.22		

(g) Continued.

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Glutamate receptor, ionotropic, AMPA 3, transcript variant 2	NM_000828	GRIA3	5.05E − 01	1.26		
Glutamate receptor, ionotropic, AMPA 3 [Source: HGNC Symbol; Acc: 4573]	ENST00000371264	GRIA3	1.14E − 01	1.50		
Glutamate receptor, ionotropic, AMPA 3, transcript variant 2	NM_000828	GRIA3	9.05E − 01	−1.05		
Glutamate receptor, ionotropic, AMPA 4, transcript variant 1	NM_000829	GRIA4	2.30E − 02	1.80	3.73	
Glutamate receptor, ionotropic, delta 1	NM_017551	GRID1	1.82E − 01	1.50		
Glutamate receptor, ionotropic, delta 1	NM_017551	GRID1	5.08E − 05	3.63		
Glutamate receptor, ionotropic, delta 2	NM_001510	GRID2	7.64E − 01	−1.11		
Glutamate receptor, ionotropic, delta 2	NM_001510	GRID2	5.55E − 09	55.66		
Glutamate receptor, ionotropic, kainate 1, transcript variant 1	NM_000830	GRIK1	1.64E − 01	1.29	2.58	
Glutamate receptor, ionotropic, kainate 2, transcript variant 3	NM_001166247	GRIK2	1.42E − 01	1.56	2.53	
Glutamate receptor, ionotropic, kainate 2, transcript variant 1	NM_021956	GRIK2	1.14E − 01	−2.01		
Glutamate receptor, ionotropic, kainate 2, transcript variant 2	NM_175768	GRIK2	7.80E − 01	1.08		
Glutamate receptor, ionotropic, kainate 3	NM_000831	GRIK3	1.58E − 01	1.64		
Glutamate receptor, ionotropic, kainate 3	NM_000831	GRIK3	6.15E − 10	22.64		
Glutamate receptor, ionotropic, kainate 4 [Source: HGNC Symbol; Acc: 4582]	ENST00000527524	GRIK4	1.77E − 11	18.10		
Glutamate receptor, ionotropic, kainate 4	NM_014619	GRIK4	5.73E − 02	2.54		
Glutamate receptor, ionotropic, kainate 5	NM_002088	GRIK5	2.65E − 03	1.79		−2.41
Glutamate receptor, ionotropic, N-methyl D-aspartate 1, transcript variant NRI-3	NM_007327	GRIN1	1.93 E−02	−3.02	2.69	
Glutamate receptor, ionotropic, N-methyl D-aspartate 1, transcript variant NRI-3	NM_007327	GRIN1	6.01E − 02	−2.45		
Glutamate receptor, ionotropic, N-methyl D-aspartate 1, transcript variant NRI-3	NM_007327	GRIN1	3.86E − 01	−1.65		
Glutamate receptor, ionotropic, N-methyl D-aspartate 2A, transcript variant 2	NM_000833	GRIN2A	6.54E − 01	1.18		
Glutamate receptor, ionotropic, N-methyl D-aspartate 2A, transcript variant 2	NM_000833	GRIN2A	9.21E − 01	−1.02		
Glutamate receptor, ionotropic, N-methyl D-aspartate 2A, transcript variant 1	NM_001134407	GRIN2A	3.32E − 10	29.85		
Glutamate receptor, ionotropic, N-methyl D-aspartate 2A, transcript variant 2	NM_000833	GRIN2A	2.63E − 01	1.55		
Glutamate receptor, ionotropic, N-methyl D-aspartate 2B	NM_000834	GRIN2B	6.64E − 01	−1.26		
Glutamate receptor, ionotropic, N-methyl D-aspartate 2C	NM_000835	GRIN2C	8.01E − 01	−1.08		
Glutamate receptor, ionotropic, N-methyl D-aspartate 2C	NM_000835	GRIN2C	5.01E − 01	1.13		
Glutamate receptor, ionotropic, N-methyl D-aspartate 2D	NM_000836	GRIN2D	4.69E − 02	−1.82		
Glutamate receptor, ionotropic, N-methyl-D-aspartate 3A	NM_133445	GRIN3A	3.28E − 01	−1.41		

(g) Continued.

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Glutamate receptor, ionotropic, N-methyl-D-aspartate 3B	NM_138690	GRIN3B	$9.36E-01$	-1.02		
Glutamate receptor, ionotropic, N-methyl D-aspartate-associated protein 1 (glutamate binding), transcript variant 1	NM_000837	GRINA	$3.68E-04$	-2.05	2.05	
NMDA receptor glutamate-binding chain (hnrwg), partial	U44954	GRINA	$3.06E-04$	-1.81		

(h) Hyperpolarization-activated cyclic nucleotide-gated channels

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Hyperpolarization-activated cyclic nucleotide-gated potassium channel 1	NM_021072	HCN1	$3.47E-11$	52.85		
Hyperpolarization-activated cyclic nucleotide-gated potassium channel 1	NM_021072	HCN1	$1.01E-08$	9.91		
Hyperpolarization-activated cyclic nucleotide-gated potassium channel 2	NM_001194	HCN2	$5.36E-02$	-1.63		
Hyperpolarization-activated cyclic nucleotide-gated potassium channel 3	NM_020897	HCN3	$1.98E-02$	2.12	6.56	
Hyperpolarization-activated cyclic nucleotide-gated potassium channel 4	NM_005477	HCN4	$1.01E-01$	-1.66		

(i) Serotonin receptors

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
5-hydroxytryptamine (serotonin) receptor 3A, transcript variant 1	NM_213621	HTR3A	$1.99E-03$	2.77	-2.20	
5-hydroxytryptamine (serotonin) receptor 3A, transcript variant 1	NM_213621	HTR3A	$2.63E-11$	11.11		
5-hydroxytryptamine (serotonin) receptor 3B	NM_006028	HTR3B	$1.99E-01$	-1.54		
5-hydroxytryptamine (serotonin) receptor 3, family member C	NM_130770	HTR3C	$4.81E-01$	1.34		
5-hydroxytryptamine (serotonin) receptor 3 family member D, transcript variant 2	NM_182537	HTR3D	$4.75E-01$	-1.37		
5-hydroxytryptamine (serotonin) receptor 3, family member E	NM_182589	HTR3E	$1.52E-01$	-1.44		

(j) Voltage-gated potassium channels

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Potassium voltage-gated channel, shaker-related subfamily, member 1 (episodic ataxia with myokymia)	NM_000217	KCNA1	$1.64E-01$	2.00		
Potassium voltage-gated channel, shaker-related subfamily, member 1 (episodic ataxia with myokymia)	NM_000217	KCNA1	$1.31E-01$	1.87		
Potassium voltage-gated channel, shaker-related subfamily, member 2, transcript variant 2	NM_001204269	KCNA2	$8.55E-02$	1.33		

(j) Continued.

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Potassium voltage-gated channel, shaker-related subfamily, member 2, transcript variant 1	NM_004974	KCNA2	8.33E – 01	1.06		
Potassium voltage-gated channel, shaker-related subfamily, member 3	NM_002232	KCNA3	2.43E – 01	1.76		
Potassium voltage-gated channel, shaker-related subfamily, member 4	NM_002233	KCNA4	5.75E – 01	1.11		
Potassium voltage-gated channel, shaker-related subfamily, member 5	NM_002234	KCNA5	1.76E – 04	4.11		5.63
Potassium voltage-gated channel, shaker-related subfamily, member 6	NM_002235	KCNA6	7.20E – 02	2.28		
Potassium voltage-gated channel, shaker-related subfamily, member 7	NM_031886	KCNA7	8.53E – 03	3.67		
Potassium voltage-gated channel, shaker-related subfamily, member 7	NM_031886	KCNA7	1.04E – 01	2.39		
Potassium voltage-gated channel, shaker-related subfamily, member 10	NM_005549	KCNA10	3.32E – 01	–1.62		
Potassium voltage-gated channel, shaker-related subfamily, beta member 1, transcript variant 2	NM_003471	KCNAB1	2.70E – 01	1.46		
Potassium voltage-gated channel, shaker-related subfamily, beta member 1, transcript variant 2	NM_003471	KCNAB1	3.60E – 02	2.42		
Potassium voltage-gated channel, shaker-related subfamily, beta member 1, transcript variant 2	NM_003471	KCNAB1	7.87E – 02	3.06		
Potassium voltage-gated channel, shaker-related subfamily, beta member 2, transcript variant 1	NM_003636	KCNAB2	5.35E – 01	–1.35		
Potassium voltage-gated channel, shaker-related subfamily, beta member 2, transcript variant 1	NM_003636	KCNAB2	2.20E – 01	1.26		
Potassium voltage-gated channel, shaker-related subfamily, beta member 3 [Source: HGNC Symbol; Acc: 6230]	ENST00000303790	KCNAB3	8.14E – 02	1.62		
Potassium voltage-gated channel, shaker-related subfamily, beta member 3	NM_004732	KCNAB3	3.66E – 01	1.35		
Potassium voltage-gated channel, Shab-related subfamily, member 1	NM_004975	KCNB1	7.42E – 08	16.32	2.59	
Potassium voltage-gated channel, Shab-related subfamily, member 2	NM_004770	KCNB2	1.54E – 05	4.58		
Potassium voltage-gated channel, Shaw-related subfamily, member 1, transcript variant A	NM_001112741	KCNC1	1.47E – 01	1.75		
Potassium voltage-gated channel, Shaw-related subfamily, member 1, transcript variant B	NM_004976	KCNC1	4.86E – 08	13.97		
Potassium voltage-gated channel, Shaw-related subfamily, member 1, transcript variant A	NM_001112741	KCNC1	2.17E – 01	1.56		
Potassium voltage-gated channel, Shaw-related subfamily, member 2, transcript variant 1	NM_139136	KCNC2	8.53E – 01	1.06		
Potassium voltage-gated channel, Shaw-related subfamily, member 2, transcript variant 2	NM_139137	KCNC2	4.04E – 01	1.34		
Potassium voltage-gated channel, Shaw-related subfamily, member 3	NM_004977	KCNC3	3.56E – 01	1.20		
Potassium voltage-gated channel, Shaw-related subfamily, member 4, transcript variant 3	NM_001039574	KCNC4	7.70E – 02	–1.40		
Potassium voltage-gated channel, Shal-related subfamily, member 1	NM_004979	KCND1	1.48E – 04	7.11		
Potassium voltage-gated channel, Shal-related subfamily, member 2	NM_012281	KCND2	1.62E – 12	91.08	2.76	–2.57

(j) Continued.

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Potassium voltage-gated channel, Shal-related subfamily, member 2	NM.012281	KCND2	2.76E – 09	21.83		
Potassium voltage-gated channel, Shal-related subfamily, member 3 [Source: HGNC Symbol; Acc: 6239]	ENST00000369697	KCND3	5.36E – 01	–1.36		
Potassium voltage-gated channel, Shal-related subfamily, member 3, transcript variant 1	NM.004980	KCND3	1.05E – 01	1.55		
Potassium voltage-gated channel, Shal-related subfamily, member 3, transcript variant 1	NM.004980	KCND3	4.41E – 01	1.37		
Potassium voltage-gated channel, Shal-related subfamily, member 3 [Source: HGNC Symbol; Acc: 6239]	ENST00000369697	KCND3	3.86E – 04	–1.84		
Potassium voltage-gated channel, Isk-related family, member 1, transcript variant 2	NM.000219	KCNE1	3.53E – 01	–1.45		
Potassium voltage-gated channel, Isk-related family, member 1, transcript variant 2	NM.000219	KCNE1	3.59E – 01	1.25		
KCNE1-like	NM.012282	KCNE1L	1.34E – 08	4.44		–2.18
KCNE1-like	NM.012282	KCNE1L	1.11E – 08	3.65		
KCNE1-like	NM.012282	KCNE1L	6.41E – 03	1.88		
Potassium voltage-gated channel, Isk-related family, member 2	NM.172201	KCNE2	1.41E – 04	3.93		
Potassium voltage-gated channel, Isk-related family, member 3	NM.005472	KCNE3	8.65E – 15	52.36		
Potassium voltage-gated channel, Isk-related family, member 4	NM.080671	KCNE4	5.04E – 01	1.20		
Potassium voltage-gated channel, subfamily F, member 1	NM.002236	KCNF1	2.25E – 05	6.48	2.84	
Potassium voltage-gated channel, subfamily G, member 1	NM.002237	KCNG1	8.51E – 03	–1.82		
Potassium voltage-gated channel, subfamily G, member 1	NM.002237	KCNG1	1.55E – 01	–1.65		
Potassium voltage-gated channel, subfamily G, member 2	NM.012283	KCNG2	6.21E – 02	1.32		
Potassium voltage-gated channel, subfamily G, member 3, transcript variant 1	NM.133329	KCNG3	3.74E – 11	58.95		–2.93
Potassium voltage-gated channel, subfamily G, member 4	NM.172347	KCNG4	3.36 E–03	–3.33		
Potassium voltage-gated channel, subfamily G, member 4, (cDNA clone IMAGE: 3028985)	BC008969	KCNG4	2.24 E–04	–3.29		
Potassium voltage-gated channel, subfamily H (eag-related), member 1, transcript variant 1	NM.172362	KCNH1	8.05E – 01	–1.15		
Potassium voltage-gated channel, subfamily H (eag-related), member 2, transcript variant 1	NM.000238	KCNH2	5.94E – 04	37.53		
Potassium voltage-gated channel, subfamily H (eag-related), member 2, transcript variant 2	NM.172056	KCNH2	5.70E – 03	1.73		
Potassium voltage-gated channel, subfamily H (eag-related), member 3	NM.012284	KCNH3	9.65E – 01	1.01		
Potassium voltage-gated channel, subfamily H (eag-related), member 4	NM.012285	KCNH4	3.18E – 01	–1.24		

(j) Continued.

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Potassium voltage-gated channel, subfamily H (eag-related), member 5, transcript variant 2	NM.172376	KCNH5	3.04E – 03	2.96		
Potassium voltage-gated channel, subfamily H (eag-related), member 5, transcript variant 1	NM.139318	KCNH5	5.41E – 01	1.47		
Potassium voltage-gated channel, subfamily H (eag-related), member 6, transcript variant 2	NM.173092	KCNH6	2.93E – 02	3.04		
cDNA FLJ33650 fis, clone BRAMY2024514, highly similar to Rattus norvegicus Potassium channel (erg2)	AK090969	KCNH6	2.08E – 05	12.42		
Potassium voltage-gated channel, subfamily H (eag-related), member 6, transcript variant 2	NM.173092	KCNH6	2.55E – 02	2.29		
Potassium voltage-gated channel, subfamily H (eag-related), member 7, transcript variant 2	NM.173162	KCNH7	4.63E – 01	–1.40		
Potassium voltage-gated channel, subfamily H (eag-related), member 7, transcript variant 1	NM.033272	KCNH7	7.58E – 01	–1.09		
Potassium voltage-gated channel, subfamily H (eag-related), member 8	NM.144633	KCNH8	3.00E – 09	24.57	2.97	
Kv channel interacting protein 1, transcript variant 1	NM.001034837	KCNIP1	8.33E – 02	1.96	4.24	
Kv channel interacting protein 2, transcript variant 7	NM.173197	KCNIP2	6.43E – 01	1.21		
Kv channel interacting protein 2, transcript variant 1	NM.014591	KCNIP2	2.53E – 03	1.96		
Kv channel interacting protein 3, calsenilin, transcript variant 1	NM.013434	KCNIP3	1.15 E–03	–3.47		
Kv channel interacting protein 4, transcript variant 5	NM.001035003	KCNIP4	6.73E – 01	1.16		2.01
Kv channel interacting protein 4, transcript variant 5	NM.001035003	KCNIP4	9.26E – 01	1.05		
Potassium voltage-gated channel, KQT-like subfamily, member 1, transcript variant 1	NM.000218	KCNQ1	3.31E – 09	79.52		
Potassium voltage-gated channel, KQT-like subfamily, member 2, transcript variant 5	NM.172109	KCNQ2	5.68E – 19	3408.43		
Potassium voltage-gated channel, KQT-like subfamily, member 2, transcript variant 3	NM.004518	KCNQ2	1.29E – 02	1.82		
Potassium voltage-gated channel, KQT-like subfamily, member 2, (cDNA clone IMAGE: 4154700)	BC020384	KCNQ2	1.09E – 08	37.12		
Potassium voltage-gated channel, KQT-like subfamily, member 3, transcript variant 1	NM.004519	KCNQ3	1.10E – 04	–1.69		
Potassium voltage-gated channel, KQT-like subfamily, member 4, transcript variant 1	NM.004700	KCNQ4	9.77E – 01	1.02		
Potassium voltage-gated channel, KQT-like subfamily, member 5, transcript variant 1	NM.019842	KCNQ5	9.81E – 02	–1.79		
Potassium voltage-gated channel, delayed-rectifier, subfamily S, member 1	NM.002251	KCNS1	3.39E – 01	–1.27		–2.64
Potassium voltage-gated channel, delayed-rectifier, subfamily S, member 2	NM.020697	KCNS2	6.19E – 02	–2.45	2.11	
Potassium voltage-gated channel, delayed-rectifier, subfamily S, member 3	NM.002252	KCNS3	1.04E – 04	2.22	–2.61	–2.39
Potassium channel, subfamily T, member 1 (sodium activated)	NM.020822	KCNT1	8.51E – 02	–2.17		

(j) Continued.

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Potassium channel, subfamily V, member 1	NM_014379	KCNV1	$7.84E - 02$	1.58		
Potassium channel, subfamily V, member 2	NM_133497	KCNV2	$7.63E - 01$	-1.16		

(k) Inwardly rectifying potassium channels

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Potassium inwardly-rectifying channel, subfamily J, member 1, transcript variant rom-k5	NM_153767	KCNJ1	$2.54E - 02$	2.66		
Potassium inwardly-rectifying channel, subfamily J, member 1, transcript variant rom-k5	NM_153767	KCNJ1	$8.51E - 01$	-1.05		
Potassium inwardly-rectifying channel, subfamily J, member 2	NM_000891	KCNJ2	$5.14E - 04$	4.38		
Potassium inwardly-rectifying channel, subfamily J, member 3	NM_002239	KCNJ3	$4.56E - 01$	1.34		
Potassium inwardly-rectifying channel, subfamily J, member 4, transcript variant 1	NM_152868	KCNJ4	$4.80E - 02$	2.30		
Potassium inwardly-rectifying channel, subfamily J, member 5	NM_000890	KCNJ5	$1.88E - 08$	-45.40		
Potassium inwardly-rectifying channel, subfamily J, member 5	NM_000890	KCNJ5	$4.42E - 12$	-109.11		
Potassium inwardly-rectifying channel, subfamily J, member 6	NM_002240	KCNJ6	$3.29E - 05$	6.75	5.54	
Potassium inwardly-rectifying channel, subfamily J, member 8	NM_004982	KCNJ8	$1.98E - 04$	5.97		2.74
Potassium inwardly-rectifying channel, subfamily J, member 9 [Source: HGNC Symbol; Acc: 6270]	ENST00000368088	KCNJ9	$3.17E - 01$	1.49		
Potassium inwardly-rectifying channel, subfamily J, member 9	NM_004983	KCNJ9	$3.07E - 01$	1.79		
Potassium inwardly-rectifying channel, subfamily J, member 10	NM_002241	KCNJ10	$8.85E - 04$	4.14		
Potassium inwardly-rectifying channel, subfamily J, member 10	NM_002241	KCNJ10	$9.56E - 01$	1.01		
Potassium inwardly-rectifying channel, subfamily J, member 11, transcript variant 1	NM_000525	KCNJ11	$4.00E - 04$	2.93		
Potassium inwardly-rectifying channel, subfamily J, member 12	NM_021012	KCNJ12	$1.00E - 01$	1.47		
Potassium inwardly-rectifying channel, subfamily J, member 12	NM_021012	KCNJ12	$3.53E - 06$	5.81		
Potassium inwardly-rectifying channel, subfamily J, member 13, transcript variant 1	NM_002242	KCNJ13	$7.75E - 01$	1.20		
Potassium inwardly-rectifying channel, subfamily J, member 14, transcript variant 2	NM_170720	KCNJ14	$9.05E - 02$	-1.40		
Potassium inwardly-rectifying channel, subfamily J, member 15, transcript variant 1	NM_170736	KCNJ15	$3.41E - 08$	-150.95		
Potassium inwardly-rectifying channel, subfamily J, member 16, transcript variant 2	NM_170741	KCNJ16	$2.22E - 01$	-1.93		

(l) Two-P potassium channels

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Potassium channel, subfamily K, member 1	NM_002245	KCNK1	3.02E – 04	–1.68		
Potassium channel, subfamily K, member 2, transcript variant 1	NM_001017424	KCNK2	3.46E – 01	1.57		
Potassium channel, subfamily K, member 3	NM_002246	KCNK3	9.98E – 01	1.00	2.36	
Potassium channel, subfamily K, member 3	NM_002246	KCNK3	8.75E – 01	–1.10		
Potassium channel, subfamily K, member 3	NM_002246	KCNK3	5.34E – 01	–1.30		
Potassium channel, subfamily K, member 3 [Source: HGNC Symbol; Acc: 6278]	ENST00000302909	KCNK3	2.53E – 01	1.52		
Potassium channel, subfamily K, member 4	NM_033310	KCNK4	6.21E – 01	–1.14	2.05	
Potassium channel, subfamily K, member 5	NM_003740	KCNK5	1.16E – 14	29.46	–2.35	–2.03
Potassium channel, subfamily K, member 6	NM_004823	KCNK6	1.22E – 06	–3.48	–2.20	
Potassium channel, subfamily K, member 6	NM_004823	KCNK6	2.05E – 08	–5.11		
Potassium channel, subfamily K, member 6	NM_004823	KCNK6	7.04E – 06	–5.22		
Potassium channel, subfamily K, member 7, transcript variant A	NM_033347	KCNK7	1.17E – 01	–1.52		
Potassium channel, subfamily K, member 9	NM_016601	KCNK9	6.67E – 01	–1.29		
Potassium channel, subfamily K, member 10, transcript variant 1	NM_021161	KCNK10	2.63E – 01	–1.30		
Potassium channel, subfamily K, member 10, transcript variant 2	NM_138317	KCNK10	1.41E – 02	1.43		
Potassium channel, subfamily K, member 12	NM_022055	KCNK12	4.58E – 10	23.40		–5.58
Potassium channel, subfamily K, member 13	NM_022054	KCNK13	7.65E – 01	–1.10		
Potassium channel, subfamily K, member 15	NM_022358	KCNK15	4.96E – 02	–1.50		
Potassium channel, subfamily K, member 15	NM_022358	KCNK15	9.10E – 01	–1.09		
Pancreatic potassium channel TALK-1d; alternatively spliced	AY253147	KCNK16	2.17E – 02	–2.43		
Potassium channel, subfamily K, member 17, transcript variant 1	NM_031460	KCNK17	1.87E – 11	28.60		
Potassium channel, subfamily K, member 17, transcript variant 1	NM_031460	KCNK17	1.52E – 05	19.62		
Potassium channel, subfamily K, member 18	NM_181840	KCNK18	1.77E – 01	1.61		

(m) Calcium-activated potassium channels

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Potassium large conductance calcium-activated channel, subfamily M, alpha member 1, transcript variant 2	NM_002247	KCNMA1	1.01E – 02	1.84		2.01
Potassium large conductance calcium-activated channel, subfamily M, alpha member 1, transcript variant 1	NM_001014797	KCNMA1	3.19E – 01	1.32		
Potassium large conductance calcium-activated channel, subfamily M, alpha member 1, transcript variant 2	NM_002247	KCNMA1	3.67E – 03	2.27		
Potassium large conductance calcium-activated channel, subfamily M, alpha member 1, transcript variant 2	NM_002247	KCNMA1	8.53E – 02	1.44		
Maxi-K channel HSLO	AF349445	KCNMA1	3.04E – 01	1.44		

(m) Continued.

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Potassium large conductance calcium-activated channel, subfamily M, alpha member 1, transcript variant 2	NM.002247	KCNMA1	$2.15E - 01$	-1.23		
Potassium large conductance calcium-activated channel, subfamily M, beta member 1	NM.004137	KCNMB1	$7.12E - 02$	-1.41		
Potassium large conductance calcium-activated channel, subfamily M, beta member 1	NM.004137	KCNMB1	$2.23E - 06$	7.82		
Potassium large conductance calcium-activated channel, subfamily M, beta member 2, transcript variant 1	NM.181361	KCNMB2	$3.36E - 01$	1.56		
Potassium large conductance calcium-activated channel, subfamily M, beta member 2, transcript variant 1	NM.181361	KCNMB2	$1.17E - 01$	1.47		
Potassium large conductance calcium-activated channel, subfamily M beta member 3, transcript variant 1	NM.171828	KCNMB3	$1.41E - 05$	2.63		
Potassium large conductance calcium-activated channel, subfamily M, beta member 4	NM.014505	KCNMB4	$8.85E - 12$	103.99		
Potassium large conductance calcium-activated channel, subfamily M, beta member 4 [Source: HGNC Symbol; Acc: 6289]	ENST00000258111	KCNMB4	$9.22E - 10$	18.40		
Potassium intermediate/small conductance calcium-activated channel, subfamily N, member 1	NM.002248	KCNN1	$3.40E - 05$	4.67		
Potassium intermediate/small conductance calcium-activated channel, subfamily N, member 2, transcript variant 1	NM.021614	KCNN2	$1.88E - 13$	22.23	-3.63	
Potassium intermediate/small conductance calcium-activated channel, subfamily N, member 3, transcript variant 1	NM.002249	KCNN3	$2.58E - 06$	8.25		
Potassium intermediate/small conductance calcium-activated channel, subfamily N, member 3, transcript variant 3	NM.001204087	KCNN3	$1.11E - 07$	15.29		
Potassium intermediate/small conductance calcium-activated channel, subfamily N, member 4	NM.002250	KCNN4	$1.06E - 02$	-1.84		

(n) P2X receptors

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Purinergic receptor P2X, ligand-gated ion channel, 1	NM.002558	P2RX1	$7.71E - 01$	1.14		
Purinergic receptor P2X, ligand-gated ion channel, 2, transcript variant 4	NM.170683	P2RX2	$5.25E - 02$	-1.66		
Purinergic receptor P2X, ligand-gated ion channel, 3	NM.002559	P2RX3	$2.43E - 02$	-1.76		
Purinergic receptor P2X, ligand-gated ion channel, 4	NM.002560	P2RX4	$8.68E - 01$	-1.03		
Purinergic receptor P2X, ligand-gated ion channel, 5, transcript variant 1	NM.002561	P2RX5	$7.68E - 08$	6.97		

(n) Continued.

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Purinergic receptor P2X, ligand-gated ion channel, 5, transcript variant 2	NM_175080	P2RX5	6.13E – 02	1.54		
Purinergic receptor P2X, ligand-gated ion channel, 6 [Source: HGNC Symbol; Acc: 8538]	ENST00000413302	P2RX6	4.13E – 01	1.68		
Purinergic receptor P2X, ligand-gated ion channel, 7, transcript variant 1	NM_002562	P2RX7	6.13E – 01	1.23		
Purinergic receptor P2X, ligand-gated ion channel, 7, transcript variant 1	NM_002562	P2RX7	9.17E – 01	1.04		

(o) Transient receptor potential channels

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Polycystic kidney disease 2 (autosomal dominant)	NM_000297	PKD2	7.48E – 07	1.79		
Polycystic kidney disease 2 (autosomal dominant)	NM_000297	PKD2	2.38E – 05	1.98		
Polycystic kidney disease 2-like 1	NM_016112	PKD2L1	1.29E – 03	4.66		
Polycystic kidney disease 2-like 2	NM_014386	PKD2L2	1.94E – 01	1.49		
Polycystic kidney disease 2-like 2	NM_014386	PKD2L2	7.43E – 01	1.19		
Polycystic kidney disease 2-like 2	NM_014386	PKD2L2	6.74E – 01	1.09		
Transient receptor potential cation channel, subfamily A, member 1	NM_007332	TRPA1	2.99E – 01	1.74		
Transient receptor potential cation channel, subfamily A, member 1	NM_007332	TRPA1	5.64E – 02	2.70		
Transient receptor potential cation channel, subfamily C, member 1	NM_003304	TRPC1	4.30E – 06	2.01		
Transient receptor potential cation channel, subfamily C, member 3, transcript variant 2	NM_003305	TRPC3	1.12E – 07	5.78		2.45
Transient receptor potential cation channel, subfamily C, member 4, transcript variant alpha	NM_016179	TRPC4	3.25E – 07	25.38		
Transient receptor potential cation channel, subfamily C, member 5	NM_012471	TRPC5	1.20E – 01	1.28		
Transient receptor potential cation channel, subfamily C, member 6	NM_004621	TRPC6	1.21E – 01	2.33		
Transient receptor potential cation channel, subfamily C, member 7, transcript variant 1	NM_020389	TRPC7	6.18E – 01	–1.36		
Transient receptor potential cation channel, subfamily M, member 1	NM_002420	TRPM1	2.29E – 01	1.31		
Transient receptor potential cation channel, subfamily M, member 2, transcript variant 1	NM_003307	TRPM2	1.22E – 01	1.97		
Transient receptor potential cation channel, subfamily M, member 2, transcript variant 1	NM_003307	TRPM2	5.04E – 02	2.53		
Transient receptor potential cation channel, subfamily M, member 3, transcript variant 7	NM_206948	TRPM3	3.04E – 09	10.69		
Transient receptor potential cation channel, subfamily M, member 3 [Source: HGNC Symbol; Acc: 17992]	ENST00000354500	TRPM3	2.34E – 03	3.74		
Transient receptor potential cation channel, subfamily M, member 3, transcript variant 7	NM_206948	TRPM3	3.41E – 06	6.23		
Transient receptor potential cation channel, subfamily M, member 3, transcript variant 9	NM_001007471	TRPM3	7.23E – 01	1.13		
Transient receptor potential cation channel, subfamily M, member 3, transcript variant 9	NM_001007471	TRPM3	1.54E – 05	2.60		

(o) Continued.

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Transient receptor potential cation channel, subfamily M, member 4, transcript variant 1	NM_017636	TRPM4	$3.72E - 02$	-1.89		
Transient receptor potential cation channel, subfamily M, member 5	NM_014555	TRPM5	$8.21E - 01$	-1.08		
Transient receptor potential cation channel, subfamily M, member 6, transcript variant a	NM_017662	TRPM6	$3.80E - 01$	1.64		
Transient receptor potential cation channel, subfamily M, member 6, transcript variant a	NM_017662	TRPM6	$9.10E - 01$	1.05		
Transient receptor potential cation channel, subfamily M, member 6, transcript variant a	NM_017662	TRPM6	$5.32E - 07$	5.59		
Transient receptor potential cation channel, subfamily M, member 6, transcript variant a	NM_017662	TRPM6	$3.32E - 01$	1.36		
Transient receptor potential cation channel, subfamily M, member 7	NM_017672	TRPM7	$2.16E - 02$	-1.25		
Transient receptor potential cation channel, subfamily M, member 7	NM_017672	TRPM7	$3.64E - 03$	-1.38		
Transient receptor potential cation channel, subfamily M, member 8	NM_024080	TRPM8	$2.04E - 01$	1.89		
Transient receptor potential cation channel, subfamily V, member 1, transcript variant 3	NM_080706	TRPV1	$3.59E - 05$	1.92		
Transient receptor potential cation channel, subfamily V, member 1, transcript variant 3	NM_080706	TRPV1	$4.14E - 05$	2.25		
Transient receptor potential cation channel, subfamily V, member 2	NM_016113	TRPV2	$8.02E - 08$	4.68		
Transient receptor potential cation channel, subfamily V, member 3	NM_145068	TRPV3	$4.64E - 01$	1.35		
Transient receptor potential cation channel, subfamily V, member 3	NM_145068	TRPV3	$1.35E - 01$	-2.39		
Transient receptor potential cation channel, subfamily V, member 4, transcript variant 2	NM_147204	TRPV4	$4.33E - 01$	1.39		
Transient receptor potential cation channel, subfamily V, member 5, (cDNA clone MGC: 34269 IMAGE: 5186668)	BC034740	TRPV5	$1.00E - 01$	1.38		
Transient receptor potential cation channel, subfamily V, member 5	NM_019841	TRPV5	$3.14E - 01$	-1.40		
Transient receptor potential cation channel, subfamily V, member 6	NM_018646	TRPV6	$1.57E - 01$	-1.38		

(p) Voltage-gated sodium channels

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Sodium channel, voltage-gated, type I, alpha subunit, transcript variant 2	NM_006920	SCN1A	$1.98E - 01$	1.54		
Sodium channel, voltage-gated, type I, beta, transcript variant b	NM_199037	SCN1B	$9.95E - 01$	-1.00		
Sodium channel, voltage-gated, type II, alpha subunit, transcript variant 1	NM_021007	SCN2A	$4.16E - 01$	1.62	5.00	
Sodium channel, voltage-gated, type II, beta	NM_004588	SCN2B	$3.61E - 01$	-1.58		
Sodium channel, voltage-gated, type II, beta	NM_004588	SCN2B	$8.16E - 04$	-2.41		

(p) Continued.

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Sodium channel, voltage-gated, type III, alpha subunit, transcript variant 1	NM_006922	SCN3A	$2.87E - 01$	-2.00	2.85	
Sodium channel, voltage-gated, type III, alpha subunit, transcript variant 1	NM_006922	SCN3A	$1.89E - 01$	1.49		
Sodium channel, voltage-gated, type III, beta, transcript variant 1	NM_018400	SCN3B	$8.74E - 01$	-1.06	4.70	
Sodium channel, voltage-gated, type III, beta, transcript variant 1	NM_018400	SCN3B	$1.57E - 01$	-1.46		
Sodium channel, voltage-gated, type III, beta, transcript variant 1	NM_018400	SCN3B	$1.56E - 01$	1.96		
Sodium channel, voltage-gated, type IV, alpha subunit	NM_000334	SCN4A	$1.62E - 08$	9.91		
Sodium channel, voltage-gated, type IV, beta, transcript variant 1	NM_174934	SCN4B	$3.25E - 04$	-3.66		
Sodium channel, voltage-gated, type V, alpha subunit, transcript variant 1	NM_198056	SCN5A	$2.43E - 08$	14.10		
Sodium channel, voltage-gated, type V, alpha subunit, transcript variant 2	NM_000335	SCN5A	$4.93E - 02$	1.95		
Sodium channel, voltage-gated, type VII, alpha	NM_002976	SCN7A	$4.49E - 03$	1.63		
Sodium channel, voltage-gated, type VII, alpha	NM_002976	SCN7A	$2.07E - 01$	1.72		
Sodium channel, voltage-gated, type VIII, alpha subunit, transcript variant 1	NM_014191	SCN8A	$1.89E - 09$	15.61		
Sodium channel, voltage-gated, type VIII, alpha subunit, transcript variant 1	NM_014191	SCN8A	$4.06E - 07$	11.23		
Sodium channel, voltage-gated, type IX, alpha subunit	NM_002977	SCN9A	$4.08E - 08$	11.51		
Sodium channel, voltage-gated, type IX, alpha subunit	NM_002977	SCN9A	$1.62E - 01$	1.66		
Sodium channel, voltage-gated, type IX, alpha subunit	NM_002977	SCN9A	$3.38E - 03$	2.57		
Sodium channel, voltage-gated, type X, alpha subunit	NM_006514	SCN10A	$4.06E - 01$	1.35		
Sodium channel, voltage-gated, type XI, alpha subunit	NM_014139	SCN11A	$3.02E - 01$	1.57		
Sodium channel, voltage-gated, type XI, alpha subunit [Source: HGNC Symbol; Acc: 10583]	ENST00000444237	SCN11A	$8.83E - 03$	1.94		

(q) Nonvoltage-gated sodium channels

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Sodium channel, nonvoltage-gated 1 alpha, transcript variant 1	NM_001038	SCNN1A	$1.33E - 09$	51.49	-21.13	-10.71
Sodium channel, nonvoltage-gated 1, beta	NM_000336	SCNN1B	$2.11E - 01$	1.64		
Sodium channel, nonvoltage-gated 1, beta	NM_000336	SCNN1B	$5.52E - 01$	1.09		
Sodium channel, nonvoltage-gated 1, delta, transcript variant 1	NM_001130413	SCNN1D	$8.43E - 02$	1.49		
Sodium channel, nonvoltage-gated 1, gamma	NM_001039	SCNN1G	$2.33E - 01$	1.59		
Gamma subunit of epithelial amiloride-sensitive sodium channel	X87160	SCNN1G	$3.99E - 01$	1.27		

(r) Two-pore channels

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Two-pore segment channel 1, transcript variant 1	NM_001143819	TPCN1	6.20E − 04	−1.76		
Two-pore segment channel 1, transcript variant 2	NM_017901	TPCN1	1.71 E − 03	−2.10		
Two-pore segment channel 1, transcript variant 2	NM_017901	TPCN1	3.55 E − 04	−2.44		
Two-pore segment channel 1, transcript variant 1	NM_001143819	TPCN1	8.06E − 03	−1.83		
Two-pore segment channel 2	NM_139075	TPCN2	5.06E − 01	−1.15		

(s) Zinc-activated ligand-gated ion channels

Gene name	Accession number	Gene symbol	<i>P</i>	Fold change Ker → hiPSC	Fold change hiPSC → Neuron	Fold change hiPSC → CM
Zinc-activated ligand-gated ion channel	NM_180990	ZACN	9.24E − 02	1.45		

During reprogramming 14 (33%) of them are significantly upregulated while only 1 (2%) is significantly downregulated. From iPSCs to neurons, as expected, neuronal alpha-subunits *CACNAID*, *CACNAIE* are upregulated together with several neuronal channel subunits ($\alpha_2\delta$, β , and γ). In cardiomyocytes, solely the alpha subunit of the cardiac *CACNAIC* is upregulated. Of note, none of the differentiated progeny downregulated any voltage-gated calcium channels.

3.3. Sperm-Associated Cation Channels. Sperm-associated cation channels or CatSper channels are calcium ion channels. They are flagellar proteins involved in sperm motility and therefore affect fertility [27]. During reprogramming to iPSCs, from seven used probes just one showed a significant downregulation and none had a significant upregulation. We observed nothing noteworthy in the differentiated progeny.

3.4. Nicotinic Acetylcholine Receptors. Nicotinic acetylcholine receptors play a role in interneuronal synapses and neuromuscular junctions. They are composed of five subunits as homomeric or heteromeric receptors. They are located at the postsynaptic site and upon binding of acetylcholine they allow the transmission of cations, especially sodium and potassium ions, in some versions also calcium ions. This leads to a depolarization of the membrane and triggers further signaling pathways [28]. Several acetylcholine receptors and subunits are thought to play roles in a variety of pathomechanisms, for example, psychiatric disorders, cardiovascular diseases, or cancer [29–31]. From 21 probes, 5 (24%) showed a significant upregulation while 3 (14%) showed a significant downregulation from keratinocytes to iPSCs. In neurons, solely the neuronal nicotinic acetylcholine receptor *CHRNA6* alpha subunit was upregulated, none in cardiomyocytes.

3.5. Cyclic Nucleotide-Gated Channels. Cyclic nucleotide-gated channels form tetrameric channels which—upon binding of cGMP—allow a flow of cations. For that, these channels track the intracellular concentration of cNMPs to produce a

voltage response [32]. Their major role is the depolarization of rod photoreceptors, but they are also found in other tissues like olfactory sensory neurons [33], testis, kidney, or heart [34] and play a role in cellular development such as neuronal growth cone guidance [35]. Defects in these genes are reported to cause retinitis pigmentosa [36]. From 7 used probes one showed a significant upregulation and one showed a significant downregulation (both 14%) during reprogramming, while none was differentially regulated in differentiated neurons or cardiomyocytes.

3.6. GABA Receptors. GABA (gamma-aminobutyric acid) receptors are ligand-gated chloride channels. Since GABA is the main inhibitory neurotransmitter in the central nervous system GABA receptors play an important role for the brain function. The receptors are composed of five subunits which form heteromers. The GABA receptors are a drug target for anesthetics and other psychoactive drugs. We have analyzed 29 probes within GABA receptor subunits. From keratinocytes to iPSCs, 6 (21%) of them are significantly upregulated while just one (3%) is significantly downregulated. Interestingly, none was up- or downregulated in differentiated neurons, while *GABRB3* was downregulated and *GABRP* was upregulated in cardiomyocytes. Up to now, more or less nothing is known about GABA receptors or their subunits in cardiac cells.

3.7. Glycine Receptors. Glycine receptors are inhibitory receptors of the postsynaptic site. They are activated by glycine and mediate an influx of chloride ions. Accordingly, GlyRs regulate not only the excitability of motor and sensory neurons but are also essential for the processing of photoreceptor signals, neuronal development, and inflammatory pain sensitization [37]. The heteromeric pore is formed by five subunits. Concerning their role in pathomechanisms, it was reported that mutations are causing hyperekplexia (also known as startle disease) [38]. We have analyzed 7 probes and none of them was significantly altered in hiPSCs.

Transcript levels for the alpha2 subunit and the beta subunit were highly upregulated in differentiated neurons, pointing to their functional role in the nervous system.

3.8. Ionotropic Glutamate Receptors. Glutamate is the predominant excitatory neurotransmitter in the central nervous system. Therefore, ionotropic glutamate receptors play a key role for learning and memory processes. They are located in the postsynaptic membrane and are composed of several heteromeric subunits. Ionotropic glutamate receptors are further divided into AMPA (alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate), NMDA (N-Methyl-D-aspartate) or kainate receptors depending on their sensitivity for the agonists. From 36 probes, 5 (14%) showed a significantly increased expression and 2 (6%) showed a decreased expression in iPSCs compared to keratinocytes. As expected, several subunits were upregulated in neurons (none downregulated). None was upregulated in cardiomyocytes, while the kainate subtype *GRIK5* was downregulated.

3.9. Hyperpolarization-Activated Cyclic Nucleotide-Gated Channels. Hyperpolarization-activated cyclic nucleotide-gated channels are homodimers or heterodimers and form a hyperpolarization-activated potassium channel. CNG channels display a very complex heteromeric structure with various subunits and domains that play a critical role in their function [39]. They contribute to pacemaker currents in the heart [40] but are also found in neurons [41]. We have analyzed 5 probes and 3 (60%) of them showed a significant upregulation while none was downregulated after reprogramming. In neurons the *HCN3* channel transcript was upregulated, which plays a role in several neuronal functions including excitability of basal ganglia output neurons [42]. Although *HCN4* for example plays a critical role in the conduction system of the heart, we did not observe a specific regulation in cardiomyocytes.

3.10. Serotonin Receptors. Serotonin (or 5-hydroxytryptamine) receptors are ligand-gated receptors mainly found presynaptically in neurons. The type 3 receptor is the only ion channel while the other serotonin receptors are G-protein-coupled receptors. It forms a heteropentameric pore which upon activation by serotonin allows the flow of sodium and potassium, leading to a depolarization. Serotonin receptors modulate neuronal function and are therefore involved in various brain functions. The analysis of 6 probes for type 3 serotonin receptors showed a significant upregulation of 2 (33%, both for *HTR3A*) probes while none was downregulated in hiPSCs. Although we expected serotonin receptors to be upregulated during neuronal differentiation, no subunit was upregulated in neurons or cardiomyocytes, while only the receptor transcript for *HTR3A* was downregulated in neurons.

3.11. Voltage-Gated Potassium Channels. Voltage-gated potassium channels are composed of a large group of subunits with different characteristics concerning for example their inactivation speed. Functional channels

are formed by heterotetramers. The channels are highly specific for potassium with a low affinity for sodium or other cations. Voltage-gated potassium channels are responsible for the repolarization of excitable cells following the sodium-mediated excitation of an action potential and are therefore found in neurons and other cells displaying action potentials. We have analyzed 81 probes. From keratinocytes to hiPSCs, 25 (31%) were significantly upregulated while only 3 (4%) were significantly downregulated. In neurons, several subunits were upregulated including *KCNB1*, coding for Kv2.1 and *KCND2*, both are better known for their role in cardiac cell excitability as well as *KCNF1*, *KCNH8*, and *KCNIP1*, all known for their contribution in neuronal excitability [43]. It should be noted that *KCNT1* is actually sodium activated but is included in the alphabetical list for a better overview. Further, several “S” subunits were upregulated (*KCNS2* in neurons) or downregulated (*KCNS3* in neurons; *KCNS1* and 3 in cardiomyocytes). These subunits are unable to form functional channels as homotetramers but instead heterotetramerize with other alpha-subunits to form conductive channels. These subunits are involved in modifying the channels response and conductivity [44]. Few is known about distinct roles in other tissues, but they were associated with for example, pain modulation [45] or airway responsiveness [46].

3.12. Inwardly Rectifying Potassium Channels. Inwardly rectifying potassium channels have a higher tendency to allow the flow of potassium ions into the cell rather than to the outside of the cell. Therefore, they play an important role in the maintenance of the resting membrane potential. Their activation is constitutive or controlled by ATP binding and G-proteins [47]. Functional channels are formed as homo- or heterotetramers. These channels can be found predominantly in neurons, cardiac myocytes, the pancreas, or the kidneys. From 20 analyzed probes in hiPSCs compared to keratinocytes, 8 (40%) showed a significant upregulation and 3 (15%) were significantly downregulated. The *GIRK2* channel encoded by *KCNJ6* is upregulated in neurons playing multiple roles in various tissues including the pancreas and brain [48, 49] and is associated with epileptic seizures in mice lacking the gene [50]. On the other hand, *Kir6.1*, encoded by *KCNJ8*, is upregulated in cardiomyocytes and has been reported to be involved in the pathogenesis of cardiac arrest in the early repolarization syndrome [51].

3.13. Two-P Potassium Channels. Two-P potassium channels contain two pore-forming P domains. After dimerization they form an outward rectifying potassium channel. They can be found in several tissues and are activated by various chemical or physical means (TRAAK channels). We have analyzed 23 probes. 4 (17%) of them were significantly upregulated while 4 (17%) were significantly downregulated. In neurons *KCNK3*, vital for setting the resting membrane potential and primary target for volatile anesthetics [52] as well as *KCNK4*, which is mechanically gated and contributes to axonal pathfinding, growth cone motility, and neurite elongation, as well as possibly having a role in touch or

pain detection [53, 54], were upregulated. *KCNK5* and 6 were downregulated. In cardiomyocytes, downregulation of *KCNK5* and 12 was observed.

3.14. Calcium-Activated Potassium Channels. Calcium-activated potassium channels are mostly activated by intracellular calcium; some family members are also voltage gated. The family consists of large, intermediate, and small conductance family members. Channels are formed by two units (*KCNM* family) or most commonly four units (*KCNN* family). They are involved in, for example, afterhyperpolarization following the action potential and are predominantly found in neurons. Additionally, they are known to play different roles in cellular mechanisms, including stem cell biology [4–6, 55]. We have analyzed 18 probes of which 9 (50%) were significantly upregulated and none was significantly downregulated after reprogramming. In neurons, solely *KCNN2* was downregulated, while in cardiomyocytes *KCNMA1*, encoding the large conductance BK-channel, involved in heart rate regulation [56], was upregulated.

3.15. P2X Receptors. P2X receptors are receptors for extracellular ATP and upon activation open a channel for ions, predominantly calcium. The channel is formed by homo- or heterotrimers. They are found in several tissues, mainly in the nervous system and muscle tissue. They are involved in a range of physiological processes such as modulation of synaptic transmission, vascular tone, cardiac rhythm, and contractility and immune response [57–61]. In stem cells an influence of P2X receptors on embryonic stem cell proliferation was reported [62]. From the 9 analyzed probes 1 (11%) was significantly upregulated and none downregulated. None of these receptors/channels was regulated in differentiated progeny.

3.16. Transient Receptor Potential Channels. Transient receptor potential channels (TRP channels) are nonselective cation channels. They show different preferences for cations, as well as different activation mechanisms and functions. TRP channels are broadly expressed throughout the organism and mediate multiple functions. These include amongst others sensor activity for a wide range of hypertrophic stimuli and mutations in *TRPM4* are now recognized as causes of human cardiac conduction disorders (reviewed in [63]). Furthermore, TRP channels are related to the onset or progression of several diseases, and defects in the genes encoding TRP channels (so-called “TRP channelopathies”) underlie certain neurodegenerative disorders due to their abnormal Ca^{2+} signaling properties (reviewed in [64]). Additionally, TRP channels influence stem cell differentiation and survival [65, 66] and are involved in neuronal-stem-cell-derived development [67]. We have analyzed 40 probes. After reprogramming, 11 (28%) of them were significantly upregulated and none was significantly downregulated. In neurons, none of these channels was noteworthy regulated while solely *TRPC3* was upregulated in cardiomyocytes, reported to be involved in conduction disturbances induced

by adenosine receptor A1AR by enhanced Ca^{2+} entry through the *TRPC3* channel [68].

3.17. Voltage-Gated Sodium Channels. Voltage-gated sodium channels consist of a main alpha unit and some optional modulating or regulatory subunits. They are highly selective for sodium and are involved in a variety of cellular functions including action potential formation [69]. From 24 probes we have analyzed, 6 (25%) were significantly upregulated and two (8%) were significantly downregulated from keratinocytes to iPS cells. These channels seem to be expressed in different kinds of stem cells and during development [70–72]. It was further reported that, for example, *SCN5A*, highly upregulated in iPS cells, is involved in cancer stem cell invasion [73]. *SCN2A*, *SCN3A*, and *SCN3B* were upregulated in neurons, described to be involved in neuronal excitation and epilepsy pathogenesis [74]. Although voltage-gated sodium channels play multiple roles also in the cardiac system [70, 75, 76], we observed no changes of this channel family in cardiomyocytes.

3.18. Nonvoltage-Gated Sodium Channels. Epithelial nonvoltage-gated sodium channels are amiloride sensitive. They form heterotrimers and are involved in ion and fluid transport across epithelia in several organs. We have analyzed 6 probes and one (17%) of them was significantly upregulated while none was significantly downregulated. Interestingly, the upregulated *SCNN1A* in iPS cells was subsequently downregulated both in neurons and cardiomyocytes, pointing to a possible function in stem cells. Of note, it was shown already that repression of pluripotency by retinoic acid represses the *SCNN1A* gene together with several other pluripotency factors [77].

3.19. Two-Pore Channels. Two-pore channels are cation-selective ion channels activated by the second messenger nicotinic acid adenine dinucleotide phosphate (NAADP). Upon activation calcium is released from intracellular stores [78]. We have analyzed 5 probes and while none of them was significantly upregulated two (40%) were significantly downregulated from keratinocytes to iPS cells. None of these channels was regulated in neurons or cardiomyocytes.

3.20. Zinc-Activated Ligand-Gated Ion Channels. The zinc-activated ligand-gated ion channel is activated upon binding of zinc. Until now just one family member is known that is expressed in several tissues [79]. Its exact function is not known. The expression was not significantly altered as shown by one probe in all cells.

4. Discussion

Although ion channels are mainly known for their role in electrically excitable cells they can be found in almost all tissues and are additionally involved in various processes such as cell differentiation and maturation [3, 4, 6, 12, 24, 47, 70]. These large groups of channel proteins are still underestimated concerning their role during embryonic

development and cell fate determination. One of the most interesting *in vitro* models for the elucidation of both developmental processes and disease-specific cellular impairments is represented by pluripotent stem cells. Therefore we were interested in the set of ion channels expressed in hiPSCs after reprogramming and compared this set with iPSC-derived differentiated progeny, namely, neurons and cardiomyocytes. As little is known about ion channels in hiPSCs we aimed to start the analysis with gene expression microarray data. For a comparison, we chose the cells from which they were produced—namely, keratinocytes—as the reference cell type. The comparison of 6 keratinocyte samples with 9 hiPSC samples should minimize the often observed variances between hiPSC lines. We found out that almost a third (32%) of the ion channel probes we investigated showed a significant change in gene expression. Of note, this was mostly an upregulation. Additionally, while many ion channel genes were not expressed in keratinocytes they were present in hiPSCs. This indicates that several of the analyzed channel groups might play unknown roles in stem cell biology, for example, homeostasis, proliferation, or differentiation. Interestingly, after differentiation into neurons or cardiomyocytes, relatively small groups were subsequently regulated. This includes ion channel transcripts playing important roles in the respective tissues. Nevertheless, we compared already published sets of data from different experimental setups and additionally limited the analysis to a strong fold regulation. This might lead to high dropout rates of regulated genes during measurement. Still, various channel transcripts were “logically” up- or downregulated during differentiation into neurons or cardiomyocytes, following embryonic development. iPSC cells and especially patient-specific iPSC cells from persons suffering from genetic mutations leading to hereditary syndromes are a very valuable tool to investigate pathogenetic mechanisms and disease associated molecular and cellular changes [80–82]. As various channel subtypes are involved in multiple pathogenetic mechanisms it would be further interesting to analyze channel transcript regulation in patient specific iPSC cells and their differentiated progeny to elucidate possible disease-specific pathways.

Concerning the presented study it is clear that gene regulations on transcript level do not explicitly mimic either protein levels and posttranslational modifications or protein activity. This set of data is sought to describe a global overview on transcript regulation of ion channels during distinct steps of development. It should be noted that in cases where several probes bind within one gene they do not indicate the same upregulation. Sometimes they show the same trend but miss significance, but for some cases there are considerable differences. This could hint for some yet unknown splicing variants. More detailed studies of these hypothesized splice variants could give insights into their function and broaden the still scarce knowledge.

This work is intended to be a guide and start point for future work focusing on single channels and their composition, localization, and function in hiPSCs and their differentiated progeny. These studies could lead to better *in vitro* differentiation protocols but also explain some of the many disease pathomechanisms related to mutations in ion channel genes.

Conflict of Interests

There is no conflict of interests to declare.

Acknowledgments

The authors thank Sabine Seltenheim for excellent technical assistance enabling research conducted in our laboratories. The research that enabled this paper was funded by a fellowship from the Medical Faculty of Ulm University (Bausteinprogramm, L. SBR.0011) and by the Deutsche Forschungsgemeinschaft (Alexander Kleger KL2544/1-1, Stefan Liebau BO1718/4-1), the German Foundation for Heart Research (F/34/11; to Alexander Kleger and Stefan Liebau), the Else-Kröner-Fresenius-Stiftung (2011_A200; to Alexander Kleger and Stefan Liebau), Boehringer-Ingelheim BIU (N5 to Stefan Liebau; C1 to Alexander Kleger), and the Helmholtz Gesellschaft (VH-VI-510 to Tobias M. Boeckers and Stefan Liebau). Alexander Kleger is indebted to the Baden-Württemberg Stiftung for the financial support of this research project by the Eliteprogramm for Postdocs.

References

- [1] B. Malissen, “A voltage-gated sodium channel mediates positive selection of T cells,” *Nature Immunology*, vol. 13, pp. 810–812, 2012.
- [2] P. Rorsman and M. Braun, “Regulation of insulin secretion in human pancreatic islets,” *Annual Review of Physiology*, vol. 75, pp. 155–179, 2012.
- [3] A. O. Grant, “Cardiac ion channels,” *Circulation*, vol. 2, no. 2, pp. 185–194, 2009.
- [4] S. Liebau, J. Steinestel, L. Linta et al., “An SK3 channel/nWASP/Abi-1 complex is involved in early neurogenesis,” *PLoS ONE*, vol. 6, no. 3, Article ID e18148, 2011.
- [5] S. Liebau, B. Vaida, C. Proepper et al., “Formation of cellular projections in neural progenitor cells depends on SK3 channel activity,” *Journal of Neurochemistry*, vol. 101, no. 5, pp. 1338–1350, 2007.
- [6] A. Kleger, T. Seufferlein, D. Malan et al., “Modulation of calcium-activated potassium channels induces cardiogenesis of pluripotent stem cells and enrichment of pacemaker-like cells,” *Circulation*, vol. 122, no. 18, pp. 1823–1836, 2010.
- [7] K. Takahashi, K. Tanabe, M. Ohnuki et al., “Induction of pluripotent stem cells from adult human fibroblasts by defined factors,” *Cell*, vol. 131, no. 5, pp. 861–872, 2007.
- [8] K. Takahashi and S. Yamanaka, “Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors,” *Cell*, vol. 126, no. 4, pp. 663–676, 2006.
- [9] X. Carvajal-Vergara, A. Sevilla, S. L. Dsouza et al., “Patient-specific induced pluripotent stem-cell-derived models of LEOPARD syndrome,” *Nature*, vol. 465, no. 7299, pp. 808–812, 2010.
- [10] M. Stockmann, L. Linta, K. J. Fohr et al., “Developmental and functional nature of human iPSC derived motoneurons,” *Stem Cell Reviews and Reports*, 2011.
- [11] M. D. Harrell, S. Harbi, J. F. Hoffman, J. Zavadil, and W. A. Coetzee, “Large-scale analysis of ion channel gene expression in the mouse heart during perinatal development,” *Physiological Genomics*, vol. 28, no. 3, pp. 273–283, 2007.

- [12] M. P. Davies, R. H. An, P. Doevendans, S. Kubalak, K. R. Chien, and R. S. Kass, "Developmental changes in ionic channel activity in the embryonic murine heart," *Circulation Research*, vol. 78, no. 1, pp. 15–25, 1996.
- [13] H. Jäger, T. Dreker, A. Buck, K. Giehl, T. Gress, and S. Grissmer, "Blockage of intermediate-conductance Ca^{2+} -Activated K^{+} channels inhibit human pancreatic cancer cell growth in vitro," *Molecular Pharmacology*, vol. 65, no. 3, pp. 630–638, 2004.
- [14] M. Müller, M. Stockmann, D. Malan et al., " Ca^{2+} activated K channels-new tools to induce cardiac commitment from pluripotent stem cells in mice and men," *Stem Cell Reviews and Reports*, vol. 8, no. 3, pp. 720–740, 2012.
- [15] S. P. Wang, J. A. Wang, R. H. Luo, W. Y. Cui, and H. Wang, "Potassium channel currents in rat mesenchymal stem cells and their possible roles in cell proliferation," *Clinical and Experimental Pharmacology and Physiology*, vol. 35, no. 9, pp. 1077–1084, 2008.
- [16] K. J. Brennand, A. Simone, J. Jou et al., "Modelling schizophrenia using human induced pluripotent stem cells," *Nature*, vol. 473, pp. 221–225, 2011.
- [17] G. H. Liu, B. Z. Barkho, S. Ruiz et al., "Recapitulation of premature ageing with iPSCs from Hutchinson-Gilford progeria syndrome," *Nature*, vol. 472, no. 7342, pp. 221–225, 2011.
- [18] R. Maehr, S. Chen, M. Snitow et al., "Generation of pluripotent stem cells from patients with type 1 diabetes," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 106, no. 37, pp. 15768–15773, 2009.
- [19] N. Hofman, A. A. M. Wilde, S. Käb et al., "Diagnostic criteria for congenital long QT syndrome in the era of molecular genetics: do we need a scoring system?" *European Heart Journal*, vol. 28, no. 5, pp. 575–580, 2007.
- [20] M. E. Curran, I. Splawski, K. W. Timothy, G. M. Vincent, E. D. Green, and M. T. Keating, "A molecular basis for cardiac arrhythmia: HERG mutations cause long QT syndrome," *Cell*, vol. 80, no. 5, pp. 795–803, 1995.
- [21] T. Aasen, A. Raya, M. J. Barrero et al., "Efficient and rapid generation of induced pluripotent stem cells from human keratinocytes," *Nature Biotechnology*, vol. 26, no. 11, pp. 1276–1284, 2008.
- [22] T. Aasen and J. C. I. Belmonte, "Isolation and cultivation of human keratinocytes from skin or plucked hair for the generation of induced pluripotent stem cells," *Nature Protocols*, vol. 5, no. 2, pp. 371–382, 2010.
- [23] L. Linta, M. Stockmann, K. N. Kleinhans et al., "Rat embryonic fibroblasts improve reprogramming of human keratinocytes into induced pluripotent stem cells," *Stem Cells and Development*, vol. 21, no. 6, pp. 965–976, 2011.
- [24] W. A. Catterall, "Voltage-gated calcium channels," *Cold Spring Harbor Perspectives in Biology*, vol. 3, no. 8, Article ID a003947, 2011.
- [25] A. C. Dolphin, "A short history of voltage-gated calcium channels," *British Journal of Pharmacology*, vol. 147, supplement 1, pp. S56–S62, 2006.
- [26] D. Magis, E. Boon, G. Coppola, A. Daron, and J. Schoenen, "A novel CACNA1A mutation results in episodic ataxia with migrainous features without headache," *Cephalalgia*, vol. 32, pp. 1147–1149, 2012.
- [27] H. Qi, M. M. Moran, B. Navarro et al., "All four CatSper ion channel proteins are required for male fertility and sperm cell hyperactivated motility," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 104, no. 4, pp. 1219–1223, 2007.
- [28] J. P. Changeux, "The nicotinic acetylcholine receptor: the founding father of the pentameric ligand-gated ion channel superfamily," *The Journal of Biological Chemistry*, vol. 287, pp. 40207–40215, 2012.
- [29] F. M. Benes, "Nicotinic receptors and functional regulation of GABA cell microcircuitry in bipolar disorder and schizophrenia," in *Handbook of Experimental Pharmacology*, pp. 401–417, Springer, Berlin, Germany, 2012.
- [30] C. Liu and D. Su, "Nicotinic acetylcholine receptor $\alpha 7$ subunit: a novel therapeutic target for cardiovascular diseases," *Frontiers of Medicine*, vol. 6, pp. 35–40, 2012.
- [31] H. M. Schuller, "Regulatory role of the $\alpha 7$ nAChR in cancer," *Current Drug Targets*, vol. 13, pp. 680–687, 2012.
- [32] A. Cukkemane, R. Seifert, and U. B. Kaupp, "Cooperative and uncooperative cyclic-nucleotide-gated ion channels," *Trends in Biochemical Sciences*, vol. 36, no. 1, pp. 55–64, 2011.
- [33] U. B. Kaupp and R. Seifert, "Cyclic nucleotide-gated ion channels," *Physiological Reviews*, vol. 82, no. 3, pp. 769–824, 2002.
- [34] M. Biel, X. Zong, M. Distler et al., "Another member of the cyclic nucleotide-gated channel family, expressed in testis, kidney, and heart," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 91, no. 9, pp. 3505–3509, 1994.
- [35] K. Togashi, M. J. von Schimmelmann, M. Nishiyama et al., "Cyclic GMP-gated CNG channels function in Sema3A-induced growth cone repulsion," *Neuron*, vol. 58, no. 5, pp. 694–707, 2008.
- [36] Q. Zhang, F. Zulfiqar, S. A. Riazuddin et al., "Autosomal recessive retinitis pigmentosa in a Pakistani family mapped to CNGA1 with identification of a novel mutation," *Molecular Vision*, vol. 10, pp. 884–889, 2004.
- [37] H. Betz and B. Laube, "Glycine receptors: recent insights into their structural organization and functional diversity," *Journal of Neurochemistry*, vol. 97, no. 6, pp. 1600–1610, 2006.
- [38] J. W. Lynch, "Molecular structure and function of the glycine receptor chloride channel," *Physiological Reviews*, vol. 84, no. 4, pp. 1051–1095, 2004.
- [39] M. Biel and S. Michalakakis, "Function and dysfunction of CNG channels: insights from channelopathies and mouse models," *Molecular Neurobiology*, vol. 35, no. 3, pp. 266–277, 2007.
- [40] S. Herrmann, B. Layh, and A. Ludwig, "Novel insights into the distribution of cardiac HCN channels: an expression study in the mouse heart," *Journal of Molecular and Cellular Cardiology*, vol. 51, pp. 997–1006, 2011.
- [41] J. Klueva, A. D. Lima, S. Meis, T. Voigt, and T. Munsch, "Hyperpolarization-activated cation current contributes to spontaneous network activity in developing neocortical cultures," *Neurosignals*, vol. 20, pp. 35–47, 2012.
- [42] B. H. Meurers, G. Dziewczapolski, A. Bittner, T. Shi, F. Kamme, and C. W. Shults, "Dopamine depletion induced up-regulation of HCN3 enhances rebound excitability of basal ganglia output neurons," *Neurobiology of Disease*, vol. 34, no. 1, pp. 178–188, 2009.
- [43] G. A. Gutman, K. G. Chandy, S. Grissmer et al., "International Union of Pharmacology. LIII. Nomenclature and molecular relationships of voltage-gated potassium channels," *Pharmacological Reviews*, vol. 57, no. 4, pp. 473–508, 2005.
- [44] F. C. Richardson and L. K. Kaczmarek, "Modification of delayed rectifier potassium currents by the Kv9.1 potassium channel subunit," *Hearing Research*, vol. 147, no. 1–2, pp. 21–30, 2000.
- [45] M. Costigan, I. Belfer, R. S. Griffin et al., "Multiple chronic pain states are associated with a common amino acid-changing allele in KCNS1," *Brain*, vol. 133, no. 9, pp. 2519–2527, 2010.

- [46] K. Hao, T. Niu, X. Xu, Z. Fang, and X. Xu, "Single-nucleotide polymorphisms of the KCNS3 gene are significantly associated with airway hyperresponsiveness," *Human Genetics*, vol. 116, no. 5, pp. 378–383, 2005.
- [47] H. Hibino, A. Inanobe, K. Furutani, S. Murakami, I. Findlay, and Y. Kurachi, "Inwardly rectifying potassium channels: their structure, function, and physiological roles," *Physiological Reviews*, vol. 90, no. 1, pp. 291–366, 2010.
- [48] N. Patil, D. R. Cox, D. Bhat, M. Faham, R. M. Myers, and A. S. Peterson, "A potassium channel mutation in weaver mice implicates membrane excitability in granule cell differentiation," *Nature Genetics*, vol. 11, no. 2, pp. 126–129, 1995.
- [49] J. Ferrer, C. G. Nichols, E. N. Makhina et al., "Pancreatic islet cells express a family of inwardly rectifying K⁺ channel subunits which interact to form G-protein-activated channels," *The Journal of Biological Chemistry*, vol. 270, no. 44, pp. 26086–26091, 1995.
- [50] S. Signorini, Y. J. Liao, S. A. Duncan, L. Y. Jan, and M. Stoffel, "Normal cerebellar development but susceptibility to seizures in mice lacking G protein-coupled, inwardly rectifying K⁺ channel GIRK2," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 94, no. 3, pp. 923–927, 1997.
- [51] J. B. Gourraud, S. le Scouarnec, F. Sacher et al., "Identification of large families in early repolarization syndrome," *Journal of the American College of Cardiology*, vol. 61, pp. 164–172, 2013.
- [52] L. Gabriel, A. Lvov, D. Orthodoxou et al., "The acid-sensitive, anesthetic-activated potassium leak channel, KCNK3, is regulated by 14-3-3beta-dependent, protein kinase C, (PKC)-mediated endocytic trafficking," *The Journal of Biological Chemistry*, vol. 287, pp. 32354–32366, 2012.
- [53] D. H. Vandorpe and C. E. Morris, "Stretch activation of the *Aplysia* S-channel," *Journal of Membrane Biology*, vol. 127, no. 3, pp. 205–214, 1992.
- [54] F. Maingret, M. Fosset, F. Lesage, M. Lazdunski, and E. Honoré, "TRAAK is a mammalian neuronal mechano-gated K⁺ channel," *The Journal of Biological Chemistry*, vol. 274, no. 3, pp. 1381–1387, 1999.
- [55] A. Kleger and S. Liebau, "Calcium-activated potassium channels, cardiogenesis of pluripotent stem cells, and enrichment of pacemaker-like cells," *Trends in Cardiovascular Medicine*, vol. 21, pp. 74–83, 2011.
- [56] W. L. Imlach, S. C. Finch, J. H. Miller, A. L. Meredith, and J. E. Dalziel, "A role for BK channels in heart rate regulation in rodents," *PLoS ONE*, vol. 5, no. 1, Article ID e8698, 2010.
- [57] W. Guo, X. Xu, X. Gao, G. Burnstock, C. He, and Z. Xiang, "Expression of P2X₅ receptors in the mouse CNS," *Neuroscience*, vol. 156, no. 3, pp. 673–692, 2008.
- [58] M. Ryten, S. Y. Yang, P. M. Dunn, G. Goldspink, and G. Burnstock, "Purinoreceptor expression in regenerating skeletal muscle in the mdx mouse model of muscular dystrophy and in satellite cell cultures," *The FASEB Journal*, vol. 18, no. 12, pp. 1404–1406, 2004.
- [59] M. Ryten, P. M. Dunn, J. T. Neary, and G. Burnstock, "ATP regulates the differentiation of mammalian skeletal muscle by activation of a P2X₅ receptor on satellite cells," *Journal of Cell Biology*, vol. 158, no. 2, pp. 345–355, 2002.
- [60] D. Sonin, S. Y. Zhou, C. Cronin et al., "Role of P2X purinergic receptors in the rescue of ischemic heart failure," *American Journal of Physiology*, vol. 295, no. 3, pp. H1191–H1197, 2008.
- [61] S. Feske, E. Y. Skolnik, and M. Prakriya, "Ion channels and transporters in lymphocyte function and immunity," *Nature Reviews Immunology*, vol. 12, pp. 532–547, 2012.
- [62] S. H. Jung and J. H. Ho, "ATP stimulates mouse embryonic stem cell proliferation via protein kinase C, phosphatidylinositol 3-kinase/Akt, and mitogen-activated protein kinase signaling pathways," *Stem Cells*, vol. 24, no. 12, pp. 2637–2648, 2006.
- [63] H. Watanabe, K. Iino, T. Ohba, and H. Ito, "Possible involvement of TRP channels in cardiac hypertrophy and arrhythmia," *Current Topics in Medicinal Chemistry*. In press.
- [64] Y. Takada, T. Numata, and Y. Mori, "Targeting TRPs in neurodegenerative disorders," *Current Topics in Medicinal Chemistry*. In press.
- [65] H. Cheng, J. M. Feng, M. L. Figueiredo et al., "Transient receptor potential melastatin type 7 channel is critical for the survival of bone marrow derived mesenchymal stem cells," *Stem Cells and Development*, vol. 19, no. 9, pp. 1393–1403, 2010.
- [66] E. den Dekker, D. G. M. Molin, G. Breikers et al., "Expression of transient receptor potential mRNA isoforms and Ca²⁺ influx in differentiating human stem cells and platelets," *Biochimica et Biophysica Acta*, vol. 1539, no. 3, pp. 243–255, 2001.
- [67] J. P. Weick, M. A. Johnson, and S. C. Zhang, "Developmental regulation of human embryonic stem cell-derived neurons by calcium entry via transient receptor potential channels," *Stem Cells*, vol. 27, no. 12, pp. 2906–2916, 2009.
- [68] J. Sabourin, F. Antigny, E. Robin, M. Frieden, and E. Raddatz, "Activation of transient receptor potential canonical 3 (TRPC3)-mediated Ca²⁺ entry by A₁ adenosine receptor in cardiomyocytes disturbs atrioventricular conduction," *The Journal of Biological Chemistry*, vol. 287, pp. 26688–26701, 2012.
- [69] A. Haythornthwaite, S. Stoelzle, A. Hasler et al., "Characterizing human ion channels in induced pluripotent stem cell-derived neurons," *Journal of Biomolecular Screening*, vol. 17, pp. 1264–1272, 2012.
- [70] S. S. Chopra, D. M. Stroud, H. Watanabe et al., "Voltage-gated sodium channels are required for heart development in zebrafish," *Circulation Research*, vol. 106, no. 8, pp. 1342–1350, 2010.
- [71] G. R. Li, H. Sun, X. Deng, and C. P. Lau, "Characterization of ionic currents in human mesenchymal stem cells from bone marrow," *Stem Cells*, vol. 23, no. 3, pp. 371–382, 2005.
- [72] P. Jiang, S. N. Rushing, C. W. Kong et al., "Electrophysiological properties of human induced pluripotent stem cells," *American Journal of Physiology*, vol. 298, no. 3, pp. C486–C495, 2010.
- [73] C. D. House, C. J. Vaske, A. M. Schwartz et al., "Voltage-gated Na⁺ channel SCN5A is a key regulator of a gene transcriptional network that controls colon cancer invasion," *Cancer Research*, vol. 70, no. 17, pp. 6957–6967, 2010.
- [74] M. H. Meisler, J. E. O'Brien, and L. M. Sharkey, "Sodium channel gene family: epilepsy mutations, gene interactions and modifier effects," *The Journal of Physiology*, vol. 588, no. 11, pp. 1841–1848, 2010.
- [75] D. E. Arking and N. Sotoodehnia, "The genetics of sudden cardiac death," *Annual Review of Genomics and Human Genetics*, vol. 13, pp. 223–239, 2012.
- [76] R. P. Davis, S. Casini, C. W. van den Berg et al., "Cardiomyocytes derived from pluripotent stem cells recapitulate electrophysiological characteristics of an overlap syndrome of cardiac sodium channel disease," *Circulation*, vol. 125, pp. 3079–3091, 2012.
- [77] C. J. Giuliano, J. S. Kerley-Hamilton, T. Bee et al., "Retinoic acid represses a cassette of candidate pluripotency chromosome 12p genes during induced loss of human embryonal carcinoma tumorigenicity," *Biochimica et Biophysica Acta*, vol. 1731, no. 1, pp. 48–56, 2005.

- [78] V. Rybalchenko, M. Ahuja, J. Coblenz et al., "Membrane potential regulates nicotinic acid adenine dinucleotide phosphate (NAADP) dependence of the pH- and Ca^{2+} -sensitive organellar two-pore channel TPC1," *The Journal of Biological Chemistry*, vol. 287, pp. 20407–20416, 2012.
- [79] P. A. Davies, W. Wang, T. G. Hales, and E. F. Kirkness, "A novel class of ligand-gated ion channel is activated by Zn^{2+} ," *The Journal of Biological Chemistry*, vol. 278, no. 2, pp. 712–717, 2003.
- [80] J. A. Kaye and S. Finkbeiner, "Modeling Huntington's disease with induced pluripotent stem cells," *Molecular and Cellular Neurosciences*, 2013.
- [81] S. Nedelec, B. Onteniente, M. Peschanski, and C. Martinat, "Genetically-modified stem cells for the treatment of neurological diseases," *Current Gene Therapy*, vol. 4, pp. 1170–1181, 2013.
- [82] H. Okano, M. Nakamura, K. Yoshida et al., "Steps toward safe cell therapy using induced pluripotent stem cells," *Circulation Research*, vol. 112, pp. 523–533, 2013.