Review Article

Stress Response and Pathogenicity of the Necrotrophic Fungal Pathogen *Alternaria alternata*

Kuang-Ren Chung^{1,2}

¹ Citrus Research and Education Center, Institute of Food and Agricultural Sciences (IFAS), University of Florida, 700 Experiment Station Road, Lake Alfred, FL 33850, USA

² Department of Plant Pathology, IFAS, University of Florida, Gainesville, FL 32611, USA

Correspondence should be addressed to Kuang-Ren Chung; krchung@ufl.edu

Received 12 September 2012; Accepted 3 October 2012

Academic Editors: J. R. Blazquez and M. Shoda

Copyright © 2012 Kuang-Ren Chung. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The production of host-selective toxins by the necrotrophic fungus *Alternaria alternata* is essential for the pathogenesis. *A. alternata* infection in citrus leaves induces rapid lipid peroxidation, accumulation of hydrogen peroxide (H_2O_2), and cell death. The mechanisms by which *A. alternata* avoids killing by reactive oxygen species (ROS) after invasion have begun to be elucidated. The ability to coordinate of signaling pathways is essential for the detoxification of cellular stresses induced by ROS and for pathogenicity in *A. alternata*. A low level of H_2O_2 , produced by the NADPH oxidase (NOX) complex, modulates ROS resistance and triggers conidiation partially via regulating the redox-responsive regulators (YAP1 and SKN7) and the mitogen-activated protein (MAP) kinase (HOG1) mediated pathways, which subsequently regulate the genes required for the biosynthesis of siderophore, an iron-chelating compound. Siderophore-mediated iron acquisition plays a key role in ROS detoxification because of the requirement of iron for the activities of antioxidants (e.g., catalase and SOD). Fungal strains impaired for the ROS-detoxifying system severely reduce the virulence on susceptible citrus cultivars. This paper summarizes the current state of knowledge of signaling pathways associated with cellular responses to multidrugs, oxidative and osmotic stress, and fungicides, as well as the pathogenicity/virulence in the tangerine pathotype of *A. alternata*.

1. Introduction

Alternaria species have different lifestyles, ranging from saprophytes to endophytes to pathogens [1]. *Alternaria* species are a highly successful group of fungal pathogens that cause diseases in a wide variety of economically important crops, including apple, broccoli, cauliflower, carrot, citrus, pear, rice, strawberry, tomato, potato, and tobacco, as well as many ornamental and weed species. Due to their wide host range and worldwide distribution, *Alternaria* species cause severe economic problems. *Alternaria* species have been reported to cause diseases in nearly 400 plant species; *A. alternata* alone can infect more than 100 plant species [2–4]. One reason for the success of these pathogens may be attributed to their production of diverse phytotoxins [5, 6]. The host-selective toxins (HSTs) produced by many members of the genus *Alternaria* have unique modes of action and

toxicity to their respective host plants. Production of HST is critical for successful pathogenesis because HST-deficient mutants are incapable of attacking their host plants [7–11]. In addition to HSTs, many *Alternaria* species produce nonhost selective phytotoxins, such as brefeldin A, altertoxin, and tentoxin [1]. Others can produce mycotoxins that are harmful to humans and other animals [12]. Several *Alternaria* species can also cause upper respiratory tract infections and asthma in humans [13].

Alternaria alternata (Fr.) Keissler has several pathogenic variants, each producing a unique HST and causing disease in different host plants [5, 9, 10, 14, 15]. HSTs produced by *Alternaria* pathotypes are chemically diverse, ranging from low-molecular-weight compounds to cyclic peptides. The genes encoding polypeptides for biosynthesis of *Alternaria* HSTs have been shown to reside on a dispensable chromosome [9]. In citrus, *A. alternata* has two major

FIGURE 1: Life cycle of *Alternaria alternata*, the causal agent of citrus brown spot. ACT toxin produced by the tangerine pathotype of *A. alternata* is transported via the vascular system and formation of necrotic lesions on a detached calamondin leaf (bottom right).

pathotypes—the tangerine pathotype and the rough lemon type [16]. The citrus pathotypes are morphologically similar and can be differentiated only by pathological and genetic analyses [17]. The rough lemon pathotype, producing the host-selective ACRL toxin, is pathogenic exclusively to lemon (*Citrus jambhiri* Lush) and Rangpur lime (*Citrus x limonia* Osbeck). ACRL toxin affects mitochondrial function, disrupting posttranscriptional RNA splicing and causing metabolite leakage and malfunction of oxidative phosphorylation in susceptible host cells [18, 19]. In contrast, the tangerine pathotype of *A. alternata* produces the host-selective ACT toxin with a core 9,10-epoxy-8-hydroxy-9-methyl-decatienoic acid structure [20] and causes brown spots on citrus leaves and fruit. ACT toxin is highly toxic to tangerines (*C. reticulata* Blanco) and grapefruit (*C. paradisi* Macfad.), as well as hybrids from grapefruit and tangerine, or tangerine and sweet orange (*C. sinensis* Osbeck). ACT toxin does not affect rough lemon or Rangpur lime [20]. The toxin is quickly translocated outward through the vascular system, causing rapid electrolyte leakage and necrotic lesions along the veins (Figure 1). *A. alternata* infection in citrus leaves induces rapid lipid peroxidation and accumulation of hydrogen peroxide (H_2O_2) [21]. Studies show that *A. alternata* has evolved a dramatic flexibility and uniqueness in the signaling pathways in order to respond to diverse environmental stimuli and to thrive within host plants. This paper discusses signaling pathways related to oxidative and osmotic stress resistance, fungicide sensitivity, conidia formation, and pathogenesis of *A. alternata*.

2. Roles of Reactive Oxygen Species in Plant-Fungal Interactions

All organisms with an aerobic lifestyle inevitably generate toxic reactive oxygen species (ROS), primarily superoxide (O_2^-), and hydrogen peroxide (H_2O_2) during physiological metabolisms [22–26]. During the course of host colonization, fungal pathogens of plants need to overcome a wide range of potentially harmful environmental challenges, particularly an oxidative burst, which could result in the production and accumulation of highly toxic ROS. In addition to the direct toxicity of ROS to cells, when produced in abundance, ROS can also serve as secondary messengers in the pathogen-response signal transduction pathways [23, 27]. Among ROS, H_2O_2 is relatively stable and able to pass freely through membranes, serving as a signaling cue for defense responses in surrounding cells and as a substrate for oxidative cross-linking in the plant cell wall [27–32]. Hydrogen peroxide can react with O_2^- via the Haber-Weiss reaction or with metal ions via the Fenton pathway [33–35] to generate the extremely toxic hydroxyl radical. It has been well known that plants produce toxic ROS as a defense against pathogens [36–41]. In response to the microbe invasion, plant cells often produce excessive amounts of H_2O_2 by a specific plasma membrane NADPH oxidase, termed as the hypersensitive reaction (HR), which leads to programmed cell death and cellular defense against pathogen attack [42–46]. The HR plays a vital role in plant defenses against saprophytes and noncompatible or biotrophic pathogens; however, HR has been shown less

effective against necrotrophic phytopathogens [47–51]. ROS have been shown to be involved in nonhost resistance in plants as well [52, 53].

The burst of the HR may ironically provide an advantage for necrotrophic phytopathogens, as they acquire nutrients exclusively from dead cells [54]. ROS have been thought to enhance plant colonization by necrotrophic pathogens such as *Botrytis cinerea* and *Sclerotinia sclerotiorum* [50, 51, 55]. Because many necrotrophic pathogens are able to produce a wide array of HSTs or cell-wall-degrading enzymes that kill host cells before colonization, leading to the accumulation of ROS, the pathogens must have evolved effective mechanisms to cope with the toxicity of ROS [47, 49].

3. YAP1-Mediated ROS Detoxification in *A. alternata*

ROS damage a wide range of biological molecules, including fatty acids, proteins/enzymes, sugars, and nucleic acids; thus, exposure to ROS may result in cell death [56–58]. The relative sensitivity of the fungal pathogen to ROS is likely determined by the effectiveness of its own ROS detoxification ability. In order to survive under aerobic conditions, fungi must have detoxification systems that can effectively scavenge ROS, maintain reduced redox states within subcellular microenvironments, and repair ROS-triggered damage [22, 59, 60]. Molecular and genetic studies aimed at understanding the mechanisms by which cells cope with the oxidative stresses and are protected from the deleterious effects of ROS have been intensively studied in both prokaryotes and eukaryotes. In the budding yeast *Saccharomyces cerevisiae*, the YAP1 transcription regulator plays a central role in the cellular pathways associated with the oxidative stress response [61, 62]. YAP1 is responsible for transcriptional activation of genes involved in multidrug resistance as well. YAP1, resembling mammalian AP-1, has a basic leucine zipper (bZIP) domain and has been shown to be activated by H_2O_2 and various ROS-generating oxidants, as well as heavy metals [63–66]. In the absence of oxidative challenges, YAP1 can be found in the cytoplasm at low levels. Upon perceiving oxidative or chemical stimuli, YAP1 quickly forms disulfide bonds, changes conformation, and is translocated into the nucleus where YAP1 regulates the expression of genes responsible for stress alleviation [67–71]. Conserved cysteine residues in both the amino and carboxyl terminal domains are essential for the formation of the disulfide bonds, nuclear relocalization and transcriptional regulation of YAP1 [72].

Although all microorganisms employ complex mechanisms, both enzymatic and nonenzymatic to avoid ROS toxicity [47, 61, 73], the pathological roles of oxidative stress mitigation remain uncertain in pathogenic fungal species. The role of ROS in host resistance and pathogen invasion is likely dictated by the physiological conditions of the host, the lifestyle of the pathogen, and the combination of different stimuli [38, 41, 74–76]. Hence, ROS produced by plants may have different effects against different pathogens [49, 75]. YAP1-mediated detoxification of ROS is an essential virulence determinant in the opportunistic human pathogen

Candida albicans and the biotrophic maize pathogen *Ustilago maydis* [77, 78]. However, YAP1 is not required for virulence in the plant pathogens *Cochliobolus heterostrophus* and *B. cinerea* and in the animal pathogen *Aspergillus fumigatus*, even though the disrupted mutants exhibit increased sensitivity to H_2O_2 [79, 80]. The necrotrophic fungus *Sc. sclerotiorum* produces oxalic acid that suppresses host-generated ROS, and thus facilitates fungal evasion [73]. The *Magnaporthe oryzae* *MoHYR1* gene encoding a glutathione peroxidase (GSHPx) is required for detoxifying plant-generated ROS and full virulence [81]. In *S. cerevisiae*, HYR1 forms a disulfide bond with YAP1, inducing a conformational alteration and nuclear localization of YAP1 upon exposure to ROS [82].

To thrive within host plants, *A. alternata* must be able to detoxify or obviate the ROS-mediated plant defense barriers. Our studies have demonstrated that cellular detoxification of ROS regulated by the redox-responsive YAP1 transcription regulator is important for pathogenesis of *A. alternata* to citrus [83, 84]. Inactivation of the *A. alternata* *API* gene (designated *AaAPI*), encoding a YAP1-like transcription factor, resulted in fungal mutants that are hypersensitive to H_2O_2 , menadione, and potassium superoxide (KO_2). The promoter of *AaAPI* contains a putative stress responsive element (STRE: AGAGGGG). Upon activation by H_2O_2 , the *AaAPI::sGFP* fusion protein became localized in the nucleus. Fungal mutants lacking *AaAPI* ($\Delta yap1$) are weakly virulent on susceptible citrus cultivars even though they synthesize HST toxins normally. However, $\Delta yap1$ mutant is not sensitive to osmotic and salt stress-related compounds (e.g., sorbitol, mannitol, NaCl, and KCl) (Figure 2). $\Delta yap1$ mutant produces wild-type level of conidia that germinate at a rate and magnitude similar to the wild-type strain. The nonpathogenic phenotype of $\Delta yap1$ mutant was accompanied with reduced activities of fungal antioxidants, including catalase, peroxidase, superoxide dismutase (SOD), and glutathione reductase. The inability of $\Delta yap1$ mutants to incite necrotic lesions is likely due to the mutants' inability to detoxify ROS because coapplication of $\Delta yap1$ mutants with the NADPH oxidase inhibitor, apocynin, or diphenylene iodonium, partially restored lesion-forming capability to the mutants. $\Delta yap1$ mutant is impaired in the penetration and colonization stages because the impaired mutant did not cause any visible necrotic lesions on wounded or unwounded leaves of the citrus cultivar Minneola. All mutant phenotypes were completely restored to the wild type in fungal strains expressing a functional copy of *AaAPI*. $\Delta yap1$ mutant displayed severe defects in antioxidant activities and was unable to detoxify H_2O_2 effectively. Our studies concluded that effective detoxification of ROS via the *AaAPI*-mediated pathway is absolutely required for successful colonization of citrus by *A. alternata* [83, 84]. The tobacco pathotype of *A. alternata* impaired for the biosynthesis of mannitol, an antioxidant and quencher of the hydroxyl radical, also greatly reduces virulence [85, 86], consistent with the importance of ROS detoxification in the pathogenesis of *A. alternata*.

Furthermore, *AaAPI* was found to be required for full resistance to 2,3,5-triodobenzoic acid (TIBA), 2-chloro-5-hydroxypyridine (CHP), diethyl maleate (DEM), and many pyridine-containing compounds [87]. Diethyl maleate is

a glutathione-depleting agent that has been shown to generate a nonreversible modification of cysteine residues in the *Schizosaccharomyces pombe* Pap1 protein. As a result, Pap1 is constitutively localized in nucleus and activates the genes required for ROS tolerance [88]. TIBA is often used as herbicides or as an inhibitor of indole-3-acetic acid (IAA) transportation [89, 90]. Pyridine is a heteroaromatic compound composed of five carbons and one nitrogen atom. Pyridine could accelerate the production of superoxide and hydroxyl radicals when Cu^{2+} and H_2O_2 are present [91, 92]. Pyridine and its derivatives serve as constituents of RNA and DNA, as electron carriers such as NADP/NADPH and flavin nucleotides (FAD/FADH) and as energy storage molecules such as ATP and GTP.

4. The “Two-Component” Histidine Kinase (HSK) Signaling Pathway

All living cells have a complicated yet well-regulated network often comprising different signaling transduction pathways to perceive changes in their environments and to adjust physiological and developmental processes [93–99]. “Two-component” histidine kinase (HSK) signaling transfer systems are commonly present in bacteria, slime molds, fungi, and plants; however, these systems have not yet been identified in animals [100, 101]. In bacteria, HSK signaling systems contain a histidine kinase (HSK) and a response regulator (RR); each is encoded by a separate gene [102, 103]. In contrast, all fungal HSKs have both the HSK and RR domains [101, 104, 105]. In response to environmental changes, a series of phosphate transfers between histidine (His) and aspartate (Asp) residues occurs in a pattern of His-Asp-His-Asp to regulate downstream signaling pathways such as mitogen-activated protein kinase (MAPK) cascades and eventually leads to a change in gene expression [100, 106, 107].

The *S. cerevisiae* histidine kinase, designated SLN1p, is required for osmotic adaption via the SLN1p-YPD1p (a protein containing a His phosphotransfer domain)-SSK1p or SKN7p cascade [108, 109]. SSK1p is the major regulator for osmolarity response; SKN7p plays only a minor role in osmosensing. Under normal osmolarity, the SLN1p kinase is phosphorylated and able to activate YPD1p and SSK1p with a phosphorelay mechanism (Figure 3). The phosphorylated SSK1p is inactive and incapable of activating the High Osmolarity-Glycerol 1 (HOG1) MAP kinase pathway (see below for details). In contrast, SLN1p is not phosphorylated under conditions of high osmolarity; therefore SSK1p is able to activate the HOG1-signaling cascade. The activated HOG1 pathway is responsible for glycerol accumulation, allowing the yeast to cope with the high osmolarity. *S. cerevisiae* also utilizes a non-HSK-related protein SHO1p to cope with osmotic stress [108, 110]. However, deletion of an *SHO1* homolog in *A. alternata* did not impact cellular tolerance to oxidative and osmotic stress, fungicide sensitivity or fungal virulence (L.-H. Chen, unpublished).

The budding yeast *S. cerevisiae* has only one HSK; all other fungi have multiple HSK signaling genes [111]. Fungal HSKs

are divided into 11 groups based on phylogenetic relationships inferred from the conserved HSK and RR domains. Among them, Group III HSK is one of the best characterized HSKs in the filamentous fungi. Collectively, Group III HSK has been implicated in osmotic and oxidative responses, toxin biosynthesis, hyphal development, conidia formation, and virulence, as well as sensitivity to dicarboximide and phenylpyrrole fungicides in different fungal species [112–121].

Signals sensed by HSK are often transduced down to the HOG1 MAP kinase pathway. Fungi lacking Group III HSK or HOG1 often became resistant to dicarboximide and phenylpyrrole fungicides and exhibited an elevated sensitivity to osmotic stress [117, 122, 123]. Although the HSK-HOG1 signaling pathway is conserved, it may be recruited for divergent functions in different fungal species. As discussed above, the *S. cerevisiae* SLN1p negatively regulates HOG1 phosphorylation under conditions of high osmolarity [100, 106, 107]. The filamentous fungus *Co. heterostrophus* Group III HSK (Dic-1) positively regulates phosphorylation of the HOG1 MAP kinase, which subsequently activates expression of genes responsible for osmotic resistance and fungicide sensitivity [117]. In *B. cinerea*, the HOG1-like MAP kinase is not required for fungicide sensitivity even though it is negatively regulated by the “two-component” HSK. Furthermore, the salt-tolerant yeast species, *Hortaea werneckii*, copes with osmotic stress using a Group VII HSK-HOG1 pathway [124]. Those studies indicate that the HSK-HOG1 signaling pathways can be operated in very different regulatory mechanisms in various species.

The *AaHSK1* gene, encoding a putative histidine kinase, was cloned from the tangerine pathotype of *A. alternata* [87]. *AaHSK1*, containing no transmembrane regions, is required for adaption to osmotic stress induced by sugars but not by salts (Figure 2). $\Delta hsk1$ displayed increased sensitivity to glucose, sucrose, sorbitol, or mannitol, but not to H_2O_2 , KCl, or NaCl [125]. Similarly, the *M. grisea* HSK is required for resistance to sugar, but not salt, osmotic stress [116]. The *F. oxysporum* histidine kinase Fhk1 is responsible for resistance to osmotic stress, menadione, but not H_2O_2 [126]. Similar to $\Delta yap1$ mutant, *AaHSK1* disruption mutants displayed an elevated sensitivity to TIBA and CHP, suggesting a possible link between YAP1 and HSK. $\Delta hsk1$ mutants displayed an elevated resistance to dicarboximide (iprodione and vinclozolin) and phenylpyrrole (fludioxonil) fungicides, suggesting that *AaHSK1* is one of the primary targets of these fungicides. Similarly, resistance to dicarboximide and phenylpyrrole fungicides has been demonstrated to be associated with a mutation within the gene encoding a Group III HSK and/or an HOG1 MAP kinase in a number of filamentous fungi [113–118, 121, 122, 126, 127]. However, HSK is not involved in dicarboximide susceptibility in *A. longipes*.

The *A. alternata* HSK1 is not required for response to oxidative stress. The *AaHSK1* gene product is not involved in pathogenicity or virulence because the *AaHSK1*-impaired mutants ($\Delta hsk1$) induced necrotic lesions at rates and magnitudes similar to the wild-type strain or the genetically reverted strain on wounded or unwounded leaves of citrus. In contrast, Group III HSK is a virulence determinant in

FIGURE 2: Phenotypic changes in *A. alternata* mutants lacking the transcription regulator (*YAP1*), the MAP kinase (*HOG1*), the “two-component” histidine kinase (*HSK1*), the response regulator (*SKN7*), or the NADPH oxidase (*NOXA*). Open rectangles denote wild-type phenotypes.

FIGURE 3: Schematic illustration and comparison of signaling pathways leading to osmotic stress resistance in the budding yeast, *S. cerevisiae*, and the pathways leading to ROS resistance, osmotic stress response, fungicide sensitivity, and conidia formation in the tangerine pathotype of *A. alternata*.

the phytopathogenic fungi *B. cinerea*, *Claviceps purpurea*, and *Fusarium oxysporum* and in the human pathogen *Cryptococcus neoformans* [118, 126, 128, 129].

5. The HOG1 Mitogen-Activated Protein kinase (MAPK-) Mediated Signaling Pathway

The HOG1 MAPK-mediated signaling cascades in eukaryotic cells are vital for sensing environmental stimuli and for transmitting these signals to the nucleus to modulate gene expression [130, 131]. MAPK-mediated cascade pathways are composed of three serine/threonine protein kinases—MAP kinase kinase kinase (MAPKKK), MAP kinase kinase (MAPKK), and MAP kinase (MAPK). This signal transduction pathway, in conjunction with HSK, is well conserved in all eukaryotes and functions in perceiving environmental stimuli via phosphorylation and gene activation [132, 133]. The phosphorylated MAPK activates a set of genes via regulating appropriate transcription factors.

The *A. alternata* ortholog (AaHOG1) contains a distinct phosphorylation motif (TGY) involved in the osmotic stress response [134]. Inactivation of the HOG1 ortholog by targeted gene disruption in the tangerine pathotype of *A. alternata* resulted in mutants that are highly sensitive to the oxidants *tert*-butyl-hydroxyperoxide, H_2O_2 , and menadione, salts (Figure 2), as well as TIBA and CHP [87]. Because fungal strain lacking the FUS3 MAP kinase ($\Delta fus3$) grew faster than wild type in the presence of KCl or NaCl [125]. Thus, AaHOG1 and FUS3 play an opposite role in KCl and NaCl tolerance. HOG1 has been shown to suppress the FUS3/KSS1 signaling cascade during hyperosmotic stress in *S. cerevisiae* [135, 136]. *A. alternata* strains impaired at AaHOG1 ($\Delta hog1$) displayed wild-type levels of sensitivity to high concentrations of glucose, sucrose, sorbitol, or mannitol (Figure 2) even though sugar osmoticants increased AaHOG1 phosphorylation and subsequently nuclear localization in the $\Delta hsk1$ mutant background. In the wild-type background, sugar osmoticant had less effect on AaHOG1 phosphorylation and did not facilitate nuclear localization of AaHOG1.

The wild-type isolate of *A. alternata* is extremely sensitive to dicarboximide and phenylpyrrole fungicides, whereas fungal strain lacking AaHSK1 is highly resistant to them. Compared to the resistance seen in mutants defective at AaHSK1, $\Delta hog1$ mutant displayed only slightly increased resistance to these fungicides. In the wild-type strain of *A. alternata*, the AaHOG1 protein was phosphorylated at low levels under normal conditions. Exposure to iprodione or fludioxonil fungicide, NaCl, or H_2O_2 elevated AaHOG1 phosphorylation to varying degrees. Although impairment of AaHSK1 reduces AaHOG1 phosphorylation, *A. alternata* apparently recruits AaHSK1 and AaHOG1 to exert a unique function in resistance to sugar osmoticants and salt stress, respectively (Figure 3).

Under unchallenged conditions, expression of AaHOG1::sGFP fusion protein under control of the endogenous AaHOG1 promoter in the wild-type strain resulted in green fluorescence uniformly diffused along the hyphal cytoplasm. However, the green fluorescence became dense

patches after exposure to H_2O_2 , iprodione and fludioxonil fungicides, or NaCl. Thus, nuclear localization is important for proper functions of HOG1. Compared to $\Delta hsk1$ or $\Delta yap1$ mutant, $\Delta hog1$ mutant was highly resistant to cell wall-degrading enzymes (lyticase, driselase, β -D-glucanase, and β -glucuronidase), thereby failing to generate any protoplasts. As judged from distinct phenotypes in fungal mutants impaired in the AaHSK1 or AaHOG1 gene in *A. alternata*, it appears that AaHSK1 functions in osmotic tolerance and fungicide sensitivity via AaHOG1 and other gene (e.g., SKN7) activation branches.

AaHSK1 plays no role in pathogenesis of *A. alternata*. On the other hand, pathogenicity assays revealed that the AaHOG1-impaired mutants are nonpathogenic, producing no necrotic lesions on *Minneola* leaves that were unwounded or prewounded before inoculation. Similar to AaAPI disruption, the AaHOG1-impaired mutant is defective at the penetration and colonization steps. Inactivation of the AaHOG1 gene did not impact the production of host-selective ACT toxin by *A. alternata*. HOG1 is required for virulence/pathogenicity in various fungal pathogens. These include *Co. heterostrophus*, *Cryphonectria parasitica*, *B. cinerea*, *Mycosphaerella graminicola*, *Ca. albicans*, and *C. neoformans* [137–142]. However, HOG1 is not a virulence determinant in *M. grisea*, *Colletotrichum lagenarium*, *Bipolaris oryzae* and *As. fumigatus* [122, 143–145]. Again, a conserved protein may have very different functions in fungi.

6. SKN7-Mediated ROS Detoxification

“Two-component” HSK-mediated signal transduction is vital for sensing and adapting to environmental changes in microorganisms. In *S. cerevisiae*, SLN1 histidine kinase transmits signals via a phosphotransfer process down to two response regulators, SSK1p and SKN7p, in response to osmotic stress. However, SKN7p is not regulated by the SLN1p-mediated phosphorylation in the oxidative stress response [146], indicating that there are two different activation mechanisms in response to osmotic and oxidative stress. Under oxidative stress, SKN7p is phosphorylated at serine or threonine residue, forming a heterodimer with YAP1; together they transcriptionally activate the genes involved in the oxidative stress response [146–152]. YAP1 regulates cadmium resistance independent of SKN7p [153]. SKN7p can also form a heterodimer with the heat-shock transcription factor, the cell cycle transcription regulator, the calcium responsive activator, or the Rho1 GTPase [154–157].

In filamentous fungi, SKN7 is required for oxidative stress adaptation, hypoosmotic stress response, cell cycle, sexual mating, sporulation, cell wall biosynthesis, and fungicide sensitivity [120, 146, 155, 158–163]. The *A. alternata* SKN7 homolog (AaSKN7) was cloned and characterized in the tangerine pathotype [164]. The promoter of AaSKN7 contains a putative stress responsive element (STRE: AGAGGGG) that is often present in genes induced by various stresses such as oxidative damage in yeasts. AaSKN7 has a heat-shock transcription factor- (HSF-) type helix-turn-helix DNA-binding domain signature and a response regulatory (RR) domain.

Genetic mutation analysis revealed that *AaSKN7* is required for resistance to osmotic and oxidative stress and fungicide sensitivity, as well as conidiation and conidia morphology. *AaSKN7* is primarily localized in the nucleus, whereas *YAP1* and *HOG1* are quickly transported into the nucleus upon sensing oxidative stress. *AaSKN7* may interact directly with *AaAP1* in nucleus in response to oxidative stress as demonstrated in the budding yeast [151]. Both *A. alternata* *AP1* and *HOG1* are required for resistance to different types of ROS including hydrogen peroxide, superoxide, and singlet oxygen. On the other hand, *AaSKN7* is required for resistance to H_2O_2 , *tert*-butyl hydroperoxide, and cumyl peroxide, but not to superoxide-generating compounds—diamide, menadione and potassium superoxide (Figure 2). It appears that *AaSKN7* and *AaHOG1* contribute independently to oxidative stress in *A. alternata*. Because *AaHSK1* is not required for ROS resistance [87]; activation of *AaSKN7* and *AaHOG1* in response to ROS is likely mediated by unknown regulatory sensors other than *AaHSK1*. Furthermore, *A. alternata* *HSK1* apparently can recruit *SKN7* and *HOG1* to deal with sugar and salt osmoticants, respectively. Although *AaHSK1* and *AaSKN7* play no roles in resistance to salt-induced stress, *AaSKN7* is involved in resistance to sugar osmoticants likely via the *AaHSK1*—mediated signaling pathway (Figure 3). $\Delta skn7/\Delta hog1$ double mutants were hypersensitive to both salts and sugars, indicating that *A. alternata* is capable of sensing different environmental stimuli using distinct or shared signaling pathways. Furthermore, *AaSKN7*, independent of *AaHSK1*, is involved in conidia formation. Our studies also revealed that formation of conidia by *A. alternata* is closely regulated by the *FUS3* and *SLT2* MAP kinases-mediated signaling pathways, as well as by the G-protein and the NOX complex [125, 165–167]. However, recent studies revealed that cAMP-dependent protein kinase A (PKA) suppresses conidia formation by the tangerine pathotype of *Alternaria alternata* [168]. It remains uncertain if these signaling pathways actually interact during conidia formation.

A. alternata strains deleted for *HSK1* or *HOG1* showed an elevated resistance to dicarboximide and phenylpyrrole fungicides. $\Delta skn7$ mutant displayed an elevated resistance to those fungicides at levels between $\Delta hsk1$ and $\Delta hog1$ mutant strains, indicating that the involvement of *SKN7* in fungicide sensitivity is likely mediated by the *HSK1* signaling pathway. Fungal strain carrying *skn7/hog1* double mutations exhibited fungicide resistance, similar to the strain carrying a single *AaHSK1* gene mutation. The results indicated that the signals associated with fungicide sensitivity are transduced from *AaHSK1* simultaneously down to both *AaSKN7*- and *AaHOG*-mediated pathways. The *HSK*-*HOG* signaling pathways are associated with fungicide susceptibility in *A. brassicicola* and *N. crassa* [121]; yet *SKN7* is not involved in fungicide sensitivity in *N. crassa*. *HSK* governs *Ssk1p* (an upstream regulator of *HOG1*) and *Skn7p* for osmolarity adaption and fungicide sensitivity in the phytopathogenic fungus *Co. heterostrophus* and the human pathogen *C. neoformans* [128, 169, 170].

The roles of *SKN7* in pathogenicity/virulence vary among fungal pathogens. *A. alternata* *SKN7* is required for fungal colonization and lesion development in susceptible cultivars

of citrus. Similar to $\Delta yap1$ and $\Delta hog1$, mutational inactivation of *AaSKN7* in *A. alternata* resulted in reduced activities of catalase, SOD, and peroxidase, confirming further that the ability to detoxify host-generating H_2O_2 by *A. alternata* is crucial for successful pathogenesis in citrus. The *SKN7* homologs are required for virulence in the human pathogens *C. neoformans*, *Ca. albicans*, and *Ca. glabrata* [150, 171, 172]. Again, in contrast, *SKN7* is not a virulence determinant in the plant pathogens *Co. heterostrophus* and *M. oryzae* and in the human pathogen *As. fumigatus* [116, 162, 173].

7. The NADPH Oxidase (NOX): Mediated Signaling Pathway

The NADPH-dependent oxidase transfers electrons from NADPH to the oxygen molecule, leading to the production of a superoxide that is further metabolized to H_2O_2 by SOD [174, 175]. NOX complex is commonly found in animals, plants, and many multicellular microorganisms, but completely absent in prokaryotes [176]. Functionally, NOX complex plays a crucial role in cellular differentiation and defence response. In humans, the phagocytic NOX complex, involved in the production of superoxide and immunity, contains two major catalytic components $gp22^{phox}$ and $gp91^{phox}$ and multiple regulatory subunits *Rac* (a small GTPase), $p40^{phox}$, $p47^{phox}$, and $p67^{phox}$ [177]. NOX complex is also required for the regulation of hormone responses, cell proliferation, and apoptosis in animals ([178–180]. Activation of $gp91^{phox}$ is primarily regulated by $p67^{phox}$ and *Rac2* [181]. Plants also have oxidases analogous to $gp91^{phox}$, designated respiratory burst homologs (*Rboh*), which are required for physiological metabolisms and for ROS generation in response to pathogen invasion [175, 182, 183].

Many fungi have NADPH oxidase orthologs, NOXA, NOXB and NOXC that have been documented by genetic analysis to be required for developmental, physiological and pathological functions [25, 76, 184, 185]. Both NOXA, and NOXB are analogous with mammalian $gp91^{phox}$. Expression of the NOXA/NOXB coding genes is closely regulated by the regulatory subunit, NOXR ($p67^{phox}$ homolog), and the small GTPase (*Rac* homolog) [185]. Fungal NOXC contains a calcium-binding EF-hand motif and is analogous to the mammalian NOX5 and the plant *Rboh* enzymes.

The functions of NOX complex in the regulation of multicellular development and pathogenicity vary markedly among fungal species that possess it [186]. Both NOXA and NOXB are involved in the regulation of sclerotia formation in both *B. cinerea* and *Sc. sclerotiorum* [141, 187]. In *B. cinerea*, only NOXB is required for the formation of the penetration structure, even though both NOX isoforms have a role in pathogenicity. In *M. grisea*, NOXA, and NOXB play a role in pathogenesis because both isoforms are required for the formation of penetration peg under the appressorium [188]. While NOXA is required for the development of sexual fruiting body in fungi, only NOXB is required for ascospore germination in *N. crassa* and *Podospira anserina* [189–191]. NOXA, but not NOXB, is required for establishing

the mutualistic association between the fungal endophyte *Epichloë festucae* and perennial ryegrass [76]. When inoculated into its grass host, *E. festucae* strain lacking NOXA or NOXR becomes pathogenic, showing increased branching and causing severe stunting and premature senescence of the host [192, 193]. NOXA is coordinately regulated by the small GTPase Rac and NOXR as evidenced by yeast two-hybrid and pull-down analyses [194]. Furthermore, the yeast polarity protein orthologs, Bem1 and Cdc24, have recently been proven to be parts of fungal NADPH oxidase complex [195].

The tangerine pathotype of *A. alternata* has NOXA, NOXB, and NOXR homologs. *A. alternata* NOXA (AaNOXA) contains a NADPH-binding domain and six transmembrane domains and a ferredoxin synthase-type FAD-binding domain, commonly found in the NOXA-like family. Genetic analysis revealed that AaNOXA is responsible for producing superoxide and H₂O₂. $\Delta noxA$ mutants accumulated less ROS within hyphae than the wild type, as judged by nitroblue tetrazolium (NBT), 3,3'-diaminobenzidine (DAB), and dichlorodihydrofluorescein diacetate (H₂DCFDA) staining for the presence of superoxide and H₂O₂. Moreover, deletion of AaNOXA in *A. alternata* resulted in an elevated sensitivity to H₂O₂, superoxide-generating compounds (menadione and KO₂), diamide, SDS, CHP, TIBA, and potent singlet oxygen-generating compounds (hematoporphyrin and rose Bengal) (Figure 2). These deficiencies are similar to the phenotypes previously seen for $\Delta yap1$ or $\Delta hog1$ mutant. Expression of the AaAPI and AaHOG1 genes is likely regulated by AaNOXA, as deletion of AaNOXA decreased the accumulation of the AaAPI and AaHOG1 gene transcripts. Reintroducing and expressing a wild-type AaNOXA in a $\Delta noxA$ mutant restored ROS resistance and expression of both AaAPI and AaHOG1 genes. $\Delta noxA$ mutants also displayed increased sensitivity to NADPH oxidase inhibitors [diphenylene iodonium (DPI) and apocynin], NO[•]-generating compounds [sodium nitroprusside (SNP) and hydroxyl amine HCl (HAD)], NO[•] synthase substrate (L-arginine) and NO[•] synthase inhibitor [nitroarginine methyl ester (nitro-arg)]. Similar to $\Delta yap1$ and $\Delta hog1$, $\Delta noxA$ mutants, producing normal ACT toxin, induced significantly smaller and fewer necrotic lesions than the wild type on detached Minneola or calamondin leaves 3 days postinoculation, indicating that NOXA is an important virulence determinant in *A. alternata*.

NOXA, NOXB, and NOXR are core components of the NOX complex, responsible for the production of H₂O₂. All three NOX components are required for vegetative growth, conidiation, resistance to oxidative and nitrosative stress, and full virulence. However, each isoform may independently and cooperatively interact with other yet unidentified components under different environmental conditions and during different developmental stages because the degree of impairment varied considerably among individual Δnox mutants. $\Delta noxA$ mutant was more sensitive to H₂O₂, KO₂, and diamide than $\Delta noxB$ or $\Delta noxR$. In contrast, $\Delta noxB$ or $\Delta noxR$ mutation strain was more sensitive to cumyl H₂O₂ and SDS than $\Delta noxA$. The elevated sensitivity of $\Delta noxB$ and $\Delta noxR$ mutants to ROS was also accompanied by a reduced expression of two redox-responsive genes AaAPI

and AaHOG1. Although expression of AaAPI and HOG1 was upregulated by the NOX system; both AaAPI and HOG1 negatively regulate the expression of NoxB and NoxR. This transcriptional feedback loop might allow fungus to avoid excessive production of toxic ROS. In *A. nidulans*, NOXA is regulated by an HOG1 homolog [189]. Expression of the NOX complex coding genes has been shown to be regulated by FUS3/KSS1 and SLT2 MAP kinases in *N. crassa* and *B. cinerea* [141, 191]. Mammalian p47^{phox} and p67^{phox} are phosphorylated by the p38 HOG1 MAP kinase.

In fungi, expression of NOXA and NOXB is regulated by NOXR and Rac [141, 191, 194]. However, *A. alternata* NOXR negatively regulated the expression of NOXA and had no effects for the expression of NOXB. $\Delta noxB$ mutant was highly resistant to calcofluor white, Congo red and dicarboximide and phenylpyrrole fungicides compared to the wild type or the mutant strain lacking NOXA and NOXR. NOXB seemingly plays a negatively regulatory role in the biosynthesis of chitin because $\Delta noxB$, but not $\Delta noxA$ and $\Delta noxR$, had higher chitin content than the wild type (S.L. Yang, personal communication). As stated above, fungal strains disrupted at any of the AaAPI, AaHSK1, and AaHOG1 genes were all hypersensitive to CHP and TIBA. $\Delta noxA$, $\Delta noxB$, and $\Delta noxR$ mutants also displayed increased sensitivity to these two compounds, suggesting the existence of essential cross-talks between different signaling pathways in the context of multidrug resistance.

The NOX complex has been shown to be required for pathogenicity/virulence in a number of fungal species [141, 167, 188, 196]. Pathogenicity assays revealed that $\Delta noxB$ or $\Delta noxR$ mutant is unable to produce necrotic lesions on unwounded citrus leaves. Both mutants induced wild-type lesions on citrus leaves that were wounded before inoculation, indicating that Δnox mutants are primarily arrested in the penetration stage. $\Delta yap1$ and $\Delta hog1$ mutants are blocked in both penetration and colonization stages.

8. Nonribosomal Peptide Synthetase- (NPS)- Mediated ROS Detoxification

Deletion of an *A. alternata* gene (AaNPS6), encoding a polypeptide analogous to fungal nonribosomal peptide synthetases (NPSs) resulted in fungi that reduced accumulation of host-selective toxin and melanin and displayed increased sensitivity to H₂O₂, superoxide-generating compounds (KO₂ and menadione), and iron depletion (L.-H. Chen, personal communication). $\Delta nps6$ failed to produce siderophore, a low-molecular organic compound involved in acquiring iron from the environment [197, 198]. In nearly all living cells, iron required for numerous metabolic functions and electron transfer processes plays a vital role for cell proliferation and survival [199]. When starved for iron, microorganisms secrete siderophores to solubilize and extract iron. All fungal siderophores (rhodotorulic acid, fusarinines, coprogens, and ferrichromes) that have been identified contain hydroxamates and are synthesized from an unconventional amino acid, L-ornithine [197, 200, 201]. NPSs function to synthesize linear or cyclic peptides without the aid of ribosomes, adding

FIGURE 4: Schematic illustration of signaling pathways leading to ROS detoxification in the tangerine pathotype of *A. alternata*. H_2O_2 produced by the membrane-bound NADPH oxidase (NOX) complex plays a central role in the activation of genes responsible for ROS resistance. Upon exposure to ROS, YAP1 forms disulfide bonds between two conserved cysteine residues, undergoes conformation changes, and is transported into the nucleus where YAP1 regulates the expression of numerous genes associated with environmental stress. The YAP1 and SKN7 redox-responsive regulators, the HOG1 mitogen-activated protein (MAP) kinase, the NPS6-mediated siderophore biosynthesis, and the NOX complex are required for ROS detoxification. *NPS6* encoding a nonribosomal peptide synthetase is required for the biosynthesis of siderophores, which can extract environmental iron. Iron is stored in vacuoles. Siderophore-mediated iron acquisition plays a critical role if ROS resistance because iron is a major cofactor for the activities of catalase and SOD. HOG1, in cooperation with unknown regulators, is also required for salt resistance. The two-component histidine kinase (HSK1), likely interacting with SKN7, is primarily used for cellular resistance to sugars. Fungicide sensitivity involves HSK1, HOG1, and SKN7.

D- or L-amino acids, proteins, nonproteins, hydroxyl acids and ornithine into nonribosomal peptides. Many of these peptides have medicinal, pharmaceutical, or agricultural values [202]. AM-toxin produced by the apple pathotype of *A. alternata*, HC-toxin produced by race 1 of the maize pathogen, *Co. carbonum*, and enniatin produced by *Fusarium* spp. are all nonribosomal peptides [7, 203, 204].

Many *Alternaria* species produce and excrete dimethyl coprogen siderophores [205–208]. Coprogen contains a diketopiperazine ring (dimerium acid), in which two N^5 -acyl- N^5 -hydroxy-ornithine units are joined by a peptide bond [198]. The third acyl ornithine unit is linked to the ring via an ester bond. NPSs are involved in assembling three N^5 -acyl- N^5 -hydroxy-ornithine units, which are

the immediate precursors of hydroxamate siderophores. Fungi often have multiple *NPS* genes; each encodes a polypeptide with discrete domains—AMP-binding adenylation (A), thiolation (T) or peptidyl carrier protein (ACP), and condensation (C) domains that are organized as a module [209–212]. The adenylation domain is required for recognition and activation of amino acid substrate. The thiolation domain is involved in 4'-phosphopanthine binding. The condensation domain is involved in the formation of a peptide bond and elongation and release of the newly synthesized peptide. The number and order of modules in an *NPS* affect the length and structure of nonribosomal peptide.

$\Delta yap1$ and $\Delta nps6$ mutants of *A. alternata* displayed varying levels of hypersensitivity to H_2O_2 and superoxide-generating compounds. $\Delta nps6$ mutant is less sensitive to ROS than $\Delta yap1$, displaying an increased sensitivity to the test oxidants only when applied at higher concentrations: H_2O_2 ($\geq 0.2\%$), KO_2 (≥ 20 mM), and menadione (≥ 5 mM). However, elevated sensitivity to H_2O_2 seen in $\Delta nps6$ or $\Delta yap1$ mutant was alleviated by adding ferric iron into the medium, implicating an important role of iron and siderophore-mediated iron acquisition in the ROS resistance. We have observed that expression of the *AaNPS6* gene was significantly downregulated in fungal mutant lacking *YAP1*. Deletion of *YAP1* in *A. alternata* also reduced the production of siderophores. Moreover, the rescued strain expressing a functional copy of *YAP1* accumulated wild-type level of siderophores and *AaNPS6* gene transcript. Expression of *AaNPS6* and production of siderophores were also downregulated in fungal strain lacking *HOG1* or *NOXA* (L.-H. Chen and S.L. Yang, personal communication), confirming a close linkage between iron acquisition and ROS resistance. The wild-type strain of *A. alternata*, when grown under iron-depleted conditions, produced barely measurable catalase and SOD activities (L.-H. Chen, personal communication). Both antioxidant activities were detectable in *A. alternata* cultured under iron-rich conditions. Hence, we concluded that the increased sensitivity to oxidative stress and the reduced pathogenicity seen in $\Delta yap1$, $\Delta hog1$, $\Delta noxA$, or $\Delta nps6$ were seemingly due to the decreased ability of oxidative stress-detoxifying enzymes.

Genetic analyses revealed that siderophore produced by *NPS6* is required for full virulence of the tangerine pathotype of *A. alternata*. This is likely due to the inability of $\Delta nps6$ mutant to detoxify toxic ROS efficiently. Siderophores are also required for fungal pathogenesis in *A. brassicicola*, *As. fumigatus*, *Co. heterostrophus*, *Co. miyabeanus*, *F. graminearum*, and *M. grisea* [208, 210, 213, 214]. However, siderophore is not required for pathogenesis of the basidiomycete maize pathogen, *U. maydis* [215]. It will be of great interest to determine if *NPS6* is also regulated by *YAP1* and *HOG1* in other fungal species.

9. Conclusions

Based on the observed phenotypes derived from mutants lacking *YAP1*, *HOG1*, *SKN7*, *NOX*, or *NPS6*, a regulatory network is assembled to underscore the intricate interplays

among these signaling pathways in *A. alternata* (Figure 4). The *NOX* complex appears to have an important role in the production of ROS, which may act as secondary messages to regulate various metabolic processes in *A. alternata*. The *NOX* complex is required for transcriptional activation of two important regulators, *YAP1* and *HOG1*, which subsequently regulate the expression of genes encoding the nonribosomal peptide synthetase (*NPS6*) and perhaps other enzymes involved in the biosynthesis of siderophores as well. *SKN7* physically interacts with *YAP1*, regulating the genes involved in ROS detoxification. Maintaining iron homeostasis is critical for ROS detoxification because of the requirement of iron for antioxidant enzymatic activities. Impairment of the *NOX* complex, the *YAP1* regulator, the *HOG1* kinase, or the siderophore-mediated iron acquisition in *A. alternata* impacts its ability to detoxify ROS and to colonize host plant, implicating the importance of ROS detoxification in the successful pathogenesis of *A. alternata*. In addition to detoxifying ROS, *A. alternata* utilizes specialized or synergistically regulated signaling pathways, involved in *HSK1*, *HOG1*, and/or *SKN7*, in response to osmotic stress, fungicides, and other toxic compounds. This cross-interaction between different signaling pathways may have eccentric advantages for integrating cellular responses to a broader spectrum of environmental stimuli.

Acknowledgments

The author would like to thank the current and former Chung Laboratory Members Siwy Ling Yang, Li-Hung Chen, Hsieh-Chin Tsai, and Dr. Ching-Hsuan Lin for their contributions to this work and Wen-Chun Wu for her assistance in taking photographs. This paper was supported by the Florida Agricultural Experiment Station.

References

- [1] B. P. H. J. Thomma, "Alternaria spp.: from general saprophyte to specific parasite," *Molecular Plant Pathology*, vol. 4, no. 4, pp. 225–236, 2003.
- [2] E. G. Simmons, "Alternaria taxonomy: current status, viewpoint, change," in *Alternaria Biology, Plant Disease and Metabolites*, J. Chelkowski and A. Visconti, Eds., pp. 1–35, Elsevier, Amsterdam, The Netherlands, 1992.
- [3] M. Kusaba and T. Tsuge, "Phylogeny of Alternaria fungi known to produce host specific toxins on the basis of variation in internal transcribed spacers of ribosomal DNA," *Current Genetics*, vol. 28, no. 5, pp. 491–498, 1995.
- [4] J. Rotem, *The Genus Alternaria: Biology, Epidemiology and Pathogenicity*, APS Press, St. Paul, Minn, USA, 1994.
- [5] S. Nishimura and K. Kohmoto, "Host-specific toxins and chemical structures from Alternaria species," *Annual Review of Phytopathology*, vol. 21, pp. 87–116, 1983.
- [6] N. Montemurro and A. Visconti, "Alternaria metabolites—chemical and biological data," in *Alternaria Biology, Plant Disease and Metabolites*, J. Chelkowski and A. Visconti, Eds., pp. 449–557, Elsevier, Amsterdam, The Netherlands, 1992.
- [7] R. D. Johnson, L. Johnson, Y. Itoh, M. Kodama, H. Otani, and K. Kohmoto, "Cloning and characterization of a cyclic

- peptide synthetase gene from *Alternaria alternata* apple pathotype whose product is involved in AM-toxin synthesis and pathogenicity," *Molecular Plant-Microbe Interactions*, vol. 13, no. 7, pp. 742–753, 2000.
- [8] L. J. Johnson, R. D. Johnson, H. Akamatsu et al., "Spontaneous loss of a conditionally dispensable chromosome from the *Alternaria alternata* apple pathotype leads to loss of toxin production and pathogenicity," *Current Genetics*, vol. 40, no. 1, pp. 65–72, 2001.
 - [9] R. Hatta, K. Ito, Y. Hosaki et al., "A conditionally dispensable chromosome controls host-specific pathogenicity in the fungal plant pathogen *Alternaria alternata*," *Genetics*, vol. 161, no. 1, pp. 59–70, 2002.
 - [10] K. Ito, T. Tanaka, R. Hatta, M. Yamamoto, K. Akimitsu, and T. Tsuge, "Dissection of the host range of the fungal plant pathogen *Alternaria alternata* by modification of secondary metabolism," *Molecular Microbiology*, vol. 52, no. 2, pp. 399–411, 2004.
 - [11] Y. Harimoto, R. Hatta, M. Kodama, M. Yamamoto, H. Otani, and T. Tsuge, "Expression profiles of genes encoded by the supernumerary chromosome controlling AM-toxin biosynthesis and pathogenicity in the apple pathotype of *Alternaria alternata*," *Molecular Plant-Microbe Interactions*, vol. 20, no. 12, pp. 1463–1476, 2007.
 - [12] M. Orved, P. Haggblom, and K. Soderhall, "Nitrogen inhibition of mycotoxin production by *Alternaria alternata*," *Applied and Environmental Microbiology*, vol. 54, no. 10, pp. 2361–2364, 1988.
 - [13] R. K. Bush and J. J. Prochnau, "Alternaria-induced asthma," *Journal of Allergy and Clinical Immunology*, vol. 113, no. 2, pp. 227–234, 2004.
 - [14] K. Kohmoto, K. Akimitsu, and H. Otani, "Correlation of resistance and susceptibility of citrus to *Alternaria alternata* with sensitivity to host-specific toxins," *Phytopathology*, vol. 81, pp. 719–722, 1991.
 - [15] H. Otani, K. Kohmoto, and M. Kodama, "Alternaria toxins and their effects on host plants," *Canadian Journal of Botany*, vol. 73, pp. S453–S458, 1996.
 - [16] T. L. Peever, G. Su, L. Carpenter-Boggs, and L. W. Timmer, "Molecular systematics of citrus-associated *Alternaria* species," *Mycologia*, vol. 96, no. 1, pp. 119–134, 2004.
 - [17] T. L. Peever, Y. Canihos, L. Olsen, A. Ibañez, Y. C. Liu, and L. W. Timmer, "Population genetic structure and host specificity of *Alternaria* spp. causing brown spot of Minneola tangelo and rough lemon in Florida," *Phytopathology*, vol. 89, no. 10, pp. 851–860, 1999.
 - [18] K. Akimitsu, K. Kohmoto, H. Otani, and S. Nishimura, "Host-specific effect of toxin from the rough lemon pathotype of *Alternaria alternata* on mitochondria," *Plant Physiology*, vol. 89, pp. 925–931, 1989.
 - [19] K. Ohtani, H. Yamamoto, and K. Akimitsu, "Sensitivity to *Alternaria alternata* toxin in citrus because of altered mitochondrial RNA processing," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 99, no. 4, pp. 2439–2444, 2002.
 - [20] K. Kohmoto, Y. Itoh, N. Shimomura et al., "Isolation and biological activities of two host-specific toxins from tangerine pathotype of *Alternaria alternata*," *Phytopathology*, vol. 83, pp. 495–502, 1993.
 - [21] C. H. Lin, S. L. Yang, and K. R. Chung, "Cellular responses required for oxidative stress tolerance, colonization, and lesion formation by the necrotrophic fungus *Alternaria alternata* in citrus," *Current Microbiology*, vol. 62, no. 3, pp. 807–815, 2011.
 - [22] R. Mittler, "Oxidative stress, antioxidants and stress tolerance," *Trends in Plant Science*, vol. 7, no. 9, pp. 405–410, 2002.
 - [23] S. Neill, R. Desikan, and J. Hancock, "Hydrogen peroxide signalling," *Current Opinion in Plant Biology*, vol. 5, no. 5, pp. 388–395, 2002.
 - [24] K. Apel and H. Hirt, "Reactive oxygen species: metabolism, oxidative stress, and signal transduction," *Annual Review of Plant Biology*, vol. 55, pp. 373–399, 2004.
 - [25] J. Aguirre, M. Ríos-Momberg, D. Hewitt, and W. Hansberg, "Reactive oxygen species and development in microbial eukaryotes," *Trends in Microbiology*, vol. 13, no. 3, pp. 111–118, 2005.
 - [26] E. Herrero, J. Ros, G. Bellí, and E. Cabisco, "Redox control and oxidative stress in yeast cells," *Biochimica et Biophysica Acta*, vol. 1780, no. 11, pp. 1217–1235, 2008.
 - [27] E. A. Veal, A. M. Day, and B. A. Morgan, "Hydrogen peroxide sensing and signaling," *Molecular Cell*, vol. 26, no. 1, pp. 1–14, 2007.
 - [28] M. Orozco-Cardenas and C. A. Ryan, "Hydrogen peroxide is generated systemically in plant leaves by wounding and systemin via the octadecanoid pathway," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 96, no. 11, pp. 6553–6557, 1999.
 - [29] M. R. Branco, H. S. Marinho, L. Cyrne, and F. Antunes, "Decrease of H₂O₂ plasma membrane permeability during Adaptation to H₂O₂ in *Saccharomyces cerevisiae*," *Journal of Biological Chemistry*, vol. 279, no. 8, pp. 6501–6506, 2004.
 - [30] J. R. Stone and S. Yang, "Hydrogen peroxide: a signaling messenger," *Antioxidants and Redox Signaling*, vol. 8, no. 3–4, pp. 243–270, 2006.
 - [31] B. D'Autrèaux and M. B. Toledano, "ROS as signalling molecules: mechanisms that generate specificity in ROS homeostasis," *Nature Reviews Molecular Cell Biology*, vol. 8, no. 10, pp. 813–824, 2007.
 - [32] I. Ślesak, M. Libik, B. Karpinska, S. Karpinski, and Z. Miszalski, "The role of hydrogen peroxide in regulation of plant metabolism and cellular signalling in response to environmental stresses," *Acta Biochimica Polonica*, vol. 54, no. 1, pp. 39–50, 2007.
 - [33] B. Halliwell and J. M. C. Gutteridge, "Biologically relevant metal ion-dependent hydroxyl radical generation. An update," *FEBS Letters*, vol. 307, no. 1, pp. 108–112, 1992.
 - [34] D. A. Wink, R. W. Nims, J. E. Saavedra, W. E. Utermahlen Jr., and P. C. Ford, "The fenton oxidation mechanism: reactivities of biologically relevant substrates with two oxidizing intermediates differ from those predicted for the hydroxyl radical," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 91, no. 14, pp. 6604–6608, 1994.
 - [35] J. A. Imlay, "Pathways of oxidative damage," *Annual Review of Microbiology*, vol. 57, pp. 395–418, 2003.
 - [36] M. C. Mehdy, "Active oxygen species in plant defense against pathogens," *Plant Physiology*, vol. 105, no. 2, pp. 467–472, 1994.
 - [37] E. Lam, N. Kato, and M. Lawton, "Programmed cell death, mitochondria and the plant hypersensitive response," *Nature*, vol. 411, no. 6839, pp. 848–853, 2001.
 - [38] D. G. Mellersh, I. V. Foulds, V. J. Higgins, and M. C. Heath, "H₂O₂ plays different roles in determining penetration failure in three diverse plant-fungal interactions," *Plant Journal*, vol. 29, no. 3, pp. 257–268, 2002.
 - [39] J. L. Dangl and J. D. G. Jones, "Plant pathogens and integrated defence responses to infection," *Nature*, vol. 411, no. 6839, pp. 826–833, 2001.

- [40] R. Mahalingam and N. Fedoroff, "Stress response, cell death and signalling: the many faces of reactive oxygen species," *Physiologia Plantarum*, vol. 119, no. 1, pp. 56–68, 2003.
- [41] N. P. Shetty, R. Mehrabi, H. Lütken et al., "Role of hydrogen peroxide during the interaction between the hemibiotrophic fungal pathogen *Septoria tritici* and wheat," *New Phytologist*, vol. 174, no. 3, pp. 637–647, 2007.
- [42] N. Doke, Y. Miura, L. M. Sanchez et al., "The oxidative burst protects plants against pathogen attack: mechanism and role as an emergency signal for plant bio-defence—a review," *Gene*, vol. 179, no. 1, pp. 45–51, 1996.
- [43] C. Lamb and R. A. Dixon, "The oxidative burst in plant disease resistance," *Annual Review of Plant Biology*, vol. 48, pp. 251–275, 1997.
- [44] J. T. Greenberg, "Programmed cell death in plant-pathogen interactions," *Annual Review of Plant Biology*, vol. 48, pp. 525–545, 1997.
- [45] P. Wojtaszek, "Oxidative burst: an early plant response to pathogen infection," *Biochemical Journal*, vol. 322, no. 3, pp. 681–692, 1997.
- [46] J. T. Greenberg and N. Yao, "The role of regulation of programmed cell death in plant-pathogen interactions," *Cellular Microbiology*, vol. 6, no. 3, pp. 201–211, 2004.
- [47] A. M. Mayer, R. C. Staples, and N. L. Gil-ad, "Mechanisms of survival of necrotrophic fungal plant pathogens in hosts expressing the hypersensitive response," *Phytochemistry*, vol. 58, no. 1, pp. 33–41, 2001.
- [48] A. J. Able, "Role of reactive oxygen species in the response of barley to necrotrophic pathogens," *Protoplasma*, vol. 221, no. 1–2, pp. 137–143, 2003.
- [49] J. Glazebrook, "Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens," *Annual Review of Phytopathology*, vol. 43, pp. 205–227, 2005.
- [50] C. Unger, S. Kleta, G. Jandl, and A. V. Tiedemann, "Suppression of the defence-related oxidative burst in bean leaf tissue and bean suspension cells by the necrotrophic pathogen *Botrytis cinerea*," *Journal of Phytopathology*, vol. 153, no. 1, pp. 15–26, 2005.
- [51] A. Walz, I. Zingen-Sell, S. Theisen, and A. Kortekamp, "Reactive oxygen intermediates and oxalic acid in the pathogenesis of the necrotrophic fungus *Sclerotinia sclerotiorum*," *European Journal of Plant Pathology*, vol. 120, no. 4, pp. 317–330, 2008.
- [52] M. Trujillo, K. H. Kogel, and R. Hückelhoven, "Superoxide and hydrogen peroxide play different roles in the nonhost interaction of barley and wheat with inappropriate formae speciales of *Blumeria graminis*," *Molecular Plant-Microbe Interactions*, vol. 17, no. 3, pp. 304–312, 2004.
- [53] Y. Narusaka, M. Narusaka, M. Seki et al., "Cytological and molecular analyses of non-host resistance of *Arabidopsis thaliana* to *Alternaria alternata*," *Molecular Plant Pathology*, vol. 6, no. 6, pp. 615–627, 2005.
- [54] H. H. Divon and R. Fluhr, "Nutrition acquisition strategies during fungal infection of plants," *FEMS Microbiology Letters*, vol. 266, no. 1, pp. 65–74, 2007.
- [55] E. M. Govrin and A. Levine, "The hypersensitive response facilitates plant infection by the necrotrophic pathogen *Botrytis cinerea*," *Current Biology*, vol. 10, no. 13, pp. 751–757, 2000.
- [56] M. E. Daub and M. Ehrenschaft, "The photoactivated *Cercospora* toxin cercosporin: contributions to plant disease and fundamental biology," *Annual Review of Phytopathology*, vol. 38, pp. 461–490, 2000.
- [57] V. J. Thannickal and B. L. Fanburg, "Reactive oxygen species in cell signaling," *American Journal of Physiology—Lung Cellular and Molecular Physiology*, vol. 279, no. 6, pp. L1005–L1028, 2000.
- [58] M. E. Daub, S. Herrero, and K. R. Chung, "Photoactivated perylenequinone toxins in fungal pathogenesis of plants," *FEMS Microbiology Letters*, vol. 252, no. 2, pp. 197–206, 2005.
- [59] R. A. Miller and B. E. Britigan, "Role of oxidants in microbial pathophysiology," *Clinical Microbiology Reviews*, vol. 10, no. 1, pp. 1–18, 1997.
- [60] B. Thammavongs, E. Denou, G. Missous, M. Guéguen, and J. M. Panoff, "Response to environmental stress as a global phenomenon in biology: the example of microorganisms," *Microbes and Environments*, vol. 23, no. 1, pp. 20–23, 2008.
- [61] W. S. Moye-Rowley, "Regulation of the transcriptional response to oxidative stress in fungi: similarities and differences," *Eukaryotic Cell*, vol. 2, no. 3, pp. 381–389, 2003.
- [62] M. B. Toledano, A. Delaunay, L. Monceau, and F. Tacnet, "Microbial H₂O₂ sensors as archetypical redox signaling modules," *Trends in Biochemical Sciences*, vol. 29, no. 7, pp. 351–357, 2004.
- [63] A. Delaunay, A. D. Isnard, and M. B. Toledano, "H₂O₂ sensing through oxidation of the Yap1 transcription factor," *The EMBO Journal*, vol. 19, no. 19, pp. 5157–5166, 2000.
- [64] W. M. Toone and N. Jones, "Stress-activated signalling pathways in yeast," *Genes to Cells*, vol. 3, no. 8, pp. 485–498, 1998.
- [65] W. M. Toone and N. Jones, "AP-1 transcription factors in yeast," *Current Opinion in Genetics and Development*, vol. 9, no. 1, pp. 55–61, 1999.
- [66] W. M. Toone, B. A. Morgan, and N. Jones, "Redox control of AP-1-like factors in yeast and beyond," *Oncogene*, vol. 20, no. 19, pp. 2336–2346, 2001.
- [67] C. Yan, L. H. Lee, and L. I. Davis, "Crm1p mediates regulated nuclear export of a yeast AP-1-like transcription factor," *The EMBO Journal*, vol. 17, no. 24, pp. 7416–7429, 1998.
- [68] S. T. Coleman, E. A. Epping, S. M. Steggerda, and W. S. Moye-Rowley, "Yap1p activates gene transcription in an oxidant-specific fashion," *Molecular and Cellular Biology*, vol. 19, no. 12, pp. 8302–8313, 1999.
- [69] M. J. Wood, E. C. Andrade, and G. Storz, "The redox domain of the Yap1p transcription factor contains two disulfide bonds," *Biochemistry*, vol. 42, no. 41, pp. 11982–11991, 2003.
- [70] K. Gulshan and W. S. Moye-Rowley, "Multidrug resistance in fungi," *Eukaryotic Cell*, vol. 6, no. 11, pp. 1933–1942, 2007.
- [71] S. Okazaki, T. Tachibana, A. Naganuma, N. Mano, and S. Kuge, "Multistep disulfide bond formation in Yap1 is required for sensing and transduction of H₂O₂ stress signal," *Molecular Cell*, vol. 27, no. 4, pp. 675–688, 2007.
- [72] S. Kuge, M. Arita, A. Murayama et al., "Regulation of the yeast Yap1p nuclear export signal is mediated by redox signal-induced reversible disulfide bond formation," *Molecular and Cellular Biology*, vol. 21, no. 18, pp. 6139–6150, 2001.
- [73] S. G. Cessna, V. E. Sears, M. B. Dickman, and P. S. Low, "Oxalic acid, a pathogenicity factor for *Sclerotinia sclerotiorum*, suppresses the oxidative burst of the host plant," *Plant Cell*, vol. 12, no. 11, pp. 2191–2199, 2000.
- [74] R. Hückelhoven and K. H. Kogel, "Reactive oxygen intermediates in plant-microbe interactions: Who is who in powdery mildew resistance?" *Planta*, vol. 216, no. 6, pp. 891–902, 2003.
- [75] S. H. Spoel, J. S. Johnson, and X. Dong, "Regulation of trade-offs between plant defenses against pathogens with different

- lifestyles," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 104, no. 47, pp. 18842–18847, 2007.
- [76] D. Takemoto, A. Tanaka, and B. Scott, "NADPH oxidases in fungi: diverse roles of reactive oxygen species in fungal cellular differentiation," *Fungal Genetics and Biology*, vol. 44, no. 11, pp. 1065–1076, 2007.
- [77] B. Enjalbert, D. M. MacCallum, F. C. Odds, and A. J. P. Brown, "Niche-specific activation of the oxidative stress response by the pathogenic fungus *Candida albicans*," *Infection and Immunity*, vol. 75, no. 5, pp. 2143–2151, 2007.
- [78] L. Molina and R. Kahmann, "An *Ustilago maydis* gene involved in H₂O₂ detoxification is required for virulence," *Plant Cell*, vol. 19, no. 7, pp. 2293–2309, 2007.
- [79] S. Lev, R. Hadar, P. Amedeo, S. E. Baker, O. C. Yoder, and B. A. Horwitz, "Activation of an AP1-like transcription factor of the maize pathogen *Cochliobolus heterostrophus* in response to oxidative stress and plant signals," *Eukaryotic Cell*, vol. 4, no. 2, pp. 443–454, 2005.
- [80] F. Lessing, O. Kniemeyer, I. Wozniok et al., "The *Aspergillus fumigatus* transcriptional regulator AfYap1 represents the major regulator for defense against reactive oxygen intermediates but is dispensable for pathogenicity in an intranasal mouse infection model," *Eukaryotic Cell*, vol. 6, no. 12, pp. 2290–2302, 2007.
- [81] K. Huang, K. J. Czymmek, J. L. Caplan, J. A. Sweigard, and N. M. Donofrio, "HYR1-mediated detoxification of reactive oxygen species is required for full virulence in the rice blast fungus," *PLoS Pathogens*, vol. 7, no. 4, Article ID e1001335, 2011.
- [82] K. H. Kim, S. D. Willger, S. W. Park et al., "Tmpl, a transmembrane protein required for intracellular redox homeostasis and virulence in a plant and an animal fungal pathogen," *PLoS Pathogens*, vol. 5, no. 11, Article ID e1000653, 2009.
- [83] C. H. Lin, S. L. Yang, and K. R. Chung, "The YAP1 homolog-mediated oxidative stress tolerance is crucial for pathogenicity of the necrotrophic fungus *Alternaria alternata* in citrus," *Molecular Plant-Microbe Interactions*, vol. 22, no. 8, pp. 942–952, 2009.
- [84] S. L. Yang, C. H. Lin, and K. R. Chung, "Coordinate control of oxidative stress tolerance, vegetative growth, and fungal pathogenicity via the AP1 pathway in the rough lemon pathotype of *Alternaria alternata*," *Physiological and Molecular Plant Pathology*, vol. 74, no. 2, pp. 100–110, 2009.
- [85] H. Véléz, N. J. Glassbrook, and M. E. Daub, "Mannitol metabolism in the phytopathogenic fungus *Alternaria alternata*," *Fungal Genetics and Biology*, vol. 44, no. 4, pp. 258–268, 2007.
- [86] H. Véléz, N. J. Glassbrook, and M. E. Daub, "Mannitol biosynthesis is required for plant pathogenicity by *Alternaria alternata*," *FEMS Microbiology Letters*, vol. 285, no. 1, pp. 122–129, 2008.
- [87] C. H. Lin and K. R. Chung, "Specialized and shared functions of the histidine kinase- and *HOG1* MAP kinase-mediated signaling pathways in *Alternaria alternata*, a filamentous fungal pathogen of citrus," *Fungal Genetics and Biology*, vol. 47, no. 10, pp. 818–827, 2010.
- [88] E. A. Castillo, J. Ayté, C. Chiva et al., "Diethylmaleate activates the transcription factor Pap1 by covalent modification of critical cysteine residues," *Molecular Microbiology*, vol. 45, no. 1, pp. 243–254, 2002.
- [89] S. Tsurumi and Y. Ohwaki, "Transport of ¹⁴C-labeled indoleacetic acid in *Vicia* root segments," *Plant and Cell Physiology*, vol. 19, no. 7, pp. 1195–1206, 1978.
- [90] K. A. Lahey, R. Yuan, J. K. Burns, P. P. Ueng, L. W. Timmer, and K. R. Chung, "Induction of phytohormones and differential gene expression in citrus flowers infected by the fungus *Colletotrichum acutatum*," *Molecular Plant-Microbe Interactions*, vol. 17, no. 12, pp. 1394–1401, 2004.
- [91] T. Watanabe, K. Koller, and K. Messner, "Copper-dependent depolymerization of lignin in the presence of fungal metabolite, pyridine," *Journal of Biotechnology*, vol. 62, no. 3, pp. 221–230, 1998.
- [92] F. Nerud, P. Baldrian, J. Gabriel, and D. Ogbeifun, "Decolorization of synthetic dyes by the Fenton reagent and the Cu/pyridine/H₂O₂ system," *Chemosphere*, vol. 44, no. 5, pp. 957–961, 2001.
- [93] B. N. Kunkel and D. M. Brooks, "Cross talk between signaling pathways in pathogen defense," *Current Opinion in Plant Biology*, vol. 5, no. 4, pp. 325–331, 2002.
- [94] J. L. Martindale and N. J. Holbrook, "Cellular response to oxidative stress: signaling for suicide and survival," *Journal of Cellular Physiology*, vol. 192, no. 1, pp. 1–15, 2002.
- [95] E. Vranová, D. Inzé, and F. Van Breusegem, "Signal transduction during oxidative stress," *Journal of Experimental Botany*, vol. 53, no. 372, pp. 1227–1236, 2002.
- [96] H. P. Harding, Y. Zhang, H. Zeng et al., "An integrated stress response regulates amino acid metabolism and resistance to oxidative stress," *Molecular Cell*, vol. 11, no. 3, pp. 619–633, 2003.
- [97] A. Ikner and K. Shiozaki, "Yeast signaling pathways in the oxidative stress response," *Mutation Research*, vol. 569, no. 1–2, pp. 13–27, 2005.
- [98] N. Chauhan, J. P. Latge, and R. Calderone, "Signalling and oxidant adaptation in *Candida albicans* and *Aspergillus fumigatus*," *Nature Reviews Microbiology*, vol. 4, no. 6, pp. 435–444, 2006.
- [99] A. Pitzschke, C. Forzani, and H. Hirt, "Reactive oxygen species signaling in plants," *Antioxidants and Redox Signaling*, vol. 8, no. 9–10, pp. 1757–1764, 2006.
- [100] P. Thomason and R. Kay, "Eukaryotic signal transduction via histidine-aspartate phosphorelay," *Journal of Cell Science*, vol. 113, no. 18, pp. 3141–3150, 2000.
- [101] P. M. Wolanin, P. A. Thomason, and J. B. Stock, "Histidine protein kinases: key signal transducers outside the animal kingdom," *Genome Biology*, vol. 3, no. 10, article 3013, pp. 3013.1–3013.8, 2002.
- [102] J. S. Parkinson and E. C. Kofoid, "Communication modules in bacterial signaling proteins," *Annual Review of Genetics*, vol. 26, pp. 71–112, 1992.
- [103] J. A. Hoch, "Two-component and phosphorelay signal transduction," *Current Opinion in Microbiology*, vol. 3, no. 2, pp. 165–170, 2000.
- [104] J. L. Santos and K. Shiozaki, "Fungal histidine kinases," *Science's STKE*, vol. 2001, no. 98, p. re1, 2001.
- [105] A. H. West and A. M. Stock, "Histidine kinases and response regulator proteins in two-component signaling systems," *Trends in Biochemical Sciences*, vol. 26, no. 6, pp. 369–376, 2001.
- [106] S. M. Wurgler-Murphy and H. Saito, "Two-component signal transducers and MAPK cascades," *Trends in Biochemical Sciences*, vol. 22, no. 5, pp. 172–176, 1997.
- [107] M. Kruppa and R. Calderone, "Two-component signal transduction in human fungal pathogens," *FEMS Yeast Research*, vol. 6, no. 2, pp. 149–159, 2006.

- [108] T. Maeda, S. M. Wurgler-Murphy, and H. Saito, "A two-component system that regulates an osmosensing MAP kinase cascade in yeast," *Nature*, vol. 369, no. 6477, pp. 242–245, 1994.
- [109] F. Posas, S. M. Wurgler-Murphy, T. Maeda, E. A. Witten, T. C. Thai, and H. Saito, "Yeast HOG1 MAP kinase cascade is regulated by a multistep phosphorelay mechanism in the SLN1-YPD1-SSK1 'two-component' osmosensor," *Cell*, vol. 86, no. 6, pp. 865–875, 1996.
- [110] P. J. Westfall, D. R. Ballon, and J. Thorner, "When the stress of your environment makes you go HOG wild," *Science*, vol. 306, no. 5701, pp. 1511–1512, 2004.
- [111] N. L. Catlett, O. C. Yoder, and B. G. Turgeon, "Whole-genome analysis of two-component signal transduction genes in fungal pathogens," *Eukaryotic Cell*, vol. 2, no. 6, pp. 1151–1161, 2003.
- [112] L. A. Alex, K. A. Borkovich, and M. I. Simon, "Hyphal development in *Neurospora crassa*: involvement of a two-component histidine kinase," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 93, no. 8, pp. 3416–3421, 1996.
- [113] W. Cui, R. E. Beever, S. L. Parkes, P. L. Weeds, and M. D. Templeton, "An osmosensing histidine kinase mediates dicarboximide fungicide resistance in *Botryotinia fuckeliana* (*Botrytis cinerea*)," *Fungal Genetics and Biology*, vol. 36, no. 3, pp. 187–198, 2002.
- [114] I. B. Dry, K. H. Yuan, and D. G. Hutton, "Dicarboximide resistance in field isolates of *Alternaria alternata* is mediated by a mutation in a two-component histidine kinase gene," *Fungal Genetics and Biology*, vol. 41, no. 1, pp. 102–108, 2004.
- [115] H. Avenot, P. Simoneau, B. Iacomi-Vasilescu, and N. Bataillé-Simoneau, "Characterization of mutations in the two-component histidine kinase gene *AbNIK1* from *Alternaria brassicicola* that confer high dicarboximide and phenylpyrrole resistance," *Current Genetics*, vol. 47, no. 4, pp. 234–243, 2005.
- [116] T. Motoyama, K. Kadokura, T. Ohira et al., "A two-component histidine kinase of the rice blast fungus is involved in osmotic stress response and fungicide action," *Fungal Genetics and Biology*, vol. 42, no. 3, pp. 200–212, 2005.
- [117] A. Yoshimi, K. Kojima, Y. Takano, and C. Tanaka, "Group III histidine kinase is a positive regulator of HOG1-type mitogen-activated protein kinase in filamentous fungi," *Eukaryotic Cell*, vol. 4, no. 11, pp. 1820–1828, 2005.
- [118] M. Viaud, S. Fillinger, W. Liu et al., "A class III histidine kinase acts as a novel virulence factor in *Botrytis cinerea*," *Molecular Plant-Microbe Interactions*, vol. 19, no. 9, pp. 1042–1050, 2006.
- [119] N. Ochiai, T. Tokai, T. Nishiuchi, N. Takahashi-Ando, M. Fujimura, and M. Kimura, "Involvement of the osmosensor histidine kinase and osmotic stress-activated protein kinases in the regulation of secondary metabolism in *Fusarium graminearum*," *Biochemical and Biophysical Research Communications*, vol. 363, no. 3, pp. 639–644, 2007.
- [120] I. Vargas-Pérez, O. Sánchez, L. Kawasaki, D. Georgellis, and J. Aguirre, "Response regulators SrrA and SskA are central components of a phosphorelay system involved in stress signal transduction and asexual sporulation in *Aspergillus nidulans*," *Eukaryotic Cell*, vol. 6, no. 9, pp. 1570–1583, 2007.
- [121] A. Dongo, N. Bataillé-Simoneau, C. Champion et al., "The group III two-component histidine kinase of filamentous fungi is involved in the fungicidal activity of the bacterial polyketide ambruticin," *Applied and Environmental Microbiology*, vol. 75, no. 1, pp. 127–134, 2009.
- [122] K. Kojima, Y. Takano, A. Yoshimi, C. Tanaka, T. Kikuchi, and T. Okuno, "Fungicide activity through activation of a fungal signalling pathway," *Molecular Microbiology*, vol. 53, no. 6, pp. 1785–1796, 2004.
- [123] K. Furukawa, Y. Hoshi, T. Maeda, T. Nakajima, and K. Abe, "*Aspergillus nidulans* HOG pathway is activated only by two-component signalling pathway in response to osmotic stress," *Molecular Microbiology*, vol. 56, no. 5, pp. 1246–1261, 2005.
- [124] M. Lenassi and A. Plemenitaš, "Novel group VII histidine kinase HwHhk7B from the halophilic fungi *Hortaea werneckii* has a putative role in osmosensing," *Current Genetics*, vol. 51, no. 6, pp. 393–405, 2007.
- [125] C. H. Lin, S. L. Yang, N. Y. Wang, and K. R. Chung, "The FUS3 MAPK signaling pathway of the citrus pathogen *Alternaria alternata* functions independently or cooperatively with the fungal redox-responsive AP1 regulator for diverse developmental, physiological and pathogenic processes," *Fungal Genetics and Biology*, vol. 47, no. 4, pp. 381–391, 2010.
- [126] N. Rispaill and A. di Pietro, "The two-component histidine kinase Fhk1 controls stress adaptation and virulence of *Fusarium oxysporum*," *Molecular Plant Pathology*, vol. 11, no. 3, pp. 395–407, 2010.
- [127] C. J. Eaton, I. Jourdain, S. J. Foster, J. S. Hyams, and B. Scott, "Functional analysis of a fungal endophyte stress-activated MAP kinase," *Current Genetics*, vol. 53, no. 3, pp. 163–174, 2008.
- [128] Y. S. Bahn, K. Kojima, G. M. Cox, and J. Heitman, "A unique fungal two-component system regulates stress responses, drug sensitivity, sexual development, and virulence of *Cryptococcus neoformans*," *Molecular Biology of the Cell*, vol. 17, no. 7, pp. 3122–3135, 2006.
- [129] E. Nathues, C. Jörgens, N. Lorenz, and P. Tudzynski, "The histidine kinase CpHK2 has impact on spore germination, oxidative stress and fungicide resistance, and virulence of the ergot fungus *Claviceps purpurea*," *Molecular Plant Pathology*, vol. 8, no. 5, pp. 653–665, 2007.
- [130] M. C. Gustin, J. Albertyn, M. Alexander, and K. Davenport, "Map kinase pathways in the yeast *Saccharomyces cerevisiae*," *Microbiology and Molecular Biology Reviews*, vol. 62, no. 4, pp. 1264–1300, 1998.
- [131] M. Qi and E. A. Elion, "MAP kinase pathways," *Journal of Cell Science*, vol. 118, no. 16, pp. 3569–3572, 2005.
- [132] S. L. Pelech and J. S. Sanghera, "MAP kinases: charting the regulatory pathways," *Science*, vol. 257, no. 5075, pp. 1355–1356, 1992.
- [133] M. J. Robinson and M. H. Cobb, "Mitogen-activated protein kinase pathways," *Current Opinion in Cell Biology*, vol. 9, no. 2, pp. 180–186, 1997.
- [134] D. Kültz, "Phylogenetic and functional classification of mitogen—and stress—activated protein kinases," *Journal of Molecular Evolution*, vol. 46, no. 5, pp. 571–588, 1998.
- [135] S. M. O'Rourke and I. Herskowitz, "The HOG1 MAPK prevents cross talk between the HOG and pheromone response MAPK pathways in *Saccharomyces cerevisiae*," *Genes and Development*, vol. 12, no. 18, pp. 2874–2886, 1998.
- [136] T. R. Shock, J. Thompson, J. R. Yates, and H. D. Madhani, "HOG1 mitogen-activated protein kinase (MAPK) interrupts signal transduction between the Kss1 MAPK and the Tec1 transcription factor to maintain pathway specificity," *Eukaryotic Cell*, vol. 8, no. 4, pp. 606–616, 2009.
- [137] S. M. Park, E. S. Choi, M. J. Kim, B. J. Cha, M. S. Yang, and D. H. Kim, "Characterization of HOG1 homologue, *CpMK1*, from *Cryphonectria parasitica* and evidence for hypovirus-mediated perturbation of its phosphorylation in response to hypertonic

- stress," *Molecular Microbiology*, vol. 51, no. 5, pp. 1267–1277, 2004.
- [138] Y. S. Bahn, K. Kojima, G. M. Cox, and J. Heitman, "Specialization of the HOG pathway and its impact on differentiation and virulence of *Cryptococcus neoformans*," *Molecular Biology of the Cell*, vol. 16, no. 5, pp. 2285–2300, 2005.
- [139] R. Mehrabi, L. H. Zwiers, M. A. de Waard, and G. H. J. Kema, "MgHOG1 regulates dimorphism and pathogenicity in the fungal wheat pathogen *Mycosphaerella graminicola*," *Molecular Plant-Microbe Interactions*, vol. 19, no. 11, pp. 1262–1269, 2006.
- [140] V. Menon, D. Li, N. Chauhan et al., "Functional studies of the Ssk1p response regulator protein of *Candida albicans* as determined by phenotypic analysis of receiver domain point mutants," *Molecular Microbiology*, vol. 62, no. 4, pp. 997–1013, 2006.
- [141] N. Segmüller, L. Kokkelink, S. Giesbert, D. Odinius, J. van Kan, and P. Tudzynski, "NADPH oxidases are involved in differentiation and pathogenicity in *Botrytis cinerea*," *Molecular Plant-Microbe Interactions*, vol. 21, no. 6, pp. 808–819, 2008.
- [142] A. Igbaria, S. Lev, M. S. Rose et al., "Distinct and combined roles of the MAP kinases of *Cochliobolus heterostrophus* in virulence and stress responses," *Molecular Plant-Microbe Interactions*, vol. 21, no. 6, pp. 769–780, 2008.
- [143] K. P. Dixon, J. R. Xu, N. Smirnov, and N. J. Talbot, "Independent signaling pathways regulate cellular turgor during hyperosmotic stress and appressorium-mediated plant infection by *Magnaporthe grisea*," *Plant Cell*, vol. 11, no. 10, pp. 2045–2058, 1999.
- [144] C. Du, J. Sarfati, J. P. Latge, and R. Calderone, "The role of the sakA (*HOG1*) and tcsB (*SLN1*) genes in the oxidant adaptation of *Aspergillus fumigatus*," *Medical Mycology*, vol. 44, no. 3, pp. 211–218, 2006.
- [145] A. Moriwaki, J. Kihara, C. Mori, and S. Arase, "A MAP kinase gene, *BMK1*, is required for conidiation and pathogenicity in the rice leaf spot pathogen *Bipolaris oryzae*," *Microbiological Research*, vol. 162, no. 2, pp. 108–114, 2007.
- [146] B. A. Morgan, G. R. Banks, W. M. Toone, D. Raitt, S. Kuge, and L. H. Johnston, "The Skn7 response regulator controls gene expression in the oxidative stress response of the budding yeast *Saccharomyces cerevisiae*," *The EMBO Journal*, vol. 16, no. 5, pp. 1035–1044, 1997.
- [147] J. M. Y. Lu, R. J. Deschenes, and J. S. Fassler, "*Saccharomyces cerevisiae* histidine phosphotransferase Ypd1p shuttles between the nucleus and cytoplasm for *SLN1*-dependent phosphorylation of Ssk1p and Skn7p," *Eukaryotic Cell*, vol. 2, no. 6, pp. 1304–1314, 2003.
- [148] J. M. Y. Lu, R. J. Deschenes, and J. S. Fassler, "Role for the ran binding protein, Mog1p, in *Saccharomyces cerevisiae* *SLN1*-*SKN7* signal transduction," *Eukaryotic Cell*, vol. 3, no. 6, pp. 1544–1556, 2004.
- [149] X. J. He and J. S. Fassler, "Identification of novel Yap1p and Skn7p binding sites involved in the oxidative stress response of *Saccharomyces cerevisiae*," *Molecular Microbiology*, vol. 58, no. 5, pp. 1454–1467, 2005.
- [150] F. L. Wormley Jr., G. Heinrich, J. L. Miller, J. R. Perfect, and G. M. Cox, "Identification and characterization of an *SKN7* homologue in *Cryptococcus neoformans*," *Infection and Immunity*, vol. 73, no. 8, pp. 5022–5030, 2005.
- [151] X. J. He, K. E. Mulford, and J. S. Fassler, "Oxidative stress function of the *Saccharomyces cerevisiae* skn7 receiver domain," *Eukaryotic Cell*, vol. 8, no. 5, pp. 768–778, 2009.
- [152] J. S. Fassler and A. H. West, "Fungal Skn7 stress responses and their relationship to virulence," *Eukaryotic Cell*, vol. 10, no. 2, pp. 156–167, 2011.
- [153] J. Lee, C. Godon, G. Lagniel et al., "Yap1 and Skn7 control two specialized oxidative stress response regulons in yeast," *Journal of Biological Chemistry*, vol. 274, no. 23, pp. 16040–16046, 1999.
- [154] A. S. Alberts, N. Bouquin, L. H. Johnston, and R. Treisman, "Analysis of RhoA-binding proteins reveals an interaction domain conserved in heterotrimeric G protein β subunits and the yeast response regulator protein Skn7," *Journal of Biological Chemistry*, vol. 273, no. 15, pp. 8616–8622, 1998.
- [155] N. Bouquin, A. L. Johnson, B. A. Morgan, and L. H. Johnston, "Association of the cell cycle transcription factor Mbp1 with the Skn7 response regulator in budding yeast," *Molecular Biology of the Cell*, vol. 10, no. 10, pp. 3389–3400, 1999.
- [156] D. C. Raitt, A. L. Johnson, A. M. Erkin et al., "The Skn7 response regulator of *Saccharomyces cerevisiae* interacts with Hsf1 in vivo and is required for the induction of heat shock genes by oxidative stress," *Molecular Biology of the Cell*, vol. 11, no. 7, pp. 2335–2347, 2000.
- [157] K. E. Williams and M. S. Cyert, "The eukaryotic response regulator Skn7p regulates calcineurin signaling through stabilization of Crz1p," *The EMBO Journal*, vol. 20, no. 13, pp. 3473–3483, 2001.
- [158] J. L. Brown, S. North, and H. Bussey, "SKN7, a yeast multicopy suppressor of a mutation affecting cell wall β -glucan assembly, encodes a product with domains homologous to prokaryotic two-component regulators and to heat shock transcription factors," *Journal of Bacteriology*, vol. 175, no. 21, pp. 6908–6915, 1993.
- [159] J. L. Brown, H. Bussey, and R. C. Stewart, "Yeast Skn7p functions in a eukaryotic two-component regulatory pathway," *The EMBO Journal*, vol. 13, no. 21, pp. 5186–5194, 1994.
- [160] W. Tao, R. J. Deschenes, and J. S. Fassler, "Intracellular glycerol levels modulate the activity of *SLN1p*, a *Saccharomyces cerevisiae* two-component regulator," *Journal of Biological Chemistry*, vol. 274, no. 1, pp. 360–367, 1999.
- [161] N. Nakamichi, H. Yanada, H. Aiba, K. Aoyama, R. Ohmiya, and T. Mizuno, "Characterization of the Prr1 response regulator with special reference to sexual development in *Schizosaccharomyces pombe*," *Bioscience, Biotechnology and Biochemistry*, vol. 67, no. 3, pp. 547–555, 2003.
- [162] S. Oide, J. Liu, S. H. Yun et al., "Histidine kinase two-component response regulator proteins regulate reproductive development, virulence, and stress responses of the fungal cereal pathogens *Cochliobolus heterostrophus* and *Gibberella zeae*," *Eukaryotic Cell*, vol. 9, no. 12, pp. 1867–1880, 2010.
- [163] D. Hagiwara, T. Mizuno, and K. Abe, "Characterization of the conserved phosphorylation site in the *Aspergillus nidulans* response regulator SrrA," *Current Genetics*, vol. 57, no. 2, pp. 103–114, 2011.
- [164] L.-H. Chen, C.-H. Lin, and K.-R. Chung, "Roles for SKN7 response regulator in stress resistance, conidiation and virulence in the citrus pathogen *Alternaria alternata*," *Fungal Genetics and Biology*, vol. 49, no. 10, pp. 802–813, 2012.
- [165] N. Y. Wang, C. H. Lin, and K. R. Chung, "A $G\alpha$ subunit gene is essential for conidiation and potassium efflux but dispensable for pathogenicity of *Alternaria alternata* on citrus," *Current Genetics*, vol. 56, no. 1, pp. 43–51, 2010.
- [166] J. I. Yago, C. H. Lin, and K. R. Chung, "The SLT2 mitogen-activated protein kinase-mediated signalling pathway governs

- conidiation, morphogenesis, fungal virulence and production of toxin and melanin in the tangerine pathotype of *Alternaria alternata*,” *Molecular Plant Pathology*, vol. 12, no. 7, pp. 653–665, 2011.
- [167] S. L. Yang and K. R. Chung, “The NADPH oxidase-mediated production of H_2O_2 and resistance to oxidative stress in the necrotrophic pathogen *Alternaria alternata* of citrus,” *Molecular Plant Pathology*, vol. 13, pp. 900–914, 2012.
- [168] H.-C. Tsai, S. L. Yang, and K.-R. Chung, “Cyclic AMP-dependent protein kinase A negatively regulates conidia formation by the tangerine pathotype of *Alternaria alternata*,” *World Journal of Microbiology and Biotechnology*. In press.
- [169] K. Izumitsu, A. Yoshimi, and C. Tanaka, “Two-component response regulators Ssk1p and Skn7p additively regulate high-osmolarity adaptation and fungicide sensitivity in *Cochliobolus heterostrophus*,” *Eukaryotic Cell*, vol. 6, no. 2, pp. 171–181, 2007.
- [170] K. Izumitsu, A. Yoshimi, S. Hamada, A. Morita, Y. Saitoh, and C. Tanaka, “*Dic2* and *Dic3* loci confer osmotic adaptation and fungicidal sensitivity independent of the HOG pathway in *Cochliobolus heterostrophus*,” *Mycological Research*, vol. 113, no. 10, pp. 1208–1215, 2009.
- [171] P. Singh, N. Chauhan, A. Ghosh, F. Dixon, and R. Calderone, “*SKN7* of *Candida albicans*: mutant construction and phenotype analysis,” *Infection and Immunity*, vol. 72, no. 4, pp. 2390–2394, 2004.
- [172] T. Saijo, T. Miyazaki, K. Izumikawa et al., “Skn7p is involved in oxidative stress response and virulence of *Candida glabrata*,” *Mycopathologia*, vol. 169, no. 2, pp. 81–90, 2010.
- [173] C. Lamarre, O. Ibrahim-Granet, C. Du, R. Calderone, and J. P. Latgé, “Characterization of the *SKN7* ortholog of *Aspergillus fumigatus*,” *Fungal Genetics and Biology*, vol. 44, no. 7, pp. 682–690, 2007.
- [174] J. D. Lambeth, “NOX enzymes and the biology of reactive oxygen,” *Nature Reviews Immunology*, vol. 4, no. 3, pp. 181–189, 2004.
- [175] R. Fluhr, “Reactive oxygen-generating HADPH oxidases in plants,” in *Reactive Oxygen Species in Plant Signaling, Signaling and Communication in Plants*, L. A. del Río and A. Puppo, Eds., pp. 1–23, Springer, Berlin, Germany, 2009.
- [176] B. T. Kawahara, M. T. Quinn, and J. D. Lambeth, “Molecular evolution of the reactive oxygen-generating NADPH oxidase (Nox/Duox) family of enzymes,” *BMC Evolutionary Biology*, vol. 7, article 109, 2007.
- [177] D. Diekmann, A. Abo, C. Johnston, A. W. Segal, and A. Hall, “Interaction of Rac with $P67^{phox}$ and regulation of phagocytic NADPH oxidase activity,” *Science*, vol. 265, no. 5171, pp. 531–533, 1994.
- [178] Y. A. Suh, R. S. Arnold, B. Lassegue et al., “Cell transformation by the superoxide-generating oxidase Mox1,” *Nature*, vol. 401, no. 6748, pp. 79–82, 1999.
- [179] J. D. Lambeth, G. Cheng, R. S. Arnold, and W. A. Edens, “Novel homologs of $gp91^{phox}$,” *Trends in Biochemical Sciences*, vol. 25, no. 10, pp. 459–461, 2000.
- [180] B. Lardy, M. Bof, L. Aubry et al., “NADPH oxidase homologs are required for normal cell differentiation and morphogenesis in *Dictyostelium discoideum*,” *Biochimica et Biophysica Acta*, vol. 1744, no. 2, pp. 199–212, 2005.
- [181] B. A. Diebold and G. M. Bokoch, “Molecular basis for Rac2 regulation of phagocyte NADPH oxidase,” *Nature Immunology*, vol. 2, no. 3, pp. 211–215, 2001.
- [182] M. A. Torres, J. L. Dangel, and J. D. G. Jones, “*Arabidopsis gp91^{phox}* homologues *AtrbohD* and *AtrbohF* are required for accumulation of reactive oxygen intermediates in the plant defense response,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 99, no. 1, pp. 517–522, 2002.
- [183] H. Yoshioka, N. Numata, K. Nakajima et al., “*Nicotiana benthamiana gp91^{phox}* homologs *NbrbohA* and *NbrbohB* participate in H_2O_2 accumulation and resistance to *Phytophthora infestans*,” *Plant Cell*, vol. 15, no. 3, pp. 706–718, 2003.
- [184] H. Lalucque and P. Silar, “NADPH oxidase: an enzyme for multicellularity?” *Trends in Microbiology*, vol. 11, no. 1, pp. 9–12, 2003.
- [185] B. Scott and C. J. Eaton, “Role of reactive oxygen species in fungal cellular differentiations,” *Current Opinion in Microbiology*, vol. 11, no. 6, pp. 488–493, 2008.
- [186] J. Heller and P. Tudzynski, “Reactive oxygen species in phytopathogenic fungi: signaling, development, and disease,” *Annual Review of Phytopathology*, vol. 49, pp. 369–390, 2011.
- [187] H.-J. Kim, C. Chen, M. Kabbage, and M. B. Dickman, “Identification and characterization of *Sclerotinia sclerotiorum* NADPH oxidases,” *Applied and Environmental Microbiology*, vol. 77, no. 21, pp. 7721–7729, 2011.
- [188] M. J. Egan, Z. Y. Wang, M. A. Jones, N. Smirnov, and N. J. Talbot, “Generation of reactive oxygen species by fungal NADPH oxidases is required for rice blast disease,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 104, no. 28, pp. 11772–11777, 2007.
- [189] T. Lara-Ortiz, H. Riveros-Rosas, and J. Aguirre, “Reactive oxygen species generated by microbial NADPH oxidase NoxA regulate sexual development in *Aspergillus nidulans*,” *Molecular Microbiology*, vol. 50, no. 4, pp. 1241–1255, 2003.
- [190] F. Malagnac, H. Lalucque, G. Lepère, and P. Silar, “Two NADPH oxidase isoforms are required for sexual reproduction and ascospore germination in the filamentous fungus *Podospora anserina*,” *Fungal Genetics and Biology*, vol. 41, no. 11, pp. 982–997, 2004.
- [191] N. Cano-Domínguez, K. Álvarez-Delfín, W. Hansberg, and J. Aguirre, “NADPH oxidases NOX-1 and NOX-2 require the regulatory subunit NOR-1 to control cell differentiation and growth in *Neurospora crassa*,” *Eukaryotic Cell*, vol. 7, no. 8, pp. 1352–1361, 2008.
- [192] D. Takemoto, A. Tanaka, and B. Scott, “A $P67^{phox}$ -like regulator is recruited to control hyphal branching in a fungal-grass mutualistic symbiosis,” *Plant Cell*, vol. 18, no. 10, pp. 2807–2821, 2006.
- [193] A. Tanaka, M. J. Christensen, D. Takemoto, P. Park, and B. Scott, “Reactive oxygen species play a role in regulating a fungus-perennial ryegrass mutualistic interaction,” *Plant Cell*, vol. 18, no. 4, pp. 1052–1066, 2006.
- [194] A. Tanaka, D. Takemoto, G. S. Hyon, P. Park, and B. Scott, “NoxA activation by the small GTPase RacA is required to maintain a mutualistic symbiotic association between *Epichloë festucae* and perennial ryegrass,” *Molecular Microbiology*, vol. 68, no. 5, pp. 1165–1178, 2008.
- [195] D. Takemoto, S. Kamakura, S. Saikia et al., “Polarity proteins Bem1 and Cdc24 are components of the filamentous fungal NADPH oxidase complex,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 108, no. 7, pp. 2861–2866, 2011.
- [196] S. Giesbert, T. Schürg, S. Scheele, and P. Tudzynski, “The NADPH oxidase Cpnox1 is required for full pathogenicity of

- the ergot fungus *Claviceps purpurea*,” *Molecular Plant Pathology*, vol. 9, no. 3, pp. 317–327, 2008.
- [197] H. Haas, “Molecular genetics of fungal siderophore biosynthesis and uptake: the role of siderophores in iron uptake and storage,” *Applied Microbiology and Biotechnology*, vol. 62, no. 4, pp. 316–330, 2003.
- [198] H. Haas, M. Eisendle, and B. G. Turgeon, “Siderophores in fungal physiology and virulence,” *Annual Review of Phytopathology*, vol. 46, pp. 149–187, 2008.
- [199] C. D. Kaplan and J. Kaplan, “Iron acquisition and transcriptional regulation,” *Chemical Reviews*, vol. 109, no. 10, pp. 4536–4552, 2009.
- [200] J. C. Renshaw, G. D. Robson, A. P. J. Trinci et al., “Fungal siderophores: structures, functions and applications,” *Mycological Research*, vol. 106, no. 10, pp. 1123–1142, 2002.
- [201] A. Stintzi and K. N. Raymond, “Siderophore chemistry,” in *Molecular and Cellular Iron Transport*, D. M. Templeton, Ed., pp. 273–319, Marcel Dekker, New York, NY, USA, 2002.
- [202] H. Kleinkauf and H. von Döhren, “A nonribosomal system of peptide biosynthesis,” *European Journal of Biochemistry*, vol. 236, no. 2, pp. 335–351, 1996.
- [203] D. G. Panaccione, J. S. Scott-Craig, J. A. Pocard, and J. D. Walton, “A cyclic peptide synthetase gene required for pathogenicity of the fungus *Cochliobolus carbonum* on maize,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 89, no. 14, pp. 6590–6594, 1992.
- [204] A. Haese, M. Schubert, M. Herrmann, and R. Zocher, “Molecular characterization of the enniatin synthetase gene encoding a multifunctional enzyme catalysing *N*-methyldepsipeptide formation in *Fusarium scirpi*,” *Molecular Microbiology*, vol. 7, no. 6, pp. 905–914, 1993.
- [205] M. A. F. Jalal, S. K. Love, and D. van der Helm, “ $N\alpha$ -Dimethylcoprogens Three novel trihydroxamate siderophores from pathogenic fungi,” *Biology of Metals*, vol. 1, no. 1, pp. 4–8, 1988.
- [206] M. A. F. Jalal and D. van der Helm, “Siderophores of highly phytopathogenic *Alternaria longipes*: structures of hydroxycoprogens,” *Biology of Metals*, vol. 2, no. 1, pp. 11–17, 1989.
- [207] J. Ohra, K. Morita, Y. Tsujino et al., “Production of two phytotoxic metabolites by the fungus *Alternaria cassia*,” *Bioscience, Biotechnology, and Biochemistry*, vol. 59, pp. 1782–1783, 1995.
- [208] S. Oide, W. Moeder, S. Krasnoff et al., “NPS6, encoding a nonribosomal peptide synthetase involved in siderophore-mediated iron metabolism, is a conserved virulence determinant of plant pathogenic ascomycetes,” *Plant Cell*, vol. 18, no. 10, pp. 2836–2853, 2006.
- [209] H. von Döhren, “Biochemistry and general genetics of nonribosomal peptide synthetases in fungi,” *Advances in Biochemical Engineering/Biotechnology*, vol. 88, pp. 217–264, 2004.
- [210] B. N. Lee, S. Kroken, D. Y. T. Chou, B. Robbertse, O. C. Yoder, and B. G. Turgeon, “Functional analysis of all nonribosomal peptide synthetases in *Cochliobolus heterostrophus* reveals a factor, NPS6, involved in virulence and resistance to oxidative stress,” *Eukaryotic Cell*, vol. 4, no. 3, pp. 545–555, 2005.
- [211] K. E. Bushley, D. R. Ripoll, and B. G. Turgeon, “Module evolution and substrate specificity of fungal nonribosomal peptide synthetases involved in siderophore biosynthesis,” *BMC Evolutionary Biology*, vol. 8, no. 1, article 328, 2008.
- [212] K. E. Bushley and B. G. Turgeon, “Phylogenomics reveals subfamilies of fungal nonribosomal peptide synthetases and their evolutionary relationships,” *BMC Evolutionary Biology*, vol. 10, no. 1, article 26, 2010.
- [213] A. H. T. Hissen, A. N. C. Wan, M. L. Warwas, L. J. Pinto, and M. M. Moore, “The *Aspergillus fumigatus* siderophore biosynthetic gene *sidA*, encoding L-ornithine N^5 -oxygenase, is required for virulence,” *Infection and Immunity*, vol. 73, no. 9, pp. 5493–5503, 2005.
- [214] M. Schrettl, E. Bignell, C. Kragl et al., “Distinct roles for intra- and extracellular siderophores during *Aspergillus fumigatus* infection,” *PLoS Pathogens*, vol. 3, no. 9, article e128, 2007.
- [215] B. Mei, A. D. Budde, and S. A. Leong, “*sid1*, a gene initiating siderophore biosynthesis in *Ustilago maydis*: molecular characterization, regulation by iron, and role in phytopathogenicity,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 90, no. 3, pp. 903–907, 1993.