Signal Transduction of Sphingosine-1-Phosphate G Protein–Coupled Receptors

Nicholas Young¹ and James R. Van Brocklyn^{2,*}

¹Integrated Biomedical Science Graduate Program and ²Department of Pathology,
The Ohio State University, Columbus

E-mail: young.651@osu.edu; james.vanbrocklyn@osumc.edu

Received June 14, 2006; Accepted July 31, 2006; Published August 11, 2006

Sphingosine-1-phosphate (S1P) is a bioactive lipid capable of eliciting dramatic effects in a variety of cell types. Signaling by this molecule is by a family of five G protein–coupled receptors named S1P_{1–5} that signal through a variety of pathways to regulate cell proliferation, migration, cytoskeletal organization, and differentiation. These receptors are expressed in a wide variety of tissues and cell types, and their cellular effects contribute to important biological and pathological functions of S1P in many processes, including angiogenesis, vascular development, lymphocyte trafficking, and cancer. This review will focus on the current progress in the field of S1P receptor signaling and biology.

KEYWORDS: sphingosine-1-phosphate, S1P, EDG, cell proliferation, cell migration, angiogenesis, lymphocyte trafficking, cancer

INTRODUCTION

The bioactive lipid sphingosine-1-phosphate (S1P) is an agonist for a subset of G protein–coupled receptors (GPCRs) known as the endothelial differentiation gene (EDG) family. S1P is present at high concentrations in the blood[1], roughly 0.1–1 μ M, where it is bound to albumin and to lipoproteins[2]. It is secreted by platelets on activation[3] and by mast cells[4], as well as by a variety of other cell types. It is generally thought to be ubiquitously present at low levels outside of the bloodstream within tissues.

S1P is formed by phosphorylation of sphingosine, the long-chain base that serves as the backbone of sphingolipids, by the ubiquitously expressed enzyme sphingosine kinase (SphK). SphK displays basal activity in unstimulated cells, however, it also can be activated by a variety of signals, including growth factors[5,6,7,8], immunoglobulin receptors[9,10], tumor necrosis factor[11], and various GPCRs[12,13]. As SphK is a cytoplasmic-signaling enzyme, it was originally thought that S1P functioned within cells as a second messenger. Interestingly, although S1P is now known to function at the cell surface by activating several GPCRs, significant evidence exists that S1P also has receptor-independent, intracellular functions[14], although definitive proof of this awaits the identification of a direct intracellular target.

Since the discovery of the S1P family of receptors in the late 1990s, the modulation of cellular and biological functions by S1P receptor stimulation has been under intense investigation. Five related GPCRs that specifically bind S1P with high affinity have been identified, including S1P₁/EDG-1/LP_{B1}, S1P₂/EDG-5/H218/AGR16/LP_{B2}, S1P₃/EDG-3/LP_{B3}, S1P₄/EDG-6/LP_{B4}, and S1P₅/EDG-8/NGR-1. Although a variety

*Corresponding author.

©2006 with author.

Published by TheScientificWorld, Ltd.; www.thescientificworld.com

of naming systems for these receptors have been used, the currently accepted system, based on International Union of Pharmacology (IUPHAR) guidelines, is the S1P nomenclature[15] and, therefore, that system will be used in this review, regardless of which names were used in the original papers. Each of the five S1P receptors has significant homology displaying 40–50% amino acid identity and approximately 60% amino acid similarity[16,17,18]. Several other orphan receptors have been implicated as potential S1P receptors, including GPR3, 6, 12, and 63, although these remain unconfirmed and little is known regarding the significance of these receptors (reviewed in [19]).

S1P₁, S1P₂, and S1P₃ were originally orphan receptors. The S1P₁ receptor subtype was the first of the family to be cloned and was identified as an immediate early transcript in phorbol ester-induced endothelial cell differentiation[20], leading to the designation endothelial differentiation gene (EDG)-1. S1P₂ was cloned from rat brain and rat vascular smooth muscle cells(VSMC)[21,22], while S1P₃ was cloned from a human genomic library[23]. The identification of a related receptor, EDG-2/vzg-1, as a receptor for lysophosphatidic acid (LPA)[24], another bloodborne lipid that is structurally similar to S1P and has several overlapping cellular effects, suggested that S1P may be a ligand for other EDG family receptors. This led to the identification of S1P₁ as a specific S1P receptor[25]. Shortly after the identification of EDG-2/LPA₁ as the first LPA receptor, S1P₂ and S1P₃ were shown to mediate S1P-induced transcription from a serum response, element-driven promoter[26]. S1P₄ was later cloned from *in vitro* differentiated murine and human dendritic cells by a PCR-based strategy using degenerate primers based on other EDG receptors[27]. S1P₅ was identified as a nerve growth factor down-regulated gene based on EST profiling and cloned from a rat PC12 cell cDNA library[28].

S1P regulates a wide range of cellular effects and biological responses. S1P stimulates proliferation of several cell types, although it can inhibit proliferation in some cases. S1P affects cell migration, again both positively and negatively depending on the receptors expressed. S1P can also block apoptosis induced by a variety of stimuli. S1P can affect cellular differentiation and is important in some aspects of development. S1P has also recently been shown to be a critical regulator of immune cell trafficking and an important regulator of tumor angiogenesis. These responses have been linked to the various GPCRs that S1P regulates and will be discussed in detail below.

Each S1P receptor subtype signals through a unique set of G proteins that result in various downstream cellular effects. Therefore, it is critical to know what receptors are expressed in order to predict cellular behavior on S1P stimulation. Of the four major families of heterotrimeric G proteins, G_s, G_{i/o}, G_{q/11}, and G_{12/13} S1P receptors couple to G_i, G_q, and G_{12/13}[29]. Thus, S1P leads to activation of the classical pathways downstream of these G proteins, adenylyl cyclase inhibition as well as the Ras/MAP kinase cascade and phosphatidylinositol 3-phosphate kinase (PI3 kinase)/Akt for G_i, phospholipase C (PLC) for G_q, and small Rho GTPases for G_{12/13}. The particular pathways activated by various receptors and the biology that they regulate will be discussed.

LIGAND AFFINITIES AND SPECIFICITIES OF S1P RECEPTORS

All five S1P receptors bind S1P with nanomolar affinities although precise values vary somewhat between experiments. S1P₁ has been shown to bind S1P with K_Ds ranging from 8–50 nM[25,30,31,32,33]. The mean value from the affinity measurements for S1P₁ reported in these papers is 24.7 ± 17.8 nM. Similar values are obtained for S1P₂ (18.4 ± 7.2 nM) and S1P₃ (19.7 ± 8.6 nM)[25,30,31,32,33,34]. S1P₅ appears to bind S1P with slightly higher affinity, as this has been reported to be 2[17] and 6 nM[35]. Reports of S1P₄ binding affinity for S1P have varied more widely, including 12.8[18], 63[36], 95, and even 156 nM[37]. Notably, all of these studies except the Yamazaki et al. paper found that S1P₄ has the lowest binding affinity for S1P of the five S1P receptors. Interestingly, S1P₄ has a much higher affinity (approximately 2 nM) for phytosphingosine-1-phosphate (phS1P), which has a 4-hydroxyl group and is commonly found in plants[37]. As this is a minor lipid in mammalian cells, the biological significance of this finding is uncertain. Interestingly, one study showed significantly higher-affinity binding of S1P to membranes from S1P₁ S1P₂, S1P₃, and S1P₅ overexpressing cells, all ranging from 0.17–0.61 nM[38].

The reason for this discrepancy is unclear. Nevertheless, as concentrations of S1P in plasma and serum range from approximately 100 nM to 1 μ M[1,39], it is apparent that all the above numbers represent physiologically relevant affinities.

Several other sphingolipids, related in structure to S1P, have been tested as potential ligands for S1P receptors. The majority of these lipids have shown no binding or agonist properties for these receptors. The exceptions are dihydrosphingosine-1-phosphate (dhS1P), which lacks the trans-double bond at the 4 position, and sphingosylphosphorylcholine (SPC), which is similar to S1P, but contains a choline head group, i.e., lysosphingomyelin. dhS1P is an agonist for all five S1P receptors. For S1P₁ and S1P₃, dhS1P binds and activates the receptors with a similar affinity to S1P[33]. Interestingly, at S1P₂, dhS1P is less potent, competing for S1P binding with an IC₅₀ of approximately 200 nM[33]. One study showed an IC₅₀ for dhS1P for S1P₄ of approximately 9 nM[37], significantly higher than its affinity for S1P. Similar numbers were obtained in another study that also showed that dhS1P bound S1P₄ with an affinity similar to that of phS1P, approximately 4 nM, while S1P bound with 10- to 20-fold lower affinity[40]. SPC also functions as an agonist for the five S1P receptors, however, for all five, it is a low-affinity agonist with IC₅₀s for binding and EC₅₀s for various signaling responses generally occurring in the low micromolar range[17,30,36,41,42,43].

G PROTEIN COUPLING OF S1P RECEPTORS AND PATHWAYS ACTIVATED

Prior to the identification of S1P as its ligand, S1P₁ was shown to couple to G_i and the extracellular signal regulated kinase (ERK) mitogen-activated protein (MAP) kinase pathway[44]. GTP γ S binding studies showed that S1P₁ couples exclusively to G_i proteins[29] and, thus, responses mediated through this receptor are sensitive to pertussis toxin (PTX), which ADP-ribosylates and inactivates G_{i/o} proteins. S1P₁ activation by S1P has been shown to lead to the activation of several signal transduction pathways (including ERK[25,45]), inhibition of forskolin-stimulated cAMP accumulation[46], and increased [Ca²⁺]_i in CHO cells[30], although the calcium response was not detected in HEK293 cells overexpressing S1P₁[46] and in *Xenopus* oocytes, S1P₁ required coexpression of a chimeric G $\alpha_{q/i}$ protein, allowing S1P₁ to couple to phospholipase C in order to affect [Ca²⁺]_i[43]. The increased [Ca²⁺]_i seen with S1P₁ signaling in some cell types may be mediated by G_i-induced activation of SphK, leading to intracellular production of S1P- and IP₃-independent calcium release from intracellular stores[47]. On the other hand, S1P₁ signaling can inhibit calcium responses induced by various other signals[48].

S1P₁ activates PI3 kinase, leading to Akt activation and endothelial nitric oxide synthase (eNOS) phosphorylation in endothelial cells[49]. The activation of this pathway may be affected by the subcellular localization of S1P₁. In this regard, S1P stimulation of S1P₁ causes its accumulation in caveolae[50]. Interaction of S1P₁ with caveolin leads to a decrease in its ability to activate eNOS[50] as well as Akt and the small GTPase Rac. Interestingly, caveolin interaction does not affect S1P₁-induced ERK activation[51]. Caveolar localization of S1P₁ requires N-glycosylation, as does receptor internalization[52].

Several studies have also shown interesting cooperative interactions of S1P₁ and platelet-derived growth factor (PDGF) signaling. Spiegel's group demonstrated that S1P₁ expression was necessary for PDGF-induced Rac activation and chemotaxis in HEK293 cells and mouse embryonic fibroblasts (MEFs)[53]. Since PDGF activates SphK, and led to S1P₁ phosphorylation, they suggested an autocrine model in which PDGF induces production of S1P that in turn activates S1P₁ leading to Rac activation and cell migration. The same group also showed that PDGF required S1P₁ to induce membrane ruffles and that PDGF causes SphK to translocate to membrane ruffles, suggesting that local S1P formation at the leading edge might be involved in directed cell migration[54]. S1P₁ was also involved in PDGF-induced FAK, src, and p38 activation[54]. In addition, Pyne's group has shown that the PDGF receptor and S1P₁ are physically linked. In their "integrative" model, the PDGF receptor phosphorylates G α_{12} , which is recruited by S1P₁, and this step enhances signaling by both S1P₁ and the PDGF receptor[55,56]. The same group further found that PDGF-induced cell migration was blocked by an inverse agonist of

S1P₁[57]. It is unclear which of the above models is more correct, nor are the two ideas necessarily mutually exclusive. However, it is possible that this system may function differently in different cell types. In this regard, S1P₁ was found not to be required for PDGF-induced chemotaxis of rat VSMC and MEFs from S1P₁ knockout embryos[58]. To further add to the complexity of the S1P/PDGF interplay, S1P₁ signaling was recently found to induce expression of PDGF-A and PDGF-B[59], and S1P has been shown to transactivate epidermal growth factor receptor (EGFR) and PDGFR in VSMC[60] although the S1P receptor involved was not identified.

S1P₂ and S1P₃ both couple to G_i, G_q, and G₁₃[29], however, it appears that they do so with different preferences, leading to different biological responses. As both of these receptors link to G_i activation, they activate ERK MAP kinase in a PTX-sensitive fashion, as does S1P₁. Interestingly, while the potency of ERK activation by S1P₃ is similar to that of S1P₁, it is higher than that mediated by S1P₂[41]. Both are also capable of activating PLC to induce IP₃ formation and Ca²⁺ mobilization in a partially PTX-sensitive fashion[31,41], suggesting involvement of both G_i and G_q. However, S1P₃ more potently stimulates inositol phosphate production and increased [Ca²⁺]_i[31]. On the other hand, S1P₂ couples strongly to activation of the small GTPase Rho and stress fiber formation[42]. S1P₂ also is a potent inducer of cell rounding and process retraction in PC12 cells[34], a process linked to Rho activation[61]. Results using cells from S1P₂ and S1P₃ knockout mice support this distinction. Analysis of S1P signaling in MEFs from S1P₂^{-/-} embryos revealed a defect in Rho activation, but no change in calcium mobilization, adenylyl cyclase inhibition, or inositol phosphate production[62]. In contrast, MEFs from S1P₃^{-/-} embryos displayed a loss of PLC activation and calcium mobilization, but with only a minor effect on Rho activation[62]. Thus, it appears that S1P₂ couples preferentially to G_{12/13}, leading to potent Rho activation, while S1P₃ prefers G_q, thus favoring PLC activation. Nevertheless, S1P₂ couples to G_{αq}, G_{α13}, and G_{αi2} in smooth muscle cells and leads to PLC as well as Rho activation[63]. Thus, the extent of coupling to various G protein and signaling pathways activated by the various receptors will likely vary in different cellular contexts.

S1P₂ has also been shown to activate p38 and Jun N-terminal kinase (JNK) MAP kinases[42] and phospholipase D activation[32], all by PTX-insensitive mechanisms. Both S1P₂ and S1P₃ also activate NF-κB, probably via G_q[64]. As expected for G_i-coupled receptors, S1P₂ and S1P₃ inhibit forskolin-induced cAMP accumulation[31]. Interestingly, however, in the absence of forskolin, signaling by these receptors leads to increased cAMP[31]. S1P₁ causes this effect as well, although it is much weaker in this capacity.

Recently, S1P₃ has also been shown to participate in cross-talk with a receptor tyrosine kinase. S1P₃ causes PDGFR transactivation, leading to Akt activation[65]. S1P₃ also causes transactivation of EGFR via a Src family kinase and matrix metalloproteinase-dependent release of heparin-binding EGF-like growth factor (HB-EGF)[66]. Interestingly, via this pathway, S1P₃ mediates EGFR transactivation by estrogen receptors that activate SphK, leading to S1P secretion[66].

S1P₄ couples to both G_i and G_{12/13}, but not G_q[67]. This specific G protein coupling has distinct downstream signaling effects that influence a variety of cellular activities. S1P₄ can lead to activation of PLC, ERK, and the Rho family small GTPase Cdc42 through G_i coupling[18,36,68]. Cdc42 activation by S1P₄ positively regulates cellular motility[68]. Coupling of S1P₄ to G_{12/13} leads to Rho activation, formation of peripheral stress fibers, and cell rounding[67].

S1P₅ couples to both G_i and G₁₂, but not G_q[35]. G_i signaling leads to inhibition of cAMP accumulation in both CHO and RH7777 cells in a PTX-sensitive manner[17,35]. Surprisingly, however, S1P₅ is unusual compared to other S1P receptors as it causes a decrease in ERK1/2 phosphorylation and activity[35]. Repression of ERK by S1P₅ is PTX insensitive, but is prevented by the phosphatase inhibitors orthovanadate and okadaic acid, indicating a role for a phosphatase[35]. Furthermore, S1P₅ activates JNK MAP kinase, but not p38[35].

REGULATION OF CELL PROLIFERATION AND SURVIVAL

The first biological effect defined for S1P, in the early 1990s, was mitogenesis[69]. PTX sensitivity of the mitogenic effect of S1P on fibroblasts suggested that GPCRs were involved in this response[70]. Nevertheless, it was, and still is, unclear to what degree receptors vs. intracellular receptor-independent effects of S1P contribute to mitogenesis. However, significant evidence exists that S1P receptors do, in some cases, stimulate cell proliferation.

S1P has been shown to stimulate proliferation of a variety of cell types, including fibroblasts[69], VSMC[71], osteoblasts[72], breast cancer cells[73], hepatic stellate cells[74], endothelial cells[75], airway smooth muscle[76], astrocytes[77], mesangial cells[78], intestinal epithelial cells[79], glioma cells[80], neural progenitor cells[81], and chondrocytes[82].

Multiple S1P receptors and signaling pathways can contribute to the mitogenic effect of S1P. S1P-induced proliferation of VSMC was augmented by overexpression of S1P₁ through activation of p70 S6 kinase and cyclin D1 expression[71]. S1P stimulated proliferation of glioma cells that express S1P₁, S1P₂, and S1P₃ through ERK and PI3-kinase β activation[80]. S1P-induced proliferation of breast cancer cells, expressing S1P₂ and S1P₃, but not S1P₁, was mediated by direct activation of serum-response element signaling and by induction of IGF-II expression[73]. Furthermore, overexpression of S1P₂ or S1P₃ in hepatoma cells allowed S1P to stimulate proliferation[83].

S1P can also exert antiproliferative effects in several cell types. S1P has an antiproliferative effect in hepatocytes[84]. S1P also blocks insulin and TSH-stimulated proliferation of thyroid cells[85]. In hepatic myofibroblasts, S1P blocks proliferation through induction of cyclooxygenase-2 (COX-2) and production of prostaglandin E₂ and cAMP[86]. Although the receptor(s) involved were not identified, this effect was PTX insensitive. S1P also suppresses proliferation of T lymphocytes[87,88], keratinocytes[89], and gastric cancer cells[90].

S1P₂ may be a major regulator of the antiproliferative effects of S1P, as S1P₂^{-/-} MEFs proliferate more rapidly than wild type MEFs[91]. Restoration of S1P₂ expression decreases proliferation[91]. Furthermore, S1P inhibition of hepatocyte proliferation was mediated by S1P₂-induced Rho activation[84]. On the other hand, suppression of T cell proliferation by S1P is mediated by S1P₁[87]. In addition, S1P₅ overexpression inhibits proliferation of CHO cells[35]. Thus, as with positive proliferation responses, antiproliferative responses to S1P can be mediated by various receptors. Moreover, in some cases, the same S1P receptors can stimulate proliferation in some cell types, while inhibiting it in others.

S1P inhibits apoptosis induced by a variety of signals and insults[13]. Inhibition of apoptosis by S1P may also be mediated by several pathways, however, the PI3 kinase/Akt pathway appears to be particularly important. As S1P₁ has most commonly been linked with this pathway, this may be the primary receptor involved in the survival response, although S1P₂ and S1P₃ have also been implicated in some cases. For example, S1P-induced Akt activation promotes survival of ovarian cancer cells[92]. Protection of endothelial cells from apoptosis by S1P is mediated by S1P₁ and S1P₃ signaling through Akt to activate eNOS[93]. Although S1P blocks proliferation of melanocytes, it enhances cell survival through activation of ERK and Akt[94]. Interestingly, S1P can activate both pro- and antiapoptotic effects in the same cell type. In this regard, it was found that S1P stimulates apoptosis of hepatic myofibroblasts by a receptor-independent mechanism, while activating PI3 kinase to inhibit apoptotic signaling[95]. S1P also enhances neutrophil survival through the p38 pathway[96]. In addition, protection of T lymphoblastoma cells from apoptosis by S1P appears to involve S1P₂ and S1P₃ signaling and decreased expression of the proapoptotic protein Bax[97].

REGULATION OF CELL MIGRATION

A great deal of work has been done in the analysis of S1P signaling in the cellular migratory response. Early studies showed that S1P inhibited migration of several cell types including melanoma, fibrosarcoma[98], and breast cancer cells[99]. In contrast, S1P potently stimulates motility of endothelial

cells[100,101,102], some glioma cells[103] and, at low concentrations, T cells[104]. In some cases, S1P enhances VSMC migration[105,106] while inhibiting movement of these cells in others[107].

The differential influence of S1P on cell migration is dependent on the receptor subtypes expressed by the cell. Overexpression of S1P₁ or S1P₃ in CHO cells allowed for S1P-induced cell migration, while S1P₂ overexpression had no effect[31]. Moreover, S1P₂ overexpression caused inhibition of Rac activation, membrane ruffling, and cell migration[108]. Furthermore, S1P₂ overexpression decreases stimulation of endothelial cell migration, while S1P₁ overexpression in smooth muscle cells reduces inhibition of migration[107] or, in some cases, allows S1P-stimulation of migration[71]. The inhibitory regulation of Rac by S1P₂ was shown to be downstream of Rho, which is potently activated by S1P₂ through G_{12/13}[109]. Inhibition of G_i signaling by S1P₃ using PTX allowed S1P to inhibit migration through this receptor as well. Thus, the overall picture is that S1P₂ preferentially couples to G_{12/13} to inhibit Rac and cell migration, while only weakly activating G_i, and S1P₃ activates G_i and G_q more potently, thus stimulating migration. S1P₁, coupling only to G_i, enhances cell migration.

In contrast, a more recent study has shown that S1P stimulation of S1P₂ in glioma cells actually increases Rac activation and that the decrease in observed migration in these cells is due to Rho signaling through Rho-associated protein kinase (ROCK), as ROCK inhibition reversed the effect[110]. Additionally, this study showed stress fiber formation by S1P₂ signaling and cell rounding on overexpression of S1P₂. Thus, the S1P₂ receptor consistently opposes migration, but does so through different signaling mechanisms, i.e., inhibition of Rac in some cell types and ROCK-mediated stress fiber formation in others. Through Rho signaling, S1P₂ can also activate PTEN, a phosphatase acting on 3-phosphorylated phosphoinositides and thus counteracting PI3 kinase signaling, to inhibit cell migration[111].

S1P₄ may also be able to stimulate cell migration. Overexpression of S1P₄ allowed S1P to stimulate migration of CHO cells through activation of another small GTPase, Cdc42[68]. Also, mutation of a zebrafish S1P receptor Miles Apart (Mil) prevented migration of heart precursors to the midline[112]. Interestingly, Mil expression was not required in the heart precursors themselves, suggesting that Mil established a permissive environment for heart precursor migration.

S1P RECEPTOR EXPRESSION

S1P₁, S1P₂, and S1P₃ are widely expressed in most tissues at various levels[16], suggesting that these receptors influence many cellular processes. S1P₁ is expressed at high levels in brain, heart, lung, liver, and spleen; moderate levels in kidney, muscle, and thymus; and at low levels in several other tissues[16,113]. On a cellular level, S1P₁ is expressed in endothelial cells of most tissues, although not in spleen, kidney, and testis[113]. Expression was also detected in pulmonary smooth muscle, cardiomyocytes, hepatocytes, neurons, especially Purkinje cells, astrocytes, spermatids, spleen marginal zone, and the collecting ducts of the kidney[113]. Mouse splenic T cells also express S1P₁[104]. During embryogenesis, S1P₁ was induced late in development (after embryonic day 15.5) at centers of ossification[114]. S1P₁ is also expressed in areas of neurogenesis adjacent to ventricles and in areas of angiogenesis in developing brain[115].

S1P₂ is also ubiquitously expressed in adult mice, but is present at the highest levels in heart and lung. In embryonic rat, S1P₂ was expressed in lung, kidney, skin, gut, spleen, brain, and liver[22], however, in E14–18 mouse embryos, S1P₂ was only detected in the brain[16]. Brain expression decreased significantly postnatally[22]. S1P₂ is expressed in young neuronal cell bodies and later is found in growing axons[116]. In addition, NGF treatment of PC12 cells led to decreased S1P₂ expression[22].

S1P₃ is expressed at high levels in adult mouse heart, lung, kidney, and spleen with lower levels seen in most other tissues[16]. Northern analysis of human RNA confirmed high S1P₃ expression in heart and kidney, but showed lower expression in lung[23]. Expression was also high in human liver, pancreas, and placenta[23], while liver expression was low in the mouse[16].

In contrast to S1P₁₋₃, S1P₄ and S1P₅ are more restricted in their expression. S1P₄ expression is limited principally to lymphoid tissue and hematopoietic cells, and lung in both adults and embryos[27]. With the exception of S1P₄, all S1P receptor subtypes have been observed to be expressed in the central nervous system (CNS). This appears to be the predominant location of S1P₅ expression. Northern analysis further revealed expression of S1P₅ prominently throughout the rat brain[17,28]. More specifically, this receptor subtype was strongly detected by *in situ* hybridization in the white matter tracts throughout the CNS[17]. In postnatal day 21, mouse brain S1P₅ expression was detected exclusively in oligodendrocyte precursors and mature oligodendrocytes, although in the mature cells S1P₅ was associated with the myelin sheath and not the cell body[117]. Radial glial cells in human fetal brain also express S1P₅[118]. In addition, S1P₅ expression has also been detected in the spleen[17] and skin[119].

Clearly, many tissues and individual cell types coexpress several S1P receptor subtypes. In addition, as noted above, the various receptors affect overlapping, but not identical, sets of signaling pathways. As a result, it is likely that many of the biological responses to S1P are mediated by the coordinated effects of multiple S1P receptor subtypes. In addition, S1P receptors heterodimerize among the various receptor subtypes[120], suggesting possible functional interactive effects of coexpressed receptors.

BIOLOGICAL ROLES OF S1P RECEPTORS DETERMINED FROM KNOCKOUT MICE

Results from S1P₁ knockout mice revealed an important role for this receptor in embryonic development. S1P₁^{-/-} embryos died between E12.5 and E14.5[121]. The embryonic lethality was attributed to massive hemorrhage due to failure of VSMC to cover the dorsal surface of blood vessels. Although this highlights a clear role for S1P₁ in vascular maturation, the endothelium formed a normal vascular network as shown by immunohistochemical staining for endothelial cell markers. Therefore, normal vasculogenesis and angiogenesis apparently occurred. Nevertheless, a conditional mutant, in which S1P₁ was knocked out only in endothelial cells, displayed the same phenotype, indicating that although the defect lies in smooth muscle migration to cover blood vessels, S1P₁ expression is required in the endothelial cells[122]. S1P₁ expression in endothelial cells is also necessary for proper limb development and digit formation[123]. Thus S1P₁ expression in endothelial cells may be important for creating a permissive environment for several developmental processes. Furthermore, S1P₁^{-/-} embryos displayed a significant loss of cells in the developing brain due to increased apoptosis and decreased mitosis, indicating an important role for S1P₁ in neurogenesis[124].

S1P₂ knockout mice have been reported from two independent investigations. The S1P₂^{-/-} mice had a much milder phenotype than the S1P₁ knockout in both instances, being born apparently normal, which suggests a possible redundant signaling role with other S1P receptors. Modest decreases in litter size were observed in one study in otherwise normal mice[62]. In contrast, another group detected seizures due to increased excitability of neocortical pyramidal neurons, which were occasionally fatal[125]. These receptor null mice were normal in all other phenotypic areas investigated.

S1P₃ receptor knockout mice are also born apparently normal[119]. The litter sizes from homozygous S1P₃^{-/-} mice crosses were slightly smaller than normal. S1P₃ deletion led to a compensatory increase in expression of S1P₂ in the heart and brain[119].

The double knockout mouse for S1P₂ and S1P₃ showed a dramatically reduced litter size and marked perinatal lethality, although surviving double knockout mice appeared normal[62]. The reduced litter size in S1P₂/S1P₃ double knockout mice appears to be the result of similar vascular defects in many embryos due to endothelial cells with abnormally thin cell bodies leading to bleeding[126]. On a cellular level, double knockout MEFs displayed a complete lack of Rho activation and severely reduced PLC activation with no loss of adenylyl cyclase inhibition[62]. S1P₁/S1P₂ double knockouts show a more severe phenotype than S1P₁ knockout alone, dying approximately 2 days sooner, with S1P₁/S1P₂/S1P₃ triple knockouts slightly more severe[126]. This was due to a less-well-formed vasculature containing fewer

capillary branches in the head. Thus, it seems that S1P₁, S1P₂ and S1P₃ cooperate in the process of vascular formation.

S1P₄ knockout mice have not yet been reported, but an S1P₅ receptor knockout mouse has been reported[117]. Although S1P₅ expression was found in both oligodendrocyte precursors and mature oligodendrocytes, myelination in brains of S1P₅^{-/-} mice appears normal[117]. S1P₅ was shown to influence both survival and process retraction of oligodendrocytes depending on the status of cellular differentiation[117]. Preoligodendrocytes stimulated with S1P resulted in process retraction, while the mature cells activated Akt to promote survival in a PTX-sensitive manner.

ANGIOGENESIS

As noted above, the cloning of S1P₁ from differentiating endothelial cells suggested a potential role for this receptor in angiogenesis. Indeed, one of the first biological responses that was demonstrated downstream of S1P₁ signaling was induction of morphogenetic differentiation of endothelial cells, i.e., the formation of networks resembling vascular tubes[25]. S1P induction of endothelial morphogenetic differentiation requires S1P₁-dependent enhancement of cell survival, S1P₁ and S1P₃ regulated cytoskeletal changes, and adherens junction assembly[127]. Enhanced endothelial cell survival is mediated by activation of eNOS[93], which is activated through PI3-kinase β [128], whereas cytoskeletal responses required S1P₁-induced Rac activation and S1P₃-induced Rho activation, leading to adherens junction assembly[127].

S1P is also potentially chemotactic for endothelial cells[100,101,102], and this response requires S1P₁ and S1P₃ signaling[129,130], leading to Rho-dependent clustering of integrins at focal adhesions and Rac activation[130]. Inhibition of Rho or p38 MAP kinase blocks S1P-induced endothelial cell migration[131]. Furthermore, S1P₁-induced endothelial cell migration requires phosphorylation of the S1P₁ receptor itself by Akt, and this step is required for Rac activation and angiogenesis[132]. This Akt-mediated S1P₁ phosphorylation/Rac pathway, regulates the formation of a cortactin-actin-related protein (Arp)2/3 complex, which ultimately results in membrane ruffling, formation of lamellipodia, and endothelial migration[133]. These profound effects of S1P signaling on endothelial cells lead to potent stimulation of angiogenesis. S1P stimulates angiogenesis *in vivo* in the matrigel plug assay[134], and synergizes with FGF-2 and VEGF to induce *in vivo* angiogenesis by an S1P₁- and S1P₃-dependent mechanism[102,127]. Synergism with VEGF may be related to the induction of S1P₁ expression by VEGF[135]. Moreover, S1P transactivates VEGFR in endothelial cells and the VEGFR-2/flt-1/KDR was required for S1P-induced Akt and eNOS activation[136], and membrane ruffling and motility[137].

The importance of S1P₁ for endothelial cell function was emphasized by one study showing that siRNA-mediated S1P₁ knock down eliminated S1P induction of actin ruffles and angiogenesis, sensitized cells to oxidant-mediated injury, and decreased expression of several adhesion molecules[138]. S1P₃ also appears to be important in the angiogenic response to S1P, as a peptide design to mimic the second intracellular loop of S1P₃, specifically caused S1P₃ internalization, and mimicked the effects of S1P on stimulating angiogenesis[139]. In addition, S1P₁ and S1P₃ mediate HDL-induced stimulation of endothelial cell migration, while S1P₁ mediates HDL-induced endothelial cell survival[140]. S1P₂, on the other hand, appears to antagonize the proangiogenic effects of S1P₁ and S1P₃, as inhibition of S1P₂ signaling with a specific antagonist enhances S1P-induced endothelial cell migration and angiogenic response[141].

CARDIOVASCULAR EFFECTS OF S1P

Several studies have investigated the effects of S1P on vascular smooth muscle and vascular tone. Intravenous injection of S1P reduces renal and mesenteric blood flow[142]. Many of these studies have shown that S1P causes contraction of isolated arteries *in vitro* and vasoconstriction *in vivo*, although in

some arteries S1P causes dilation, particularly in aorta when precontracted (for reviews see [143,144,145]). For example, S1P causes contraction of renal and mesenteric[146], canine basilar[147], spiral modiolar[148], pulmonary[149], and placental[150] vessels, but dilates precontracted aorta[149].

S1P may affect vascular tone through a number of signaling pathways. S1P stimulated contraction of cerebral artery that was mediated by ROCK downstream of Rho activation and calcium mobilization from intracellular stores downstream of PLC[151]. On the other hand, in the same study, S1P did not affect aorta and this was correlated with low levels of S1P₂ and S1P₃ expression in aorta and higher levels in cerebral artery[151]. S1P also caused contraction of coronary artery smooth muscle cells through S1P₂ activation[152] and increased portal pressure via Rho-mediated contraction of hepatic stellate cells[153]. Thus, it is likely that both S1P₂ and S1P₃ contribute to S1P-induced vascular smooth muscle contraction and arterial constriction through Rho and PLC pathways, respectively. Moreover, S1P₂ also induced increased COX-2 expression and activation, and cAMP accumulation in coronary artery smooth muscle cells, leading to prostaglandin synthesis[154]. Various mediators produced downstream of COX-2 can cause both vasoconstriction and vasodilation[155].

Dilation of precontracted aorta was mediated by S1P₃, since the effect was absent in vessels from S1P knockout mice[156]. Furthermore, S1P mediates HDL-induced vasodilation via S1P₃ and nitric oxide production from endothelial cells in aorta[157]. In addition, S1P₁ has been shown to activate eNOS in endothelial cells[50] and, thus, it may also contribute to this response. Indeed, loss of S1P₁ expression in vessels of aged rats allows enhanced sensitivity to S1P-induced vasoconstriction[158], presumably due to S1P₃ signaling.

Another important role for S1P in the vasculature is endothelial barrier enhancement. This is an important step late in angiogenesis as well as a part of vascular homeostasis. S1P enhances endothelial barrier integrity through S1P₁ and partially S1P₃ signaling, leading to Rho activation and actin rearrangement[159]. Activation by S1P recruits S1P₁ to caveolae[50]. This step promotes activation of PI3 kinase, Tiam1-mediated Rac activation, and cortical actin rearrangement, leading to endothelial barrier enhancement[160]. Furthermore, S1P₁ is also transactivated by activated protein C to mediate endothelial barrier enhancement[161]. In contrast, in alveolar epithelial cells, S1P₃ signaling leads increased epithelial permeability by opening tight junctions[162]. The targeted deletion of S1P₃ prevents disruption of pulmonary epithelial tight junctions by S1P[162]. Thus, S1P₃ is being investigated with regard to mediating enhanced epithelial permeability observed in pulmonary edema.

S1P also regulates heart rate. Intravenously administered S1P decreases heart rate, ventricular contraction, and blood pressure in rats[163]. S1P induction of bradycardia was found to be mediated by S1P₃, as this effect was absent in S1P₃^{-/-} mice[164,165].

ROLES OF S1P RECEPTORS IN IMMUNE CELL REGULATION

A major current focus of research on S1P, in particular through S1P₁ signaling, is its effects on lymphocyte trafficking. This topic has been the subject of several recent comprehensive reviews[166,167,168,169,170] and, therefore, it will be discussed only briefly here. These studies were initiated by the discovery that the immunosuppressant FTY720 is a sphingosine analog that, when phosphorylated to FTY720-P by SphK, acts as an agonist at four of the five S1P receptors, excluding S1P₂[38,171]. However, FTY720 causes a down-regulation of S1P₁ surface expression on lymphocytes, thus acting as a functional antagonist of this receptor[172,173]. This ultimately leads to sequestration of lymphocytes in lymph nodes, as an S1P gradient is responsible for chemotactically calling lymphocytes out of nodes into the lymph. FTY720-P also induces inhibition of thymic egress of T cells through down-regulation of surface S1P₁ expression, as S1P₁ is necessary to induce thymic egress of mature thymocytes[173,174]. Furthermore, S1P₁ signaling is also required for retention of B cells in the splenic marginal zone[175,176].

S1P at low concentrations is chemotactic for lymphocytes through S1P₁ signaling[104], while at high concentrations, S1P inhibits lymphocyte chemotaxis towards chemokines[177]. This may modulate

lymphocyte circulation, as the high concentrations of S1P in blood repress chemokine-driven lymphocyte movement into lymph nodes. Activation by antigen leads to down-regulation of S1P₁ expression, allowing lymphocytes to be called into lymph nodes. Subsequent re-expression of S1P₁ in the lymph nodes allows the S1P gradient to call lymphocytes back into lymph[166]. In addition, cyclic modulation of S1P₁ expression on lymphocytes regulates their transit through secondary lymphoid organs such that low levels of S1P₁, when the cells enter lymph nodes, allow cells to remain there until up-regulation of S1P₁ expression calls T cells into the lymph or splenic B cells into the circulation[178]. S1P₁ function on T cells may also be regulated by post-translation modification, as tyrosine sulfation of S1P₁ was shown to be necessary for the chemotactic response of T cells to S1P[179]. Furthermore, lymphocytes can also regulate S1P₁ by modulating expression of the lectin CD69, which interacts with and inhibits S1P₁. Thus, interferon (IFN)- α/β up-regulates CD69 to promote lymphocyte retention in lymphoid organs[180].

S1P and FTY720 can also affect trafficking of macrophages[181], eosinophils[182], dendritic cells[183,184], and hematopoietic progenitor cells[185]. Thus, S1P₁ is a major regulator of immune cell trafficking. In addition, S1P₁ has also been shown to suppress proliferation of naïve T cell[87,88]. T cell activation leads to S1P₁ down-regulation, allowing proliferation[104].

T cells also express S1P₄ as well as S1P₁. While S1P₁ controls T cell migration, S1P₄ plays roles in T cell proliferation and cytokine secretion[186]. Secretion of interleukin-4 (IL-4), IL-2, and IFN- γ , cytokines with positive regulatory effects on immune responses, was decreased in T cells through S1P₄ signaling, while secretion of the inhibitory cytokine IL-10 was increased. Other S1P receptors modulate cytokine secretion as well. S1P induces release of IL-6 and IL-8 from immature dendritic cells primed with lipopolysaccharide, which express S1P₁ and S1P₃, although the specific receptor involved was not identified[187]. Furthermore, S1P treatment of maturing dendritic cells decreased secretion of TNF- α and IL-12, but increased IL-10 secretion, thus favoring a Th2 response from T cells[188]. Thus, S1P is an important regulator of the immune system, not only by affecting lymphocyte trafficking, but also by regulating cytokine secretion (reviewed in [189]).

S1P₄ signaling inhibits T cell proliferation[186]. On T cell activation, S1P₄ is down-regulated, thus allowing proliferation[104]. Furthermore, it is interesting to note that FTY720-P selectively down-regulates surface S1P₁ expression, while leaving S1P₄, for which FTY720-P functions as an agonist, available on the cell surface[172]. This may enhance the immunosuppressive activity of FTY720, which may involve a combination of S1P₁ internalization and, hence, lymphocyte sequestration and S1P₄-mediated inhibition of proliferation.

S1P plays an additional role in immune cell regulation by controlling both chemotaxis and degranulation of mast cells. IgE binding to Fc ϵ RI triggers SphK activation and autocrine signaling of S1P through S1P₁, which enhances chemotaxis toward antigen. In addition, S1P₂ expression is up-regulated by Fc ϵ RI signaling, and S1P₂ signaling inhibits migration and causes mast cell degranulation[190,191].

FTY720 has tremendous therapeutic potential and is currently being studied in clinical trials. FTY720 administration to cause lymphocyte sequestration through surface down-regulation of S1P₁ is being investigated to treat transplant rejection and multiple sclerosis[192,193,194,195]. Since S1P₁ influences a variety of biological activities, additional therapeutic roles for FTY720 and other compounds, with perhaps enhanced selectivity for various S1P receptors, will likely arise with applications to cancer and other immune disorders.

S1P AND ITS RECEPTORS IN CANCER

Significant evidence has linked S1P to several aspects of malignancy. Much of this has come through studies on the enzyme that synthesizes S1P, SphK. SphK1 overexpression transforms NIH 3T3 fibroblasts, suggesting a potential oncogenic role[196]. Overexpression of SphK1 was found in several human tumor types, including breast, lung, and colon tumors, compared to matched normal tissue, suggesting that this enzyme may play a role in a wide variety of tumor types[197]. SphK1 is also

overexpressed in chemically induced, rat colon adenocarcinomas[198] and a mouse leukemia model[199]. Overexpression of SphK1 in MCF-7 breast cancer cells caused enhanced proliferation, decreased apoptosis, and led to formation of larger tumors in nude mice in an estrogen-dependent manner[200]. Conversely, a dominant-negative form of SphK1 inhibited estrogen-mediated mitogenic signaling in MCF-7 cells and decreased tumor formation in nude mice[201]. In addition, SphK overexpression also prevents apoptosis induced by serum withdrawal or chemotherapeutic drugs in several cancer types[199,202,203].

As discussed above, S1P₂ is known to inhibit cell migration. This effect has been found in cancer cells as well. S1P₂ inhibited migration and invasion through Matrigel of melanoma cells by inhibiting Rac activation[204]. Furthermore, S1P treatment prevented lung metastasis of S1P₂-expressing melanoma cells in mice, thus demonstrating an *in vivo* significance of this effect[205].

Our group has focused on the roles played by SphK and S1P in regulating the malignant behavior of human gliomas. We recently found that high levels of SphK1 expression correlate with poor survival of patients with the most malignant form of brain tumor, glioblastoma multiforme[206]. Thus, patients with low SphK1 levels survived more than three times as long as those with high levels. Moreover, SphK knock down by RNA interference in glioma cell lines decreased cell proliferation by preventing entry of cells into the cell cycle[206].

Using a panel of human glioma cell lines, we determined that S1P stimulated proliferation of 50%, and motility and invasiveness of 60% of human glioma cell lines tested[103]. S1P-induced mitogenesis was mediated in part by G_i signaling as it was approximately 50% inhibited by PTX, but required ERK and PI3 kinase activation, as inhibitors of these pathways completely blocked this response[80]. On the other hand, stimulation of motility by S1P required G_i signaling, as it was completely inhibited by PTX. The motility response was partially dependent on ERK and PI 3-kinase signaling. In addition, S1P enhanced glioma cell motility regardless of whether or not it was present in a concentration gradient, indicating that S1P stimulates a chemokinetic response in these cells[103]. Thus, S1P derived from the glioma cells themselves and stimulating in an autocrine fashion would be expected to enhance cell migration and invasion. In this regard, it has been shown that S1P is released extracellularly by astrocytes[207,208] and C6 glioma cells[209].

Whether a glioma cell line responds to S1P with proliferation or motility, or both or neither, is due to the profile of S1P receptor expression. Thus, although the glioma cell lines used all express the three S1P receptors, S1P₁, S1P₂, and S1P₃, the relative levels of expression of these receptors vary. Cell lines that do not respond mitogenically to S1P expressed extremely low levels of the receptor S1P₁[103], suggesting that this receptor is crucial for mediating S1P-stimulated glioma cell proliferation. In agreement, recent data indicate that S1P₁ overexpression in glioma cells stimulates their proliferation more potently than overexpression of S1P₂ or S1P₃, while S1P₁ knock down by RNA interference is most efficacious at blocking growth of glioma cells (NY and JVB, manuscript in preparation).

Conversely, glioma cells in which S1P stimulates motility express high proportions of S1P₁ and S1P₃, relative to S1P₂[103]. By overexpressing or knocking down S1P receptor expression in glioma cells, Lepley et al. showed that S1P₂ mediates inhibition of glioma migration, while S1P₁ mediates enhanced migration in response to S1P[110]. Our recent results confirm that S1P₁ or S1P₃ overexpression enhances glioma cell motility and invasiveness, while S1P₂ overexpression decreases motility. Surprisingly, however, S1P₂ overexpression enhances invasion of glioma cells through Matrigel, and this response appears to be mediated through an enhanced ability of S1P₂-overexpressing glioma cells to adhere to Matrigel (NY and JVB, manuscript in preparation).

Malchinkhuu et al. confirmed that S1P inhibits migration of some glioma cell lines through S1P₂ signaling[210]. They also suggested that S1P₂ is up-regulated in astrocytoma cells in comparison to normal astrocytes based on receptor expression in glioma cell lines and glioblastoma multiforme (GBM) tissue[210]. However, their analysis of GBM tissue utilized only two cases. We recently examined expression levels of S1P₁, S1P₂, and S1P₃ by real-time PCR analysis in 48 cases of GBM in comparison to 20 cases of the relatively benign pilocytic astrocytoma. We found no significant difference in expression of S1P₁, S1P₂, or S1P₃ between these two tumor types[206]. However, S1P₂ expression in GBMs was

lower than that of S1P₁ or S1P₃. Thus, although its expression level is high in some long-term glioma cell lines, S1P₂ is not likely to be a dominant S1P receptor in gliomas *in vivo*. This suggests that the promigratory effect of S1P may be dominant in glioma cells *in vivo*.

S1P may also be very important as a stimulator of tumor angiogenesis and as a therapeutic target to block this crucial aspect of cancer behavior. FTY720, which down-regulates S1P₁ expression, blocked tumor angiogenesis in a melanoma mouse model and inhibited growth of metastatic tumors[211]. Furthermore, a monoclonal antibody specific for S1P was used in animal models of several different types of tumors[212]. Without exception, the antibody decreased tumor growth potently and this was shown to be due to blocking of tumor angiogenesis. Interestingly, this was true even for a tumor cell line that is not responsive to S1P in terms of proliferation, migration, or prevention of apoptosis, suggesting that angiogenesis is a critical aspect of S1P function in malignancy.

SUMMARY

The discovery of S1P receptors has led to an explosion of information regarding the biological significance of this bioactive lipid. In a relatively short time, numerous signaling pathways regulated by these receptors, as well as their involvement in a variety of physiologic and pathologic processes particularly within the immune and cardiovascular systems as well as in neoplastic disease, have come to light. Nevertheless, the ubiquitous nature of S1P and its receptors within most cells and tissues of the body suggest that other roles played by this signaling system remain to be discovered. Knockout mouse models have provided clear functions for at least one of the receptors, S1P₁, as well as some information regarding the remaining receptors. However, it seems likely that functional redundancies, certainly among the S1P receptor family and possibly even with other receptors such as the LPA receptors, are masking other important functions of this system.

Given the diversity of dramatic effects described above for these GPCRs, the eventual understanding of the roles played by each subtype will likely have great medical impact. In 2001, 50% of current, commercially available pharmaceuticals targeted GPCRs and were a \$30 billion industry[213]. The potential therapeutic application of S1P receptor-G-protein signaling can be applied to a variety of settings, most notably cancer and immune system disorders. Many of the aspects of malignant behavior of cancers are regulated by S1P receptor activation, including proliferation, cell survival by resistance to apoptosis, invasion, and angiogenesis. Additionally, S1P migratory effects are an integral influence on lymphocyte trafficking. Thus, the near future will undoubtedly see a continuation of the intense interest in this field.

ACKNOWLEDGMENTS

Work in the author's laboratory was supported by Grant # R01 NS41517 from the National Institute of Neurological Disorders and Stroke (NINDS) to JRVB.

REFERENCES

1. Yatomi, Y., Igarashi, Y., Yang, L., Hisano, N., Qi, R., Asazuma, N., Satoh, K., Ozaki, Y., and Kume, S. (1997) Sphingosine 1-phosphate, a bioactive sphingolipid abundantly stored in platelets, is a normal constituent of human plasma and serum. *J. Biochem.* **121**, 969–973.
2. Murata, N., Sato, K., Kon, J., Tomura, H., Yanagita, M., Kuwabara, A., Ui, M., and Okajima, F. (2000) Interaction of sphingosine 1-phosphate with plasma components, including lipoproteins, regulates the lipid receptor-mediated actions. *Biochem. J.* **352**, 809–815.
3. Yatomi, Y., Ruan, F., Hakomori, S., and Igarashi, Y. (1995) Sphingosine-1-phosphate: a platelet-activating sphingolipid released from agonist-stimulated human platelets. *Blood* **86**, 193–202.
4. Olivera, A. and Rivera, J. (2005) Sphingolipids and the balancing of immune cell function: lessons from the mast cell.

- J. Immunol.* **174**, 1153–1158.
5. Olivera, A. and Spiegel, S. (1993) Sphingosine-1-phosphate as a second messenger in cell proliferation induced by PDGF and FCS mitogens. *Nature* **365**, 557–560.
6. Edsall, L.C., Pirianov, G.G., and Spiegel, S. (1997) Involvement of sphingosine 1-phosphate in nerve growth factor-mediated neuronal survival and differentiation. *J. Neurosci.* **17**, 6952–6960.
7. Duan, H.F., Wu, C.T., Lu, Y., Wang, H., Liu, H.J., Zhang, Q.W., Jia, X.X., Lu, Z.Z., and Wang, L.S. (2004) Sphingosine kinase activation regulates hepatocyte growth factor induced migration of endothelial cells. *Exp. Cell Res.* **298**, 593–601.
8. Sarkar, S., Maceyka, M., Hait, N.C., Paugh, S.W., Sankala, H., Milstien, S., and Spiegel, S. (2005) Sphingosine kinase 1 is required for migration, proliferation and survival of MCF-7 human breast cancer cells. *FEBS Lett.* **579**, 5313–5317.
9. Choi, O.H., Kim, J.-H., and Kinet, J.-P. (1996) Calcium mobilization via sphingosine kinase in signalling by the FcεRI antigen receptor. *Nature* **380**, 634–636.
10. Melendez, A., Floto, R.A., Gillooly, D.J., Harnett, M.M., and Allen, J.M. (1998) FcγRI coupling to phospholipase D initiates sphingosine kinase-mediated calcium mobilization and vesicular trafficking. *J. Biol. Chem.* **273**, 9393–9402.
11. Xia, P., Gamble, J.R., Rye, K.A., Wang, L., Hii, C.S.T., Cockerill, P., Khew-Goodall, Y., Bert, A.G., Barter, P.J., and Vadas, M.A. (1998) Tumor necrosis factor-α induces adhesion molecule expression through the sphingosine kinase pathway. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 14196–14201.
12. Meyer zu Heringdorf, D., Lass, H., Alemany, R., Laser, K.T., Neumann, E., Zhang, C., Schmidt, M., Rauen, U., Jakobs, K.H., and van Koppen, C.J. (1998) Sphingosine kinase-mediated Ca²⁺ signalling by G-protein-coupled receptors. *EMBO J.* **17**, 2830–2837.
13. Maceyka, M., Payne, S.G., Milstien, S., and Spiegel, S. (2002) Sphingosine kinase, sphingosine-1-phosphate, and apoptosis. *Biochim. Biophys. Acta* **1585**, 193–201.
14. Spiegel, S. and Milstien, S. (2003) Exogenous and intracellularly generated sphingosine 1-phosphate can regulate cellular processes by divergent pathways. *Biochem. Soc. Trans.* **31**, 1216–1219.
15. Chun, J., Goetzl, E.J., Hla, T., Igarashi, Y., Lynch, K.R., Moolenaar, W., Pyne, S., and Tigyi, G. (2002) International Union of Pharmacology. XXXIV. Lysophospholipid Receptor Nomenclature. *Pharmacol. Rev.* **54**, 265–269.
16. Zhang, G., Contos, J.J.A., Weiner, J.A., Fukushima, N., and Chun, J. (1999) Comparative analysis of three murine G-protein coupled receptors activated by sphingosine-1-phosphate. *Gene* **227**, 89–99.
17. Im, D.-S., Heise, C.E., Ancellin, N., O'Dowd, B.F., Shei, G.-J., Heavens, R.P., Rigby, M.R., Hla, T., Mandala, S., McAllister, G., George, S.R., and Lynch, K.R. (2000) Characterization of a novel sphingosine 1-phosphate receptor, Edg-8. *J. Biol. Chem.* **275**, 14281–14286.
18. Yamazaki, Y., Kon, J., Sato, K., Tomura, H., Sato, M., Yoneya, T., Okazaki, H., Okajima, F., and Ohta, H. (2000) Edg-6 as a putative sphingosine 1-phosphate receptor coupling to Ca²⁺ signaling pathway. *Biochem. Biophys. Res. Commun.* **268**, 583–589.
19. Kostenis, E. (2004) Novel clusters of receptors for sphingosine-1-phosphate, sphingosylphosphorylcholine, and (lyso)-phosphatidic acid: new receptors for "old" ligands. *J. Cell. Biochem.* **92**, 923–936.
20. Hla, T. and Maciag, T. (1990) An abundant transcript induced in differentiating human endothelial cells encodes a polypeptide with structural similarities to G-protein coupled receptors. *J. Biol. Chem.* **265**, 9308–9313.
21. Okazaki, H., Ishizaka, N., Sakurai, T., Kurokawa, K., Goto, K., Kumada, M., and Takuwa, Y. (1993) Molecular cloning of a novel putative G protein-coupled receptor expressed in the cardiovascular system. *Biochem. Biophys. Res. Commun.* **190**, 1104–1109.
22. MacLennan, A.J., Browe, C.S., Gaskin, A.A., Lado, D.C., and Shaw, G. (1994) Cloning and characterization of a putative G-protein coupled receptor potentially involved in development. *Mol. Cell Neurosci.* **5**, 201–209.
23. Yamaguchi, F., Tokuda, M., Hatase, O., and Brenner, S. (1996) Molecular cloning of the novel human G protein-coupled receptor (GPCR) gene mapped on chromosome 9. *Biochem. Biophys. Res. Commun.* **227**, 608–614.
24. Hecht, J.H., Weiner, J.A., Post, S.R., and Chun, J. (1996) Ventricular zone gene-1 (vzg-1) encodes a lysophosphatidic acid receptor expressed in neurogenic regions of the developing cerebral cortex. *J. Cell Biol.* **135**, 1071–1083.
25. Lee, M.-J., Van Brocklyn, J.R., Thangada, S., Liu, C.H., Hand, A.R., Menzeleev, R., Spiegel, S., and Hla, T. (1998) Sphingosine-1-phosphate as a ligand for the G protein-coupled receptor EDG-1. *Science* **279**, 1552–1555.
26. An, S., Bleu, T., Huang, W., Hallmark, O.G., Coughling, S.R., and Goetzl, E.J. (1997) Identification of cDNAs encoding two G protein-coupled receptors for lysosphingolipids. *FEBS Lett.* **417**, 279–282.
27. Gräler, M.H., Bernhardt, G., and Lipp, M. (1998) EDG6, a novel G-protein-coupled receptor related to receptors for bioactive lysophospholipids, is specifically expressed in lymphoid tissue. *Genomics* **53**, 164–169.
28. Glickman, M., Malek, R.L., Kwitek-Black, A.E., Jacob, H.J., and Lee, N.H. (1999) Molecular cloning, tissue-specific expression, and chromosomal localization of a novel nerve growth factor-regulated G-protein-coupled receptor, nrg-1. *Mol. Cell. Neurosci.* **14**, 141–152.
29. Windh, R.T., Lee, M.J., Hla, T., An, S., Barr, A.J., and Manning, D.R. (1999) Differential coupling of the sphingosine 1-phosphate receptors Edg-1, Edg-3, and H218/Edg-5 to the G_i, G_q, and G₁₂ families of heterotrimeric G proteins. *J. Biol. Chem.* **274**, 27351–27358.
30. Okamoto, H., Takuwa, N., Gonda, K., Okazaki, H., Chang, K., Yatomi, Y., Shigematsu, H., and Takuwa, Y. (1998) EDG1 is a functional sphingosine-1-phosphate receptor that is linked via a Gi/o to multiple signaling pathways,

- including phospholipase C activation, Ca^{2+} mobilization, ras-mitogen-activated protein kinase activation, and adenylate cyclase inhibition. *J. Biol. Chem.* **273**, 27104–27110.
31. Kon, J., Sato, K., Watanabe, T., Tomura, H., Kuwabara, A., Kimura, T., Tamama, K., Ishizuka, T., Murata, N., Kanda, T., Kobayashi, I., Ohta, H., Ui, M., and Okajima, F. (1999) Comparison of intrinsic activities of the putative sphingosine 1-phosphate receptor subtypes to regulate several signaling pathways in their cDNA-transfected chinese hamster ovary cells. *J. Biol. Chem.* **274**, 23940–23947.
32. Sato, K., Ui, M., and Okajima, F. (2000) Differential roles of Edg-1 and Edg-5, sphingosine 1-phosphate receptors, in the signaling pathways in C6 glioma cells. *Brain Res. Mol. Brain Res.* **85**, 151–160.
33. Tamama, K., Kon, J., Sato, K., Tomura, H., Kuwabara, A., Kimura, T., Kanda, T., Ohta, H., Ui, M., Kobayashi, I., and Okajima, F. (2001) Extracellular mechanism through the Edg family of receptors might be responsible for sphingosine-1-phosphate-induced regulation of DNA synthesis and migration of rat aortic smooth-muscle cells. *Biochem. J.* **353**, 139–146.
34. Van Brocklyn, J.R., Tu, Z., Edsall, L., Schmidt, R.R., and Spiegel, S. (1999) Sphingosine 1-phosphate-induced cell rounding and neurite retraction are mediated by the G protein-coupled receptor H218. *J. Biol. Chem.* **274**, 4626–4632.
35. Malek, R.L., Toman, R.E., Edsall, L.C., Wong, S., Chiu, J., Letterle, C.A., Van Brocklyn, J.R., Milstein, S., Spiegel, S., and Lee, N.H. (2001) Nrg-1 belongs to the endothelial differentiation gene family of G protein-coupled sphingosine-1-phosphate receptors. *J. Biol. Chem.* **276**, 5692–5699.
36. Van Brocklyn, J.R., Gräler, M.H., Bernhardt, G., Lipp, M., and Spiegel, S. (2000) Sphingosine-1-phosphate is a ligand for the G protein-coupled receptor EDG-6. *Blood* **95**, 2624–2629.
37. Candelore, M.R., Wright, M., Tota, L., Milligan, J., Shei, G., Bergstrom, J., and Mandala, S. (2002) Phytosphingosine 1-phosphate: a high affinity ligand for the S1P₄/Edg-6 receptor. *Biochem. Biophys. Res. Commun.* **297**, 600–606.
38. Mandala, S., Hajdu, R., Bergstrom, J., Quackenbush, E., Xie, J., Milligan, J., Thornton, R., Shei, G.J., Card, D., Keohane, C., Rosenbach, M., Hale, J., Lynch, C.L., Rupprecht, K., Parsons, W., and Rosen, H. (2002) Alteration of lymphocyte trafficking by sphingosine 1-phosphate receptor agonists. *Science* **296**, 346–349.
39. Murata, N., Sato, K., Kon, J., Tomura, H., and Okajima, F. (2000) Quantitative measurement of sphingosine 1-phosphate by radioreceptor-binding assay. *Anal. Biochem.* **282**, 115–120.
40. Fossetta, J., Deno, G., Gonsiorek, W., Fan, X., Lavey, B., Das, P., Lunn, C., Zavodny, P.J., Lundell, D., and Hipkin, R.W. (2004) Pharmacological characterization of human S1P₄ using a novel radioligand, [4,5-³H]-dihydrosphingosine-1-phosphate. *Br. J. Pharmacol.* **142**, 851–860.
41. Okamoto, H., Takuwa, N., Yatomi, Y., Gonda, K., Shigematsu, H., and Takuwa, Y. (1999) EDG3 is a functional receptor specific for sphingosine 1-phosphate and sphingosylphosphorylcholine with signaling characteristics distinct from EDG1 and AGR16. *Biochem. Biophys. Res. Commun.* **260**, 203–208.
42. Gonda, K., Okamoto, H., Takuwa, N., Yatomi, Y., Okazaki, H., Sakurai, T., Kimura, S., Sillard, R., Harii, K., and Takuwa, Y. (1999) The novel sphingosine 1-phosphate receptor AGR16 is coupled via pertussis toxin-sensitive and -insensitive G-proteins to multiple signalling pathways. *Biochem. J.* **337**, 67–75.
43. Ancellin, N. and Hla, T. (1999) Differential pharmacological properties and signal transduction of the sphingosine 1-phosphate receptors EDG-1, EDG-3, and EDG-5. *J. Biol. Chem.* **274**, 18997–19002.
44. Lee, M.-J., Evans, M., and Hla, T. (1996) The inducible G protein-coupled receptor *edg-1* signals via the G_i/mitogen-activated protein kinase pathway. *J. Biol. Chem.* **271**, 11272–11282.
45. Zondag, G.C.M., Postma, F.R., Etten, I.V., Verlaan, I., and Moolenaar, W.H. (1998) Sphingosine 1-phosphate signalling through the G-protein-coupled receptor Edg-1. *Biochem. J.* **330**, 605–609.
46. Van Brocklyn, J.R., Lee, M.J., Menzeleev, R., Olivera, A., Edsall, L., Cuvillier, O., Thomas, D.M., Coopman, P.J.P., Thangada, S., Hla, T., and Spiegel, S. (1998) Dual actions of sphingosine-1-phosphate: extracellular through the G_i-coupled orphan receptor *edg-1* and intracellular to regulate proliferation and survival. *J. Cell Biol.* **142**, 229–240.
47. Meyer zu Heringdorf, D., Lass, H., Kuchar, I., Lipinski, M., Alemany, R., Rumenapp, U., and Jakobs, K.H. (2001) Stimulation of intracellular sphingosine-1-phosphate production by G-protein-coupled sphingosine-1-phosphate receptors. *Eur. J. Pharmacol.* **414**, 145–154.
48. Meyer zu Heringdorf, D., Vincent, M.E., Lipinski, M., Danneberg, K., Stropp, U., Wang de, A., Tigyi, G., and Jakobs, K.H. (2003) Inhibition of Ca^{2+} signalling by the sphingosine 1-phosphate receptor S1P₁. *Cell. Signal.* **15**, 677–687.
49. Morales-Ruiz, M., Lee, M.J., Zoellner, S., Gratton, J.P., Scotland, R., Shiojima, I., Walsh, K., Hla, T., and Sessa, W.C. (2001) Sphingosine-1-phosphate activates Akt, nitric oxide production and chemotaxis through a G_i-protein/phosphoinositide 3-kinase pathway in endothelial cells. *J. Biol. Chem.* **276**, 19672–19677.
50. Igarashi, J. and Michel, T. (2000) Agonist-modulated targeting of the EDG-1 receptor to plasmalemmal caveolae: eNOS activation by sphingosine 1-phosphate and the role of caveolin-1 in sphingolipid signal transduction. *J. Biol. Chem.* **275**, 32363–32370.
51. Gonzalez, E., Nagiel, A., Lin, A.J., Golan, D.E., and Michel, T. (2004) Small interfering RNA-mediated down-regulation of caveolin-1 differentially modulates signaling pathways in endothelial cells. *J. Biol. Chem.* **279**, 40659–40669.
52. Kohno, T., Wada, A., and Igarashi, Y. (2002) N-glycans of sphingosine 1-phosphate receptor Edg-1 regulate ligand-induced receptor internalization. *FASEB J.* **16**, 983–992.
53. Hobson, J.P., Rosenfeldt, H.M., Barak, L.S., Olivera, A., Poulton, S., Caron, M.G., Milstien, S., and Spiegel, S.

- (2001) Role of the sphingosine-1-phosphate receptor EDG-1 in PDGF-induced cell motility. *Science* **291**, 1800–1803.
54. Rosenfeldt, H.M., Hobson, J.P., Maceyka, M., Olivera, A., Nava, V.E., Milstien, S., and Spiegel, S. (2001) EDG-1 links the PDGF receptor to Src and focal adhesion kinase activation leading to lamellipodia formation and cell migration. *FASEB J.* **15**, 2649–2659.
55. Alderton, F., Rakhit, S., Choi, K.K., Palmer, T., Sambi, B., Pyne, S., and Pyne, N.J. (2001) Tethering of the platelet-derived growth factorb receptor to G-protein coupled receptors: a novel platform for integrative signaling by these receptor classes in mammalian cells. *J. Biol. Chem.* **276**, 28578–28585.
56. Waters, C., Sambi, B.S., Kong, K.C., Thompson, D., Pitson, S.M., Pyne, S., and Pyne, N.J. (2003) Sphingosine 1-phosphate and platelet-derived growth factor (PDGF) act via PDGFb receptor-sphingosine 1-phosphate receptor complexes in airway smooth muscle cells. *J. Biol. Chem.* **278**, 6282–6290.
57. Waters, C.M., Long, J., Gorshkova, I., Fujiwara, Y., Connell, M., Belmonte, K.E., Tigyi, G., Natarajan, V., Pyne, S., and Pyne, N.J. (2006) Cell migration activated by platelet-derived growth factor receptor is blocked by an inverse agonist of the sphingosine 1-phosphate receptor-1. *FASEB J.* **20**(3), 509–511.
58. Kluk, M.J., Colmont, C., Wu, M.T., and Hla, T. (2003) Platelet-derived growth factor (PDGF)-induced chemotaxis does not require the G protein-coupled receptor S1P₁ in murine embryonic fibroblasts and vascular smooth muscle cells. *FEBS Lett.* **533**, 25–28.
59. Usui, S., Sugimoto, N., Takuwa, N., Sakagami, S., Takata, S., Kaneko, S., and Takuwa, Y. (2004) Blood lipid mediator sphingosine-1-phosphate potently stimulates platelet-derived growth factor-A and -B chain expression through S1P₁-Gi-Ras-MAPK-dependent induction of Krupel-like factor 5. *J. Biol. Chem.* **279**, 12300–12311.
60. Tanimoto, T., Lungu, A.O., and Berk, B.C. (2004) Sphingosine 1-phosphate transactivates the platelet-derived growth factor b receptor and epidermal growth factor receptor in vascular smooth muscle cells. *Circ. Res.* **94**, 1505–1058.
61. Postma, F.R., Jalink, K., Hengeveld, T., and Moolenaar, W.H. (1996) Sphingosine-1-phosphate rapidly induces rho-dependent neurite retraction: action through a specific cell surface receptor. *EMBO J.* **15**, 2388–2392.
62. Ishii, I., Ye, X., Friedman, B., Kawamura, S., Contos, J.J., Kingsbury, M.A., Yang, A.H., Zhang, G., Brown, J.H., and Chun, J. (2002) Marked perinatal lethality and cellular signaling deficits in mice null for the two sphingosine 1-phosphate receptors, S1P₂/LPB₂/EDG-5 and S1P₃/LPB₃/EDG-3. *J. Biol. Chem.* **277**, 25152–25159.
63. Hu, W., Mahavadi, S., Huang, J., Li, F., and Murthy, K.S. (2006) Characterization of S1P₁ and S1P₂ receptor function in smooth muscle by receptor silencing and receptor protection. *Am. J. Physiol. Gastrointest. Liver Physiol.* [E-pub]
64. Siehler, S., Wang, Y., Fan, X., Windh, R.T., and Manning, D.R. (2001) Sphingosine 1-phosphate activates nuclear factor-kB through Edg receptors: activation through Edg-3 and Edg-5, but not Edg-1, in human embryonic kidney 293 cells. *J. Biol. Chem.* **276**, 48733–48739.
65. Baudhuin, L.M., Jiang, Y., Zaslavsky, A., Ishii, I., Chun, J., and Xu, Y. (2004) S1P₃-mediated Akt activation and crosstalk with platelet-derived growth factor receptor (PDGFR). *FASEB J.* **18**, 341–343.
66. Sukocheva, O., Wadham, C., Holmes, A., Albanese, N., Verrier, E., Feng, F., Bernal, A., Derian, C.K., Ullrich, A., Vadas, M.A., and Xia, P. (2006) Estrogen transactivates EGFR via the sphingosine 1-phosphate receptor Edg-3: the role of sphingosine kinase-1. *J. Cell Biol.* **173**, 301–310.
67. Gräler, M.H., Grosse, R., Kusch, A., Kremmer, E., Gudermann, T., and Lipp, M. (2003) The sphingosine 1-phosphate receptor S1P₄ regulates cell shape and motility via coupling to G_i and G_{12/13}. *J. Cell. Biochem.* **89**, 507–519.
68. Kohno, T., Matsuyuki, H., Inagaki, Y., and Igarashi, Y. (2003) Sphingosine 1-phosphate promotes cell migration through the activation of Cdc42 in Edg-6/S1P₄-expressing cells. *Genes Cells* **8**, 685–697.
69. Zhang, H., Desai, N.N., Olivera, A., Seki, T., Brooker, G., and Spiegel, S. (1991) Sphingosine-1-phosphate, a novel lipid, involved in cellular proliferation. *J. Cell Biol.* **114**, 155–167.
70. Goodemote, K.A., Mattie, M.E., Berger, A., and Spiegel, S. (1995) Involvement of a pertussis toxin sensitive G protein in the mitogenic signaling pathways of sphingosine 1-phosphate. *J. Biol. Chem.* **270**, 10272–10277.
71. Kluk, M.J. and Hla, T. (2001) Role of the sphingosine 1-phosphate receptor edg-1 in vascular smooth muscle cell proliferation and migration. *Circ. Res.* **89**, 496–502.
72. Carpio, L.C., Stephan, E., Kamer, A., and Dziak, R. (1999) Sphingolipids stimulate cell growth via MAP kinase activation in osteoblastic cells. *Prostaglandins Leukot. Essent. Fatty Acids* **61**, 267–273.
73. Goetzl, E.J., Dolezalova, H., Kong, Y., and Zeng, L. (1999) Dual mechanisms for lysophospholipid induction of proliferation of human breast carcinoma cells. *Cancer Res.* **59**, 4732–4737.
74. Ikeda, H., Yatomi, Y., Yanase, M., Satoh, H., Maekawa, H., Ogata, I., Ozaki, Y., Takuwa, Y., Mochida, S., and Fujiwara, K. (2000) Biological activities of novel lipid mediator sphingosine 1-phosphate in rat hepatic stellate cells. *Am. J. Physiol. Gastrointest. Liver Physiol.* **279**, G304–G310.
75. Kimura, T., Watanabe, T., Sato, K., Kon, J., Tomura, H., Tamama, K., Kuwabara, A., Kanda, T., Kobayashi, I., Ohta, H., Ui, M., and Okajima, F. (2000) Sphingosine 1-phosphate stimulates proliferation and migration of human endothelial cells possibly through the lipid receptors, Edg-1 and Edg-3. *Biochem. J.* **348**, 71–76.
76. Ammit, A.J., Hastie, A.T., Edsall, L.C., Hoffman, R.K., Amrani, Y., Krymskaya, V.P., Kane, S.A., Peters, S.P., Penn, R.B., Spiegel, S., and Panettieri, R.A., Jr. (2001) Sphingosine 1-phosphate modulates human airway smooth muscle cell functions that promote inflammation and airway remodeling in asthma. *FASEB J.* **15**, 1212–1214.
77. Pébay, A., Toutant, M., Premont, J., Calvo, C.F., Venance, L., Cordier, J., Glowinski, J., and Tence, M. (2001) Sphingosine-1-phosphate induces proliferation of astrocytes: regulation by intracellular signalling cascades. *Eur. J. Neurosci.* **13**, 2067–2076.

78. Hanafusa, N., Yatomi, Y., Yamada, K., Hori, Y., Nangaku, M., Okuda, T., Fujita, T., Kurokawa, K., and Fukagawa, M. (2002) Sphingosine 1-phosphate stimulates rat mesangial cell proliferation from outside the cells. *Nephrol. Dial. Transplant.* **17**, 580–586.
79. Thamilselvan, V., Li, W., Sumpio, B.E., and Basson, M.D. (2002) Sphingosine-1-phosphate stimulates human Caco-2 intestinal epithelial proliferation via p38 activation and activates ERK by an independent mechanism. *In Vitro Cell Dev. Biol. Anim.* **38**, 246–253.
80. Van Brocklyn, J.R., Letterle, C.A., Snyder, P.J., and Prior, T.W. (2002) Sphingosine-1-phosphate stimulates human glioma cell proliferation through G_i-coupled receptors: role of ERK MAP kinase and phosphatidylinositol 3-kinase b. *Cancer Lett.* **181**, 195–204.
81. Harada, J., Foley, M., Moskowitz, M.A., and Waeber, C. (2004) Sphingosine-1-phosphate induces proliferation and morphological changes of neural progenitor cells. *J. Neurochem.* **88**, 1026–1039.
82. Kim, M.K., Lee, H.Y., Kwak, J.Y., Park, J.I., Yun, J., and Bae, Y.S. (2006) Sphingosine-1-phosphate stimulates rat primary chondrocyte proliferation. *Biochem. Biophys. Res. Commun.* **345**, 67–73.
83. An, S., Zheng, Y., and Bleu, T. (2000) Sphingosine 1-phosphate-induced cell proliferation, survival, and related signaling events mediated by G protein-coupled receptors edg3 and edg5. *J. Biol. Chem.* **275**, 288–296.
84. Ikeda, H., Satoh, H., Yanase, M., Inoue, Y., Tomiya, T., Arai, M., Tejima, K., Nagashima, K., Maekawa, H., Yahagi, N., Yatomi, Y., Sakurada, S., Takuwa, Y., Ogata, I., Kimura, S., and Fujiwara, K. (2003) Antiproliferative property of sphingosine 1-phosphate in rat hepatocytes involves activation of Rho via Edg-5. *Gastroenterology* **124**, 459–469.
85. Björklund, S., Palmberg, S., Rask, S., Westerdahl, A.C., and Törnquist, K. (2005) Effects of sphingosine 1-phosphate on calcium signaling, proliferation and S1P₂ receptor expression in PC Cl₃ rat thyroid cells. *Mol. Cell. Endocrinol.* **231**, 65–74.
86. Davaille, J., Gallois, C., Habib, A., Li, L., Mallat, A., Tao, J., Levade, T., and Lotersztajn, S. (2000) Antiproliferative effects of sphingosine-1-phosphate in human hepatic myofibroblasts: a cyclooxygenase-2-mediated pathway. *J. Biol. Chem.* **275**, 34628–34633.
87. Dorsam, G., Graeler, M.H., Seroogy, C., Kong, Y., Voice, J.K., and Goetzl, E.J. (2003) Transduction of multiple effects of sphingosine 1-phosphate (S1P) on T cell functions by the S1P₁ G protein-coupled receptor. *J. Immunol.* **171**, 3500–3507.
88. Jin, Y., Knudsen, E., Wang, L., Bryceson, Y., Damaj, B., Gessani, S., and Maghazachi, A.A. (2003) Sphingosine 1-phosphate is a novel inhibitor of T cell proliferation. *Blood* **101**, 4909–4915.
89. Kim, D.S., Kim, S.Y., Kleuser, B., Schafer-Korting, M., Kim, K.H., and Park, K.C. (2004) Sphingosine-1-phosphate inhibits human keratinocyte proliferation via Akt/protein kinase B inactivation. *Cell. Signal.* **16**, 89–95.
90. Yamashita, H., Kitayama, J., Shida, D., Yamaguchi, H., Mori, K., Osada, M., Aoki, S., Yatomi, Y., Takuwa, Y., and Nagawa, H. (2006) Sphingosine 1-phosphate receptor expression profile in human gastric cancer cells: differential regulation on the migration and proliferation. *J. Surg. Res.* **130**, 80–87.
91. Goparaju, S.K., Jolly, P.S., Watterson, K.R., Bektas, M., Alvarez, S., Sarkar, S., Mel, L., Ishii, I., Chun, J., Milstien, S., and Spiegel, S. (2005) The S1P₂ receptor negatively regulates platelet-derived growth factor-induced motility and proliferation. *Mol. Cell. Biol.* **25**, 4237–4249.
92. Baudhuin, L.M., Cristina, K.L., Lu, J., and Xu, Y. (2002) Akt activation induced by lysophosphatidic acid and sphingosine-1-phosphate requires both mitogen-activated protein kinase kinase and p38 mitogen-activated protein kinase and is cell-line specific. *Mol. Pharmacol.* **62**, 660–671.
93. Kwon, Y.G., Min, J.K., Kim, K.M., Lee, D.J., Billiar, T.R., and Kim, Y.M. (2001) Sphingosine 1-phosphate protects human umbilical vein endothelial cells from serum-deprived apoptosis by nitric oxide production. *J. Biol. Chem.* **276**, 10627–10633.
94. Kim, D.S., Hwang, E.S., Lee, J.E., Kim, S.Y., and Park, K.C. (2003) Sphingosine-1-phosphate promotes mouse melanocyte survival via ERK and Akt activation. *Cell. Signal.* **15**, 919–926.
95. Davaille, J., Li, L., Mallat, A., and Lotersztajn, S. (2002) Sphingosine-1-phosphate triggers both apoptotic and survival signals for human hepatic myofibroblasts. *J. Biol. Chem.* **277**, 37323–37330.
96. Chihab, R., Porn-Ares, M.I., Alvarado-Kristensson, M., and Andersson, T. (2003) Sphingosine 1-phosphate antagonizes human neutrophil apoptosis via p38 mitogen-activated protein kinase. *Cell. Mol. Life Sci.* **60**, 776–785.
97. Goetzl, E.J., Kong, Y., and Mei, B. (1999) Lysophosphatidic acid and sphingosine 1-phosphate protection of T cells from apoptosis in association with suppression of Bax. *J. Immunol.* **162**, 2049–2056.
98. Sadahira, Y., Ruan, F., Hakomori, S., and Igarashi, Y. (1992) Sphingosine 1-phosphate, a specific endogenous signaling molecule controlling cell motility and tumor cell invasiveness. *Proc. Natl. Acad. Sci. U. S. A.* **89**, 9686–9690.
99. Wang, F., Van Brocklyn, J.R., Edsall, L., Nava, V.E., and Spiegel, S. (1999) Sphingosine-1-phosphate inhibits motility of human breast cancer cells independently of cell surface receptors. *Cancer Res.* **59**, 6185–6191.
100. Wang, F., Van Brocklyn, J.R., Hobson, J.P., Movafagh, S., Zukowska-Grojec, Z., Milstien, S., and Spiegel, S. (1999) Sphingosine 1-phosphate stimulates cell migration through a G_i-coupled cell surface receptor: potential involvement in angiogenesis. *J. Biol. Chem.* **274**, 35343–35350.
101. Okamoto, H., Yatomi, Y., Ohmori, T., Satoh, K., Matsumoto, Y., and Ozaki, Y. (2000) Sphingosine 1-phosphate stimulates G_i- and Rho-mediated vascular endothelial cell spreading and migration. *Thromb. Res.* **99**, 259–265.
102. English, D., Welch, Z., Kovala, A.T., Harvey, K., Volpert, O.V., Brindley, D.N., and Garcia, J.G. (2000) Sphingosine

- 1-phosphate released from platelets during clotting accounts for the potent endothelial cell chemotactic activity of blood serum and provides a novel link between hemostasis and angiogenesis. *FASEB J.* **14**, 2255–2265.
103. Van Brocklyn, J.R., Young, N., and Roof, R. (2003) Sphingosine-1-phosphate stimulates motility and invasiveness of human glioblastoma multiforme cells. *Cancer Lett.* **199**, 53–60.
 104. Graeler, M. and Goetzl, E.J. (2002) Activation-regulated expression and chemotactic function of sphingosine 1-phosphate receptors in mouse splenic T cells. *FASEB J.* **16**, 1874–1878.
 105. Boguslawski, G., Grogg, J.R., Welch, Z., Ciechanowicz, S., Sliva, D., Kovala, A.T., McGlynn, P., Brindley, D.N., Rhoades, R.A., and English, D. (2002) Migration of vascular smooth muscle cells induced by sphingosine 1-phosphate and related lipids: potential role in the angiogenic response. *Exp. Cell Res.* **274**, 264–274.
 106. Tanski, W., Roztocil, E., and Davies, M.G. (2002) Sphingosine-1-phosphate induces G_{i1} -coupled, PI3K/ras-dependent smooth muscle cell migration. *J. Surg. Res.* **108**, 98–106.
 107. Ryu, Y., Takuwa, N., Sugimoto, N., Sakurada, S., Usui, S., Okamoto, H., Matsui, O., and Takuwa, Y. (2002) Sphingosine-1-phosphate, a platelet-derived lysophospholipid mediator, negatively regulates cellular Rac activity and cell migration in vascular smooth muscle cells. *Circ. Res.* **90**, 325–332.
 108. Okamoto, H., Takuwa, N., Yokomizo, T., Sugimoto, N., Sakurada, S., Shigematsu, H., and Takuwa, Y. (2000) Inhibitory regulation of Rac activation, membrane ruffling, and cell migration by the G protein-coupled sphingosine-1-phosphate receptor EDG5 but not EDG1 or EDG3. *Mol. Cell. Biol.* **20**, 9247–9261.
 109. Sugimoto, N., Takuwa, N., Okamoto, H., Sakurada, S., and Takuwa, Y. (2003) Inhibitory and stimulatory regulation of rac and cell motility by the $G_{12/13}$ -Rho and G_i pathways integrated downstream of a single G protein-coupled sphingosine-1-phosphate receptor isoform. *Mol. Cell. Biol.* **23**, 1534–1545.
 110. Lepley, D., Paik, J.H., Hla, T., and Ferrer, F. (2005) The G protein-coupled receptor S1P₂ regulates Rho/Rho kinase pathway to inhibit tumor cell migration. *Cancer Res.* **65**, 3788–3795.
 111. Sanchez, T., Thangada, S., Wu, M.T., Kontos, C.D., Wu, D., Wu, H., and Hla, T. (2005) PTEN as an effector in the signaling of antimigratory G protein-coupled receptor. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 4312–4317.
 112. Kupperman, E., An, S., Osborne, N., Waldron, S., and Stainier, D.Y. (2000) A sphingosine-1-phosphate receptor regulates cell migration during vertebrate heart development. *Nature* **406**, 192–195.
 113. Chae, S.S., Proia, R.L., and Hla, T. (2004) Constitutive expression of the S1P₁ receptor in adult tissues. *Prostaglandins Other Lipid Mediat.* **73**, 141–150.
 114. Liu, C.H. and Hla, T. (1997) The mouse gene for the inducible G-protein-coupled receptor *edg-1*. *Genomics* **43**, 15–24.
 115. McGiffert, C., Contos, J.J., Friedman, B., and Chun, J. (2002) Embryonic brain expression analysis of lysophospholipid receptor genes suggests roles for s1p₁ in neurogenesis and s1p₁₋₃ in angiogenesis. *FEBS Lett.* **531**, 103–108.
 116. MacLennan, A.J., Marks, L., Gaskin, A.A., and Lee, N. (1997) Embryonic expression pattern of H218, a G-protein coupled receptor homolog, suggests roles in early mammalian nervous system development. *Neuroscience* **79**, 217–224.
 117. Jaillard, C., Harrison, S., Stankoff, B., Aigrot, M.S., Calver, A.R., Duddy, G., Walsh, F.S., Pangalos, M.N., Arimura, N., Kaibuchi, K., Zalc, B., and Lubetzki, C. (2005) Edg8/S1P5: an oligodendroglial receptor with dual function on process retraction and cell survival. *J. Neurosci.* **25**, 1459–1469.
 118. Ulfig, N. and Briesse, M. (2004) Evidence for the presence of the sphingosine-1-phosphate receptor Edg-8 in human radial glial fibers. *Acta Histochem.* **106**, 373–378.
 119. Ishii, I., Friedman, B., Ye, X., Kawamura, S., McGiffert, C., Contos, J.J., Kingsbury, M.A., Zhang, G., Brown, J.H., and Chun, J. (2001) Selective loss of sphingosine 1-phosphate signaling with no obvious phenotypic abnormality in mice lacking its G protein-coupled receptor, LPB3/EDG-3. *J. Biol. Chem.* **276**, 33697–33704.
 120. Van Brocklyn, J.R., Behbahani, B., and Lee, N.H. (2002) Homodimerization and heterodimerization of S1P/EDG sphingosine-1-phosphate receptors. *Biochim. Biophys. Acta* **1582**, 89–93.
 121. Liu, Y., Wada, R., Yamashita, T., Mi, Y., Deng, C.X., Hobson, J.P., Rosenfeldt, H.M., Nava, V.E., Chae, S.S., Lee, M.J., Liu, C.H., Hla, T., Spiegel, S., and Proia, R.L. (2000) Edg-1, the G protein-coupled receptor for sphingosine-1-phosphate, is essential for vascular maturation. *J. Clin. Invest.* **106**, 951–961.
 122. Allende, M.L., Yamashita, T., and Proia, R.L. (2003) G-protein coupled receptor S1P₁ acts within endothelial cells to regulate vascular maturation. *Blood* **102**, 3665–3667.
 123. Chae, S.S., Paik, J.H., Allende, M.L., Proia, R.L., and Hla, T. (2004) Regulation of limb development by the sphingosine 1-phosphate receptor *S1p₁/EDG-1* occurs via the hypoxia/VEGF axis. *Dev. Biol.* **268**, 441–447.
 124. Mizugishi, K., Yamashita, T., Olivera, A., Miller, G.F., Spiegel, S., and Proia, R.L. (2005) Essential role for sphingosine kinases in neural and vascular development. *Mol. Cell. Biol.* **25**, 11113–11121.
 125. MacLennan, A.J., Carney, P.R., Zhu, W.J., Chaves, A.H., Garcia, J., Grimes, J.R., Anderson, K.J., Roper, S.N., and Lee, N. (2001) An essential role for the H218/AGR16/Edg-5/LPB₂ sphingosine 1-phosphate receptor in neuronal excitability. *Eur. J. Neurosci.* **14**, 203–209.
 126. Kono, M., Mi, Y., Liu, Y., Sasaki, T., Allende, M.L., Wu, Y.P., Yamashita, T., and Proia, R.L. (2004) The sphingosine-1-phosphate receptors S1P₁, S1P₂ and S1P₃ receptors coordinately function during embryonic angiogenesis. *J. Biol. Chem.* **279**, 29367–29373.
 127. Lee, M.J., Thangada, S., Claffey, K.P., Ancellin, N., Liu, C.H., Kluk, M., Volpi, M., Sha'afi, R.I., and Hla, T. (1999)

- Vascular endothelial cell adherens junction assembly and morphogenesis induced by sphingosine-1-phosphate. *Cell* **99**, 301–312.
128. Igarashi, J. and Michel, T. (2001) Sphingosine 1-phosphate and isoform-specific activation of phosphoinositide 3-kinase b: evidence for divergence and convergence of receptor-regulated endothelial nitric oxide synthase signaling pathways. *J. Biol. Chem.* **276**, 36281–36288.
 129. English, D., Kovala, A.T., Welch, Z., Harvey, K.A., Siddiqui, R.A., Brindley, D.N., and Garcia, J.G. (1999) Induction of endothelial cell chemotaxis by sphingosine 1-phosphate and stabilization of endothelial monolayer barrier function by lysophosphatidic acid, potential mediators of hematopoietic angiogenesis. *J. Hematother. Stem Cell Res.* **8**, 627–634.
 130. Paik, J.H., Chae, S.S., Lee, M.J., Thangada, S., and Hla, T. (2001) Sphingosine-1-phosphate induced endothelial cell migration requires the expression of EDG-1 and EDG-3 receptors and Rho-dependent activation of $\alpha_v\beta_3$ and β_1 -containing integrins. *J. Biol. Chem.* **276**, 11830–11837.
 131. Liu, F., Verin, A.D., Wang, P., Day, R., Wersto, R.P., Chrest, F.J., English, D.K., and Garcia, J.G. (2001) Differential regulation of sphingosine-1-phosphate- and VEGF-induced endothelial cell chemotaxis. involvement of G_{i2} -linked rho kinase activity. *Am. J. Respir. Cell. Mol. Biol.* **24**, 711–719.
 132. Lee, M., Thangada, S., Paik, J., Sapkota, G.P., Ancellin, N., Chae, S., Wu, M., Morales-Ruiz, M., Sessa, W.C., Alessi, D.R., and Hla, T. (2001) Akt-mediated phosphorylation of the g protein-coupled receptor edg-1 is required for endothelial cell chemotaxis. *Mol. Cell* **8**, 693–704.
 133. Lee, J.F., Ozaki, H., Zhan, X., Wang, E., Hla, T., and Lee, M.J. (2006) Sphingosine-1-phosphate signaling regulates lamellipodia localization of cortactin complexes in endothelial cells. *Histochem. Cell. Biol.* 1–8. [E-pub]
 134. Rikitake, Y., Hirata, K., Kawashima, S., Ozaki, M., Takahashi, T., Ogawa, W., Inoue, N., and Yokoyama, M. (2002) Involvement of endothelial nitric oxide in sphingosine-1-phosphate-induced angiogenesis. *Arterioscler. Thromb. Vasc. Biol.* **22**, 108–114.
 135. Igarashi, J., Erwin, P.A., Dantas, A.P., Chen, H., and Michel, T. (2003) VEGF induces S1P₁ receptors in endothelial cells: implications for cross-talk between sphingolipid and growth factor receptors. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 10664–10669.
 136. Tanimoto, T., Jin, Z.G., and Berk, B.C. (2002) Transactivation of vascular endothelial growth factor (VEGF) receptor Flk-1/KDR is involved in sphingosine 1-phosphate-stimulated phosphorylation of Akt and endothelial nitric-oxide synthase (eNOS). *J. Biol. Chem.* **277**, 42997–43001.
 137. Endo, A., Nagashima, K.I., Kurose, H., Mochizuki, S., Matsuda, M., and Mochizuki, N. (2002) Sphingosine 1-phosphate induces membrane ruffling and increases motility of human umbilical vein endothelial cells via vascular endothelial growth factor receptor and CrkII. *J. Biol. Chem.* **277**, 23747–23754.
 138. Krump-Konvalinkova, V., Yasuda, S., Rubic, T., Makarova, N., Mages, J., Erl, W., Vosseler, C., Kirkpatrick, C.J., Tigyi, G., and Siess, W. (2005) Stable knock-down of the sphingosine 1-phosphate receptor S1P₁ influences multiple functions of human endothelial cells. *Arterioscler. Thromb. Vasc. Biol.* **25**, 546–552.
 139. Licht, T., Tsurulnikov, L., Reuveni, H., Yarnitzky, T., and Ben-Sasson, S.A. (2003) Induction of pro-angiogenic signaling by a synthetic peptide derived from the 2nd intracellular loop of S1P₃(EDG3). *Blood* **102**, 2099–2107.
 140. Kimura, T., Sato, K., Malchinkhuu, E., Tomura, H., Tamama, K., Kuwabara, A., Murakami, M., and Okajima, F. (2003) High-density lipoprotein stimulates endothelial cell migration and survival through sphingosine 1-phosphate and its receptors. *Arterioscler. Thromb. Vasc. Biol.* **23**, 1283–1288.
 141. Inoki, I., Takuwa, N., Sugimoto, N., Yoshioka, K., Takata, S., Kaneko, S., and Takuwa, Y. (2006) Negative regulation of endothelial morphogenesis and angiogenesis by S1P₂ receptor. *Biochem. Biophys. Res. Commun.* **346**(1), 293–300.
 142. Bischoff, A., Czyborra, P., Meyer zu Heringdorf, D., Jakobs, K.H., and Michel, M.C. (2000) Sphingosine-1-phosphate reduces rat renal and mesenteric blood flow in vivo in a pertussis toxin-sensitive manner. *Br. J. Pharmacol.* **130**, 1878–1883.
 143. Alewijnse, A.E., Peters, S.L., and Michel, M.C. (2004) Cardiovascular effects of sphingosine-1-phosphate and other sphingomyelin metabolites. *Br. J. Pharmacol.* **143**, 666–684.
 144. Hemmings, D.G. (2006) Signal transduction underlying the vascular effects of sphingosine 1-phosphate and sphingosylphosphorylcholine. *Naunyn Schmiedebergs Arch. Pharmacol.* **373**, 18–29.
 145. Brinkmann, V. and Baumruker, T. (2006) Pulmonary and vascular pharmacology of sphingosine 1-phosphate. *Curr. Opin. Pharmacol.* **6**, 244–250.
 146. Bischoff, A., Czyborra, P., Fetscher, C., Meyer zu Heringdorf, D., Jakobs, K.H., and Michel, M.C. (2000) Sphingosine-1-phosphate and sphingosylphosphorylcholine constrict renal and mesenteric microvessels in vitro. *Br. J. Pharmacol.* **130**, 1871–1877.
 147. Tosaka, M., Okajima, F., Hashiba, Y., Saito, N., Nagano, T., Watanabe, T., Kimura, T., and Sasaki, T. (2001) Sphingosine 1-phosphate contracts canine basilar arteries in vitro and in vivo: possible role in pathogenesis of cerebral vasospasm. *Stroke* **32**, 2913–2919.
 148. Scherer, E.Q., Lidington, D., Oestreicher, E., Arnold, W., Pohl, U., and Bolz, S.S. (2006) Sphingosine-1-phosphate modulates spiral modiolar artery tone: A potential role in vascular-based inner ear pathologies? *Cardiovasc. Res.* **70**, 79–87.
 149. Hsiao, S.H., Constable, P.D., Smith, G.W., and Haschek, W.M. (2005) Effects of exogenous sphinganine,

- sphingosine, and sphingosine-1-phosphate on relaxation and contraction of porcine thoracic aortic and pulmonary arterial rings. *Toxicol. Sci.* **86**, 194–199.
150. Hemmings, D.G., Hudson, N.K., Halliday, D., O'Hara, M., Baker, P.N., Davidge, S.T., and Taggart, M.J. (2006) Sphingosine-1-phosphate acts via Rho-associated kinase and nitric oxide to regulate human placental vascular tone. *Biol. Reprod.* **74**, 88–94.
 151. Coussin, F., Scott, R.H., Wise, A., and Nixon, G.F. (2002) Comparison of sphingosine 1-phosphate-induced intracellular signaling pathways in vascular smooth muscles: differential role in vasoconstriction. *Circ. Res.* **91**, 151–157.
 152. Ohmori, T., Yatomi, Y., Osada, M., Kazama, F., Takafuta, T., Ikeda, H., and Ozaki, Y. (2003) Sphingosine 1-phosphate induces contraction of coronary artery smooth muscle cells via S1P₂. *Cardiovasc. Res.* **58**, 170–177.
 153. Ikeda, H., Nagashima, K., Yanase, M., Tomiya, T., Arai, M., Inoue, Y., Tejima, K., Nishikawa, T., Watanabe, N., Omata, M., and Fujiwara, K. (2004) Sphingosine 1-phosphate enhances portal pressure in isolated perfused liver via S1P₂ with Rho activation. *Biochem. Biophys. Res. Commun.* **320**, 754–759.
 154. Damirin, A., Tomura, H., Komachi, M., Tobo, M., Sato, K., Mogi, C., Nochi, H., Tamoto, K., and Okajima, F. (2005) Sphingosine-1-phosphate receptors mediate the lipid-induced cAMP accumulation through cyclooxygenase-2/prostaglandin I₂ pathway in human coronary artery smooth muscle cells. *Mol. Pharmacol.* **67**, 1177–1185.
 155. Davidge, S.T. (2001) Prostaglandin H synthase and vascular function. *Circ. Res.* **89**, 650–660.
 156. Tolle, M., Levkau, B., Keul, P., Brinkmann, V., Giebing, G., Schonfelder, G., Schafers, M., von Wnuck Lipinski, K., Jankowski, J., Jankowski, V., Chun, J., Zidek, W., and Van der Giet, M. (2005) Immunomodulator FTY720 induces eNOS-dependent arterial vasodilatation via the lysophospholipid receptor S1P₃. *Circ. Res.* **96**, 913–920.
 157. Nofer, J.R., Van Der Giet, M., Tolle, M., Wolinska, I., Von Wnuck Lipinski, K., Baba, H.A., Tietge, U.J., Godecke, A., Ishii, I., Kleuser, B., Schafers, M., Fobker, M., Zidek, W., Assmann, G., Chun, J., and Levkau, B. (2004) HDL induces NO-dependent vasorelaxation via the lysophospholipid receptor S1P₃. *J. Clin. Invest.* **113**, 569–581.
 158. Hemmings, D.G., Xu, Y., and Davidge, S.T. (2004) Sphingosine 1-phosphate-induced vasoconstriction is elevated in mesenteric resistance arteries from aged female rats. *Br. J. Pharmacol.* **143**, 276–284.
 159. Garcia, J.G., Liu, F., Verin, A.D., Birukova, A., Dechert, M.A., Gerthoffer, W.T., Bamberg, J.R., and English, D. (2001) Sphingosine 1-phosphate promotes endothelial cell barrier integrity by Edg-dependent cytoskeletal rearrangement. *J. Clin. Invest.* **108**, 689–701.
 160. Singleton, P.A., Dudek, S.M., Chiang, E.T., and Garcia, J.G. (2005) Regulation of sphingosine 1-phosphate-induced endothelial cytoskeletal rearrangement and barrier enhancement by S1P₁ receptor, PI3 kinase, Tiam1/Rac1, and a-actinin. *FASEB J.* **19**, 1646–1656.
 161. Finigan, J.H., Dudek, S.M., Singleton, P.A., Chiang, E.T., Jacobson, J.R., Camp, S.M., Ye, S.Q., and Garcia, J.G. (2005) Activated protein C mediates novel lung endothelial barrier enhancement. Role of sphingosine 1-phosphate receptor transactivation. *J. Biol. Chem.* **280**, 17286–17293.
 162. Gon, Y., Wood, M.R., Kiesses, W.B., Jo, E., Sanna, M.G., Chun, J., and Rosen, H. (2005) S1P₃ receptor-induced reorganization of epithelial tight junctions compromises lung barrier integrity and is potentiated by TNF. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 9270–9275.
 163. Sugiyama, A., Aye, N.N., Yatomi, Y., Ozaki, Y., and Hashimoto, K. (2000) Effects of sphingosine 1-phosphate, a naturally occurring biologically active lysophospholipid, on the rat cardiovascular system. *Jpn. J. Pharmacol.* **82**, 338–342.
 164. Sanna, M.G., Liao, J., Jo, E., Alfonso, C., Ahn, M.Y., Peterson, M.S., Webb, B., Lefebvre, S., Chun, J., Gray, N., and Rosen, H. (2004) S1P receptor subtypes S1P₁ and S1P₃, respectively, regulate lymphocyte recirculation and heart rate. *J. Biol. Chem.* **279**, 13839–13848.
 165. Forrest, M., Sun, S.Y., Hajdu, R., Bergstrom, J., Card, D., Hale, J., Keohane, C.A., Meyers, C., Milligan, J., Mills, S., Nomura, N., Rosenbach, M., Shei, G.J., Singer, I.I., Tian, M., West, S., White, V., Xie, J., Rosen, H., Proia, R., Doherty, G., and Mandala, S. (2004) Immune cell regulation and cardiovascular effects of sphingosine 1-phosphate receptor agonists in rodents are mediated via distinct receptor sub-types. *J. Pharmacol. Exp. Ther.* **309**, 758–768.
 166. Goetzl, E.J., Wang, W., McGiffert, C., Huang, M.C., and Graler, M.H. (2004) Sphingosine 1-phosphate and its G protein-coupled receptors constitute a multifunctional immunoregulatory system. *J. Cell. Biochem.* **92**, 1104–1114.
 167. Goetzl, E.J. and Rosen, H. (2004) Regulation of immunity by lysosphingolipids and their G protein-coupled receptors. *J. Clin. Invest.* **114**, 1531–1537.
 168. Chiba, K. (2005) FTY720, a new class of immunomodulator, inhibits lymphocyte egress from secondary lymphoid tissues and thymus by agonistic activity at sphingosine 1-phosphate receptors. *Pharmacol. Ther.* **108**, 308–319.
 169. Cyster, J.G. (2005) Chemokines, sphingosine-1-phosphate, and cell migration in secondary lymphoid organs. *Annu. Rev. Immunol.* **23**, 127–159.
 170. Rosen, H. and Goetzl, E.J. (2005) Sphingosine 1-phosphate and its receptors: an autocrine and paracrine network. *Nat. Rev. Immunol.* **5**, 560–570.
 171. Brinkmann, V., Davis, M.D., Heise, C.E., Albert, R., Cottens, S., Hof, R., Bruns, C., Prieschl, E., Baumruker, T., Hiestand, P., Foster, C.A., Zollinger, M., and Lynch, K.R. (2002) The immune modulator FTY720 targets sphingosine 1-phosphate receptors. *J. Biol. Chem.* **277**, 21453–21457.
 172. Gräler, M.H. and Goetzl, E.J. (2004) The immunosuppressant FTY720 down-regulates sphingosine 1-phosphate G protein-coupled receptors. *FASEB J.* **18**, 551–553.

173. Matloubian, M., Lo, C.G., Cinamon, G., Lesneski, M.J., Xu, Y., Brinkmann, V., Allende, M.L., Proia, R.L. and Cyster, J.G. (2004) Lymphocyte egress from thymus and peripheral lymphoid organs is dependent on S1P receptor 1. *Nature* **427**, 355–360.
174. Allende, M.L., Dreier, J.L., Mandala, S., and Proia, R.L. (2004) Expression of the sphingosine-1-phosphate receptor, S1P₁, on T-cells controls thymic emigration. *J. Biol. Chem.* **279**, 15396–15401.
175. Cinamon, G., Matloubian, M., Lesneski, M.J., Xu, Y., Low, C., Lu, T., Proia, R.L., and Cyster, J.G. (2004) Sphingosine 1-phosphate receptor 1 promotes B cell localization in the splenic marginal zone. *Nat. Immunol.* **5**, 713–720.
176. Vora, K.A., Nichols, E., Porter, G., Cui, Y., Keohane, C.A., Hajdu, R., Hale, J., Neway, W., Zaller, D., and Mandala, S. (2005) Sphingosine 1-phosphate receptor agonist FTY720-phosphate causes marginal zone B cell displacement. *J. Leukoc. Biol.* **78**, 471–480.
177. Graeler, M., Shankar, G., and Goetzl, E.J. (2002) Cutting edge: suppression of T cell chemotaxis by sphingosine 1-phosphate. *J. Immunol.* **169**, 4084–4087.
178. Lo, C.G., Xu, Y., Proia, R.L., and Cyster, J.G. (2005) Cyclical modulation of sphingosine-1-phosphate receptor 1 surface expression during lymphocyte recirculation and relationship to lymphoid organ transit. *J. Exp. Med.* **201**, 291–301.
179. Fieger, C.B., Huang, M.-C., Van Brocklyn, J.R., and Goetzl, E.J. (2005) Type 1 sphingosine 1-phosphate G protein-coupled receptor signaling of lymphocyte functions requires sulfation of its extracellular amino-terminal tyrosines. *FASEB J.* **19**, 1926–1928.
180. Shiow, L.R., Rosen, D.B., Brdickova, N., Xu, Y., An, J., Lanier, L.L., Cyster, J.G., and Matloubian, M. (2006) CD69 acts downstream of interferon- α/β to inhibit S1P₁ and lymphocyte egress from lymphoid organs. *Nature* **440**, 540–544.
181. Singer, I.I., Tian, M., Wickham, L.A., Lin, J., Matheravidathu, S.S., Forrest, M.J., Mandala, S., and Quackenbush, E.J. (2005) Sphingosine-1-phosphate agonists increase macrophage homing, lymphocyte contacts, and endothelial junctional complex formation in murine lymph nodes. *J. Immunol.* **175**, 7151–7161.
182. Roviezzo, F., Del Galdo, F., Abbate, G., Bucci, M., D'Agostino, B., Antunes, E., De Dominicis, G., Parente, L., Rossi, F., Cirino, G., and De Palma, R. (2004) Human eosinophil chemotaxis and selective *in vivo* recruitment by sphingosine 1-phosphate. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 11170–11175.
183. Czeloth, N., Bernhardt, G., Hofmann, F., Genth, H., and Forster, R. (2005) Sphingosine-1-phosphate mediates migration of mature dendritic cells. *J. Immunol.* **175**, 2960–2967.
184. Lan, Y.Y., De Creus, A., Colvin, B.L., Abe, M., Brinkmann, V., Coates, P.T., and Thomson, A.W. (2005) The sphingosine-1-phosphate receptor agonist FTY720 modulates dendritic cell trafficking *in vivo*. *Am. J. Transplant.* **5**, 2649–2659.
185. Seitz, G., Boehmler, A.M., Kanz, L., and Mohle, R. (2005) The role of sphingosine 1-phosphate receptors in the trafficking of hematopoietic progenitor cells. *Ann. N Y Acad. Sci.* **1044**, 84–89.
186. Wang, W., Graeler, M.H., and Goetzl, E.J. (2005) Type 4 sphingosine 1-phosphate G protein-coupled receptor (S1P₄) transduces S1P effects on T cell proliferation and cytokine secretion without signaling migration. *FASEB J.* **19**, 1731–1733.
187. Oz-Arslan, D., Ruscher, W., Myrtek, D., Ziemer, M., Jin, Y., Damaj, B.B., Sorichter, S., Idzko, M., Norgauer, J., and Maghazachi, A.A. (2006) IL-6 and IL-8 release is mediated via multiple signaling pathways after stimulating dendritic cells with lysophospholipids. *J. Leukoc. Biol.* **80**(2), 287–297.
188. Idzko, M., Panther, E., Corinti, S., Morelli, A., Ferrari, D., Herouy, Y., Dichmann, S., Mockenhaupt, M., Gebicke-Haerter, P., Di Virgilio, F., Girolomoni, G., and Norgauer, J. (2002) Sphingosine 1-phosphate induces chemotaxis of immature dendritic cells and modulates cytokine-release in mature human dendritic cells for emergence of Th2 immune responses. *FASEB J.* **16**, 625–627.
189. Payne, S.G., Milstien, S., Barbour, S.E., and Spiegel, S. (2004) Modulation of adaptive immune responses by sphingosine-1-phosphate. *Semin. Cell Dev. Biol.* **15**, 521–527.
190. Jolly, P.S., Bektas, M., Olivera, A., Gonzalez-Espinosa, C., Proia, R.L., Rivera, J., Milstien, S., and Spiegel, S. (2004) Transactivation of sphingosine-1-phosphate receptors by Fc ϵ RI triggering is required for normal mast cell degranulation and chemotaxis. *J. Exp. Med.* **199**, 959–970.
191. Olivera, A., Urtz, N., Mizugishi, K., Yamashita, Y., Gilfillan, A.M., Furumoto, Y., Gu, H., Proia, R.L., Baumruker, T., and Rivera, J. (2006) IgE-dependent activation of sphingosine kinases 1 and 2 and secretion of sphingosine 1-phosphate requires Fyn kinase and contributes to mast cell responses. *J. Biol. Chem.* **281**, 2515–2525.
192. Tedesco-Silva, H., Mourad, G., Kahan, B.D., Boira, J.G., Weimar, W., Mulgaonkar, S., Nashan, B., Madsen, S., Charpentier, B., Pellet, P., and Vanrenterghem, Y. (2005) FTY720, a novel immunomodulator: efficacy and safety results from the first phase 2A study in de novo renal transplantation. *Transplantation* **79**, 1553–1560.
193. Boehler, T., Schuetz, M., Budde, K., Neumayer, H., and Waiser, J. (2002) FTY720 alters the composition of T-lymphocyte subpopulations in the peripheral blood compartment of renal transplant patients. *Transplant. Proc.* **34**, 2242–2243.
194. Budde, K., Schmouder, R.L., Brunkhorst, R., Nashan, B., Lucker, P.W., Mayer, T., Choudhury, S., Skerjanec, A., Kraus, G., and Neumayer, H.H. (2002) First human trial of FTY720, a novel immunomodulator, in stable renal transplant patients. *J. Am. Soc. Nephrol.* **13**, 1073–1083.

195. Webb, M., Tham, C.S., Lin, F.F., Lariosa-Willingham, K., Yu, N., Hale, J., Mandala, S., Chun, J., and Rao, T.S. (2004) Sphingosine 1-phosphate receptor agonists attenuate relapsing-remitting experimental autoimmune encephalitis in SJL mice. *J. Neuroimmunol.* **153**, 108–121.
196. Xia, P., Gamble, J.R., Wang, L., Pitson, S.M., Moretti, P.A., Wattenberg, B.W., D'Andrea, R.J., and Vadas, M.A. (2000) An oncogenic role of sphingosine kinase. *Curr. Biol.* **10**, 1527–1530.
197. French, K.J., Schrecengost, R.S., Lee, B.D., Zhuang, Y., Smith, S.N., Eberly, J.L., Yun, J.K., and Smith, C.D. (2003) Discovery and evaluation of inhibitors of human sphingosine kinase. *Cancer Res.* **63**, 5962–2969.
198. Kawamori, T., Osta, W., Johnson, K.R., Pettus, B.J., Bielawski, J., Tanaka, T., Wargovich, M.J., Reddy, B.S., Hannun, Y.A., Obeid, L.M., and Zhou, D. (2006) Sphingosine kinase 1 is up-regulated in colon carcinogenesis. *FASEB J.* **20(2)**, 386–388.
199. Le Scolan, E., Pchejetski, D., Banno, Y., Denis, N., Mayeux, P., Vainchenker, W., Levade, T., and Moreau-Gachelin, F. (2005) Overexpression of sphingosine kinase 1 is an oncogenic event in erythroleukemic progression. *Blood* **106**, 1808–1816.
200. Nava, V.E., Hobson, J.P., Murthy, S., Milstien, S., and Spiegel, S. (2002) Sphingosine kinase type 1 promotes estrogen-dependent tumorigenesis of breast cancer mcf-7 cells. *Exp. Cell Res.* **281**, 115–127.
201. Sukocheva, O.A., Wang, L., Albanese, N., Vadas, M.A., and Xia, P. (2003) Sphingosine kinase transmits estrogen signaling in human breast cancer cells. *Mol. Endocrinol.* **17**, 2002–2012.
202. Bektas, M., Jolly, P.S., Muller, C., Eberle, J., Spiegel, S., and Geilen, C.C. (2005) Sphingosine kinase activity counteracts ceramide-mediated cell death in human melanoma cells: role of Bcl-2 expression. *Oncogene* **24**, 178–187.
203. Pchejetski, D., Golzio, M., Bonhoure, E., Calvet, C., Doumerc, N., Garcia, V., Mazerolles, C., Rischmann, P., Teissie, J., Malavaud, B., and Cuvillier, O. (2005) Sphingosine kinase-1 as a chemotherapy sensor in prostate adenocarcinoma cell and mouse models. *Cancer Res.* **65**, 11667–11675.
204. Arikawa, K., Takuwa, N., Yamaguchi, H., Sugimoto, N., Kitayama, J., Nagawa, H., Takehara, K., and Takuwa, Y. (2003) Ligand-dependent inhibition of B16 melanoma cell migration and invasion via endogenous S1P₂ G protein-coupled receptor requirement of inhibition of cellular Rac activity. *J. Biol. Chem.* **278**, 32841–32851.
205. Yamaguchi, H., Kitayama, J., Takuwa, N., Arikawa, K., Inoki, I., Takehara, K., Nagawa, H., and Takuwa, Y. (2003) Sphingosine-1-phosphate receptor subtype-specific positive and negative regulation of Rac and hematogenous metastasis of melanoma cells. *Biochem. J.* **374**, 715–722.
206. Van Brocklyn, J.R., Jackson, C.A., Pearl, D.K., Kotur, M.S., Snyder, P.J., and Prior, T.W. (2005) Sphingosine kinase-1 expression correlates with poor survival of patients with glioblastoma multiforme. Roles of sphingosine kinase isoforms in growth of glioblastoma cell lines. *J. Neuropathol. Exp. Neurol.* **64**, 695–705.
207. Anelli, V., Bassi, R., Tettamanti, G., Viani, P., and Riboni, L. (2005) Extracellular release of newly synthesized sphingosine-1-phosphate by cerebellar granule cells and astrocytes. *J. Neurochem.* **92**, 1204–1215.
208. Bassi, R., Anelli, V., Giussani, P., Tettamanti, G., Viani, P., and Riboni, L. (2006) Sphingosine-1-phosphate is released by cerebellar astrocytes in response to bFGF and induces astrocyte proliferation through G_i-protein-coupled receptors. *Glia* **53**, 621–630.
209. Edsall, L.C., Cuvillier, O., Twitty, S., Spiegel, S., and Milstien, S. (2001) Sphingosine kinase expression regulates apoptosis and caspase activation in PC12 cells. *J. Neurochem.* **76**, 1573–1584.
210. Malchinkhuu, E., Sato, K., Horiuchi, Y., Mogi, C., Ohwada, S., Ishiuchi, S., Saito, N., Kurose, H., Tomura, H., and Okajima, F. (2005) Role of p38 mitogen-activated kinase and c-Jun terminal kinase in migration response to lysophosphatidic acid and sphingosine-1-phosphate in glioma cells. *Oncogene* **24**, 6676–6688.
211. Lamontagne, K., Littlewood-Evans, A., Schnell, C., O'Reilly, T., Wyder, L., Sanchez, T., Probst, B., Butler, J., Wood, A., Liau, G., Billy, E., Theuer, A., Hla, T., and Wood, J. (2006) Antagonism of sphingosine-1-phosphate receptors by FTY720 inhibits angiogenesis and tumor vascularization. *Cancer Res.* **66**, 221–231.
212. Visentin, B., Vekich, J.A., Sibbald, B.J., Cavalli, A.L., Moreno, K.M., Matteo, R.G., Garland, W.A., Lu, Y., Yu, S., Hall, H.S., Kundra, V., Mills, G.B., and Sabbadini, R.A. (2006) Validation of an anti-sphingosine-1-phosphate antibody as a potential therapeutic in reducing growth, invasion, and angiogenesis in multiple tumor lineages. *Cancer Cell* **9**, 225–238.
213. Klabunde, T. and Hessler, G. (2002) Drug design strategies for targeting G-protein-coupled receptors. *Chembiochem* **3**, 928–944.

This article should be cited as follows:

Young, N. and Van Brocklyn, J.R. (2006) Signal transduction of sphingosine-1-phosphate G protein-coupled receptors. *TheScientificWorldJOURNAL* **6**, 946–966. DOI 10.1100/tsw.2006.182.