

Research Article

Amniotic Fluid or Its Fatty Acids Produce Actions Similar to Diazepam on Lateral Septal Neurons Firing Rate

Ana G. Gutiérrez-García,^{1,2} Carlos M. Contreras,^{1,3} and Diana Idania Vásquez-Hernández¹

¹Laboratorio de Neurofarmacología, Instituto de Neuroetología, Universidad Veracruzana, Avenue Dr. Luis Castelazo S/N Col. Industrial Las Ánimas, 91190 Xalapa, VER, Mexico

²Facultad de Psicología, Universidad Veracruzana, Manantial San Cristóbal-Xalapa 2000, 91097 Xalapa, VER, Mexico

³Unidad Periférica Xalapa, Instituto de Investigaciones Biomédicas, Universidad Nacional Autónoma de México, 91190 Xalapa, VER, Mexico

Correspondence should be addressed to Carlos M. Contreras; ccontreras@uv.mx

Received 10 May 2013; Accepted 11 June 2013

Academic Editors: C. La Motta and J. Mazella

Copyright © 2013 Ana G. Gutiérrez-García et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Human amniotic fluid (AF) contains eight fatty acids (FATs), and both produce anxiolytic-like effects in adult rats and appetitive responses in human newborns. The medial amygdala and lateral septal nucleus function are related to social behavior, but the action of AF or its FATs in this circuit is known. We obtained 267 single-unit extracellular recordings in Wistar rats treated with vehicle (1 mL, s.c.; $n = 12$), human AF (1 mL, s.c.; $n = 12$), a FAT mixture (1 mL, s.c.; $n = 13$), diazepam (1 mg/kg, i.p.; $n = 11$), and fluoxetine (1 mg/kg, p.o.; $n = 12$). Compared with the vehicle group, the spontaneous septal firing rate in the AF, FAT mixture, and diazepam groups was the lowest and in the fluoxetine group the highest. Cumulative peristimulus histograms indicated that the significant change in septal firing occurred only in the AF and FAT mixture groups and exclusively in those neurons that increased their firing rate during amygdala stimulation. We conclude that human AF and its FATs produce actions comparable to anxiolytic drugs and are able to modify the responsiveness of a circuit involved in social behavior, suggesting facilitation of social recognition processes by maternal-fetal fluids.

1. Introduction

Newborn mammals, including humans, exhibit agitation and emit vocalizations when placed in an unfamiliar environment, but they calm down when they return to their nest or remain in close proximity to their mother [1, 2], an action likely related to odors that come from maternal fluids, such as amniotic fluid (AF), colostrum, and milk [3–5]. These observations open the possibility that the calming effect is an anxiolytic property of maternal fluids.

We recently reported that human AF and its fatty acids (FATs) produce anxiolytic-like effects in adult Wistar rats subjected to validated models of experimental anxiety [6]. In such a case, the fetus seemingly develops and grows in a comfortable medium that provides a protective anxiolytic action through an unknown process. Maternal fluids, such as AF,

colostrum, and milk, also produce orientating movements in some mammals [7–9], which include human newborns [10–14]. These three maternal-infant fluids that contain FATs are quite different in pigs [15] and humans [6]. Amniotic fluid or an artificial mixture of its FATs produces orientating-feeding responses in human newborns [16], illustrating early learning [17] through an unknown neural process.

The lateral septal nucleus contains γ -aminobutyric acid (GABA) and serotonin (5-hydroxytryptamine [5-HT]) terminals [18] and is part of a functional network related to social behavior processing [19]. Septal neurons increase their firing rate after the administration of fluoxetine [20], among other clinically effective antidepressant drugs [21], and decrease their firing rate after the administration of benzodiazepines [22, 23]. Anxiolytic GABAergic drugs produce a low septal firing rate, whereas serotonergic drugs

with anxiolytic actions produce a high neuronal firing rate in the lateral septal nucleus. Diazepam exerts anxiolytic effects through its well-known action on GABA_A receptors [24]. Fluoxetine is an antidepressant that selectively inhibits serotonin reuptake to also produce anxiolytic effects [25]. Therefore, these two drugs may be useful tools for exploring the anxiolytic effects of AF or its FATs on neuronal activity.

Any possible action of AF and its FATs on the neuronal activity of lateral septal neurons has not been explored. The aim of the present study was to compare the effects of AF and its FATs against diazepam and fluoxetine on the neuronal firing rate of lateral septal neurons in response to medial amygdala stimulation.

2. Materials and Methods

2.1. Ethics. For the human samples, we strictly followed international principles of confidentiality and healthcare, such as the Declaration of Helsinki, and all of the procedures in rats followed the principles of animal care based on the Guide for the Care and Use of Laboratory Animals [26]. Both protocols received authorization from the Biomedical Research Institute Ethical Committee (National Autonomous University of México).

2.2. Human Samples. All of the volunteers were given a detailed explanation of the purpose and risks of the study and signed an informed consent prior to inclusion in the study. Volunteers were invited to participate only as donors of AF, with no risk to the mother or baby. Both the mother and newborn were in optimal health, reflected by a general clinical evaluation of the mothers and proper scales for newborns. Amniotic fluid was obtained under sterile conditions by an obstetrics surgeon. For vaginal deliveries, after the amniotic membrane ruptured, approximately 5 mL of AF was collected in a sterile receptacle. For caesarean deliveries, the surgeon collected the AF with a sterile syringe (5 mL, 23 gauge). A total sample of approximately 50 mL of AF was collected from 11 healthy volunteers (15–30 years old; 37–42 weeks gestational age; first gestation, $n = 4$; vaginal delivery, $n = 5$).

2.3. Experimental Groups. Given that spontaneous lateral septal neuronal firing varies during the estrous cycle in Wistar rats [27], the present study included 60 male rats that were randomly assigned to five experimental groups. The single administration or last administration of the treatments preceded anesthesia for the surgical procedures by 30 min. After another 60 min, the first single-unit extracellular recording was obtained.

Fresh human AF (1 mL/rat, $n = 12$) was injected subcutaneously after being filtered with filter paper (no. 4, 110 mm diameter; Whatman International, Maidstone, UK).

A fatty acid mixture (1 mL/rat, $n = 13$) was injected subcutaneously. The selection of the concentrations of the FATs contained in the artificial FAT mixture was based on previous reports [6, 16, 17]. The FAT mixture group received a mixture of lauric acid (0.4 mg), myristic acid (3.0 mg), palmitic acid (15.3 mg), palmitoleic acid (7.1 mg), stearic acid (3.7 mg), oleic

acid (8.0 mg), elaidic acid (1.5 mg), and linoleic acid (4.4 mg) dissolved in 100 mL of vehicle (96% propylene glycol and 4% ethanol) at a temperature of 37–40°C. All of the FATs were analytical grade and obtained from Sigma-Aldrich (St. Louis, MO, USA).

The study included one vehicle group ($n = 12$) that received a single subcutaneous injection of the FAT mixture solvent. One active drug control group ($n = 11$) received a single dose of diazepam (Valium, 1 mg/kg; Roche, Toluca, México) dissolved in isotonic saline (0.9%) and injected intraperitoneally in a volume of 0.3 mL/rat. A second active drug control group ($n = 12$) received fluoxetine (Prozac, 1 mg/kg; Eli Lilly de México, Tlalpan, México) dissolved in isotonic saline (0.9%) and administered orally in a volume of 1.0 mL/kg daily for 21 days [20].

2.4. Animals and Housing. Wistar rats from a local strain initially supplied by Harlan (México City, México), approximately 3 months old and weighing 250–300 g, were included in the study after being housed in local facilities at a mean temperature of 25°C with a 12 h/12 h light/dark cycle (lights on 7:00 a.m.). They were housed five to six rats per cage in acrylic boxes (44 cm width × 30 cm length × 20 cm height) with free access to food (Teklad Lab Animal Diets, Harlan) and purified water. All of the experimental procedures were performed during the light period, beginning at 9:00 a.m.

2.5. Single-Unit Extracellular Recordings

2.5.1. Stereotaxic Surgery. Thirty minutes after the last injection of any treatment, the rats were profoundly anesthetized with ethyl carbamate (1 g/kg urethane, intraperitoneally; Sigma Chemical, St. Louis, MO, USA), and their head was fixed in a stereotaxic frame (Stoelting, Wood Dale, IL, USA) to proceed with surgery. Cardiac pulse and a parietal cortical surface electroencephalogram were continuously monitored on a polygraph (GRASS 79, Grass Instruments, Quincy, MA, USA). During recording, we added one-tenth of the initial dose of urethane upon detecting signs of alertness, such as respiratory acceleration, movements of the vibrissae, or blinking, or sudden changes in cardiac pulse. A midline incision uncovered the skull. Through a small trephination, we lowered a glass micropipette filled with 1 M NaCl (4–5 M Ω) using a hydraulic micromanipulator (Trent Wells, South Gate, CA, USA) toward the lateral septal nucleus (coordinates: anterior/posterior, 0.2 mm; lateral, 0.5 mm; dorsal/ventral, from –3.0 to –5.0 mm) [28]. Another trephination was made at coordinates that correspond to the medial amygdala (anterior/posterior, 2.8 mm; lateral, 3.3 mm; ventral, –8.6 mm), where a stainless-steel bipolar electrode was placed (~100 k Ω resistance, 1 mm insulation uncovered at the inner tip, 100 μ m diameter).

2.5.2. Single-Unit Extracellular Recordings. The micropipette signal was connected in series to a 7P511L Grass amplifier (Quincy, MA, USA; bandwidth pass filters: 300 Hz–3 KHz) and an oscilloscope (model 5111A, Tektronix, Beaverton, OR, USA) that received a filtered signal free from background




FIGURE 1: Location of electrodes. (a) Coronal section of the brain (Nissl technique) that shows the recording electrode tips in the dorsal aspect of the lateral septal nucleus (arrow). (b) Representative schematic of a coronal section (+0.2 mm) from a rat that illustrates the marks left by the recording electrode. (c) Coronal section (Nissl technique) that shows the marks left by the stimulation electrode in the medial amygdala (arrow). (d) Representative schematic of a coronal section (-2.8 mm) from a rat that illustrates the marks left by the stimulation electrode. CPu: striatum; LV: lateral ventricle; C: corpus callosum; Pir: piriform cortex.

noise through a window discriminator and in parallel to an audio amplifier. The absence of sudden changes in the amplitude of the firing rate over 300 s verified a stable recording. Afterward, each spike detected by the amplifier was fed to a Grass S88 stimulator (Quincy, MA, USA) that delivered a spike-corresponding square pulse of constant amplitude and duration (4 V, 0.6 ms) directed to a CED 1401 interface system (Cambridge Electronic Design, Cambridge, UK). The basal activity of lateral septal neurons was recorded

for 1 min, followed by 1 min of amygdala stimulation (20 stimuli, monophasic pulses, 0.3 Hz, 0.6 ms duration). The signals were processed (bin time, 1 ms) by Spike2 software, version 5.20, that delivered the mean and standard error of the firing rate (c/10 s) of spontaneous activity and generated 1000 ms base peristimulus histograms.

2.5.3. Histological Analysis. To mark the last recorded point in the single-unit extracellular recording of the lateral septal

nuclei, we first passed a direct current (1 min each polarity) through the recording micropipette. The stimulating electrodes were marked with electrolytic lesions (0.5 mA for 30 s for each polarity). Afterward, the rats received a lethal overdose of pentobarbital and were intracardially perfused with 200 mL of 0.9% saline solution, followed by 200 mL of 30% formaldehyde. The brains were removed and postfixed with 30% formaldehyde for 72 h. The tissues were cryoprotected with 30% sucrose for 24 h, frozen at -20°C , cut into $40\ \mu\text{m}$ -thick sections with a cryocut microtome (Leica-Jung, Nussloch, Germany), and dyed using the Nissl technique to reconstruct the path followed by the micropipette and stimulation electrodes with the aid of stereotaxic coordinates [28].

2.6. Data Collection and Statistical Analysis. We analyzed (SigmaStat version 3.5) the baseline lateral septal nucleus firing rate under basal conditions for 1 min before amygdala stimulation. One-way analysis of variance (ANOVA) was used to compare the overall effects of the treatments (vehicle, AF, FAT mixture, diazepam, and fluoxetine), followed by the Student-Neuman-Keuls (SNK) *post hoc* test. Values of $P \leq 0.05$ were considered statistically significant.

Based on the peristimulus histograms (1000 ms), we formed three groups of septal neurons: neurons that increased (\uparrow cells) or decreased (\downarrow cells) their firing rate or had no response (\emptyset cells) to amygdala stimulation. The criterion of change was based on a poststimulus firing rate difference that was higher or lower than the mean value of the prestimulus firing rate ± 1 standard deviation. After classifying neuronal activity, a complete database was constructed with all of the recorded neurons. Therefore, the cumulative peristimulus histograms and their corresponding statistics included all of the recorded neurons, classified according to their response to amygdala stimulation. For the analysis of these data, the prestimulus data were considered the basal firing rate, and the percentage of change in neuronal firing rate during amygdala stimulation was calculated and subjected to two-way ANOVA (SigmaStat, version 3.5), with the factors treatment (vehicle, AF, FAT mixture, diazepam, and fluoxetine) and type of response (\uparrow cells, \downarrow cells, and \emptyset cells), followed by the SNK *post hoc* test. Values of $P \leq 0.05$ were considered statistically significant. The data are expressed as mean \pm standard error of the mean.

3. Results

3.1. Histological Control. A total of 267 single-unit extracellular recordings were obtained from the lateral septal nucleus. The distribution of the recordings included 56 neurons in the vehicle group ($n = 12$ rats), 59 neurons in the fluoxetine group ($n = 12$ rats), 52 neurons in the diazepam group ($n = 11$ rats), 53 neurons in the FAT mixture group ($n = 13$ rats), and 47 neurons in the human AF group ($n = 12$ rats). The histological analysis allowed us to determine that the neuronal recordings were obtained from the dorsal aspect (3.0–4.2 mm beneath the cerebral cortex) of the lateral septal nucleus (Figure 1). We did not find significant differences




FIGURE 2: Overall effects of treatments. The lateral septal nucleus spontaneous neuronal firing rate is shown in the groups treated with vehicle (VEH), fluoxetine (Flx), diazepam (DZP), the fatty acid mixture (FAT), and fresh amniotic fluid (AF). The diazepam, fatty acid mixture, and fresh human amniotic fluid groups had the lowest values, and the fluoxetine group had the highest value compared with vehicle ($P < 0.05$, SNK).

between the depth of recordings (range: from 3.6 ± 0.05 mm to 3.7 ± 0.04 mm below the cortex) among groups ($F_{4,262} = 0.621$, $P = 0.648$).

3.2. Lateral Septal Nucleus Firing Rate Prior to Amygdala Stimulation. The baseline lateral septal nucleus neuronal firing rate (Figure 2) was significantly different between treatment groups ($F_{4,262} = 27.232$, $P < 0.001$). The SNK *post hoc* test revealed that the highest ($P < 0.05$) firing rate was observed in the fluoxetine group and that the lowest ($P < 0.05$) firing rates were observed in the FAT mixture, human AF, and diazepam groups, with no significant difference between them. The septal neuronal firing rate in the vehicle group was intermediate.

3.3. Classification Based on Septal Response to Amygdala Stimulation

3.3.1. Response Distribution. Table 1 shows the percent distribution of lateral septal neurons according to their type of response to electrical stimulation of the medial amygdala (base 1000 ms). Some variations were observed between groups in the distribution of the three types of responses.

3.4. Peristimulus Histograms. Figure 3 illustrates the cumulative peristimulus histograms of all septal \uparrow cells. The qualitative analysis indicated that diazepam ($n = 18$) reduced the septal firing rate compared with the vehicle group ($n = 24$). Fluoxetine ($n = 19$) produced the opposite action. In the period prior to amygdala stimulation, the septal neuronal firing rate was similar among the FAT mixture ($n = 28$), AF ($n = 22$), and diazepam ($n = 18$) groups. However, the increased firing rate in response to amygdala stimulation was higher in the AF and FAT mixture groups than in the vehicle group.

The cumulative peristimulus histogram of \downarrow cells is shown in Figure 4. The qualitative analysis of the period prior

TABLE 1: Lateral septal neuron distribution according to type of response to electrical stimulation of the medial amygdala.

	Control (<i>n</i> = 56)	Fluoxetine (<i>n</i> = 59)	Diazepam (<i>n</i> = 52)	Fatty acid mixture (<i>n</i> = 53)	Amniotic fluid (<i>n</i> = 47)
↑ Cells	42.8% (<i>n</i> = 24)	32.2% (<i>n</i> = 19)	34.6% (<i>n</i> = 18)	52.8% (<i>n</i> = 28)	46.8% (<i>n</i> = 22)
↓ Cells	21.4% (<i>n</i> = 12)	16.9% (<i>n</i> = 10)	26.9% (<i>n</i> = 14)	16.9% (<i>n</i> = 9)	31.9% (<i>n</i> = 15)
∅ Cells	35.7% (<i>n</i> = 20)	50.8% (<i>n</i> = 30)	38.4% (<i>n</i> = 20)	30.1% (<i>n</i> = 16)	21.2% (<i>n</i> = 10)

↑ Cells: cells with increased neuronal firing; ∅ Cells: cells with no change in neuronal firing; ↓ Cells: cells with decreased neuronal firing; *n*: number of cells recorded.




FIGURE 3: Cumulative peristimulus histograms. Septal neurons that responded with an increased firing rate (↑ cells) during amygdala stimulation are shown. Notice the qualitative change in firing in the FAT mixture and fresh human AF groups. Each histogram contains the mean firing rate (ordinate) from all recorded cells, 1000 ms before and 1000 ms after amygdala stimulation (abscissas).




FIGURE 4: Cumulative peristimulus histogram for septal neurons that responded with a decreased firing rate after amygdala stimulation (\downarrow cells). The response was similar among treatments. The coordinates are the same as in Figure 3.

to amygdala stimulation showed similarities between the vehicle ($n = 12$) and fluoxetine ($n = 10$) groups. The other three groups (diazepam, $n = 14$; FAT mixture, $n = 9$; AF, $n = 15$) fired at a similarly lower frequency.

As expected, \emptyset cells did not respond to amygdala stimulation. The fluoxetine group ($n = 30$) qualitatively fired at the highest rate, whereas the diazepam ($n = 20$) and FAT mixture ($n = 16$) groups fired at the lowest rate. The vehicle ($n = 20$) and AF ($n = 10$) groups fired at similar frequencies (Figure 5).

3.4.1. Percent Change in Firing Rate. The two-way ANOVA indicated a significant effect of treatment ($F_{4,262} = 2.908$, $P <$

0.02). The highest statistically significant ($P \leq 0.05$, SNK; Figure 6(a)) change in firing rate was observed in the human AF and FAT mixture groups. Similar percent changes in septal neuronal firing rate were observed after amygdala stimulation in the vehicle, fluoxetine, and diazepam groups.

The analysis also indicated a significant effect of type of response ($F_{2,262} = 55.424$, $P < 0.001$). \uparrow cells displayed the highest change in firing rate ($P \leq 0.05$) compared with \downarrow cells and \emptyset cells (Figure 6(b)).

The interaction between factors was also significant ($F_{8,262} = 2.501$, $P < 0.01$), and the *post hoc* analysis revealed that the change in the firing rate of septal neurons




FIGURE 5: Cumulative peristimulus histogram for septal neurons that did not respond to amygdala stimulation (\emptyset cells). As expected, the differences were minimal and nonsignificant. The coordinates are the same as in Figure 3.

after amygdala stimulation reached statistical significance ($P \leq 0.05$, SNK) only for \uparrow cells and exclusively in the AF and FAT mixture groups and not in the other groups (Figure 6(c)).

4. Discussion

The present study investigated the effects of fresh human AF and its FATs on lateral septal neurons identified by their connection to the medial amygdala. Regardless of their response to medial amygdala stimulation, the FAT mixture

and AF treatments decreased the spontaneous lateral septal neuronal firing rate similar to diazepam, whereas fluoxetine produced the inverse effect (i.e., an increased firing rate). An unexpected result was that the septal neurons that were most sensitive to AF or its FATs were those that responded with an increased firing rate after amygdala stimulation (\uparrow cells), an effect not observed with the other treatments.

The components used to prepare the FAT mixture correspond to intermediate-length FATs (from C6 to C18), and most of them are polyunsaturated [29]. Amniotic fluid is the natural environment for the development of the fetus [30],




FIGURE 6: (a) Global percent change in the firing rate of lateral septal neurons in the different treatment groups. Only the FAT mixture and AF produced a significant change in firing rate after amygdala stimulation. (b) Percent change in neuronal firing rate in septal neurons based on the type of response. ↑ cells displayed the highest change in firing rate. (c) Only ↑ cells significantly changed their firing rate and only in the AF and FAT groups (* $P \leq 0.05$, SNK). See Figure 2 for abbreviations.

providing a safe environment [31] and seemingly exerting some anxiolytic/protective effects during early development, given the similar neural actions of AF, its FATs, and diazepam.

Fatty acids affect neuronal function by acting on the phospholipid membrane layer, modifying membrane tension, causing conformational changes in ion channels, and altering ionic conductance [32]. Fatty acids also exert neuronal effects, even after systemic administration [33], likely through a lipocalin superfamily [34] of protein transporters [15]. These proteins have also been identified in human AF [35], colostrum [36], and olfactory epithelia [37] that are connected to deep temporal lobe structures related to emotional processing [38], suggesting the existence of a complete receptor sensorial system for such maternal fluids.

In the present study, diazepam, AF, and its FATs decreased the firing rate of lateral septal neurons. Diazepam, through its actions on GABA_A receptors, produces hyperpolarization, with the participation of chloride channels [39]. Fatty acids produce mechanical actions on the conformation of GABA_A receptors [40], modifying the opening frequency of chloride channels [32, 41] and leading to neuronal hyperpolarization [32, 42]. This has been shown for oleic acid, one of the most abundant FATs found in AF [6, 16].

The spontaneous firing rate of lateral septal neurons decrease after a single forced swim session [43]. Several antidepressants produce an increase in firing rate [20, 44], which agrees with the present results. Such antidepressants exert their main actions on the serotonergic system [45],

and we detected opposite actions between fluoxetine and diazepam, AF, and the FAT mixture, suggesting minimal participation of 5-HT in the effects of AF and its FATs. The actions of diazepam, AF, and its FATs were similar, suggesting the involvement of GABA_A receptors, a neurotransmission system already identified in the septal nucleus [46] and sensitive to treatments that reduce immobility in the forced swim test and increase the neuronal firing rate of septal neurons [20, 47], seemingly representing a neuronal correlate of the struggling behavior represented by the effort to solve an unsolvable problem.

An unexpected result arose from the cumulative peristimulus histogram analysis. Amygdala-septal neurons that responded with an increased firing rate (↑ cells) displayed a significant percent change only in the AF and FAT groups and not in the other groups. The amygdala complex contains several nuclei [48], and some specific functions of the medial amygdala have been identified. The lateral septal nucleus receives afferents from the medial amygdala [49], and this amygdala nucleus relays pheromonal and olfactory information to lateral septal neurons and participates in social behavior [50], which may be particularly relevant for maternal behavior. Our results support this possibility. Amniotic fluid is the maternal-fetal fluid in which the fetus develops and is in constant contact for months. Amniotic fluid and its FATs selectively produced a significant change in the firing rate of septal neurons that apparently receive excitatory inputs from the medial amygdala.

The septal excitatory pathway comes exclusively from the medial amygdala [51] and is involved in disinhibition processes [52], with the participation of vasopressin and oxytocin [50, 53, 54]. Amniotic fluid and its FATs may promote the activity of these connections and impinge on filial behavior.

We used different routes of administration for every treatment included in present study. It may be argued that this fact may lead to some problem for the interpretation of results. However, in every case at least one hour elapsed between treatments and tests. Consequently, all used treatments must be reaching enough plasma concentrations during tests, as confirmed by differences found against vehicles.

5. Conclusion

Amniotic fluid and a mixture of its FATs produced diazepam-like effects on lateral septal neurons, regardless of their connection with the medial amygdala, likely with the participation of GABA_A receptors. Septal neurons receive excitatory inputs from the medial amygdala and appear to participate in the integration of maternal cue recognition mediated by AF and its FATs.

Conflict of Interests

The authors declare that there is no conflict of interests.

Acknowledgments

The authors thank Michael Arends for revising and editing the English of the paper and the Hospital Escuela de Universidad Veracruzana for obtaining the human samples. This work was supported by Consejo Nacional de Ciencia y Tecnología, México (CONACyT: CB-2006-1, 61741), Universidad Nacional Autónoma de México (UNAM: DGAPA-PAPIIT IN211111-3), and Sistema Nacional de Investigadores (exp. 754 and 32755). D. I. Vázquez-Hernández received fellowships from CONACyT (reg. 235961).

References

- [1] K. Christensson, T. Cabrera, E. Christensson, K. Uvnäs-Moberg, and J. Winberg, "Separation distress call in the human neonate in the absence of maternal body contact," *Acta Paediatrica*, vol. 84, no. 5, pp. 468–473, 1995.
- [2] K. Michelsson, K. Christensson, H. Rothgänger, and J. Winberg, "Crying in separated and non-separated newborns: sound spectrographic analysis," *Acta Paediatrica*, vol. 85, no. 4, pp. 471–475, 1996.
- [3] H. Varendi, K. Christensson, R. H. Porter, and J. Winberg, "Soothing effect of amniotic fluid smell in newborn infants," *Early Human Development*, vol. 51, no. 1, pp. 47–55, 1998.
- [4] C. Rattaz, N. Goubet, and A. Bullinger, "The calming effect of a familiar odor on full-term newborns," *Journal of Developmental and Behavioral Pediatrics*, vol. 26, no. 2, pp. 86–92, 2005.
- [5] S. Nishitani, T. Miyamura, M. Tagawa et al., "The calming effect of a maternal breast milk odor on the human newborn infant," *Neuroscience Research*, vol. 63, no. 1, pp. 66–71, 2009.
- [6] C. M. Contreras, J. F. Rodríguez-Landa, A. G. Gutiérrez-García, M. R. Mendoza-López, R. I. García-Ríos, and J. Cueto-Escobedo, "Anxiolytic-like effects of human amniotic fluid and its fatty acids in Wistar rats," *Behavioural Pharmacology*, vol. 22, no. 7, pp. 655–662, 2011.
- [7] A. S. Fleming, D. H. O'Day, and G. W. Kraemer, "Neurobiology of mother-infant interactions: experience and central nervous system plasticity across development and generations," *Neuroscience and Biobehavioral Reviews*, vol. 23, no. 5, pp. 673–685, 1999.
- [8] R. H. Porter and J. Winberg, "Unique salience of maternal breast odors for newborn infants," *Neuroscience and Biobehavioral Reviews*, vol. 23, no. 3, pp. 439–449, 1999.
- [9] F. Lévy, M. Keller, and P. Poindron, "Olfactory regulation of maternal behavior in mammals," *Hormones and Behavior*, vol. 46, no. 3, pp. 284–302, 2004.
- [10] B. Schaal, L. Marlier, and R. Soussignan, "Responsiveness to the odour of amniotic fluid in the human neonate," *Biology of the Neonate*, vol. 67, no. 6, pp. 397–406, 1995.
- [11] L. Marlier, B. Schaal, and R. Soussignan, "Orientation responses to biological odours in the human newborn. Initial pattern and postnatal plasticity," *Comptes Rendus de l'Académie des Sciences. Serie III*, vol. 320, no. 12, pp. 999–1005, 1997.
- [12] L. Marlier, B. Schaal, and R. Soussignan, "Bottle-fed neonates prefer an odor experienced in utero to an odor experienced postnatally in the feeding context," *Developmental Psychobiology*, vol. 33, no. 2, pp. 133–145, 1998.
- [13] L. Marlier and B. Schaal, "Human newborns prefer human milk: conspecific milk odor is attractive without postnatal exposure," *Child Development*, vol. 76, no. 1, pp. 155–168, 2005.

- [14] B. Schaal, "Mammary odor cues and pheromones. Mammalian infant-directed communication about maternal state, mammae, and milk," *Vitamins and Hormones*, vol. 83, pp. 83–136, 2010.
- [15] G. Guiraudie-Capraz, M.-C. Slomianny, P. Pageat et al., "Biochemical and chemical supports for a transnatal olfactory continuity through sow maternal fluids," *Chemical Senses*, vol. 30, no. 3, pp. 241–251, 2005.
- [16] C. M. Contreras, A. G. Gutiérrez-García, M. R. Mendoza-López, J. R. Rodríguez-Landa, B. Bernal-Morales, and C. Díaz-Martí, "Amniotic fluid elicits appetitive responses in human newborns: fatty acids and appetitive responses," *Developmental Psychobiology*, vol. 55, no. 3, pp. 221–231, 2013.
- [17] C. M. Contreras, A. G. Gutiérrez-García, and D. I. Vásquez-Hernández, "Fatty acids and emotional behavior," in *Neurochemistry*, C. M. Contreras, Ed., pp. 109–128, INTECH, Rijeka, Croatia, 2011.
- [18] T. P. Sheehan, R. A. Chambers, and D. S. Russell, "Regulation of affect by the lateral septum: implications for neuropsychiatry," *Brain Research Reviews*, vol. 46, no. 1, pp. 71–117, 2004.
- [19] S. W. Newman, "The medial extended amygdala in male reproductive behavior. A node in the mammalian social behavior network," *Annals of the New York Academy of Sciences*, vol. 877, pp. 242–257, 1999.
- [20] C. M. Contreras, J. F. Rodríguez-Landa, A. G. Gutiérrez-García, and B. Bernal-Morales, "The lowest effective dose of fluoxetine in the forced swim test significantly affects the firing rate of lateral septal nucleus neurones in the rat," *Journal of Psychopharmacology*, vol. 15, no. 4, pp. 231–236, 2001.
- [21] C. M. Contreras, V. Alcalá-Herrera, and M. L. Marván, "Action of antidepressants on the septal nuclei of the rat," *Physiology and Behavior*, vol. 46, no. 5, pp. 793–798, 1989.
- [22] M. H. Bassant, A. Jobert, P. Dutar, and Y. Lamour, "Effect of psychotropic drugs on identified septohippocampal neurons," *Neuroscience*, vol. 27, no. 3, pp. 911–920, 1988.
- [23] H. L. Garner, M. A. Whittington, and Z. Henderson, "Induction by kainate of theta frequency rhythmic activity in the rat medial septum-diagonal band complex *in vitro*," *Journal of Physiology*, vol. 564, no. 1, pp. 83–102, 2005.
- [24] C. Campo-Soria, Y. Chang, and D. S. Weiss, "Mechanism of action of benzodiazepines on GABA_A receptors," *British Journal of Pharmacology*, vol. 148, no. 7, pp. 984–990, 2006.
- [25] Z. Rogóž and G. Skuza, "Anxiolytic-like effects of olanzapine, risperidone and fluoxetine in the elevated plus-maze test in rats," *Pharmacological Reports*, vol. 63, no. 6, pp. 1547–1552, 2011.
- [26] National Research Council, *Guide for the Care and Use of Laboratory Animals*, Publication no. 80-23, National Academy Press, Washington, DC, USA, 1996.
- [27] C. M. Contreras, M. Molina, M. Saavedra, and L. Martínez-Mota, "Lateral septal neuronal firing rate increases during proestrus-estrus in the rat," *Physiology and Behavior*, vol. 68, no. 3, pp. 279–284, 2000.
- [28] G. Paxinos and C. Watson, *The Rat Brain in Stereotaxic Coordinates*, Academic Press, San Diego, Calif, USA, 4th edition, 1998.
- [29] L. Lehninger, D. L. Nelson, and M. M. Cox, *Lehninger Principles of Biochemistry*, W.H. Freeman, New York, NY, USA, 4th edition, 2005.
- [30] M. H. Beall, J. P. H. M. van den Wijngaard, M. J. C. van Gemert, and M. G. Ross, "Amniotic fluid water dynamics," *Placenta*, vol. 28, no. 8-9, pp. 816–823, 2007.
- [31] R. Nowak, R. H. Porter, F. Lévy, P. Orgeur, and B. Schaal, "Role of mother-young interactions in the survival of offspring in domestic mammals," *Reviews of Reproduction*, vol. 5, no. 3, pp. 153–163, 2000.
- [32] A. Leaf, Y.-F. Xiao, and J. X. Kang, "Interactions of n-3 fatty acids with ion channels in excitable tissues," *Prostaglandins Leukotrienes and Essential Fatty Acids*, vol. 67, no. 2-3, pp. 113–120, 2002.
- [33] I. Lauritzen, N. Blondeau, C. Heurteaux, C. Widmann, G. Romey, and M. Lazdunski, "Polyunsaturated fatty acids are potent neuroprotectors," *The EMBO Journal*, vol. 19, no. 8, pp. 1784–1793, 2000.
- [34] D. R. Flower, "The lipocalin protein family: structure and function," *The Biochemical Journal*, vol. 318, no. 1, pp. 1–14, 1996.
- [35] S. Liberatori, L. Bini, C. de Felice et al., "A two-dimensional protein map of human amniotic fluid at 17 weeks' gestation," *Electrophoresis*, vol. 18, no. 15, pp. 2816–2822, 1997.
- [36] K. Murakami, M. Lagarde, and Y. Yuki, "Identification of minor proteins of human colostrum and mature milk by two-dimensional electrophoresis," *Electrophoresis*, vol. 19, no. 14, pp. 2521–2527, 1998.
- [37] M. Tegoni, P. Pelosi, F. Vincent et al., "Mammalian odorant binding proteins," *Biochimica et Biophysica Acta*, vol. 1482, no. 1-2, pp. 229–240, 2000.
- [38] N. Gutiérrez-Castellanos, A. Martínez-Marcos, F. Martínez-García, and E. Lanuza, "Chemosensory function of the amygdala," *Vitamins and Hormones*, vol. 83, pp. 165–196, 2010.
- [39] M. Chebib and G. A. R. Johnston, "GABA-activated ligand gated ion channels: medicinal chemistry and molecular biology," *Journal of Medicinal Chemistry*, vol. 43, no. 8, pp. 1427–1447, 2000.
- [40] R. D. Schwartz and X. Yu, "Inhibition of GABA-gated chloride channel function by arachidonic acid," *Brain Research*, vol. 585, no. 1-2, pp. 405–410, 1992.
- [41] T.-C. Hwang, S. E. Guggino, and W. B. Guggino, "Direct modulation of secretory chloride channels by arachidonic and other cis unsaturated fatty acids," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 87, no. 15, pp. 5706–5709, 1990.
- [42] R. W. Ordway, J. J. Singer, and J. V. Walsh Jr., "Direct regulation of ion channels by fatty acids," *Trends in Neurosciences*, vol. 14, no. 3, pp. 96–100, 1991.
- [43] C. M. Contreras, L. Chacón, R. J. Rodríguez-Landa, B. Bernal-Morales, A. G. Gutiérrez-García, and M. Saavedra, "Spontaneous firing rate of lateral septal neurons decreases after forced swimming test in Wistar rat," *Progress in Neuropsychopharmacology and Biological Psychiatry*, vol. 28, no. 2, pp. 343–348, 2004.
- [44] C. M. Contreras, M. L. Marván, A. Ramírez-Morales, and A. Muñoz-Méndez, "Clomipramine enhances the excitatory actions of dorsal raphe nucleus stimulation in lateral septal neurons in the rat," *Neuropsychobiology*, vol. 27, no. 2, pp. 86–90, 1993.
- [45] L. Sghendo and J. Mifsud, "Understanding the molecular pharmacology of the serotonergic system: using fluoxetine as a model," *Journal of Pharmacy and Pharmacology*, vol. 64, no. 3, pp. 317–325, 2012.
- [46] R. L. Jakab and C. Leranth, "Septum," in *The Rat Nervous System*, G. Paxinos, Ed., pp. 405–442, Academic Press, New York, NY, USA, 2nd edition, 1995.
- [47] J. F. Rodríguez-Landa, C. M. Contreras, and R. I. García-Ríos, "Allopregnanolone microinjected into the lateral septum or dorsal hippocampus reduces immobility in the forced swim test:

- participation of the GABAA receptor," *Behavioural Pharmacology*, vol. 20, no. 7, pp. 614–622, 2009.
- [48] P. Sah, E. S. L. Faber, M. L. de Armentia, and J. Power, "The amygdaloid complex: anatomy and physiology," *Physiological Reviews*, vol. 83, no. 3, pp. 803–834, 2003.
- [49] A. R. Caffé, F. W. van Leeuwen, and P. G. M. Luiten, "Vasopressin cells in the medial amygdala of the rat project to the lateral septum and ventral hippocampus," *Journal of Comparative Neurology*, vol. 261, no. 2, pp. 237–252, 1987.
- [50] T. Sheehan and M. Numan, "The septal region and social behavior," in *The Behavioral Neuroscience of the Septal Region*, R. Numan, Ed., pp. 175–209, Springer, New York, NY, USA, 2000.
- [51] M. Numan and M. J. Numan, "Importance of pup-related sensory inputs and maternal performance for the expression of fos-like immunoreactivity in the preoptic area and ventral bed nucleus of the stria terminalis of postpartum rats," *Behavioral Neuroscience*, vol. 109, no. 1, pp. 135–149, 1995.
- [52] B. R. Komisaruk, J. S. Rosenblatt, M. L. Barona et al., "Combined *c-fos* and ¹⁴C-2-deoxyglucose method to differentiate site-specific excitation from disinhibition: analysis of maternal behavior in the rat," *Brain Research*, vol. 859, no. 2, pp. 262–272, 2000.
- [53] P. van den Hooff and I. J. A. Urban, "Vasopressin facilitates excitatory transmission in slices of the rat dorso-lateral septum," *Synapse*, vol. 5, no. 3, pp. 201–206, 1990.
- [54] I. F. Bielsky and L. J. Young, "Oxytocin, vasopressin, and social recognition in mammals," *Peptides*, vol. 25, no. 9, pp. 1565–1574, 2004.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

