Research Article

Ethnoveterinary Study of Medicinal Plants in a Tribal Society of Sulaiman Range

Akash Tariq,¹ Sakina Mussarat,¹ Muhammad Adnan,¹ Naser M. AbdElsalam,² Riaz Ullah,³ and Abdul Latif Khan⁴

¹ Department of Botany, Kohat University of Science and Technology, Kohat 26000, Pakistan

² Riyadh Community College, King Saud University, Riyadh 11437, Saudi Arabia

³ Department of Chemistry, Government College Ara Khel, FR Kohat 26000, Pakistan

⁴ Department of Biological Sciences and Chemistry, University of Nizwa, 616 Nizwa, Oman

Correspondence should be addressed to Akash Tariq; akash.malik786@yahoo.com

Received 24 July 2014; Accepted 24 September 2014; Published 21 October 2014

Academic Editor: Michael J. Myers

Copyright © 2014 Akash Tariq et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The aims of the present study were (i) to document ethnoveterinary plants and their formulation techniques in an unexplored region of Pakistan and (ii) to select candidate medicinal plants with high consensus factor and fidelity value for further *in vitro* investigation. A total of 60 informants were interviewed using semistructured questionnaire. A total of 41 plants belonging to 30 families were used to treat livestock ailments in study area. Mostly leaves (47%) were used in recipes formulation mostly in the form of decoction. Gastrointestinal infections were found more common and majority of the plants were used against cow (31) and buffaloes (24) ailments. Recovery time of majority of the recipes was three to four days. Informant consensus factor (Fic) results have shown a high degree of consensus for gastrointestinal, respiratory, and reproductive (0.95 each) ailments. Fidelity level (FL) results showed that *Asparagus gracilis* ranked first with FL value 93% followed by *Rumex hastatus* ranked second (91%) and *Tinospora cordifolia* ranked third (90%). Aged farmers and nomads had more traditional knowledge as compared to younger ones. Plants with high Fic and FL values could be further investigated *in vitro* for the search of some novel bioactive compounds and young generation should be educated regarding ethnoveterinary practices.

1. Introduction

Traditional veterinary medicine was experienced as early as 1800 B.C. at the time of King Hammurabi of Babylon who devised laws on veterinary fees and charged for treating animals [1]. Many ethnoveterinary medicines were neglected due to the development of initial western drugs. Ethnoveterinary practices have gained tremendous importance for the last decade due to the discovery of some effective ethnoveterinary products [2]. Traditional veterinary medicines provide a cheap therapy and easy accessibility in comparison with western drugs [3].

Ethnoveterinary practices are more common in developing countries due to different socioeconomic factors [4]. Pakistan is an agriculture country and almost 80% of its population is dependent on agriculture and livestock. Pakistan is

the world's 5th largest milk producing country due to its high dependency on agriculture and livestock [5]. Resource-poor farmers of Pakistan greatly rely on traditional medicine due to their limited access to modern prevention health practices and particularly lack of modern health facilities in their areas [6]. Despite the fact that traditional knowledge is very much important for the livestock health and productivity, the documentation of this knowledge is very much neglected in majority of the remote areas of Pakistan. Livestock farmers all over Pakistan can draw on over 4000 years of knowledge and experience [7]. This traditional knowledge has been passed orally from generation to generation but it may be extinct or may be endangered due to the current rapid socioeconomic, environmental, and technological changes [8]. Therefore, the documentation of such knowledge is very crucial before its extinction for future developments.

values all of the ailments that were reported are grouped into major classes [10]. FL value was estimated using the formula $FL = Ip/Iu \times 100$, where Ip is the number of respondents who reported the utilization of medicinal plants for a specific main ailment and Iu is the total number of respondents who mentioned the same plant for any ailment [13]. It is assumed that those medicinal plants which are plants used in some recurring manner for the same disease category are more likely to be biologically active [14].

2.7. Collection and Preservation of the Reported Medicinal Plants. Field trips were made with local informants for the collection of the reported medicinal plants. Collected medicinal plants were brought to the laboratory of Kohat University of Science and Technology (KUST), Kohat, Pakistan, for further processing. Specimen identification and confirmation were undertaken by using Flora of Pakistan and taxonomic experts. Plants were dried and pressed on herbarium sheets and deposited at the Herbarium of Department of Botany, KUST, Kohat, Pakistan.

3. Results

The present study revealed that local farmers of Kohat region utilize 41 medicinal plants belonging to 30 families for the treatment of different types of livestock ailments (Table 1). Among all the families, Asteraceae was found to be dominant (4 species) in the study area being in use in ethnoveterinary practices in the region. Traditional farmers mostly used herbs (55%) for the preparation of ethnomedicines (Table 2) followed by shrubs (27%). Almost all plant parts were being used for medicinal purposes but leaves (47%) were found to be the most frequently used plant part followed by whole plant (32%) and roots (17%) (Table 2). Local farmers used these ethnomedicines to treat different types of domestic animals like buffaloes, cows, goats, sheep, and donkeys. A total of 31 plants were found to be used for treatment of cow ailments followed by 24 plants against buffalo's ailments and 17 for goats (Figure 2). Different types of ailments were treated which were categorized into 8 major categories. Gastrointestinal infections were found to be most common in domestic animals and a total of 13 plants were used against them followed by 7 plants which are used as antipyretic while 6 are used for wounds treatment (Table 3). Local farmers prepare different types of ethnomedicines but the most preferred techniques were decoction and powder (10 plants each) followed by crushing (7 plants) in the studied region (Figure 3). Monotherapy was most common in the study area; only few plants were found to be used in concoction form (Table 1). For example, stem of *Allium sativum* is mixed with flower of *Punica granatum* and milk and used against gastrointestinal infection; roots of *Asparagus gracilis* are mixed with leaves of *Coriandrum sativum* to make fine concoction and given with water to cattle for delivery purposes. Different types of vehicles were found to be used for preparation and administration of plant recipes like sugar, flour, water, and milk (Table 1). The most common route of administration was oral (75%) followed by dermal (24%) and only single species is administered through nasal

FIGURE 2: Number of plants used to treat different domestic animals.

FIGURE 3: Traditional medicines preparation.

pathway (Table 1). Recovery time of majority of the recipes was three to four days. Informant consensus (Fic) results have shown a high degree of consensus for gastrointestinal, respiratory, and reproductive (0.95 each) ailments, which were followed by parasitic infections and wound healing (0.90 each) (Table 3). The highest plant use citation was for gastrointestinal (260) followed by wound healing (53) and reproductive (47) ailments. The present study revealed 10 medicinal plants having high FL value (Table 4). FL values in this study varied from 1.0% to 100%. *Asparagus gracilis* ranked first with the highest FL value (93%) followed by *Rumex hastatus* ranked second (91%), *Tinospora cordifolia* ranked third (90%), and *Aloe barbadensis* ranked fourth (85%). The entire informants interviewed were aged people ranging between 40 and 70 years old. No use of ethnoveterinary medicine by women or young generation was recorded.

4. Discussion

Livestock keeping is one of the most important economic sources of rural community of study area. The farmers and nomadic people of the area not only depend on plants to get fodder for their animals but also use different medicinal plants to treat various animal diseases. The majority of the people interviewed using ethnoveterinary plants have got this knowledge from their forefathers while some have learned

TABLE 1: Ethnoveterinary plants used for the treatment of livestock ailments in Kohat region.

Plant names/families name/voucher number	Local names	Habit	Part used	Ailment treated	Animal treated	Recipe	Vehicles	Route	Recovery
<i>Acacia modesta</i> Wall. Fabaceae KUH-421	Kikar	Tree	Leaves, seeds	Delivery	Buffaloes, cow	Decoction	Water	Oral	Two days
<i>Achyranthes aspera</i> Linn. Amaranthaceae KUH-422	Gishkay	Herb	Whole plant	Anthelmintic and delivery	Buffaloes, cow, goat, sheep	Powder	Sugar	Oral	One day
<i>Allium cepa</i> L. Amaryllidaceae KUH-423	Pyaz	Herb	Whole plant	Febrifuge, tonic	Sheep, goat	Decoction	Flour	Oral	Three days
<i>Allium sativum</i> L. Liliaceae KUH-424	Thoom	Herb	Stem	Gastrointestinal	Goat	Concoction	Milk	Oral	Ten days
<i>Aloe barbadensis</i> Mill. Liliaceae KUH-425	Kunwar	Shrub	Root	Gastrointestinal	Sheep, cow, buffaloes	Powder	Milk	Oral	Six days
<i>Artemisia brevifolia</i> Wall. Asteraceae KUH-426	Jaukay	Herb	Leaves	Removal of placenta	Cow	Decoction	Sugar	Oral	
<i>Asparagus gracilis</i> Royle. Liliaceae KUH-427	Lachgawa	Herb	Root	Delivery	Goat	Concoction	Water	Oral	
<i>Brassica campestris</i> L. Brassicaceae KUH-428	Sarson	Herb	Whole plant	External lice (blood feeding)	Cow, buffaloes	Paste		Dermal	Five days
<i>Calotropis procera</i> (Willd.) R.Br. Apocynaceae KUH-429	Spulmaey	Shrub	Fruit, leaves	Intestinal worms and skin infections	Buffaloes, cow, goat, sheep	Paste, concoction	Sugar	Oral, dermal	Three days
<i>Cannabis sativa</i> L. Cannabaceae KUH-430	Bhang	Shrub	Leaves	External parasite, appetizer	Cow, donkey, buffaloes	Poultice, powder		Dermal, oral	Two days
<i>Chenopodium album</i> L. Amaranthaceae KUH-431	Samaray	Herb	Whole plant	Wound healing	Goat, sheep, cow	Paste		Dermal	One day
<i>Chrysanthemum leucanthemum</i> L. Asteraceae KUH-432	Chitti pulari	Herb	Whole plant	Increase milk production	Goat, cow, buffaloes	Powder	Flour	Oral	Continuously
<i>Citrullus colocynthis</i> (L.) Schrad. Cucurbitaceae KUH-433	Karthuma	Herb	Root	Skin infection	Buffalo, cow, goat, sheep	Juice		Dermal	Two days
<i>Convolvulus arvensis</i> L. Convolvulaceae KUH-434	Shankpuspi	Herb	Whole plant	Constipation	Cow, buffaloes, sheep	Crushed	Sugar	Oral	Four days
<i>Coriandrum sativum</i> L. Apiaceae KUH-435	Dhania	Herb	Leaves, root	Antidiuretic	Buffaloes	Decoction	Water	Oral	One week
<i>Curcuma longa</i> L. Zingiberaceae KUH-436	Haldi	Tree	Leaves	Wound healing	Cow	Decoction	Sugar	Oral	Two days
<i>Gynodon dactylon</i> L. Poaceae KUH-437	Wakha	Herb	Whole plant	Wound healing, analgesic	Cow, buffaloes	Concoction	Milk	Oral	Two days

TABLE 1: Continued.

Plant names/families name/voucher number	Local names	Habit	Part used	Ailment treated	Animal treated	Recipe	Vehicles	Route	Recovery
<i>Gynoglossum lanceolatum</i> Forssk. Boraginaceae KUH-438	Pachy	Herb	Root	Common cold	Cow, buffaloes	Crushed	Water	Oral	Two days
<i>Cyperus niveus</i> Retz. Cyperaceae KUH-439	Kulio	Herb	Whole plant	Common cold, stomach worms, joint pains	Cow, goat	Crushed	Water	Oral	Four to five days
<i>Datura innoxia</i> Mill. Solanaceae KUH-440	Mangaz	Herb	Whole plant	Antilice	Sheep, cow	Paste	Milk	Dermal	Three days
<i>Hedera nepalensis</i> K. Koch Araliaceae KUH-441	Zalai	Shrub	Leaves	To remove leeches	Sheep	Infusion		Nasal	Three days
<i>Melia azedarach</i> L. Meliaceae KUH-442	Dhrek	Tree	Leaves	Stomach flatulence	Cow, buffaloes	Powder	Sugar	Oral	Four days
<i>Meniha arvensis</i> Linn. Lamiaceae KUH-443	Pudina	Herb	Leaves	External parasite	Cow	Paste		Dermal	Four days
<i>Morus alba</i> L., Moraceae KUH-444	Toot	Tree	Leaves, fruit	Laxative	Buffaloes	Crushed	Milk	Oral	Two days
<i>Morus nigra</i> L. Moraceae KUH-445	Tor toot	Tree	Leaves	Tonic, laxative	Cow, buffaloes	Powder	Water	Oral	Three days
<i>Nerium oleander</i> L. Apocynaceae KUH-446	Ghanderay	Shrub	Whole plant	Stomachache	Sheep	Concoction	Water	Oral	One week
<i>Ocimum basilicum</i> L. Lamiaceae KUH-447	Kashmalay	Shrub	Leaves	Gastrointestinal	Buffaloes	Decoction	Water	Oral	Four days
<i>Punica granatum</i> L. Lythraceae KUH-448	Anar	Shrub	Fruit, leaves	Anthelmintic	Cow, buffaloes, goat	Decoction	Milk	Oral	Two days
<i>Ricinus communis</i> Linn. Euphorbiaceae KUH-449	Arund	Shrub	Leaves, stem	Common cold	Buffaloes, cow	Powder	Flour	Oral	Three days
<i>Rumex hastatus</i> D. Don. Polygonaceae KUH-450	Tarooky	Shrub	Root and leaves	Wound healing	Goat, cow, buffaloes	Powder	Flour	Oral	Four days
<i>Solanum surratense</i> Burm. f. Solanaceae KUH-451	Kandiari	Herb	Whole plant	Fever, cough, intestinal infections	Cow, buffaloes, goat, sheep	Crushed	Flour	Oral	One week
<i>Sonchus asper</i> (L.) Hill. Asteraceae KUH-452	Spingul	Herb	Whole plant	Milk production	Goat, cow, buffaloes	Decoction	Flour	Oral	Continuously
<i>Tagetes minuta</i> L. Asteraceae KUH-453	Ban hanjari	Herb	Leaves	Skin infections	Cow, buffaloes	Juice		Dermal	Five days
<i>Tamarix aphylla</i> (L.) H. Karst. Tamaricaceae KUH-454	Khagal	Tree	Leaves	Kill worms of wounds	Cow	Paste		Dermal	Four days

TABLE 1: Continued.

Plant names/families name/voucher number	Local names	Habit	Part used	Ailment treated	Animal treated	Recipe	Vehicles	Route	Recovery
<i>Tinospora cordifolia</i> Miers. Menispermaceae KUH-455	Giloe	Climber	Whole plant	Skin infections	Cow, goat	Poultice		Dermal	Three days
<i>Tribulus terrestris</i> Linn. Zygophyllaceae KUH-456	Markondai	Herb	Whole plant	Chronic cough	Cow, buffaloes, goat	Crushed	Sugar	Oral	One week
<i>Trifolium repens</i> L. Papilionaceae KUH-457	Shoutal	Shrub	Root	Tonic, laxative	Goat, cow	Powder	Flour	Oral	One day
<i>Triticum aestivum</i> L. Poaceae KUH-458	Gandam	Herb	Seeds	Common cold, dysentery	Cow	Powder	Flour	Oral	Two days
<i>Verbena officinalis</i> L. Verbenaceae KUH-459	Shamakay	Herb	Stem, leaves	Wound healing	Buffaloes	Decoction	Milk	Oral	Three days
<i>Vitex negundo</i> L. Verbenaceae KUH-460	Marmandi	Shrub	Stem	Mange, fever, stomach problems	Cow, goat	Crushed, decoction	Sugar	Oral	Four to six days
<i>Zizyphus nummularia</i> W. & A. Rhamnaceae KUH-461	Kurkunda	Tree	Leaves	Wound healing	Cow	Decoction	Sugar	Oral	Three days

TABLE 2: Habit and parts used of ethnoveterinary plants.

General attributes	Total plants	Percentage
Habit		
Herbs	22	55
Shrubs	11	27
Trees	06	15
Climbers	01	02
Parts used		
Leaves	19	47
Whole plant	13	32
Root	07	17
Stem	04	10
Fruit	03	07
Seed	01	02

from the other people. The majority of the farmers and nomadic pastoralists were not very well off and heavily dependent on medicinal plants due to their unaffordable potential of using modern veterinary drugs for their animals treatment.

The present study revealed that people of the region use 41 medicinal plants for their livestock health care. Similar studies have also been documented in other parts of Pakistan [5, 15]. Traditional healers of the region mostly use herbs for the treatment of their animals that might be due to the fact that herbs are available everywhere and easy to collect as compared with other growth forms. The results indicate the abundance of herbs in the study area and their high usage might also be due to the strong efficacy of herbaceous plants against livestock ailments. The same findings were also reported from other studies conducted at different parts of the world [16, 17].

The wider utilization of this Asteraceae family might be due to its higher abundance in the study area or might be due to high bioactivity. Similar studies have also been reported from other parts of world [18] and from Pakistan [19, 20] where traditional healers mostly use the member of Asteraceae family for the preparation of traditional medicines for the treatment of different livestock and human ailments. This observation is however different from that of Appidi et al. [21] and Offiah et al. [22] who in an ethnoveterinary survey reported Fabaceae family as the highest. The difference among studies might be related to the different dominant vegetation of the areas or might be associated with traditional beliefs of different cultures in using specific plants traditionally.

Most of the ethnoveterinary recipes in the study region are prepared using leaves of plants. The highest use of leaves in large number of ethnomedicinal and ethnoveterinary studies has also been documented from different parts of the world [23, 24]. Preferred use of leaves might be associated with ease of collection as compared to other plant parts. Leaves are the main site of photosynthetic apparatus and are involved in a variety of physiological processes of plants and produce numerous secondary metabolites that could be a possible reason for their effectiveness and efficacy against different

TABLE 3: Informant consensus factor.

Disease categories	Nur	Nt	Fic
Gastrointestinal	260	13	0.95
Respiratory	22	02	0.95
Reproductive	47	03	0.95
Dermatological	12	04	0.77
Wounds	53	06	0.90
Antipyretic	14	07	0.53
Parasitic	45	05	0.90
General body tonic	09	03	0.75

livestock diseases. Local people also use whole plants after leaves for herbal formulation that could be a very destructive type of harvesting for rare and slowly growing plants from conservation point of view. Harvesting of leaves does not pose any serious impact on the life cycle of plants and is considered a sustainable type of harvesting. The present results are in contradiction with other studies where roots are the most widely used plant part in ethnoveterinary practices [25, 26].

Cows and buffaloes were the most commonly treated animals followed by goats and sheep in the studied region. Similar results have also been conducted by van der Merwe et al. [27] and Benítez et al. [16]. There was almost no mention of treating dogs, cats, or donkeys. This is probably because rural people do not generally keep animals as pets and because nonproduction animals are perceived as being more resistant than humans to different kinds of ailments. Production animals are also more important because of their socioeconomic importance in the local inhabitant life. The majority of the plants in the region are used to treat different types of gastrointestinal problems of the livestock like diarrhea, expulsion of worms, constipation, and so forth. It has already been found that stomach infections are more common in lactating animals which might be due to poor quality of fodder and drinking water [17]. Informant consensus results also showed the highest informant citation for gastrointestinal, respiratory, and gynecological problems. The highest informant citation against these infections gives an indication of high prevalence of these diseases in the region. According to Heinrich et al. [11], high Fic values are very useful in the selection of specific plants for further search of bioactive compounds. Widely used medicinal plants for species ailments always score the highest fidelity level. The present study determined different plants like *Asparagus gracilis*, *Rumex hastatus*, *Tinospora cordifolia*, *Aloe barbadensis*, and so forth, scored highest fidelity values and should be further subjected to phytochemical and pharmacological investigation to prove their medicinal efficacy.

The method of drug preparation in many cases varied from individual to individual. The same plant material for the same ailment was prepared in different ways by different traditional veterinary healers. Traditional healers prepare ethnoveterinary recipes mostly in the form of powder and decoction in the study area. Deeba [28] powdering or boiling is the most common method of drugs extraction. These findings are in line with a study conducted in the Malakand

TABLE 4: Fidelity level of highly utilized species.

Number	Plant species	Disease category	Ip	Iu	FL %
01	<i>Asparagus gracilis</i>	Reproductive	29	31	93
02	<i>Rumex hastatus</i>	Wound healing	32	35	91
03	<i>Tinospora cordifolia</i>	Dermatological	28	31	90
04	<i>Aloe barbadensis</i>	Gastrointestinal	23	27	85
05	<i>Convolvulus arvensis</i>	Gastrointestinal	24	29	82
06	<i>Tribulus terrestris</i>	Respiratory	17	21	80
07	<i>Zizypus nummularia</i>	Wound healing	18	23	78
08	<i>Chenopodium album</i>	Wound healing	17	25	68
09	<i>Artemisia brevifolia</i>	Reproductive	19	30	63
10	<i>Cannabis sativa</i>	Parasitic	17	29	58

valley of Pakistan [5] while they are contradictory with the studies conducted in other parts of the world [29, 30]. Most of the recipes are prepared using single plant mixture while some of the recipes are also prepared in the form of concoction and it is generally believed that potency of the drugs can be enhanced when used in concoction form [31]. These recipes are mostly taken orally in the study area due to the high prevalence of the internal diseases. These recipes are given to the livestock with their feed along with different types of ingredients like sugar, flour, milk, and so forth, in the region. Similar findings are also reported from other regions of the world [18, 32]. The use of these vehicles might be due to their enhancing potential of taste and medicinal properties of certain plant remedies. Uniformity was lacking regarding amount of medicines to be used among informants during the interview. It was determined that a contradiction in ethnoveterinary dosage is a serious drawback of traditional medicinal plants. Informants only provided the knowledge of observed time of recovery of animals in response to given recipes. Full recovery is confirmed when the animals restart proper feeding and activities. Similar findings are also reported by other ethnoveterinary studies conducted elsewhere [5, 22].

It was confirmed from the present study that men had better knowledge regarding ethnoveterinary practices as compared to women. The reason might be due to the fact that men are mostly favored in shift of knowledge while women in the majority of the cultures are considered for family's care. This noticeable gender bias reflects the limited involvement of women in cattle production and herd health in Kohat region of Pakistan. Aged males had much more indigenous knowledge as compared to young generation which might be due to the lack of interest in such practices. Therefore, documentation of ethnoveterinary practices is an essential step toward the conservation of such knowledge before its extinction.

5. Conclusions

Local farmers and nomads of the region utilize different medicinal plants for the treatment of livestock due to their low income status and high expenses of western drugs. Traditional healers possess tremendous expertise in preparing

herbal formulations of medicinal plants. Gastrointestinal infections were most common in the studied region; therefore attention should be given to provide good quality fodder and water to the livestock. Plants with high informant consensus and fidelity level should be subjected to further *in vitro* investigation for their phytochemical analysis and pharmacological activities. Young generation should be mobilized to take interest in ethnoveterinary practices in order to conserve this knowledge.

Questionnaire of Ethnoveterinary Data Collection

Informants' Consent for the Participation in the Study

I _____ (name of informant) hereby give my full consent and conscious to participate in this study and declare that to the best of my knowledge the information that I have provided is true, accurate, and complete.

Date _____ (signature/thumb impression of informant).

Informants' Details

Name
Gender
Age
Occupation
Education
Location/residence

Data about Medicinal Plant and Its Use

Number of plants known
Names of plants (local names)
Plant part used
Name of disease(s) treated
Type of animal treated
Method of crude drug preparation
Use of single or mixture of plants

Ingredients used
 Recovery time
 Route of administration
 Dosage

Informant Consensus Factor and Fidelity Level

Name of plants used against disease category 1
 Name of plants used against disease category 2
 Name of plants used against disease category 3
 Name of plants used against disease category 4
 Name of plants used against disease category 5
 Name of plants used against disease category 6
 Name of plants used against disease category 7
 Name of plants used against disease category 8
 Name of plants used against disease category 9
 Name of plants used against disease category 10

Remarks

Plant identified as — (botanical name and family)
 Signature of researcher

Conflict of Interests

The authors declare that they have no competing interests.

Authors' Contribution

Akash Tariq and Muhammad Adnan designed the research project. Akash Tariq and Sakina Mussarat conducted the field work. Akash Tariq, Sakina Mussarat, Abdul Latif Khan, Naser M. AbdElsalam, and Riaz Ullah analyzed the data and wrote the whole paper. Muhammad Adnan and Akash Tariq supervised all the stages of research work and provided comments on the draft. All the authors have read and approved the final paper.

Acknowledgments

The authors are thankful to the Deanship of Scientific Research, King Saud University, Riyadh, Saudi Arabia, for funding the work through the Research Group Project no. RGP-210. The authors are also very thankful to the local informants for sharing their valuable knowledge.

References

- [1] S. V. Veen, "Sense or nonsense? Traditional methods of animal disease prevention and control in African savannah," in *Ethnoveterinary Research and Development*, C. M. McCorkle, E. Mathias, and T. W. Schillhorn van Veen, Eds., p. 338, Intermediate Technology Publications, London, UK, 1996.
- [2] C. Lans, N. Turner, T. Khan, G. Brauer, and W. Boepple, "Ethnoveterinary medicines used for ruminants in British Columbia, Canada," *Journal of Ethnobiology and Ethnomedicine*, vol. 3, article 11, 2007.
- [3] S. Ganesan, M. Chandhirasekaran, and A. Selvaraj, "Ethnoveterinary healthcare practices in southern districts of Tamil Nadu," *Indian Journal of Traditional Knowledge*, vol. 7, no. 2, pp. 347–354, 2008.
- [4] C. H. Rahman, A. Ghosh, and S. Mandal, "Studies on the Ethno veterinary medicinal plants used by the tribes of Birbhum district, West Bengal," *Indian Journal of Traditional Knowledge*, vol. 33, pp. 333–338, 2009.
- [5] H. Ul Hassan, W. Murad, A. Tariq, and A. Ahmad, "Ethnoveterinary study of medicinal plants in Malakand Valley, District Dir (Lower), Khyber Pakhtunkhwa, Pakistan," *Irish Veterinary Journal*, vol. 67, no. 1, article 6, 2014.
- [6] A. M. Abbasi, S. M. Khan, M. Ahmad, M. A. Khan, C. L. Quave, and A. Pieroni, "Botanical ethnoveterinary therapies in three districts of the Lesser Himalayas of Pakistan," *Journal of Ethnobiology and Ethnomedicine*, vol. 9, article 84, 2013.
- [7] Z. Farooq, Z. Iqbal, S. Mushtaq, G. Muhammad, M. Z. Iqbal, and M. Arshad, "Ethnoveterinary practices for the treatment of parasitic diseases in livestock in Cholistan desert (Pakistan)," *Journal of Ethnopharmacology*, vol. 118, no. 2, pp. 213–219, 2008.
- [8] A. Hussain, M. N. Khan, Z. Iqbal, and M. S. Sajid, "An account of the botanical anthelmintics used in traditional veterinary practices in Sahiwal district of Punjab, Pakistan," *Journal of Ethnopharmacology*, vol. 119, no. 1, pp. 185–190, 2008.
- [9] S. Shinwari, R. Qureshi, and E. Baydoun, "Ethnobotanical study of Kohat Pass (Pakistan)," *Pakistan Journal of Botany*, vol. 43, pp. 135–139, 2011.
- [10] M. Giday, Z. Asfaw, Z. Woldu, and T. Teklehaymanot, "Medicinal plant knowledge of the Bench ethnic group of Ethiopia: an ethnobotanical investigation," *Journal of Ethnobiology and Ethnomedicine*, vol. 5, article 34, 2009.
- [11] M. Heinrich, A. Ankli, B. Frei, C. Weimann, and O. Sticher, "Medicinal plants in Mexico: healers' consensus and cultural importance," *Social Science and Medicine*, vol. 47, no. 11, pp. 1859–1871, 1998.
- [12] M. Canales, T. Hernández, J. Caballero et al., "Informant consensus factor and antibacterial activity of the medicinal plants used by the people of San Rafael Coxcatlán, Puebla, México," *Journal of Ethnopharmacology*, vol. 97, no. 3, pp. 429–439, 2005.
- [13] J. Friedman, Z. Yaniv, A. Dafni, and D. Palewitch, "A preliminary classification of the healing potential of medicinal plants, based on a rational analysis of an ethnopharmacological field survey among Bedouins in the Negev Desert, Israel," *Journal of Ethnopharmacology*, vol. 16, no. 2-3, pp. 275–287, 1986.
- [14] R. T. Trotter and M. H. Logan, "Informants consensus: a new approach for identifying potentially effective medicinal plants," in *Plants in Indigenous Medicine and Diet*, N. L. Etkin, Ed., pp. 91–112, Redgrave, Bedford Hill, NY, USA, 1986.
- [15] M. A. Khan, M. A. Khan, and M. Hussain, "Ethnoveterinary medicinal uses plants of Poonch valley Azad Kashmir," *Pakistan Journal of Weed Sciences Research*, vol. 18, no. 4, pp. 495–507, 2012.
- [16] G. Benítez, M. R. González-Tejero, and J. Molero-Mesa, "Knowledge of ethnoveterinary medicine in the Province of Granada, Andalusia, Spain," *Journal of Ethnopharmacology*, vol. 139, no. 2, pp. 429–439, 2012.
- [17] D. Luseba and D. van der Merwe, "Ethnoveterinary medicine practices among Tsonga speaking people of South Africa," *Onderstepoort Journal of Veterinary Research*, vol. 73, no. 2, pp. 115–122, 2006.

- [18] H. Yineger, E. Kelbessa, T. Bekele, and E. Lulekal, "Ethnoveterinary medicinal plants at Bale Mountains National Park, Ethiopia," *Journal of Ethnopharmacology*, vol. 112, no. 1, pp. 55–70, 2007.
- [19] I. Khan, N. M. Abdelsalam, H. Fouad, A. Tariq, R. Ullah, and M. Adnan, "Application of ethnobotanical indices on the use of traditional medicines against common diseases," *Evidence-Based Complementary and Alternative Medicine*, vol. 2014, Article ID 635371, 21 pages, 2014.
- [20] N. Akhtar, A. Rashid, W. Murad, and E. Bergmeier, "Diversity and use of ethno-medicinal plants in the region of Swat, North Pakistan," *Journal of Ethnobiology and Ethnomedicine*, vol. 9, no. 1, article 25, 2013.
- [21] J. R. Appidi, D. S. Grierson, and A. J. Afolayan, "Ethnobotanical study of plants used for the treatment of diarrhoea in the Eastern Cape, South Africa," *Pakistan Journal of Biological Sciences*, vol. 11, no. 15, pp. 1961–1963, 2008.
- [22] N. V. Offiah, S. Makama, I. L. Elisha et al., "Ethnobotanical survey of medicinal plants used in the treatment of animal diarrhoea in Plateau State, Nigeria," *BMC Veterinary Research*, vol. 7, article 36, 2011.
- [23] C. P. Kala, "Ethnomedicinal botany of the Apatani in the Eastern Himalayan region of India," *Journal of Ethnobiology and Ethnomedicine*, vol. 1, article 11, 2005.
- [24] J. A. Bhat, M. Kumar, and R. W. Bussmann, "Ecological status and traditional knowledge of medicinal plants in Kedarnath Wildlife Sanctuary of Garhwal Himalaya, India," *Journal of Ethnobiology and Ethnomedicine*, vol. 9, no. 1, article 1, 2013.
- [25] J. R. S. Tabuti, S. S. Dhillon, and K. A. Lye, "Ethnoveterinary medicines for cattle (*Bos indicus*) in Bulamogi county, Uganda: plant species and mode of use," *Journal of Ethnopharmacology*, vol. 88, no. 2-3, pp. 279–286, 2003.
- [26] D. Hunde, Z. Asfaw, and E. Kelbessa, "Use and management of ethnoveterinary medicinal plants by indigenous people in "Boosat", Welenchetti area," *Ethiopian Journal of Biological Sciences*, vol. 3, pp. 113–132, 2004.
- [27] D. van der Merwe, G. E. Swan, and C. J. Botha, "Use of ethnoveterinary medicinal plants in cattle by Setswana-speaking people in the Madikwe area of the North West Province of South Africa," *Journal of the South African Veterinary Association*, vol. 72, no. 4, pp. 189–196, 2001.
- [28] F. Deebea, *Documentation of ethnoveterinary practices in urban and peri-urban areas of Faisalabad, Pakistan [Ph.D. thesis]*, University of Agriculture, Faisalabad, Pakistan, 2009.
- [29] A. P. Dold and M. L. Cocks, "Traditional veterinary medicine in the Alice district of the Eastern Cape Province, South Africa," *South African Journal of Science*, vol. 97, no. 9-10, pp. 375–379, 2001.
- [30] E. Lulekal, E. Kelbessa, T. Bekele, and H. Yineger, "An ethnobotanical study of medicinal plants in Mana Angetu District, southeastern Ethiopia," *Journal of Ethnobiology and Ethnomedicine*, vol. 4, article 10, 2008.
- [31] D. Abebe and A. Ayehu, *Medicinal Plants and Enigmatic Health Practices of Northern Ethiopia*, Berhanena Selam Printing Enterprise, Addis Ababa, Ethiopia, 1993.
- [32] A. Jabbar, M. A. Raza, Z. Iqbal, and M. N. Khan, "An inventory of the ethnobotanicals used as anthelmintics in the southern Punjab (Pakistan)," *Journal of Ethnopharmacology*, vol. 108, no. 1, pp. 152–154, 2006.