Research Article

Nasopupillary Asymmetry

Eduardo Arenas,¹ Diana Muñoz,² Evelyn Matheus,³ and Diana Morales³

¹Department of Ophthalmology, Colombia National University, Bogota, Colombia

²Department of Physiology, Colombia National University, Bogota, Colombia

³Eduardo Arenas Ophthalmology Society, Bogota, Colombia

Correspondence should be addressed to Diana Muñoz; diannymism@gmail.com

Received 2 July 2014; Revised 10 November 2014; Accepted 20 November 2014; Published 4 December 2014

Academic Editor: Dan-Ning Hu

Copyright © 2014 Eduardo Arenas et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Purpose. To establish the prevalence of nasopupillary asymmetry (difference in nasopupillary distances) in the population and its relation with the interpupillary distance. **Methods.** A retrospective descriptive study was conducted by reviewing of 1262 medical records. The values of nasopupillary asymmetry and the interpupillary distance were obtained. A statistical analysis was made and the correlation between these variables was established. **Results.** Seventy-nine percent of the population presented some degree of nasopupillary asymmetry. The interpupillary distance had a very low correlation with the nasopupillary asymmetry ($r = 0.074$, $P = 0.0$). **Conclusion.** It is advisable to use the nasopupillary distance of each eye as a standard measurement.

1. Introduction

The interpupillary distance (ID) is the distance measured between the centers of the pupils, and it is important for the creation of the stereoscopic vision, which results in a single tridimensional image [1, 2]. The ID is an important measure widely used today as part of the process of eyeglasses formulation [3, 4]. Furthermore, the nasopupillary distance (NPD) is the measure from the pupil center to the medial nasal axis [5, 6].

However, currently the nasopupillary asymmetry (NPA) (difference in nasopupillary distances) and its relation to the ID have not been studied [7]. Additionally, there have not been reported in the literature data of the percentage of the population with NPA.

In this research, the values of NPA and its relation to ID in a Colombian population were studied, evidencing that most of the population presents some degree of NPA. This is the first research reported in the literature that studies the NPA.

2. Methods

A retrospective descriptive study was conducted by reviewing 1300 medical records. Patients with any type of strabismus,

single eye and those under 18 were excluded, thereby obtaining a total sample of 1262 medical records of 631 women and an equal number of men, aged between 18 and 99 years.

The ID and NPD measurement was performed all the time by the same specialist utilizing the Essilor pupillometer calibrated by long distance. The instrument shows the ID and also the two separate nasopupillary distances (right and left eyes) (Figure 1). The NPA was obtained by calculating the difference between the NPD of the right eye (RE) and the NPD of the left eye (LE), for each patient. Data was collected and a statistical analysis for the entire population was performed.

The degree of NPA was classified as low (up to 2 mm), medium (greater than 2 mm up to 4 mm), and high (greater than 4 mm). Also, which eye was farthest from the nasal axis (greater NPD) and the distribution of this variable to the entire population were analyzed. A calculation to determine the relation between the NPA and the ID was made, by Pearson's correlation coefficient.

3. Results

The average age of the population was 46.8 ± 18 (range: 18–99) years, with a mean of 48.65 ± 17.57 (range: 18–99) years for women and 44.94 ± 18.24 (range: 18–92) years for men.

FIGURE 1: Digital pupillometer that shows in the center the ID and to the sides the NPD of each eye.

FIGURE 2: Distribution of the NPA in the population.

Of the total sample, 288 (22.82%) patients were from 18 to 30 years of age, 444 (35.18%) were from 31 to 50, 382 (30.26%) were from 51 to 70, and 148 (11.72%) were from 71 to 99.

The NPD mean of the RE was 31.65 ± 1.86 (range: 26–38) mm, median was 31.5 mm, and the mode was 31 mm. The NPD of the LE had a mean of 30.90 ± 1.8 (range: 26–37) mm, a median of 31 mm, and a mode of 31 mm.

For the ID a mean of 62.58 ± 3.35 (range: 52.5–73) mm, a median of 62 mm, and a mode of 62 mm were found. The mean of NPA for the population was 1.28 ± 1.04 (range: 0–6) mm. A median of 1 mm and a mode of 1 mm for the NPA in the population were obtained. The confidence interval for the NPA was 1.28 ± 0.057 mm.

Of all patients, 997 (79%) had some degree of NPA and 265 (21%) had a symmetrical NPD, and this difference was statistically significant ($P = 0.0$). 834 patients (83.73%) showed a low NPA, 155 (15.46%) a medium NPA, and 8 (0.8%) a high NPA.

Of the total population of patients with NPA, 76 (7.6%) patients had an NPA of 0.5 mm, 445 (44.6%) of 1 mm, 43 (4.3%) of 1.5 mm, 270 (27.08%) of 2 mm, 10 (1%) of 2.5 mm, 114 (11.43%) of 3 mm, 3 (0.3%) of 3.5 mm, 28 (2.8%) of 4 mm, 5 (0.5%) of 5 mm, and 3 (0.3%) of 6 mm (Figure 2). 729 patients (73.19%) had the RE farther than the LE (RE NPD greater

FIGURE 3: Correlation between the ID and the NPA. A very low correlation between both variables is observed ($r = 0.074$, $P = 0.0$).

than LE NPD) and the remaining 268 (26.8%) had the LE farther than the RE (LE NPD higher than the RE NPD) ($P = 0.0$). A very low correlation between ID and NPA was found ($r = 0.074$, $P = 0.0$) (Figure 3).

4. Discussion

The magnitude of ID has a great importance; however, the perfect symmetry on the face is not present in all people [8, 9]. The ID does not consider the ocular asymmetry. The nature of the NPD has elements that make it variable, such as the gender, race, and age [3, 10–12]. This variability may be influenced by the techniques and instruments used to measure it, due to parallax errors, patient care, reading accuracy, and the fixation point, especially when the millimeter ruler is used [13].

The pupillometer is a tool that can be used to measure both the NPD and the ID, in which the measurement is based on the assessment of the corneal reflex in both eyes while the patient observes a fixation point. This method provides a more accurate measurement than a metric ruler, since when the patient observes the fixation point, the corneal reflection denotes the point known as visual axis or vision line, and without the errors of parallelism of the lanterns or other systems [14].

In this paper the NPA was studied through a digital pupillometer in a Colombian population, finding that 79% of the population has some degree of NPA. 83.73% of the population with NPA presented a low NPA, which could not be clinically significant; nonetheless, the remaining 16.7% presented a greater NPA, which could be significant.

The low correlation found between the ID and the NPA ($r = 0.074$) reflects the lack of relation between the ID and the ocular asymmetry. This could be especially important for the eyeglasses prescription, since if this is not taken into account, it could cause the failure to obtain a correct centering of the ophthalmic lenses, inducing a prismatic effect and a binocular vision, especially in progressive lenses [15, 16].

All of the above underlines the importance of the NPD of each eye as an optometric measure. It should be noted that most automated refractometers and phoropters do not

have the possibility of including the NPD at the time of examination.

According to our results it would be advisable to use the NPD of each eye as standard measurement, especially in patients with medium and high NPA, when the refractive error is high and of course with the increasing use of multifocal lens for patients with presbyopia [17]. Similarly, the relation of the NPA with other variables should be established. This will be studied by the present authors in a further study.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgment

All thanks are due to Dr. Luis Artemo, Professor of the Military University, for his collaboration.

References

- [1] M. Aslankurt, L. Aslan, A. Aksoy, M. Özdemir, and Ş. Dane, "Laterality does not affect the depth perception, but interpupillary distance," *Journal of Ophthalmology*, vol. 2013, Article ID 485059, 5 pages, 2013.
- [2] Y. Eom, J.-S. Song, S.-E. Ahn et al., "Effects of interpupillary distance on stereoacuity: the Frisby Davis distance stereotest versus a 3-dimensional distance stereotest," *Japanese Journal of Ophthalmology*, vol. 57, no. 5, pp. 486–492, 2013.
- [3] C. MacLachlan and H. C. Howland, "Normal values and standard deviations for pupil diameter and interpupillary distance in subjects aged 1 month to 19 years," *Ophthalmic and Physiological Optics*, vol. 22, no. 3, pp. 175–182, 2002.
- [4] P. Thompson, "Eyes wide apart: overestimating interpupillary distance," *Perception*, vol. 31, no. 6, pp. 651–656, 2002.
- [5] C. Garrigosa and J. Perez, "La distancia interpupilar y las distancias de centros de las monturas," *Archivos de la Sociedad Oftalmológica Hispano-Americana*, vol. 28, pp. 541–555, 1968.
- [6] J. Belmonte, "La distancia naso-pupilar en las anisometropías," *ArchSocEspOftalmol*, vol. 10, pp. 923–934, 1977.
- [7] E. P. Osuobeni and M. Al-Fahdi, "Differences between anatomical and physiological interpupillary distance," *Journal of the American Optometric Association*, vol. 65, no. 4, pp. 265–271, 1994.
- [8] Y. Wang, Y. Zhao, and Y. Ai, "Survey on the growth of interpupillary distance of Chinese children aged 5 to 17 years," *Zhonghua Yan KeZaZhi*, vol. 37, no. 1, pp. 63–65, 2001.
- [9] J. S. Pointer, "The interpupillary distance in adult Caucasian subjects, with reference to "readymade" reading spectacle centration," *Ophthalmic and Physiological Optics*, vol. 32, no. 4, pp. 324–331, 2012.
- [10] H. Fesharaki, L. Rezaei, F. Farrahi, T. Banihashem, and A. Jahanbakhshi, "Normal interpupillary distance values in an Iranian population," *Journal of Ophthalmic and Vision Research*, vol. 7, no. 3, pp. 231–234, 2012.
- [11] T. Filipović, "Changes in the interpupillary distance (IPD) with ages and its effect on the near convergence/distance (NC/D) ratio," *Collegium Antropologicum*, vol. 27, no. 2, pp. 723–727, 2003.
- [12] E. P. Osuobeni and K. A. Al-Musa, "Gender differences in interpupillary distance among Arabs," *Optometry and Vision Science*, vol. 70, no. 12, pp. 1027–1030, 1993.
- [13] B. J. Holland and J. Siderov, "Repeatability of measurements of interpupillary distance," *Ophthalmic and Physiological Optics*, vol. 19, no. 1, pp. 74–78, 1999.
- [14] R&C Distribuciones, *Hoja Técnica del Pupímetro*, R&C Distribuciones, Bogotá, Colombia, 2014.
- [15] A.-H. Chen and D. J. O'Leary, "Changes in the interpupillary distance with age and its effect on the near fusion free position," *Australian and New Zealand Journal of Ophthalmology*, vol. 25, no. 1, pp. S6–S8, 1997.
- [16] R. Gockeln, "The influence of interpupillary distance on depth perception," *Klinische Monatsblätter für Augenheilkunde*, vol. 209, no. 4, pp. 205–210, 1996.
- [17] D. B. Elliott and A. Green, "Many ready-made reading spectacles fail the required standards," *Optometry & Vision Science*, vol. 89, no. 4, pp. E446–E451, 2012.