Research Article

Bifurcations of Orbit and Inclination Flips Heteroclinic Loop with Nonhyperbolic Equilibria

Fengjie Geng and Junfang Zhao

School of Science, China University of Geosciences (Beijing), Beijing, 100083, China

Correspondence should be addressed to Fengjie Geng; gengfengjie@cugb.edu.cn

Received 10 October 2013; Accepted 16 January 2014; Published 23 March 2014

Academic Editors: M. Han, Z. Jin, and Y. Xia

Copyright © 2014 F. Geng and J. Zhao. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The bifurcations of heteroclinic loop with one nonhyperbolic equilibrium and one hyperbolic saddle are considered, where the nonhyperbolic equilibrium is supposed to undergo a transcritical bifurcation; moreover, the heteroclinic loop has an orbit flip and an inclination flip. When the nonhyperbolic equilibrium does not undergo a transcritical bifurcation, we establish the coexistence and noncoexistence of the periodic orbits and homoclinic orbits. While the nonhyperbolic equilibrium undergoes the transcritical bifurcation, we obtain the noncoexistence of the periodic orbits and homoclinic orbits and the existence of two or three heteroclinic orbits.

1. Introduction

In recent years, a great deal of mathematical efforts has been devoted to the bifurcation problems of homoclinic and heteroclinic orbits with high codimension, for example, the bifurcations of homoclinic or heteroclinic loop with orbit flip, the bifurcations of homoclinic or heteroclinic loop with inclination flip, and so forth; see [1–5] and the references therein. However, most of these papers considered the bifurcation problems of orbits connecting hyperbolic equilibria, and limited work has been done in the corresponding problems with nonhyperbolic equilibria; see [6–8]. To fill this gap, we investigate the bifurcations of orbit and inclination flip heteroclinic orbits with one nonhyperbolic equilibrium and one hyperbolic saddle. The method is using the fundamental solution matrix of the linear variational system to obtain the Poincaré map, which is easier to get the bifurcation equations.

Consider the following C^r ($r \geq 5$) system

$$\dot{z} = g(z, \lambda, \mu) \quad (1)$$

and its unperturbed system

$$\dot{z} = f(z), \quad (2)$$

where $z \in \mathbb{R}^4$, the vector field g depends on the parameters (λ, μ) , $\lambda \in \mathbb{R}$, $\mu \in \mathbb{R}^l$, $l \geq 2$, $0 \leq \lambda$, $|\mu| \ll 1$, $g(z, 0, 0) = f(z)$, $g(p_1, 0, \mu) = 0$, and $g(p_2, \lambda, \mu) = 0$. Moreover, the parameter λ governs bifurcation of the nonhyperbolic equilibrium, while μ controls bifurcations of the heteroclinic orbits.

Assuming system (2) has a heteroclinic loop Γ connecting its two equilibria p_1, p_2 , where $\Gamma = \Gamma^1 \cup \Gamma^2$, $\Gamma^i = \{z = r_i(t) : t \in \mathbb{R}, r_i(+\infty) = r_{i+1}(-\infty) = p_{i+1}, i = 1, 2, r_3(t) = r_1(t), \text{ and } p_3 = p_1\}$. Furthermore, the linearization $Df(p_1)$ has real eigenvalues $0, \lambda_1^1, -\rho_1^1$, and $-\rho_1^2$ satisfying $-\rho_1^2 < -\rho_1^1 < 0 < \lambda_1^1$; $Df(p_2)$ has simple real eigenvalues $\lambda_2^1, \lambda_2^2, -\rho_2^1$, and $-\rho_2^2$ fulfilling $-\rho_2^2 < -\rho_2^1 < 0 < \lambda_2^1 < \lambda_2^2$.

The following conditions hold in the whole paper:

$$(H_1)$$

$$e_i^\pm = \lim_{t \rightarrow \pm\infty} \frac{\dot{r}_i(-t)}{|\dot{r}_i(-t)|}, \quad (3)$$

where $e_1^+ \in T_{p_1} W_1^{cu}$, $e_1^- \in T_{p_2} W_2^{ss}$, $e_2^+ \in T_{p_2} W_2^u$, $e_2^- \in T_{p_1} W_1^s$, and $e_1^- \in T_{p_2} W_2^{ss}$ mean that Γ^1 is a heteroclinic orbit with

orbit flip, W_1^{cu} is the center unstable manifold of p_1 , W_i^u (resp., W_i^s) is the unstable (resp., stable) manifold of p_i , and W_i^{uu} (resp., W_i^{ss}) is the strong unstable (resp., stable) manifold of p_i , $i = 1, 2$. Moreover,

$$\begin{aligned} \dim(T_{r_1(t)}W_1^c \cap T_{r_1(t)}W_2^s) \\ = \dim(T_{r_1(t)}W_1^{cu} \cap T_{r_1(t)}W_2^s) = 1. \end{aligned} \quad (4)$$

(H_2)

$$\begin{aligned} \lim_{t \rightarrow +\infty} T_{r_1(t)}W_1^{cu} &= \text{span}\{e_1^-, T_{p_2}W_2^{uu}\}, \\ \lim_{t \rightarrow +\infty} T_{r_2(t)}W_2^u &= \text{span}\{e_2^-, T_{p_1}W_1^u\}, \\ \lim_{t \rightarrow -\infty} T_{r_1(t)}W_2^s &= \text{span}\{e_1^+, T_{p_1}W_1^{ss}\}, \\ \lim_{t \rightarrow -\infty} T_{r_2(t)}W_1^s &= \text{span}\{e_2^+, T_{p_2}W_2^s\}, \end{aligned} \quad (5)$$

where the first three equations mean that the center unstable manifold W_1^{cu} of p_1 , the stable (resp., unstable) manifold W_2^s (resp., W_2^u) of p_2 are fulfilling the strong inclination property. And the fourth equation implies that the stable manifold W_1^s is of inclination flip as $t \rightarrow -\infty$.

It is worthy of noting that, for any integers $m \geq 1$ and $n \geq 1$, if we assume $\dim(W_1^{uu}) = \dim(W_2^{ss}) = m$ and $\dim(W_1^{ss}) = \dim(W_2^s) = n$, then all the results achieved in this paper are still valid.

Let $\lambda \in \mathbb{R}$ be a parameter to control the transcritical bifurcation of system (1), let the x -axis be the tangent space of the center manifold at p_1 , and let $\theta(x, \lambda, \mu)$ be the vector field defined on the center manifold; then by [9], we may assume

$$\begin{aligned} (H_3) \quad \theta(x_{p_1}, \lambda, \mu) &= 0, \quad (\partial\theta/\partial x)(x_{p_1}, 0, 0) = 0, \\ (\partial^2\theta/\partial x^2)(x_{p_1}, 0, 0) &> 0, \quad (\partial^2\theta/\partial x\partial\lambda)(x_{p_1}, 0, 0) < 0, \\ (\partial^2\theta/\partial x\partial\mu)(x_{p_1}, 0, \mu) &= 0, \text{ where } x_{p_1} \text{ is the } x \\ &\text{component of } p_1. \end{aligned}$$

If (H_3) is true, then system (1) exhibits the transcritical bifurcation, that is, when $\lambda > 0$ (or $\lambda < 0$; in this paper, we only consider the case $\lambda > 0$; for the case $\lambda < 0$, one may discuss it similarly); there are two hyperbolic saddles p_1^0 and p_1^1 bifurcated from p_1 . Denote by $p_1^0 = p_1 = (0, 0, 0, 0)^*$ and $p_1^1 = p_1 + (\lambda_p, 0, 0, 0)^*$, where $\lambda_p = \theta_0\lambda + O(\lambda^2) + O(\lambda\mu)$ and $\theta_0 = -(\partial^2\theta/\partial x\partial\lambda)(x_{p_1}, 0, 0)/(\partial^2\theta/\partial x^2)(x_{p_1}, 0, 0)$. Moreover, $\dim(W_{p_1^0}^s) = 3$, $\dim(W_{p_1^0}^u) = 1$, and $\dim(W_{p_1^1}^u) = \dim(W_{p_1^1}^s) = 2$.

The present paper is built up as follows. In Section 2, we devote it to deriving the successor functions by constructing a suitable Poincaré Map. The analysis to the bifurcations of system (2) is presented in Section 3, where we establish the existence of the heteroclinic loop, the homoclinic orbits, and the three or two heteroclinic orbits and the coexistence of a periodic orbit and a homoclinic loop, and the difference between the heteroclinic loop with hyperbolic equilibria and nonhyperbolic equilibria is revealed.

2. Normal Form and Poincaré Map

Let the neighborhood U_i of p_i be small enough and straight the local manifolds of $W_i^s, W_2^{uu}, W_i^{ss}$, and $i = 1, 2$ in the neighborhood U_i . And then by virtue of the invariance of these manifolds and a scale transformation $x \rightarrow \theta_{xx}^{-1}(x_{p_1}, 0, 0)x$ and $\lambda \rightarrow -\theta_{x\lambda}^{-1}(x_{p_1}, 0, 0)\lambda$, system (1) has the following expression in U_1 :

$$\begin{aligned} \dot{x} &= -\lambda_p x + x^2 + O(u)[O(y) + O(v)] \\ &\quad + O(x)[O(y) + O(u) + O(v)] + O(x)O(x^2), \\ \dot{y} &= [-\rho_1^1(\alpha) + \dots]y + O(v)[O(x) + O(u)], \\ \dot{u} &= [\lambda_1^1(\alpha) + \dots]u + O(x)[O(y) + O(v)], \\ \dot{v} &= [-\rho_1^2(\alpha) + \dots]v + O(y)[O(x) + O(y) + O(u)], \end{aligned} \quad (6)$$

and in U_2 it takes the following form:

$$\begin{aligned} \dot{x} &= [\lambda_2^1(\alpha) + \dots]x + O(u)[O(y) + O(v)], \\ \dot{y} &= [-\rho_2^1(\alpha) + \dots]y + O(v)[O(x) + O(u)], \\ \dot{u} &= [\lambda_2^2(\alpha) + \dots]u + O(x)[O(x) + O(y) + O(v)], \\ \dot{v} &= [-\rho_2^2(\alpha) + \dots]v + O(y)[O(x) + O(y) + O(u)], \end{aligned} \quad (7)$$

where $\alpha = (\lambda, \mu)$, $\lambda_p = \lambda + O(\lambda^2) + O(\lambda\mu)$, $\lambda_1^1(0) = \lambda_1^1$, $\rho_i^j(0) = \rho_i^j$, $j = 1, 2$, $i = 1, 2$, $\lambda_2^j(0) = \lambda_2^j$, $j = 1, 2$.

From the normal form (6), (7), and the condition (H_1), we may select $-T_i$ and T_i such that

$$\begin{aligned} r_1(-T_1) &= (\delta, 0, 0, 0)^*, \quad r_1(T_1) = (0, 0, 0, \delta)^*, \\ r_1(-T_2) &= (\delta, 0, 0, \delta_u)^*, \quad r_2(T_2) = (0, \delta, 0, \delta_v)^*, \end{aligned} \quad (8)$$

where $\delta > 0$ is small enough such that $\{(x, y, u, v) : |x|, |y|, |u|, |v| < 2\delta\} \subset U_i$ and $|\delta_u| = o(\delta)$, $|\delta_v| = o(\delta)$.

Consider the linear variational system

$$\dot{z} = Df(r_i(t))z \quad (9)_i$$

and its adjoint system

$$\dot{\phi} = -(Df(r_i(t)))^* \phi, \quad (10)_i$$

$i = 1, 2$, where $(Df(r_i(t)))^*$ is the transposed matrix of $Df(r_i(t))$.

Supposing $Z_i(t) = (z_i^1(t), z_i^2(t), z_i^3(t), z_i^4(t))$ is a fundamental solution matrix of $(9)_i$, then, we arrive at the following lemma.

Lemma 1. If conditions $(H_1)-(H_3)$ are satisfied, then

(1) there exists a fundamental solution matrix of $(9)_1$ satisfying

$$\begin{aligned} z_1^1(t) &\in (T_{r_1(t)} W_1^{cu})^c \cap (T_{r_1(t)} W_2^s)^c, \\ z_1^2(t) &= -\frac{\dot{r}_1(t)}{|\dot{r}_1(T_1)|} \in T_{r_1(t)} W_1^c \cap T_{r_1(t)} W_2^s, \\ z_1^3(t) &\in T_{r_1(t)} W_1^{cu} \cap (T_{r_1(t)} W_2^s)^c, \\ z_1^4(t) &\in (T_{r_1(t)} W_1^{cu})^c \cap T_{r_1(t)} W_2^s \end{aligned} \quad (11)$$

such that

$$\begin{aligned} Z_1(-T_1) &= \begin{pmatrix} w_1^{11} & w_1^{21} & 0 & w_1^{41} \\ w_1^{12} & 0 & 0 & w_1^{42} \\ w_1^{13} & w_1^{23} & 1 & w_1^{43} \\ 0 & 0 & 0 & w_1^{44} \end{pmatrix}, \\ Z_1(T_1) &= \begin{pmatrix} 1 & 0 & w_1^{31} & 0 \\ \tilde{w}_1^{12} & 0 & w_1^{32} & 1 \\ 0 & 0 & w_1^{33} & 0 \\ 0 & 1 & w_1^{34} & 0 \end{pmatrix}; \end{aligned} \quad (12)$$

(2) $(9)_2$ has a fundamental solution matrix fulfilling

$$\begin{aligned} z_2^1(t) &\in (T_{r_2(t)} W_2^u)^c \cap (T_{r_2(t)} W_1^s)^c, \\ z_2^2(t) &= -\frac{\dot{r}_2(t)}{|\dot{r}_2(T_2)|} \in T_{r_2(t)} W_2^u \cap T_{r_2(t)} W_1^s, \\ z_2^3(t) &\in T_{r_2(t)} W_2^u \cap (T_{r_2(t)} W_1^s)^c, \\ z_2^4(t) &\in (T_{r_2(t)} W_2^u)^c \cap T_{r_2(t)} W_1^s \end{aligned} \quad (13)$$

such that

$$\begin{aligned} Z_2(-T_2) &= \begin{pmatrix} w_2^{11} & w_2^{21} & 0 & w_2^{41} \\ 0 & 0 & 0 & w_2^{42} \\ w_2^{13} & w_2^{23} & 1 & w_2^{43} \\ w_2^{14} & 0 & 0 & 0 \end{pmatrix}, \\ Z_2(T_2) &= \begin{pmatrix} 1 & 0 & w_2^{31} & 0 \\ 0 & 1 & w_2^{32} & 0 \\ 0 & 0 & w_2^{33} & 0 \\ \tilde{w}_2^{14} & w_2^{24} & w_2^{34} & 1 \end{pmatrix}, \end{aligned} \quad (14)$$

where $w_i^{21} < 0$, $w_1^{12} w_i^{33} w_2^{14} w_2^{42} \neq 0$, $|(w_i^{33})^{-1} w_i^{3j}| \ll 1$, $j = 1, 2, 4$, $i = 1, 2$.

Now, let $(z_i^1(t), z_i^2(t), z_i^3(t), z_i^4(t))$ be a new local active coordinate system along Γ^i . Given $\Phi_i(t) = (\phi_i^1(t), \phi_i^2(t), \phi_i^3(t), \phi_i^4(t)) = (Z_i^{-1}(t))^*$, then $\Phi_i(t)$ is the fundamental solution matrix of $(10)_i$, $i = 1, 2$.

Let $z = r_i(t) + Z_i(t)N_i(t) \triangleq h_i(t)$, where $N_i(t) = (n_i^1, 0, n_i^3, n_i^4)^*$, $i = 1, 2$. Defining the cross sections

$$\begin{aligned} S_i^0 &= \{z = h_i(-T_i) : |x|, |y|, |u|, |v| < 2\delta\}, \\ S_i^1 &= \{z = h_i(T_i) : |x|, |y|, |u|, |v| < 2\delta\} \end{aligned} \quad (15)$$

of Γ_i at $t = -T_i$ and $t = T_i$, respectively, $i = 1, 2$.

Now that if $q_i^0 \in S_i^0$ and $q_i^1 \in S_i^1$, then

$$\begin{aligned} q_i^0 &= (x_i^0, y_i^0, u_i^0, v_i^0)^* = r_i(-T_i) + Z_1(-T_i) N_i(-T_i), \\ N_i(-T_i) &= (n_i^{0,1}, 0, n_i^{0,3}, n_i^{0,4})^*, \\ q_i^1 &= (x_i^1, y_i^1, u_i^1, v_i^1)^* = r_i(T_i) + Z_i(T_i) N_i(T_i), \\ N_i(T_i) &= (n_i^{1,1}, 0, n_i^{1,3}, n_i^{1,4})^*. \end{aligned} \quad (16)$$

Based on the expressions of $Z_i(-T_i)$ and $Z_i(T_i)$, we get their new coordinates of $q_i^0(n_i^{0,1}, 0, n_i^{0,3}, n_i^{0,4})^*$ and $q_i^1(n_i^{1,1}, 0, n_i^{1,3}, n_i^{1,4})^*$; that is,

$$\begin{aligned} n_1^{0,1} &= (w_1^{12})^{-1} [y_1^0 - w_1^{42} (w_1^{44})^{-1} v_1^0], \\ n_1^{0,3} &= u_1^0 - w_1^{13} (w_1^{12})^{-1} y_1^0 \\ &\quad + [w_1^{13} w_1^{42} (w_1^{12})^{-1} - w_1^{43}] (w_1^{44})^{-1} v_1^0, \\ n_1^{0,4} &= (w_1^{44})^{-1} v_1^0, \\ x_1^0 &= \delta + w_1^{11} n_1^{0,1} + w_1^{41} n_1^{0,4} \approx \delta, \\ n_1^{1,1} &= x_1^1 - w_1^{31} (w_1^{33})^{-1} u_1^1, \\ n_1^{1,3} &= (w_1^{33})^{-1} u_1^1, \\ n_1^{1,4} &= y_1^1 - \tilde{w}_1^{12} x_1^1 + (\tilde{w}_1^{12} w_1^{31} - w_1^{32}) (w_1^{33})^{-1} u_1^1, \\ v_1^1 &\approx \delta, \\ n_2^{0,1} &= (w_2^{14})^{-1} v_2^0, \\ n_2^{0,3} &= u_2^0 - \delta_2^u - w_2^{13} (w_2^{14})^{-1} v_2^0 - w_2^{43} (w_2^{42})^{-1} y_2^0, \\ n_2^{0,4} &= (w_2^{42})^{-1} y_2^0, \\ x_2^0 &\approx \delta, \\ n_2^{1,1} &= x_0^1 - w_2^{31} (w_2^{33})^{-1} u_0^1, \\ n_2^{1,3} &= (w_2^{33})^{-1} u_0^1, \\ n_2^{1,4} &= v_0^1 - \delta_2^v - \tilde{w}_2^{14} x_0^1 + (\tilde{w}_2^{14} w_2^{31} - w_2^{34}) (w_2^{33})^{-1} u_0^1, \\ y_0^1 &\approx \delta. \end{aligned} \quad (17)$$

Next, we divide our establishment of the Poincaré map in the new coordinate system in three steps.

First, consider the map $F_i^1 : S_i^0 \mapsto S_i^1$. Put $z = h_i(t)$ into (1); we have

$$\begin{aligned} \dot{r}_i(t) + \dot{Z}_i(t) N_i(t) + Z_i(t) \dot{N}_i(t) \\ = g(r_i(t) + Z_i(t) N_i(t), \lambda, \mu) \\ = g(r_i(t), 0, 0) + g_z(r_i(t), 0, 0) Z_i(t) N_i(t) \\ + g_\lambda(r_i(t), 0, 0) \lambda + g_\mu(r_i(t), 0, 0) \mu + \text{h.o.t.} \\ = f(r_i(t)) + Df(r_i(t)) Z_i(t) N_i(t) \\ + g_\lambda(r_i(t), 0, 0) \lambda + g_\mu(r_i(t), 0, 0) \mu + \text{h.o.t.} \end{aligned} \quad (18)$$

According to the fact $\dot{r}_i(t) = f(r_i(t))$ and $\dot{Z}_i(t) = Df(r_i(t)) Z_i(t)$, it then yields to that

$$\begin{aligned} \dot{N}_i(t) = Z_i^{-1}(t) [g_\lambda(r_i(t), 0, 0) \lambda + g_\mu(r_i(t), 0, 0) \mu] \\ + \text{h.o.t.} \end{aligned} \quad (19)$$

Integrating the above equation from $-T_i$ to T_i , we arrive at

$$\begin{aligned} N_i(T_i) = N_i(-T_i) + \int_{-T_i}^{T_i} Z_i^{-1}(t) g_\lambda(r_i(t), 0, 0) \lambda dt \\ + \int_{-T_i}^{T_i} Z_i^{-1}(t) g_\mu(r_i(t), 0, 0) \mu dt + \text{h.o.t.} \end{aligned} \quad (20)$$

Noticing that $\Phi_i^*(t) = Z_i^{-1}(t)$, then

$$n_i^{1,j} = n_i^{0,j} + M_{i\lambda}^j \lambda + M_{i\mu}^j \mu + \text{h.o.t.}, \quad j = 1, 3, 4, \quad (21)_i$$

where

$$\begin{aligned} M_{i\lambda}^j &= \int_{-T_i}^{T_i} \phi_i^{j*} g_\lambda(r_i(t), 0, 0) dt, \\ M_{i\mu}^j &= \int_{-T_i}^{T_i} \phi_i^{j*} g_\mu(r_i(t), 0, 0) dt, \quad j = 1, 3, 4. \end{aligned} \quad (22)_i$$

Together with (17) and (21)_i, (22)_i then defines the map $F_i^1 : S_i^0 \mapsto S_i^1$, $(n_i^{0,1}, 0, n_i^{0,3}, n_i^{0,4}) \mapsto (n_i^{1,1}, 0, n_i^{1,3}, n_i^{1,4})$.

Next, to construct the map $F_i^0 : S_{i-1}^1 \mapsto S_i^0$ (where $S_0^1 = S_2^1$). Let τ_i , $i = 1, 2$ be the flying time from $q_{i-1}^1(x_{i-1}^1, y_{i-1}^1, u_{i-1}^1, v_{i-1}^1)^*$ to $q_i^0(x_i^0, y_i^0, u_i^0, v_i^0)^*$; set $s_1 = e^{-\rho_1^1(\alpha)\tau_1}$ and $s_2 = e^{-\rho_2^1(\alpha)\tau_2}$. By virtue of the approximate solution of system (6) and (7), if we neglect the higher terms, then the expression of $F_i^0 : S_{i-1}^1 \mapsto S_i^0$ is

$$\begin{aligned} x_1^0 &\approx \frac{x_1^0}{h(s_1)}, & y_1^0 &\approx s_1 y_0^1, \\ u_1^0 &\approx s_1^{\lambda_1^1(\alpha)/\rho_1^1(\alpha)} u_1^0, & v_1^0 &\approx s_1^{\rho_1^2(\alpha)/\rho_1^1(\alpha)} v_0^1 \end{aligned} \quad (23)$$

and $F_2^0 : S_1^1 \mapsto S_2^1$ is

$$\begin{aligned} x_1^1 &\approx s_2^{\lambda_2^1(\alpha)/\rho_2^1(\alpha)} x_2^0, & y_2^0 &\approx s_2 y_1^1, \\ u_1^1 &\approx s_2^{\lambda_2^2(\alpha)/\rho_2^1(\alpha)} u_2^0, & v_2^0 &\approx s_2^{\rho_2^2(\alpha)/\rho_2^1(\alpha)} v_1^1, \end{aligned} \quad (24)$$

FIGURE 1

where (s_i, u_i^0, v_{i-1}^1) , $i = 1, 2$ are called Shilnikov coordinates, and

$$h(s) = \begin{cases} (\lambda_p)^{-1} [x_1^0 - (x_1^0 - \lambda_p) s^{\lambda_p/\rho_1^1(\alpha)}], & \lambda_p \neq 0, \\ 1 - (\rho_1^1(\alpha))^{-1} x_1^0 \ln s, & \lambda_p = 0. \end{cases} \quad (25)$$

Since the nonhyperbolic equilibrium p_1 undergoes a transcritical bifurcation based on the structure of orbits in U_1 , we may see that the equation $x_0^1 \approx x_1^0/h(s_1)$ holds only for $x_0^1 \geq \lambda_p$. While for $x_0^1 \in [-\beta, \lambda_p)$ ($0 < \beta \ll 1$), the map F_1^0 is well defined only if $s_1 = 0$ (see Figure 1). So, we extend the domain of F_1^0 , defining

$$x_1^0 = \delta, \quad s_1 = 0, \quad \text{if } x_0^1 \in [-\beta, \lambda_p). \quad (26)$$

The final step is to compose the maps F_i^0 and F_i^1 , and then $F_1 = F_1^1 \circ F_1^0 : S_0^1 \mapsto S_1^1$ can be expressed as

$$\begin{aligned} n_1^{1,1} &= (w_1^{12})^{-1} \delta s_1 - (w_1^{12})^{-1} w_1^{42} (w_1^{44})^{-1} s_1^{\rho_1^2(\alpha)/\rho_1^1(\alpha)} v_0^1 \\ &\quad + M_{1\lambda}^1 \lambda + M_{1\mu}^1 \mu + \text{h.o.t.}, \\ n_1^{1,3} &= u_1^0 - w_1^{13} (w_1^{12})^{-1} \delta s_1 \\ &\quad + [w_1^{13} w_1^{42} (w_1^{12})^{-1} - w_1^{43}] (w_1^{44})^{-1} s_1^{\rho_1^2(\alpha)/\rho_1^1(\alpha)} v_0^1 \\ &\quad + M_{1\lambda}^3 \lambda + M_{1\mu}^3 \mu + \text{h.o.t.}, \\ n_1^{1,4} &= (w_1^{44})^{-1} s_1^{\rho_1^2(\alpha)/\rho_1^1(\alpha)} v_0^1 + M_{1\lambda}^4 \lambda + M_{1\mu}^4 \mu + \text{h.o.t.} \end{aligned} \quad (27)$$

and $F_2 = F_2^1 \circ F_2^0 : S_1^1 \mapsto S_2^1 (= S_0^1)$ as

$$\begin{aligned} n_2^{1,1} &= (w_2^{14})^{-1} \delta s_2^{\rho_2^2(\alpha)/\rho_2^1(\alpha)} + M_{2\lambda}^1 \lambda + M_{2\mu}^1 \mu + \text{h.o.t.}, \\ n_2^{1,3} &= u_2^0 - \delta s_2^u - w_2^{13} (w_2^{14})^{-1} \delta s_2^{\rho_2^2(\alpha)/\rho_2^1(\alpha)} \\ &\quad - w_2^{43} (w_2^{42})^{-1} s_2 y_1^1 + M_{2\lambda}^3 \lambda + M_{2\mu}^3 \mu + \text{h.o.t.}, \\ n_2^{1,4} &= (w_2^{44})^{-1} s_2 y_1^1 + M_{2\lambda}^4 \lambda + M_{2\mu}^4 \mu + \text{h.o.t.} \end{aligned} \quad (28)$$

Set $G_i = F_i(q_{i-1}^1) - q_i^1$, $i = 1, 2$. Combining (21)_i, (23), (24), (27), and (28), we derive the successor functions G_i^j :

$$\begin{aligned}
 G_1^1 &= (w_1^{12})^{-1} \delta s_1 - \delta s_2^{\lambda_1^1/\rho_2^1(\alpha)} + M_{1\lambda}^1 \lambda + M_{1\mu}^1 \mu + \text{h.o.t.}, \\
 G_1^3 &= u_1^0 - w_1^{13} (w_1^{12})^{-1} \delta s_1 - (w_1^{33})^{-1} s_2^{\lambda_2^1(\alpha)/\rho_2^1(\alpha)} u_2^0 \\
 &\quad + M_{1\lambda}^3 \lambda + M_{1\mu}^3 \mu + \text{h.o.t.}, \\
 G_1^4 &= (w_1^{44})^{-1} s_1^{\rho_2^1(\alpha)/\rho_1^1(\alpha)} v_0^1 - y_1^1 + \bar{w}_1^{12} \delta s_2^{\lambda_2^1(\alpha)/\rho_2^1(\alpha)} \\
 &\quad + M_{1\lambda}^4 \lambda + M_{1\mu}^4 \mu + \text{h.o.t.}, \\
 G_2^1 &= (w_2^{14})^{-1} \delta s_2^{\rho_2^2(\alpha)/\rho_2^1(\alpha)} - \delta h(s_1) \\
 &\quad + w_2^{31} (w_2^{33})^{-1} s_1^{\lambda_1^1(\alpha)/\rho_1^1(\alpha)} u_1^0 + M_{2\lambda}^1 \lambda + M_{2\mu}^1 \mu + \text{h.o.t.}, \\
 G_2^3 &= u_2^0 - \delta_2^u - w_2^{13} (w_1^{14})^{-1} \delta s_2^{\rho_2^2(\alpha)/\rho_2^1(\alpha)} \\
 &\quad - w_2^{43} (w_2^{42})^{-1} s_2^{\rho_2^1(\alpha)/\rho_2^1(\alpha)} y_1^1 \\
 &\quad - (w_2^{33})^{-1} s_1^{\lambda_1^1(\alpha)/\rho_1^1(\alpha)} u_1^0 + M_{2\lambda}^3 \lambda + M_{2\mu}^3 \mu + \text{h.o.t.}, \\
 G_2^4 &= (w_2^{44})^{-1} s_2 y_1^1 - v_0^1 + \delta_2^v + \bar{w}_2^{14} \delta h(s_1) \\
 &\quad - (w_2^{34} - \bar{w}_2^{14} w_2^{31}) (w_2^{33})^{-1} s_1^{\lambda_1^1(\alpha)/\rho_1^1(\alpha)} u_1^0 \\
 &\quad + M_{2\lambda}^4 \lambda + M_{2\mu}^4 \mu + \text{h.o.t.}
 \end{aligned} \tag{29}$$

It is easy to see that what we need to do is considering the solutions of

$$(G_1^1, G_1^3, G_1^4, G_2^1, G_2^3, G_2^4) = 0 \tag{30}$$

with $s_1 \geq 0$ and $s_2 \geq 0$. This is because the solution of (30) with $s_1 = s_2 = 0$ (resp., $s_1 > 0, s_2 > 0$; $s_1 = 0, s_2 > 0$ or $s_1 > 0, s_2 = 0$) means that system (1) has a heteroclinic loop (resp., a periodic orbit; homoclinic loop).

3. Main Results

Based on the expressions of the successor functions and the implicit function theorem, we know that the equation $(G_1^3, G_1^4, G_2^3, G_2^4) = 0$ has a unique solution $(u_1^0, u_2^0, y_1^1, v_0^1)$. And putting it into $(G_1^1, G_2^1) = 0$, then we obtain the following bifurcation equations:

$$\begin{aligned}
 (w_1^{12})^{-1} \delta s_1 - \delta s_2^{\lambda_1^1/\rho_2^1} + M_{1\lambda}^1 \lambda + M_{1\mu}^1 \mu + \text{h.o.t.} &= 0, \\
 (w_2^{14})^{-1} \delta s_2^{\rho_2^2/\rho_2^1} - \delta h(s_1) + M_{2\lambda}^1 \lambda + M_{2\mu}^1 \mu \\
 + w_2^{31} (w_2^{33})^{-1} s_1^{\lambda_1^1/\rho_1^1} [w_1^{13} (w_1^{12})^{-1} \delta s_1 + (w_1^{33})^{-1} \\
 \times s_2^{\lambda_2^1/\rho_2^1} (\delta_2^u - M_{2\lambda}^3 \lambda - M_{2\mu}^3 \mu) \\
 - M_{1\lambda}^3 \lambda - M_{1\mu}^3 \mu] + \text{h.o.t.} &= 0.
 \end{aligned} \tag{31}$$

Firstly, we consider the case $\lambda = 0$, which means the transcritical bifurcation does not happen. By (23) and (25), (31) turns to

$$\begin{aligned}
 (w_1^{12})^{-1} \delta s_1 - \delta s_2^{\lambda_1^1/\rho_2^1} + M_{1\mu}^1 \mu + \text{h.o.t.} &= 0, \\
 (w_2^{12})^{-1} \delta s_2^{\rho_2^2/\rho_2^1} - \frac{\delta}{1 - (\rho_1^1)^{-1} \delta \ln s_1} + M_{2\mu}^1 \mu \\
 + w_2^{31} (w_2^{33})^{-1} w_1^{13} (w_1^{12})^{-1} \delta s_1^{\lambda_1^1/\rho_1^1+1} \\
 - w_2^{31} (w_2^{33})^{-1} s_1^{\lambda_1^1/\rho_1^1} M_{1\mu}^3 \mu + \text{h.o.t.} &= 0.
 \end{aligned} \tag{32}$$

Noticing that $\lambda_1^1/\rho_1^1 > 0$, which shows $\lim_{s_1 \rightarrow 0} s_1^{\lambda_1^1/\rho_1^1} (1 - (\rho_1^1)^{-1} \delta \ln s_1) = 0$, it then follows that

$$\begin{aligned}
 s_1 - w_1^{12} s_2^{\lambda_2^1/\rho_2^1} + \delta^{-1} w_1^{12} M_{1\mu}^1 \mu + \text{h.o.t.} &= 0, \\
 (w_2^{12})^{-1} s_2^{\rho_2^2/\rho_2^1} - \frac{1}{1 - (\rho_1^1)^{-1} \delta \ln s_1} + \delta^{-1} M_{2\mu}^1 \mu + \text{h.o.t.} &= 0.
 \end{aligned} \tag{33}$$

From the above bifurcation equations, we obtain the following results immediately.

Theorem 2. Let the conditions (H_1) – (H_3) be true and $M_{i\mu}^1 \neq 0$, $i = 1, 2$. Then, for $\lambda = 0$ and $0 < |\mu| \ll 1$, one has

(i) for $\text{rank}(M_{1\mu}^1, M_{2\mu}^1) = 2$, there exists a codimension 2 surface

$$L_{12} = \{\mu : M_{1\mu}^1 \mu + \text{h.o.t.} = M_{2\mu}^1 \mu + \text{h.o.t.} = 0\} \tag{34}$$

such that system (1) has a unique heteroclinic loop near Γ if and only if $\mu \in L_{12}$, where the surface L_{12} has a normal plane $\text{span}\{M_{1\mu}^1, M_{2\mu}^1\}$ at $\mu = 0$.

(ii) there exists an $(l-1)$ -dimensional surface

$$L_1^2 = \left\{ \mu : \delta^{-1} w_2^{12} M_{2\mu}^1 \mu + (\delta^{-1} M_{1\mu}^1 \mu)^{\rho_2^2/\lambda_2^1} + \text{h.o.t.} = 0, \right.$$

$$\left. M_{1\mu}^1 \mu > 0 \right\} \tag{35}$$

$$\left(\text{resp., } L_2^1 = \left\{ \mu : \frac{\delta}{1 - (\rho_1^1)^{-1} \delta \ln (-\delta^{-1} w_1^{12} M_{1\mu}^1 \mu)} \right. \right. \\
 \left. \left. - M_{2\mu}^1 \mu + \text{h.o.t.} = 0, w_1^{12} M_{1\mu}^1 \mu < 0 \right\} \right) \tag{36}$$

such that system (1) has a unique homoclinic loop connecting p_1 (resp., connecting p_2) near Γ if and only if $\mu \in L_1^2$ (resp., $\mu \in L_2^1$).

Proof. The result (i) will be proved by putting $s_1 = s_2 = 0$ into (33).

If we assume $s_1 = 0$ and $s_2 > 0$ in (33), then

$$\begin{aligned} s_2^{\lambda_2/\rho_2^1} &= \delta^{-1} M_{1\mu}^1 \mu + \text{h.o.t.}, \\ (w_2^{12})^{-1} s_2^{\rho_2^2/\rho_2^1} + \delta^{-1} M_{2\mu}^1 \mu + \text{h.o.t.} &= 0, \end{aligned} \quad (37)$$

which means

$$(w_2^{12})^{-1} s_2^{\rho_2^2/\lambda_2^1} + \delta^{-1} M_{2\mu}^1 \mu + \text{h.o.t.} = 0. \quad (38)$$

It follows that there exists an $(l-1)$ -dimensional surface L_1^2 given by (35) such that (33) has a unique solution $s_1 = 0$, $s_2 = s_2(\mu) > 0$ as $\mu \in L_1^2$ and $0 < |\mu| \ll 1$. This implies system (1) has a homoclinic loop connecting p_1 . The existence of L_1^2 can be obtained similarly.

This completes the proof. $\square$

Remark 3. There is no difficulty to see that L_1^2 has a normal vector $M_{2\mu}^1$ at $\mu = 0$ as $\rho_2^2 > \lambda_2^1$, while for $\rho_2^2 < \lambda_2^1$ (resp., $\rho_2^2 > \lambda_2^1$) it has a normal vector $M_{1\mu}^1$ (resp., $M_{1\mu}^1 + w_2^{12} M_{2\mu}^1$) at $\mu = 0$.

Theorem 4. Assume the conditions (H_1) – (H_3) hold and $M_{i\mu}^1 \neq 0$, $i = 1, 2$. Then for $\lambda = 0$, $\mu \in L_1^2$, and $0 < |\mu| \ll 1$, the periodic orbit and homoclinic loop with p_1 of system (1) cannot coexist.

Proof. Theorem 2 shows that if $\mu \in L_1^2$ and $0 < |\mu| \ll 1$, then system (1) has a homoclinic loop with p_1 . Setting $s_1 \geq 0$, $s_2^{\lambda_2^1/\rho_2^1} = (w_1^{12})^{-1} s_1 + \delta^{-1} M_{1\mu}^1 \mu + \text{h.o.t.} > 0$, and $\mu \in L_1^2$, then (33) is reduced to

$$\begin{aligned} V_1(s_1) &\triangleq \left[(w_1^{12})^{-1} s_1 + \delta^{-1} M_{1\mu}^1 \mu \right]^{\rho_2^2/\lambda_2^1} + \delta^{-1} w_2^{12} M_{2\mu}^1 \mu \\ &+ \text{h.o.t.} = \frac{w_2^{12}}{1 - (\rho_1^1)^{-1} \delta \ln s_1} \triangleq N_1(s_1). \end{aligned} \quad (39)$$

Notice that $V_1(0) = N_1(0)$ and

$$\begin{aligned} V_1'(s_1) &= \frac{\rho_2^2}{\lambda_2^1} (w_1^{12})^{-1} \left[(w_1^{12})^{-1} s_1 + \delta^{-1} M_{1\mu}^1 \mu \right]^{\rho_2^2/\lambda_2^1 - 1}, \\ N_1'(s_1) &= \frac{w_2^{12} (\rho_1^1)^{-1} \delta}{(1 - (\rho_1^1)^{-1} \delta \ln s_1)^2 s_1}. \end{aligned} \quad (40)$$

If $w_1^{12} w_2^{12} < 0$, then $V_1'(s_1) N_1'(s_1) < 0$; it is obvious that $V_1(s_1) = N_1(s_1)$ has no sufficiently small positive solutions.

While $\rho_2^2 > \lambda_2^1$, then $|V_1'(s_1)| \ll 1$ and $|N_1'(s_1)| \gg 1$ hold for $0 < s_1 \ll 1$, which shows that $V_1(s_1) = N_1(s_1)$ has no sufficiently small positive solution.

Next, we only consider the case $\rho_2^2 \leq \lambda_2^1$ and $w_1^{12} w_2^{12} > 0$. As $\mu \in L_1^2$, we have $M_{1\mu}^1 \mu > 0$, and then, for $w_i^{12} > 0$, $i = 1, 2$ we see that

$$V_1'(s_1) \leq (w_1^{12})^{-\rho_2^2/\lambda_2^1} s_1^{\rho_2^2/\lambda_2^1 - 1} < N_1'(s_1) \quad \text{for } 0 < s_1 \ll 1. \quad (41)$$

In fact, $\rho_2^2 < \lambda_2^1$ yields that $\lim_{s_1 \rightarrow 0^+} s_1^{\rho_2^2/\lambda_2^1 - 1} = +\infty$, $\lim_{s_1 \rightarrow 0^+} N_1'(s_1) = +\infty$, and $\lim_{s_1 \rightarrow 0^+} s_1^{\rho_2^2/\lambda_2^1 - 1} / N_1'(s_1) = 0$, which shows $V_1(s_1) = N_1(s_1)$ has no sufficiently small positive solutions. Obviously, the conclusion is correct as $\rho_2^2 = \lambda_2^1$.

Similarly, for $\rho_2^2 < \lambda_2^1$, $w_i^{12} < 0$, $i = 1, 2$, there does not exist a small positive solution for $V_1(s_1) = N_1(s_1)$.

The proof is then completed. $\square$

Theorem 5. Assume that the conditions (H_1) – (H_3) hold and $M_{i\mu}^1 \neq 0$, $i = 1, 2$. Let $\lambda = 0$, $\mu \in L_1^2$, and $0 < |\mu| \ll 1$; then

- (i) the periodic orbit and the homoclinic loop connecting p_2 of system (1) cannot coexist as $\rho_2^2 \geq \lambda_2^1$ or $w_1^{12} w_2^{12} < 0$;
- (ii) at least one periodic orbit and the homoclinic loop connecting p_2 of system (1) coexist as $\rho_2^2 < \lambda_2^1$, $w_1^{12} > 0$, and $w_2^{12} > 0$;
- (iii) a unique periodic orbit and the homoclinic loop connecting p_2 of system (1) coexist as $\rho_2^2 < \lambda_2^1$, $w_1^{12} < 0$, and $w_2^{12} < 0$.

Proof. By Theorem 2, the condition $\mu \in L_1^2$ for $0 < |\mu| \ll 1$ implies that system (1) has a homoclinic loop connecting p_2 .

(i) Let $s_2 = e^{-\rho_2^1 \tau_2}$ and eliminating s_1 in (33), we derive

$$\begin{aligned} V_2(s_2) &\triangleq s_2 + \delta^{-1} w_2^{12} M_{2\mu}^1 \mu + \text{h.o.t.} \\ &= \frac{w_2^{12}}{1 - (\rho_1^1)^{-1} \delta \ln \left(w_1^{12} \left(s_2^{\lambda_2^1/\rho_2^1} - \delta^{-1} M_{1\mu}^1 \mu \right) \right)} \\ &\triangleq N_2(s_2). \end{aligned} \quad (42)$$

Note that $V_2(0) = N_2(0)$ as $\mu \in L_1^2$. Moreover,

$$\begin{aligned} V_2'(s_2) &= 1, \\ N_2'(s_2) &= \frac{w_2^{12} (\rho_1^1)^{-1} \delta}{\left[1 - (\rho_1^1)^{-1} \delta \ln \left(w_1^{12} \left(s_2^{\lambda_2^1/\rho_2^1} - \delta^{-1} M_{1\mu}^1 \mu \right) \right) \right]^2} \\ &\quad \cdot \frac{(\lambda_2^1/\rho_2^1) s_2^{(\lambda_2^1 - \rho_2^1)/\rho_2^1}}{(s_2^{\lambda_2^1/\rho_2^1} - \delta^{-1} M_{1\mu}^1 \mu)}. \end{aligned} \quad (43)$$

For $\rho_2^2 \geq \lambda_2^1$ and $N_2'(s_2) \ll 1 = V_2'(s_2)$, this means $V_2(s_2) = N_2(s_2)$ has no sufficiently small positive solutions.

Now we turn to the case $w_1^{12} w_2^{12} < 0$, since we are interested in sufficiently small positive solutions of (33), it suffices to consider the sufficiently small positive solutions of $V_2(s_2) = N_2(s_2)$ satisfying $w_1^{12} (s_2^{\lambda_2^1/\rho_2^1} - \delta^{-1} M_{1\mu}^1 \mu) > 0$, which implies that $s_2^{\lambda_2^1/\rho_2^1} - \delta^{-1} M_{1\mu}^1 \mu < 0$ (resp., $s_2^{\lambda_2^1/\rho_2^1} - \delta^{-1} M_{1\mu}^1 \mu > 0$) for $w_1^{12} < 0$ (resp., $w_1^{12} > 0$). It is easy to see that $V_2(s_2) = N_2(s_2)$ has no sufficiently small positive solutions as $w_1^{12} w_2^{12} < 0$.

(ii) For $\rho_2^2 < \lambda_2^1$, we have $V_2'(0) = 1 > 0 = N_2'(0)$, which implies that there exists an $0 < \tilde{s}_2 \ll 1$ such that $V_2(s_2) > N_2(s_2)$ for $0 < s_2 < \tilde{s}_2$.

Choosing $\tilde{s}_2 = \delta^{-1} w_2^{12} M_{2\mu}^1 \mu > 0$, then

$$\begin{aligned} V_2(\tilde{s}_2) &= 2\delta^{-1} w_2^{12} M_{2\mu}^1 \mu + \text{h.o.t.}, \\ N_2(\tilde{s}_2) &= \frac{w_2^{12}}{1 - (\rho_1^1)^{-1} \delta \ln \left(w_1^{12} \left(\tilde{s}_2^{\lambda_2^1/\rho_2^2} - \delta^{-1} M_{1\mu}^1 \mu \right) \right)}. \end{aligned} \quad (44)$$

In view of $\ln(w_1^{12}(\tilde{s}_2^{\lambda_2^1/\rho_2^2} - \delta^{-1} M_{1\mu}^1 \mu)) > \ln(w_1^{12} \tilde{s}_2^{\lambda_2^1/\rho_2^2}) = \ln(w_1^{12}(\delta^{-1} w_2^{12} M_{2\mu}^1 \mu)^{\lambda_2^1/\rho_2^2})$ for $w_1^{12} > 0$, so

$$\begin{aligned} N_2(\tilde{s}_2) &> \frac{w_2^{12}}{1 - (\rho_1^1)^{-1} \delta \ln \left(w_1^{12} \left(\delta^{-1} w_2^{12} M_{2\mu}^1 \mu \right)^{\lambda_2^1/\rho_2^2} \right)} \\ &\gg 2w_1^{12} (\delta^{-1} w_2^{12} M_{2\mu}^1 \mu) = V_2(\tilde{s}_2) \end{aligned} \quad (45)$$

when $w_2^{12} > 0$. As a result, $N_2(s_2) = V_2(s_2)$ has at least one solution $\tilde{s}_2$ satisfying $0 < \tilde{s}_2 < \tilde{s}_2 < \tilde{s}_2 \ll 1$.

(iii) s_2 must fulfill $0 < s_2 < (\delta^{-1} M_{1\mu}^1 \mu)^{\rho_2^2/\lambda_2^1}$ as $w_1^{12} < 0$; with similar arguments in proof of (ii), we can prove that there exists a $0 < s_2^* \ll 1$ such that $V_2(s_2^*) = N_2(s_2^*)$ for $0 < s_2^* < (\delta^{-1} M_{1\mu}^1 \mu)^{\rho_2^2/\lambda_2^1} \ll 1$. It is easy to compute that $N_2''(s_2) > 0$ for $w_2^{12} < 0$, $0 < s_2 < (\delta^{-1} M_{1\mu}^1 \mu)^{\rho_2^2/\lambda_2^1}$, and $\mu \in L_2^1$. Combining with the fact $V_2(0) = N_2(0)$, $N_2'(s_2) > 0$, and $V_2'(s_2) = 1$, we immediately know that s_2^* is unique.

This completes the proof. $\square$

Now, we turn to discussing the bifurcations of the heteroclinic loop for $\lambda > 0$, when p_1 undergoes a transcritical bifurcation. From Figure 1, we know that when $\lambda > 0$, after the creation of the equilibria p_1^0 and p_1^1 , there always exists a straight segment orbit heteroclinic to p_1^1 and p_1^0 , its length is λ_p , and we denote this heteroclinic orbit by Γ^* . Moreover, $x_0^1 = \lambda_p$ is a critical position.

Firstly, we take into account the case $x_0^1 \geq \lambda_p$. In this case, (31) becomes

$$\begin{aligned} (w_1^{12})^{-1} \delta s_1 - \delta s_2^{\lambda_2^1/\rho_2^2} + M_{1\lambda}^1 \lambda + M_{1\mu}^1 \mu + \text{h.o.t.} &= 0, \\ (w_2^{12})^{-1} \delta s_2^{\rho_2^2/\rho_2^1} - \delta \lambda_p \left[\delta - (\delta - \lambda_p) s_1^{\lambda_p/\rho_1^1} \right]^{-1} \\ &+ M_{2\lambda}^1 \lambda + M_{2\mu}^1 \mu \\ &+ w_2^{31} (w_2^{33})^{-1} s_1^{\lambda_1^1/\rho_1^1} \left[w_1^{13} (w_1^{12})^{-1} \delta s_1 + (w_1^{33})^{-1} \delta s_2^{\lambda_2^2/\lambda_2^1} \right. \\ &\quad \left. - M_{1\lambda}^3 \lambda - M_{1\mu}^3 \mu \right] + \text{h.o.t.} = 0. \end{aligned} \quad (46)$$

Let $s = s_1^{\lambda_p/\rho_1^1}$ ($s = 0$ means $s_1 = 0$ and vice versa); by virtue of Taylor's development for $\delta \lambda_p / (\delta - (\delta - \lambda_p) s_1^{\lambda_p/\rho_1^1})$, we have

$$\begin{aligned} (w_1^{12})^{-1} s^{\rho_1^1/\lambda_p} - s_2^{\lambda_2^1/\rho_2^2} + \delta^{-1} M_{1\lambda}^1 \lambda + \delta^{-1} M_{1\mu}^1 \mu + \text{h.o.t.} &= 0, \\ (w_2^{12})^{-1} \delta s_2^{\rho_2^2/\rho_2^1} - \lambda_p - \frac{\lambda_p (\delta - \lambda_p)}{\delta} s + M_{2\lambda}^1 \lambda \\ &+ M_{2\mu}^1 \mu + \text{h.o.t.} = 0. \end{aligned} \quad (47)$$

With similar arguments to $\lambda = 0$, we may easily obtain the following results.

Theorem 6. Suppose the conditions (H_1) – (H_3) hold, $0 < \lambda \ll 1$; then

(i) if $\text{rank}(M_{1\mu}^1, M_{2\mu}^1) = 2$, there exists an $(l - 2)$ -dimensional surface

$$\begin{aligned} L_{12}^\lambda &= \{ \mu(\lambda) : M_{1\mu}^1 \mu + M_{1\lambda}^1 \lambda + \text{h.o.t.} \\ &= M_{2\mu}^1 \mu + M_{2\lambda}^1 \lambda - \lambda + \text{h.o.t.} = 0 \} \end{aligned} \quad (48)$$

such that system (1) has a unique heteroclinic loop if and only if $\mu \in L_{12}^\lambda$ and $0 < |\mu| \ll 1$;

(ii) there exists an $(l - 1)$ -dimensional surface

$$\begin{aligned} L_{1\lambda}^2 &= \left\{ \mu(\lambda) : W_1^2(\lambda, \mu) = (w_2^{12})^{-1} \left[\delta^{-1} (M_{1\mu}^1 \mu + M_{1\lambda}^1 \lambda) \right]^{\beta_2} \right. \\ &\quad \left. + \delta^{-1} (M_{2\mu}^1 \mu + M_{2\lambda}^1 \lambda) \right. \\ &\quad \left. - \delta^{-1} \lambda_p + \text{h.o.t.} = 0, \right. \\ &\quad \left. M_{1\mu}^1 \mu + M_{1\lambda}^1 \lambda > 0 \right\} \end{aligned}$$

(resp.,

$$\begin{aligned} L_{2\lambda}^1 &= \left\{ \mu(\lambda) : W_2^1(\lambda, \mu) \right. \\ &= \delta \lambda_p + \lambda_p (\delta - \lambda_p) \\ &\quad \times \left[-\delta^{-1} w_1^{12} (M_{1\lambda}^1 \lambda + M_{1\mu}^1 \mu) \right]^{\lambda_p/\rho_1^1} \\ &\quad - \delta M_{2\lambda}^1 \lambda \\ &\quad - \delta M_{2\mu}^1 \mu + \text{h.o.t.} = 0, \\ &\quad \left. w_1^{12} (M_{1\lambda}^1 \lambda + M_{1\mu}^1 \mu) < 0 \right\} \end{aligned} \quad (49)$$

such that system (1) has one homoclinic loop connecting p_1^1 (resp., connecting p_2) if and only if $\mu \in L_{1\lambda}^2$ and $0 < |\mu| \ll 1$.

Theorem 7. Suppose hypotheses (H_1) – (H_3) hold, $M_{i\mu}^1 \neq 0$, $i = 1, 2$, $0 < \lambda$, $|\mu| \ll 1$, and $w_1^{12} w_2^{12} < 0$. Then, except the

homoclinic loop connecting p_1^1 (resp., p_2), system (1) has no periodic orbits as $\mu \in L_{1\lambda}^2$ (resp., $\mu \in L_{2\lambda}^1$).

Remark 8. It is easy to see that homoclinic loop connecting p_1^0 and heteroclinic loop joining p_1^0, p_2 cannot be bifurcated from Γ , which is exactly determined by the generic condition (H_1) .

Finally, we consider the case $-\beta \leq x_0^1 < \lambda_p$. Due to Figure 1 and (25), it follows from (31) that

$$\begin{aligned} s_2 &= [\delta^{-1} (M_{1\lambda}^1 \lambda + M_{1\mu}^1 \mu)]^{\rho_2^1/\lambda_2^1} + \text{h.o.t.}, \\ x_0^1 &= (w_2^{12})^{-1} \delta s_2^{\rho_2^2/\rho_2^1} + M_{2\lambda}^1 \lambda + M_{2\mu}^1 \mu + \text{h.o.t.} \end{aligned} \quad (50)$$

Theorem 9. Assume the conditions (H_1) – (H_3) are true, $\text{rank}(M_{1\lambda}^1, M_{1\mu}^1) > 0$ and $\text{rank}(M_{2\lambda}^1, M_{2\mu}^1) > 0$. Then,

(i) there exists a surface

$$\begin{aligned} \Sigma_1(\mu, \lambda) &= \left\{ \mu(\lambda) : [\delta^{-1} (M_{1\lambda}^1 \lambda + M_{1\mu}^1 \mu)]^{\rho_2^1/\lambda_2^1} + \text{h.o.t.} = 0, \right. \\ &\quad -\beta \leq M_{2\lambda}^1 \lambda + M_{2\mu}^1 \mu + \text{h.o.t.} < \lambda_p, \\ &\quad \left. 0 < |\mu|, \lambda \ll 1 \right\}, \end{aligned} \quad (51)$$

such that system (1) has two orbits heteroclinic to p_1^1, p_2, p_1^0 as $\mu \in \Sigma_1(\mu, \lambda)$;

(ii) there exists a region in the (λ, μ) space

$$\begin{aligned} \Delta &= \left\{ (\lambda, \mu) : -\beta \leq (w_2^{12})^{-1} \delta^{(\lambda_2^1 - \rho_2^2)/\lambda_2^1} \right. \\ &\quad \times (M_{1\lambda}^1 \lambda + M_{1\mu}^1 \mu)^{\rho_2^2/\lambda_2^1} \\ &\quad + M_{2\lambda}^1 \lambda + M_{2\mu}^1 \mu \\ &\quad \left. + \text{h.o.t.} < \lambda_p, \right. \\ &\quad \left. M_{1\lambda}^1 \lambda + M_{1\mu}^1 \mu > 0, \right. \\ &\quad \left. 0 < |\mu|, \lambda \ll 1 \right\}, \end{aligned} \quad (52)$$

such that system (1) has a heteroclinic orbit connecting p_1^1 and p_1^0 for $(\lambda, \mu) \in \Delta$.

Proof. (i) If $s_2 = 0$ in (50), then

$$\begin{aligned} 0 &= [\delta^{-1} (M_{1\lambda}^1 \lambda + M_{1\mu}^1 \mu)]^{\rho_2^1/\lambda_2^1} + \text{h.o.t.}, \\ x_0^1 &= M_{2\lambda}^1 \lambda + M_{2\mu}^1 \mu + \text{h.o.t.} \end{aligned} \quad (53)$$

which shows that there exists a surface $\Sigma_1(\mu, \lambda)$ such that (50) has a solution $s_2 = 0$ and $-\beta \leq x_0^1 < \lambda_p$ for $\mu \in \Sigma_1(\mu, \lambda)$, then system (1) has two heteroclinic orbits, one is heteroclinic to p_1^1 and p_2 and the other is heteroclinic to p_2 and p_1^0 .

FIGURE 2

(ii) If $s_2 > 0$ in (50), one attains $M_{1\lambda}^1 \lambda + M_{1\mu}^1 \mu > 0$. Eliminating s_2 in (50), we achieve

$$\begin{aligned} x_0^1 &= (w_2^{12})^{-1} \delta^{(\lambda_2^1 - \rho_2^2)/\lambda_2^1} (M_{1\lambda}^1 \lambda + M_{1\mu}^1 \mu)^{\rho_2^2/\lambda_2^1} \\ &\quad + M_{2\lambda}^1 \lambda + M_{2\mu}^1 \mu + \text{h.o.t.}, \end{aligned} \quad (54)$$

which shows that there exists a region Δ such that when $(\lambda, \mu) \in \Delta$, system (1) has one heteroclinic orbit heteroclinic to p_1^1 and p_1^0 . $\square$

Remark 10. All the heteroclinic orbits joining p_1^0 will go into p_1^0 in different ways according to different fields of x_0^1 ; see Figure 2.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

This paper is supported by National Natural Science Foundation of China (nos. 11202192, 11226133), the Fundamental Research Funds for the Central Universities (no. 2652012097), and the Beijing Higher Education Young Elite Teacher Project.

References

- [1] A. J. Homburg and B. Krauskopf, "Resonant homoclinic flip bifurcations," *Journal of Dynamics and Differential Equations*, vol. 12, no. 4, pp. 807–850, 2000.
- [2] C. A. Morales and M. J. Pacifico, "Inclination-flip homoclinic orbits arising from orbit-flip," *Nonlinearity*, vol. 14, no. 2, pp. 379–393, 2001.
- [3] S. Shui, J. Li, and X. Zhang, "Nonresonant bifurcations of heteroclinic loops with one inclination flip," *International Journal of Bifurcation and Chaos*, vol. 21, no. 1, pp. 255–273, 2011.

- [4] Y. Xu and D. Zhu, "Bifurcations of heterodimensional cycles with one orbit flip and one inclination flip," *Nonlinear Dynamics*, vol. 60, no. 1-2, pp. 1-13, 2010.
- [5] T. Zhang and D. Zhu, "Codimension 3 homoclinic bifurcation of orbit flip with resonant eigenvalues corresponding to the tangent directions," *International Journal of Bifurcation and Chaos in Applied Sciences and Engineering*, vol. 14, no. 12, pp. 4161-4175, 2004.
- [6] B. Deng, "Homoclinic bifurcations with nonhyperbolic equilibria," *SIAM Journal on Mathematical Analysis*, vol. 21, pp. 693-720, 1990.
- [7] J. Klaus and J. Knobloch, "Bifurcation of homoclinic orbits to a saddle-center in reversible systems," *International Journal of Bifurcation and Chaos in Applied Sciences and Engineering*, vol. 13, no. 9, pp. 2603-2622, 2003.
- [8] J. D. M. Rademacher, "Homoclinic orbits near heteroclinic cycles with one equilibrium and one periodic orbit," *Journal of Differential Equations*, vol. 218, no. 2, pp. 390-443, 2005.
- [9] S. Wiggins, *Introduction to Applied Nonlinear Dynamical System and Chaos*, Springer, New York, NY, USA, 1990.