Research Article

Hydrogen Peroxide Is Involved in Salicylic Acid-Elicited Rosmarinic Acid Production in *Salvia miltiorrhiza* Cell Cultures

Wenfang Hao, Hongbo Guo, Jingyi Zhang, Gege Hu, Yaqin Yao, and Juane Dong

State Key Laboratory of Crop Stress Biology for Arid Areas, College of Life Sciences, Northwest A&F University, Yangling 712100, China

Correspondence should be addressed to Juane Dong; dzsys@nwsuaf.edu.cn

Received 4 April 2014; Revised 7 May 2014; Accepted 7 May 2014; Published 29 May 2014

Academic Editor: Hongbo Shao

Copyright © 2014 Wenfang Hao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Salicylic acid (SA) is an elicitor to induce the biosynthesis of secondary metabolites in plant cells. Hydrogen peroxide (H_2O_2) plays an important role as a key signaling molecule in response to various stimuli and is involved in the accumulation of secondary metabolites. However, the relationship between them is unclear and their synergetic functions on accumulation of secondary metabolites are unknown. In this paper, the roles of SA and H_2O_2 in rosmarinic acid (RA) production in *Salvia miltiorrhiza* cell cultures were investigated. The results showed that SA significantly enhanced H_2O_2 production, phenylalanine ammonia-lyase (PAL) activity, and RA accumulation. Exogenous H_2O_2 could also promote PAL activity and enhance RA production. If H_2O_2 production was inhibited by NADPH oxidase inhibitor (IMD) or scavenged by quencher (DMTU), RA accumulation would be blocked. These results indicated that H_2O_2 is secondary messenger for signal transduction, which can be induced by SA, significantly and promotes RA accumulation.

1. Introduction

Salicylic acid (SA) is often used to regulate plant growth and development, seed germination, and fruit formation and to enhance the capability of the plants to respond to abiotic and biotic stresses [1, 2]. Exogenous application of SA can promote the thermo tolerance of mustard seedlings [3], chilling tolerance of cucumber [4], salt stress of *Arabidopsis* seedlings [5], and toxicity tolerance of cadmium of barley seedlings [6]. In recent years, SA has been used as an elicitor to induce the biosynthesis of secondary metabolites in plants. Exogenous application of SA induces the biosynthesis of coumarins in *Matricaria chamomilla* [7], taxane in *Taxus chinensis* [8], saponins in ginseng [9], phenolic acids in *Salvia miltiorrhiza* [10], artemisinin in *Artemisia annua* [11], and sinapyl alcohol in *ulmus* cells [12].

The biosynthesis of secondary metabolites in plant cells is regulated by specific signal transduction pathways. Under the stimulation of elicitors, some biochemical events related to signal transduction can be activated, such as ion transmembrane transport, active oxide species (AOS) burst,

and protein phosphorylation and dephosphorylation [13, 14]. AOS are signal molecules that widely exist in plant cells [13–15]. Hydrogen peroxide (H_2O_2), one of the AOS, has been considered as the most significant signal molecules [14, 16]. Generally, when plants are under stress, H_2O_2 is produced rapidly to activate systemic acquired resistance [13, 17, 18] and acts as a signal molecule mediating the biosynthesis of secondary metabolites in plants. For example, H_2O_2 is the key signal molecule for oligosaccharides to induce the biosynthesis of taxol in *T. chinensis* cells [19], for the cell wall of *Aspergillus niger* to elicit *Catharanthus roseus* cells to synthesize catharanthine [20], for ABA to induce rice seedling leaves to synthesize anthocyanidins [21], and for the cell cultures of *Rubia tinctorum* to produce anthraquinone [22], and so forth.

Rosmarinic acid (RA) is one of effective compounds of Danshen, the root and rhizome of *Salvia miltiorrhiza* Bunge, which is used as a traditional Chinese herbal drug which removes blood stasis, stops pain, and activates blood flow and is heart-relieving [23]. RA shows a great intensity for free radical scavenging and antioxidant activity. The

phenylalanine ammonia-lyase (PAL) is the first enzyme for accumulating phenolic acid compounds in *S. miltiorrhiza* cell cultures [10], and SA could induce the production of rosmarinic acid [24]. The aim of this work is to reveal the effects of SA and H_2O_2 on the accumulation of RA in *S. miltiorrhiza* cell cultures. For this purpose, imidazole (IMD, NADPH oxidase inhibitor) is used to inhibit the enzyme activity of NADPH oxidase, and dimethylthiourea (DMTU, H_2O_2 scavenger) is employed as scavenger of H_2O_2 .

2. Results

2.1. Effects of SA on PAL Activity, H_2O_2 Production, and RA Accumulation. The PAL activity increased from 20 min after treating with 22.5 mg/L SA and reached the peak at 16 h, when the activity was 5.46 folds of that of control, followed by subsequent decrease (Figure 1(a)). The H_2O_2 production induced by SA also exhibited continuous increase at the beginning with two peaks, but decreased rapidly after 2-h elicitation. The first peak occurred at 20 min and the second did at 2 h, when 8.51 folds H_2O_2 production was found in SA-treated cells (Figure 1(b)). In terms of RA accumulation, continuous increase was found till the 2nd day when RA content reached the highest with 2.15 folds and then it decreased gradually (Figure 1(c)).

2.2. Effects of Exogenous H_2O_2 on the PAL Activation and RA Production. The effects of H_2O_2 on activation of PAL activity and RA production were investigated, after applying exogenous H_2O_2 with concentration ranging from 1 to 70 mM. The activity of PAL and RA content were enhanced with the increase of H_2O_2 concentration ranging from 1 to 10 mM, whereas both decreased if the concentration was more than 10 mM (Figure 2). This result suggests that 10 mM is an appropriate concentration to enhance PAL activity and RA accumulation.

To further estimate the suitable treatment time of H_2O_2 , continuous 6-day treatment was performed to confirm the best time point. Both PAL activity and RA accumulation were increased at early stage with time extension and then decreased at the late stage (Figure 3). The activity of PAL reached the highest peak at 8 h, followed by drastic decrease. The RA content reached the peak at 1 d and decreased drastically after 2 d.

2.3. Influences of CAT on SA-Induced RA Production. CAT is a scavenger of H_2O_2 , which can not pass through cell membrane, and thus the exogenous CAT cannot scavenge H_2O_2 within cells. Both contents of H_2O_2 (Figure 4(a)) and RA (Figure 4(b)) were slightly lower than those of control after adding CAT, whereas both contents showed no significant difference between CAT + SA-treated and SA-elicited cells. The application of CAT did not affect the production of endogenous H_2O_2 and RA accumulation in cells that were subjected to SA elicitation. However, CAT can scavenge the exogenous H_2O_2 , thereby eliminating the production of H_2O_2 and RA synthesis due to H_2O_2 regulation. This

indicates that the RA biosynthesis is mediated by intracellular H_2O_2 elicited by SA.

2.4. Influences of DMTU on SA-Induced RA Production. DMTU is a H_2O_2 quencher that can effectively remove H_2O_2 within cells. The results showed that H_2O_2 was released and RA accumulated due to SA elicitation (Figure 1(b)). In order to further explain whether endogenous H_2O_2 was involved in the activation of PAL and the synthesis of RA, DMTU was employed to quench endogenous H_2O_2 , and under such condition, both contents of H_2O_2 and RA in cells were measured (Figure 5).

The production of endogenous H_2O_2 (Figure 5(a)) and the accumulation of RA (Figure 5(b)) within cells were partially inhibited by 100 μ M DMTU, and they were significantly inhibited with the concentration of 500 μ M. If the cells were treated by 500 μ M DMTU + 22.5 mg/L SA or 500 μ M DMTU + 10 mM H_2O_2 , DMTU would significantly inhibit the production of endogenous H_2O_2 and the accumulation of RA. Application of exogenous H_2O_2 and/or SA could partially reverse the inhibition of DMTU on RA synthesis. The treatment of 700 μ M DMTU also completely inhibited the production of H_2O_2 in cells, and the inhibition could not be reversed by application of exogenous SA or H_2O_2 . These results showed DMTU treatment slightly decreased both contents of RA and H_2O_2 , when compared with the H_2O control. Meanwhile, DMTU + SA treatment would significantly block both contents if compared with those SA-elicited cells. These results indicated that H_2O_2 produced from SA elicitation is necessary for triggering the biosynthesis of RA.

2.5. Influences of IMD on SA-Induced RA Production. NADPH oxidase is one of the key enzymes for the generation of intracellular H_2O_2 [25, 26]. IMD is the specific inhibitor of NADPH oxidase on plasma membrane, which can inhibit the production of H_2O_2 through inactivating NADPH oxidase [16, 17, 25]. For instance, IMD scavenged H_2O_2 produced from NADPH oxidase elicited by ABA and decreased the synthesis of anthocyanins in rice leaves [21]. In this study, IMD was used to inhibit the activity of NADPH oxidase to examine its influence on the biosynthesis of RA elicited by exogenous H_2O_2 and/or SA.

When IMD (100 μ M) was used solely, IMD treatment slightly decreased the contents of RA and H_2O_2 when compared with the control, while IMD + SA and IMD + H_2O_2 treatments significantly decreased both contents when compared with SA-elicited cells (Figure 6). The results indicated that IMD inhibited the effects of SA elicitation, and the inhibition of IMD to the synthesis of RA in cell cultures depended on H_2O_2 generated from NADPH oxidase.

3. Discussion

This investigation shows SA can induce the burst of H_2O_2 , and the elicited H_2O_2 enhances the PAL activity and RA accumulation in *S. miltiorrhiza* cell cultures. SA signaling

FIGURE 1: Exogenous SA promotes PAL activity (a), H_2O_2 production (b) and RA synthesis (c) in the suspension cultured cells of *S. miltiorrhiza*. 6-day-old suspension cultured cells treated with 22.5 mg/L SA were harvested at the time indicated. The PAL activity and contents of H_2O_2 and RA were determined. Values are means of three independent experiments. Bars represented standard errors.

FIGURE 2: Effects of exogenous hydrogen peroxide with different concentrations on PAL activity (a) and RA accumulation (b) in the *S. miltiorrhiza* suspension cultured cells. The salicylic acid concentration is 22.5 mg/L. Values are means of three independent experiments. Bars represented standard errors. Different lowercase letters represent the significant differences between PAL activity (a) and RA content (b) after each treatment at the level of 5%.

pathways in plant defense network have been widely investigated during the past two decades [27, 28], but their role in secondary metabolism is rather limited. Zitta et al. [29] reports that 10 μM SA leads to a significant 4-fold increase in H_2O_2 concentrations, which may be attributed to the fact that SA can directly reduce the activity of catalase that decompose of H_2O_2 (Figure 4). As in our study, the content of H_2O_2

in SA-elicited cells was 8.51-fold higher than that of water control (Figure 1), which is consistent with these reports. Although low level ($<10 \mu\text{M}$) of H_2O_2 is thought to result in the induction of various intracellular signaling pathways [28, 29], application of either exogenous H_2O_2 (10 mM) or endogenous H_2O_2 elicited by SA (22 mg/L) showed its capacity to enhance both PAL activity and RA accumulation

FIGURE 3: Exogenous hydrogen peroxide promotes PAL activity and RA production in the *S. miltiorrhiza* suspension cell cultures. 6-d-old cells treated with 10 mM H_2O_2 were harvested at the time as indicated, and RA contents and PAL activities were then determined. Values are means of three independent experiments. Bars represented standard errors.

in our experiments. PAL is the first key enzyme in the phenylpropanoid pathway for the biosynthesis of RA [30], which has been reported to be enhanced by SA in the colonization of arbuscular mycorrhizal fungus [28] and in the interaction between tomato plants and *Fusarium oxysporum* f. sp. *lycopersici* [31]. H_2O_2 comes to light as a second messenger involved in SA-elicited pathway. In this paper, both PAL activity and RA accumulation were significantly increased when exogenous 10 mM H_2O_2 was applied (Figure 2). On the other hand, H_2O_2 could be elicited by SA even though it was treated by catalase or DMTU (Figures 4 and 5), and the elicited H_2O_2 also could significantly promote RA accumulation. Our results are consistent with those reports in the fact that H_2O_2 is involved in many elicitor-induced processes to produce secondary metabolites, for instance, the biosynthesis of taxol [19], catharanthine [20], anthocyanidins [21], and anthraquinone [22].

Increasing evidence proved that plant responses to elicitors, including enzymes activation and the production of secondary metabolites, are not only regulated by signaling pathway, but also a cross-talk process. The mechanism about cross-talk among SA, H_2O_2 , enzyme activity, and synthesis of secondary metabolites has gained some attention recently, but related studies are rare [32]. According to the latest progress in our lab, both nitric oxide (NO) and increase of intracellular Ca^{2+} are also involved in this cross-talk during SA induction.

Taken together, H_2O_2 production can be elicited by SA and the elicited H_2O_2 can promote PAL activity and RA accumulation, in which H_2O_2 plays an important role as a second messenger in signal transduction when SA is applied.

4. Experimental Section

4.1. Cell Culture. The detailed protocols of *S. miltiorrhiza* cell culture were provided by Dong et al. [10]. The seeds were collected from Shangluo, Shaanxi Province, and then soaked in water for 2–4 h. The wax coat was removed by gauze,

followed by washing with water. The seeds were soaked in 70% ethanol for 30 s. After washing by sterile water for 3 times, the seeds were sterilized by 0.1% $HgCl_2$ for 10–15 min. They were sown in the autoclaved MS solid media after washing with sterile water for three times, supplemented with 5.5 g/L agar (Beijing Kangbeisi Sci & Tech Company) and 30 g/L sucrose (Guangzhou Jinhua Chemical Company) to induce the germ free seedlings. The leaves of germ free seedlings were cut into 0.5 cm × 0.5 cm pieces and inoculated on the autoclaved MS solid medium supplemented with 1.0 mg/L NAA (Tianjin Bodi Chemical Company), 1.0 mg/L 6-BA (Beijing Kangbeisi Sci & Tech Company), 1.0 mg/L 2,4-D (Beijing Kangbeisi Sci & Tech Company), 5.5 g/L agar, and 30 g/L sucrose. Calli were induced and cultured at $25 \pm 2^\circ C$ with light intensity of 2000–3000 Lx for 12–16 h per day. After 15-day culture, the calli would be subcultured till two months when their morphological characteristics and growth rates were stable. The stable calli were collected and inoculated in MS liquid media (containing 30 g/L sucrose) with the ratio of callus to culture medium 1:15 (W/V). The suspension cells were cultured in the dark at $25^\circ C$ with shaking speed of 125 rpm.

4.2. Elicitation and Chemical Treatments. Stock solutions of SA, H_2O_2 , DMTU (H_2O_2 scavenger, Sigma, USA), CAT (Sigma, USA), and IMD (NADPH scavenger, Sigma, USA) were prepared in distilled water and then sterilized after filtration through 0.22 μm membrane. The resultant solutions with different concentrations were added in the media in accordance with the experimental design. In the control group, only distilled water with the same volume was added to substitute for the chemicals. DMTU, CAT, and IMD were added separately to the cell culture 30 min before SA treatment. All experiments were performed with three replications.

4.3. PAL Activity Assay. PAL activity was measured according to the method of Solecka and Kacperska [33] with some modifications. Fresh cultured calli were placed on filter paper in a Bucher funnel, filtered in vacuum, washed by distilled water, and filtered in vacuum again to remove the facial water. Two grams of calli were homogenized with 5 mL extraction buffer in a mortar. The homogenate was filtered through four-layer cheesecloth and centrifuged at 10000 rpm for 15 min at $4^\circ C$ to get supernatant that would be used as a crude enzyme. The reaction mixture (3 mL) included 0.5 mL crude enzyme, 16 mM L-phenylalanine, 50 mM Tris-HCl buffer (pH 8.9), and 3.6 mM NaCl. The mixture was incubated at $37^\circ C$ for 60 min and the reaction was stopped by 500 μL 6 M HCl. The resultant mixture was then centrifuged (12,000 ×g, 10 min). The absorbance was measured at 290 nm before and after incubation. The enzyme activity (U) was calculated by the following equation:

$$PAL(U \cdot g^{-1}FW \cdot h^{-1}) = \frac{A_{290} \times V_T \times v}{0.01 \times V_s \times FW \times t}, \quad (1)$$

where V_t represents the total volume of enzyme solution (mL); FW is the fresh weight of calli; V_s is the volume of

FIGURE 4: Influences of CAT (a scavenger of H₂O₂) on the generation of H₂O₂ (a) and the biosynthesis of RA (b) in the suspension cultured cells treated by SA and/or exogenous H₂O₂. 6-day-old suspension cultured cells treated with CAT (100 U) 30 min before SA treatment were harvested at 8 h (for H₂O₂ analysis) and 2 d (for RA analysis) later, and the H₂O₂ and rosmarinic acid contents were then determined. The SA concentration is 22.5 mg/L. The exogenous H₂O₂ concentration is 10 mM. Values are means of three independent experiments. Bars represented standard errors. Different lowercase letters represent the significant differences between the concentrations of H₂O₂ (a) and RA (b) after each treatment at the level of 5%.

FIGURE 5: Inhibition of DMTU on exogenous SA in the generation of intracellular H₂O₂ (a) and the biosynthesis of RA (b) in suspension cultured cells. 8-day-old suspension cultured cells treated with DMTU 30 min before exogenous H₂O₂ and/or SA treatment as indicated were harvested at 8 h (for H₂O₂ analysis) and 2 d (for RA analysis) later, and the H₂O₂ and Sal B contents were then determined. The Salicylic acid concentration is 22.5 mg/L. The exogenous H₂O₂ concentration is 10 mM. Values are means of three independent experiments. Bars represented standard errors. Different lowercase letters represent the significant differences between the contents of H₂O₂ (a) and Sal B (b) after each treatment at the level of 5%.

the enzyme solution sampled (mL); v is the total volume of the reaction solution (mL); and t is the reaction time (h).

4.4. Determination of Hydrogen Peroxide. Hydrogen peroxide was determined by the method of Li et al. [34]. The calli (0.2-0.3 g) were homogenized with 5 mL acetone precooled under 4°C and then ground to homogenate on the ice. The mixture was centrifuged at 12,000 rpm for 10 min at 4°C. The supernatant (2 mL) was mixed quickly with 0.5 mL 5% titanium sulfate, and ammonia solution (2 mL) was added and homogenized. The resultant mixtures were then centrifuged again; the supernatant was discarded. The precipitate (the titanium-hydro peroxide complex) was dissolved in 5 mL 2 M sulfuric acid, and the supernatant absorbance was

measured at 415 nm, and the content of hydrogen peroxide was calculated.

4.5. RA Extraction and HPLC Analysis. The *S. miltiorrhiza* cells were collected from cell cultures by centrifugation at 1200 rpm, and then dried at 47.5°C in an oven till constant weight. The dried cells (0.05 g) were put into a test tube with a stopper, and the extraction was conducted with aqueous methanol (70 : 30, methanol : water) for 45 min in an ultrasonic bath. The extract was filtered through a 0.22 μm membrane and the filtrate was obtained for detection.

The content of RA was quantified by HPLC (Shimadzu, model SCL-10AVPTM equipped with UV/Vis absorbance detector, 150 mm × 4.6 mm shim-pack column,

FIGURE 6: Inhibition of IMD on exogenous H₂O₂ and/or SA in the generation of intracellular H₂O₂ and the biosynthesis of RA in suspension cultured cells. 8-day-old suspension cultured cells treated with IMD 30 min before exogenous H₂O₂ and/or SA treatment as indicated were harvested at 8 h (for H₂O₂ analysis) and 2 d (for RA analysis) later, and the H₂O₂ and Sal B contents were then determined. The Salicylic acid concentration is 22.5 mg/L. The exogenous H₂O₂ concentration is 10 mM. The IMD concentration is 100 μ M. Values are means of three independent experiments. Bars represented standard errors.

and LC-ATVP pump). The mobile phase was acetonitrile : water : phosphate (25 : 75 : 0.1, v/v/v). The flow rate was 1 mL/min and the injection amount was 10 μ L. The detection wavelength was 285 nm and the column temperature was 25°C.

5. Conclusions

SA is an effective elicitor inducing RA accumulation in *S. miltiorrhiza* cell cultures, which can act independently or synergistically with H₂O₂. The intracellular H₂O₂ acts as a signal molecule to participate in RA accumulation in response to SA elicitation.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Authors' Contribution

Wenfang Hao and Hongbo Guo contributed equally to this work.

Acknowledgments

The authors gratefully acknowledge the funding and support from the Natural Science Foundation (no. 31170274) and from

National Forestry Public Welfare Industry Research Project (no. 20120406).

References

- [1] B. Guo, Y. C. Liang, Y. G. Zhu, and F. J. Zhao, "Role of salicylic acid in alleviating oxidative damage in rice roots (*Oryza sativa*) subjected to cadmium stress," *Environmental Pollution*, vol. 147, no. 3, pp. 743–749, 2007.
- [2] Q. Hayat, S. Hayat, M. Irfan, and A. Ahmad, "Effect of exogenous salicylic acid under changing environment: a review," *Environmental and Experimental Botany*, vol. 68, no. 1, pp. 14–25, 2010.
- [3] J. F. Dat, C. H. Foyer, and I. M. Scott, "Changes in salicylic acid and antioxidants during induced thermotolerance in mustard seedlings," *Plant Physiology*, vol. 118, no. 4, pp. 1455–1461, 1998.
- [4] H. M. Kang and M. E. Saltveit, "Chilling tolerance of maize, cucumber and rice seedling leaves and roots are differentially affected by salicylic acid," *Physiologia Plantarum*, vol. 115, no. 4, pp. 571–576, 2002.
- [5] O. Borsani, V. Valpuesta, and M. A. Botella, "Evidence for a role of salicylic acid in the oxidative damage generated by NaCl and osmotic stress in *Arabidopsis* seedlings," *Plant Physiology*, vol. 126, no. 3, pp. 1024–1030, 2001.
- [6] A. Metwally, I. Finkemeier, M. Georgi, and K.-J. Dietz, "Salicylic acid alleviates the cadmium toxicity in barley seedlings," *Plant Physiology*, vol. 132, no. 1, pp. 272–281, 2003.
- [7] A. Pastřová, M. Repčák, and A. Eliašová, "Salicylic acid induces changes of coumarin metabolites in *Matricaria chamomilla* L.," *Plant Science*, vol. 167, no. 4, pp. 819–824, 2004.
- [8] Y. D. Wang, Y. J. Yuan, and J. C. Wu, "Induction studies of methyl jasmonate and salicylic acid on taxane production in suspension cultures of *Taxus chinensis* var. mairei," *Biochemical Engineering Journal*, vol. 19, no. 3, pp. 259–265, 2004.
- [9] Y. Zhang, Y. Wang, S. C. Jiang, M. Z. Zhang, F. H. Li, and M. P. Zhang, "Cloning of the differential expression fragment from ginseng culture induced by salicylic acid," *Genomics and Applied Biology*, vol. 28, pp. 245–250, 2009.
- [10] J. Dong, G. Wan, and Z. Liang, "Accumulation of salicylic acid-induced phenolic compounds and raised activities of secondary metabolic and antioxidative enzymes in *Salvia miltiorrhiza* cell culture," *Journal of Biotechnology*, vol. 148, no. 2-3, pp. 99–104, 2010.
- [11] H. Yin, A. Kjaer, X. C. Fretté et al., "Chitosan oligosaccharide and salicylic acid up-regulate gene expression differently in relation to the biosynthesis of artemisinin in *Artemisia annua* L.," *Process Biochemistry*, vol. 47, no. 11, pp. 1559–1562, 2012.
- [12] J. A. Martín, A. Solla, M. C. García-Vallejo, and L. Gil, "Chemical changes in *Ulmus minor* xylem tissue after salicylic acid or carvacrol treatments are associated with enhanced resistance to *Ophiostoma novo-ulmi*," *Phytochemistry*, vol. 83, pp. 104–109, 2012.
- [13] M. E. Alvarez, R. I. Pennell, P.-J. Meijer, A. Ishikawa, R. A. Dixon, and C. Lamb, "Reactive oxygen intermediates mediate a systemic signal network in the establishment of plant immunity," *Cell*, vol. 92, no. 6, pp. 773–784, 1998.
- [14] S. Neill, R. Desikan, and J. Hancock, "Hydrogen peroxide signalling," *Current Opinion in Plant Biology*, vol. 5, no. 5, pp. 388–395, 2002.
- [15] A. Levine, R. Tenhaken, R. Dixon, and C. Lamb, "H₂O₂ from the oxidative burst orchestrates the plant hypersensitive disease resistance response," *Cell*, vol. 79, no. 4, pp. 583–593, 1994.

- [16] C. S. Bestwick, I. R. Brown, M. H. R. Bennett, and J. W. Mansfield, "Localization of hydrogen peroxide accumulation during the hypersensitive reaction of lettuce cells to *Pseudomonas syringae* pv phaseolicola," *Plant Cell*, vol. 9, no. 2, pp. 209–221, 1997.
- [17] Z. Chen, H. Silva, and D. F. Klessig, "Active oxygen species in the induction of plant systemic acquired resistance by salicylic acid," *Science*, vol. 262, no. 5141, pp. 1883–1886, 1993.
- [18] T. Janda, G. Szalai, K. Rios-Gonzalez, O. Veisz, and E. Páldi, "Comparative study of frost tolerance and antioxidant activity in cereals," *Plant Science*, vol. 164, no. 2, pp. 301–306, 2003.
- [19] Y. J. Yuan, C. Li, Z. D. Hu, and J. C. Wu, "Signal transduction pathway for oxidative burst and taxol production in suspension cultures of *Taxus chinensis* var. mairei induced by oligosaccharide from *Fusarium oxysprum*," *Enzyme and Microbial Technology*, vol. 29, no. 6-7, pp. 372–379, 2001.
- [20] M. Xu and J. Dong, "O²⁻ from elicitor-induced oxidative burst is necessary for triggering phenylalanine ammonia-lyase activation and catharanthine synthesis in *Catharanthus roseus* cell cultures," *Enzyme and Microbial Technology*, vol. 36, no. 2-3, pp. 280–284, 2005.
- [21] K. T. Hung, D. G. Cheng, Y. T. Hsu, and C. H. Kao, "Absciscic acid-induced hydrogen peroxide is required for anthocyanin accumulation in leaves of rice seedlings," *Journal of Plant Physiology*, vol. 165, no. 12, pp. 1280–1287, 2008.
- [22] M. Perassolo, C. V. Quevedo, V. D. Busto, A. M. Giulietti, and J. R. Talou, "Role of reactive oxygen species and proline cycle in anthraquinone accumulation in *Rubia tinctorum* cell suspension cultures subjected to methyl jasmonate elicitation," *Plant Physiology and Biochemistry*, vol. 49, no. 7, pp. 758–763, 2011.
- [23] "Chinese Pharmacopoeia Commission," Pharmacopoeia of the People's Republic of China, Beijing, China, 2010.
- [24] M. Jiao, R. Cao, H. Chen, W. Hao, and J. Dong, "Effects of salicylic acid on synthesis of rosmarinic acid and related enzymes in the suspension cultures of *Salvia miltiorrhiza*," *Shengwu Gongcheng Xuebao/Chinese Journal of Biotechnology*, vol. 28, no. 3, pp. 320–328, 2012.
- [25] C. K. Auh and T. M. Murphy, "Plasma membrane redox enzyme is involved in the synthesis of O²⁻ and H₂O₂ by *Phytophthora* elicitor-stimulated rose cells," *Plant Physiology*, vol. 107, no. 4, pp. 1241–1247, 1995.
- [26] J. Foreman, V. Demidchik, J. H. F. Bothwell et al., "Reactive oxygen species produced by NADPH oxidase regulate plant cell growth," *Nature*, vol. 422, no. 6930, pp. 442–446, 2003.
- [27] C. An and Z. Mou, "Salicylic acid and its function in plant immunity," *Journal of Integrative Plant Biology*, vol. 53, no. 6, pp. 412–428, 2011.
- [28] R.-Q. Zhang, H.-H. Zhu, H.-Q. Zhao, and Q. Yao, "Arbuscular mycorrhizal fungal inoculation increases phenolic synthesis in clover roots via hydrogen peroxide, salicylic acid and nitric oxide signaling pathways," *Journal of Plant Physiology*, vol. 170, no. 1, pp. 74–79, 2013.
- [29] K. Zitta, P. Meybohm, B. Bein et al., "Salicylic acid induces apoptosis in colon carcinoma cells grown in-vitro: influence of oxygen and salicylic acid concentration," *Experimental Cell Research*, vol. 318, no. 7, pp. 828–834, 2012.
- [30] S. Mandal, I. Kar, A. K. Mukherjee, and P. Acharya, "Elicitor-induced defense responses in *Solanum lycopersium* against *Ralstonia solanacearum*," *The Scientific World Journal*, vol. 2013, Article ID 561056, 9 pages, 2013.
- [31] S. Mandal, N. Mallick, and A. Mitra, "Salicylic acid-induced resistance to *Fusarium oxysporum* f. sp. lycopersici in tomato," *Plant Physiology and Biochemistry*, vol. 47, no. 7, pp. 642–649, 2009.
- [32] Y. Wang, C. C. Dai, Y. W. Zhao, and Y. Peng, "Fungal endophyte-induced volatile oil accumulation in *Atractylodes lancea* plantlets is mediated by nitric oxide, salicylic acid and hydrogen peroxide," *Process Biochemistry*, vol. 46, no. 3, pp. 730–735, 2011.
- [33] D. Solecka and A. Kacperska, "Phenylpropanoid deficiency affects the course of plant acclimation to cold," *Physiologia Plantarum*, vol. 119, no. 2, pp. 253–262, 2003.
- [34] S. W. Li, L. Xue, S. Xu, H. Feng, and L. An, "Hydrogen peroxide acts as a signal molecule in the adventitious root formation of mung bean seedlings," *Environmental and Experimental Botany*, vol. 65, no. 1, pp. 63–71, 2009.