Research Article

***L*-Fuzzy Fixed Points Theorems for *L*-Fuzzy Mappings via $\beta_{\mathcal{F}_L}$ -Admissible Pair**

Maliha Rashid,¹ Akbar Azam,² and Nayyar Mehmood²

¹ Department of Mathematics, International Islamic University, Sector H-10, Islamabad 44000, Pakistan

² Department of Mathematics, COMSATS Institute of Information Technology, Chack Shahzad, Islamabad 44000, Pakistan

Correspondence should be addressed to Akbar Azam; akbarazam@yahoo.com

Received 30 August 2013; Accepted 10 November 2013; Published 5 February 2014

Academic Editors: G. Bonanno and J. Wang

Copyright © 2014 Maliha Rashid et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We define the concept of $\beta_{\mathcal{F}_L}$ -admissible for a pair of *L*-fuzzy mappings and establish the existence of common *L*-fuzzy fixed point theorem. Our result generalizes some useful results in the literature. We provide an example to support our result.

1. Introduction

A large variety of the problems of analysis and applied mathematics relate to finding solutions of nonlinear functional equations which can be formulated in terms of finding the fixed points of nonlinear mappings. Heilpern [1] first introduced the concept of fuzzy mappings and established a fixed point theorem for fuzzy contraction mappings in complete metric linear spaces, which is a fuzzy extension of Banach contraction principle and Nadler's [2] fixed point theorem. Subsequently several other authors [3–17] generalized this result and studied the existence of fixed points and common fixed points of fuzzy mappings satisfying a contractive type condition.

Zadeh published his important paper “Fuzzy sets” [18], after that Goguen published the paper “*L*-Fuzzy sets” [19]. The concept of *L*-fuzzy sets is a generalization of the concept of fuzzy sets. Fuzzy set is a special case of *L*-fuzzy set when $L = [0, 1]$. There are basically two understandings of the meaning of *L*, one is when *L* is a complete lattice equipped with a multiplication $*$ operator satisfying certain conditions as shown in the initial paper [19] and the second understanding of the meaning of *L* is that *L* is a completely distributive complete lattice with an order-reversing involution (see, e.g., [20–22], etc.).

In 2012, Samet et al. [23] introduced the concept of β -admissible mapping and established fixed point theorems

via β -admissible and also showed that these results can be utilized to derive fixed point theorems in partially ordered spaces and coupled fixed point theorems. Moreover, they applied the main results to ordinary differential equations. Afterwards, Asl et al. [24] extended the concept of β -admissible for single valued mappings to multivalued mappings. Recently, Mohammadi et al. [25] introduced the concept of β -admissible for multivalued mappings which is different from the notion of β_* -admissible which has been provided in [24] and Azam and Beg [26] obtained a common α -fuzzy fixed point of a pair of fuzzy mappings on a complete metric space under a generalized contractive condition for α -level sets via Hausdorff metric for fuzzy sets.

In this paper we introduce the concept of $\beta_{\mathcal{F}_L}$ -admissible for a pair of *L*-fuzzy mappings and establish the existence of common *L*-fuzzy fixed point theorem. We also have given an example to support our main theorem.

2. Preliminaries

Let (X, d) be a metric space, and denote

$CB(X) = \{A : A \text{ is nonempty closed and bounded subset of } X\},$

$C(X) = \{A : A \text{ is nonempty compact subset of } X\}.$

For $\epsilon > 0$ and the sets $A, B \in CB(X)$ define

$$\begin{aligned} d(x, A) &= \inf_{y \in A} d(x, y), \\ d(A, B) &= \inf_{x \in A, y \in B} d(x, y), \end{aligned} \quad (1)$$

$$N(\epsilon, A) = \{x \in X : d(x, a) < \epsilon, \text{ for some } a \in A\},$$

$$E_{A,B} = \{\epsilon > 0 : A \subseteq N(\epsilon, B), B \subseteq N(\epsilon, A)\}.$$

Then the Hausdorff metric d_H on $CB(X)$ induced by d is defined as

$$d_H(A, B) = \inf E_{A,B}. \quad (2)$$

Lemma 1 (see [2]). Let (X, d) be a metric space and $A, B \in CB(X)$; then for each $a \in A$

$$d(a, B) \leq H(A, B). \quad (3)$$

Lemma 2 (see [2]). Let (X, d) be a metric space and $A, B \in CB(X)$; then for each $a \in A$, $\epsilon > 0$, there exists an element $b \in B$ such that

$$d(a, b) \leq H(A, B) + \epsilon. \quad (4)$$

Definition 3 (see [19]). A partially ordered set $(L, \preceq_L)$ is called

- (i) a lattice, if $a \vee b \in L$, $a \wedge b \in L$ for any $a, b \in L$;
- (ii) a complete lattice, if $\bigvee A \in L$, $\bigwedge A \in L$ for any $A \subseteq L$;
- (iii) distributive if $a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$, $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$ for any $a, b, c \in L$.

Definition 4 (see [19]). Let L be a lattice with top element 1_L and bottom element 0_L and let $a, b \in L$. Then b is called a complement of a , if $a \vee b = 1_L$, and $a \wedge b = 0_L$. If $a \in L$ has a complement element, then it is unique. It is denoted by $\hat{a}$.

Definition 5 (see [19]). An L -fuzzy set A on a nonempty set X is a function $A : X \rightarrow L$, where L is complete distributive lattice with 1_L and 0_L .

Remark 6. The class of L -fuzzy sets is larger than the class of fuzzy sets as an L -fuzzy set is a fuzzy set if $L = [0, 1]$.

The α_L -level set of L -fuzzy set A is denoted by A_{α_L} and is defined as follows:

$$\begin{aligned} A_{\alpha_L} &= \{x : \alpha_L \preceq_L A(x)\} \quad \text{if } \alpha_L \in L \setminus \{0_L\}, \\ A_{0_L} &= \text{cl}(\{x : 0_L \preceq_L A(x)\}). \end{aligned} \quad (5)$$

Here $\text{cl}(B)$ denotes the closure of the set B .

We denote and define the characteristic function χ_{L_A} of an L -fuzzy set A as follows:

$$\chi_{L_A} := \begin{cases} 0_L & \text{if } x \notin A, \\ 1_L & \text{if } x \in A. \end{cases} \quad (6)$$

Definition 7. Let X be an arbitrary set and Y a metric space. A mapping T is called L -fuzzy mapping if T is a mapping from X into $\mathcal{F}_L(Y)$. An L -fuzzy mapping T is an L -fuzzy subset on $X \times Y$ with membership function $T(x)(y)$. The function $T(x)(y)$ is the grade of membership of y in $T(x)$.

Definition 8. Let (X, d) be a metric space and S, T L -fuzzy mappings from X into $\mathcal{F}_L(X)$. A point $z \in X$ is called an L -fuzzy fixed point of T if $z \in [Tx]_{\alpha_L}$, where $\alpha_L \in L \setminus \{0_L\}$. The point $z \in X$ is called a common L -fuzzy fixed point of S and T if $z \in [Sx]_{\alpha_L} \cap [Tx]_{\alpha_L}$.

Definition 9 (see [23]). Let X be a nonempty set, $T : X \rightarrow X$, and $\beta : X \times X \rightarrow [0, \infty)$. We say that T is β -admissible if for all $x, y \in X$ we have

$$\beta(x, y) \geq 1 \implies \beta(Tx, Ty) \geq 1. \quad (7)$$

Definition 10 (see [24]). Let X be a nonempty set, $T : X \rightarrow 2^X$, where 2^X is a collection of subset of X , $\beta : X \times X \rightarrow [0, \infty)$ and $\beta_* : 2^X \times 2^X \rightarrow [0, \infty)$. We say that T is β_* -admissible if for all $x, y \in X$ we have

$$\beta(x, y) \geq 1 \implies \beta_*(Tx, Ty) \geq 1. \quad (8)$$

Definition 11 (see [25]). Let X be a nonempty set, $T : X \rightarrow 2^X$, where 2^X is a collection of subset of X and $\beta : X \times X \rightarrow [0, \infty)$. We say that T is β -admissible whenever for each $x \in X$ and $y \in Tx$ with $\beta(x, y) \geq 1$ we have $\beta(y, z) \geq 1$ for all $z \in Ty$.

3. Main Result

In this section, we introduce a new concept of $\beta_{\mathcal{F}_L}$ -admissible for a pair of L -fuzzy mappings and establish the existence of common L -fuzzy fixed point theorem.

Definition 12. Let (X, d) be a metric space, $\beta : X \times X \rightarrow [0, \infty)$, and S, T L -fuzzy mappings from X into $\mathcal{F}_L(X)$. The order pair (S, T) is said to be $\beta_{\mathcal{F}_L}$ -admissible if it satisfies the following conditions:

- (i) for each $x \in X$ and $y \in [Sx]_{\alpha_L(x)}$, where $\alpha_L(x) \in L \setminus \{0_L\}$, with $\beta(x, y) \geq 1$, we have $\beta(y, z) \geq 1$ for all $z \in [Ty]_{\alpha_L(y)} \neq \emptyset$, where $\alpha_L(y) \in L \setminus \{0_L\}$;
- (ii) for each $x \in X$ and $y \in [Tx]_{\alpha_L(x)}$, where $\alpha_L(x) \in L \setminus \{0_L\}$, with $\beta(x, y) \geq 1$, we have $\beta(y, z) \geq 1$ for all $z \in [Sy]_{\alpha_L(y)} \neq \emptyset$, where $\alpha_L(y) \in L \setminus \{0_L\}$.

If $S = T$ then T is called $\beta_{\mathcal{F}_L}$ -admissible.

Remark 13. It is easy to see that if (S, T) is $\beta_{\mathcal{F}_L}$ -admissible, then (T, S) is also $\beta_{\mathcal{F}_L}$ -admissible.

Next, we give a common L -fuzzy fixed point theorem for β_F -admissible pair.

Theorem 14. Let (X, d) be a complete metric space, $\beta : X \times X \rightarrow [0, \infty)$, and S, T L -fuzzy mappings from X into $\mathcal{F}_L(X)$ satisfying the following conditions.

- (a) For each $x \in X$, there exists $\alpha_L(x) \in L \setminus \{0_L\}$ such that $[Sx]_{\alpha_L(x)}, [Tx]_{\alpha_L(x)}$ are nonempty closed bounded subsets of X and for $x_0 \in X$, there exists $x_1 \in [Sx_0]_{\alpha_L(x_0)}$ with $\beta(x_0, x_1) \geq 1$.

(b) For all $x, y \in X$, we have

$$\begin{aligned} & \max \{ \beta(x, y), \beta(y, x) \} [H([Sx]_{\alpha_L(x)}, [Ty]_{\alpha_L(y)})] \\ & \leq a_1(d(x, [Sx]_{\alpha_L(x)}) + a_2(dy, [Ty]_{\alpha_L(y)}) \\ & \quad + a_3(dx, [Ty]_{\alpha_L(y)}) + a_4(dy, [Sx]_{\alpha_L(x)}) + a_5d(x, y), \end{aligned} \quad (9)$$

where a_1, a_2, a_3, a_4 and a_5 are nonnegative real numbers and $\sum_{i=1}^5 a_i < 1$ and either $a_1 = a_2$ or $a_3 = a_4$.

(c) (S, T) is $\beta_{\mathcal{F}_L}$ -admissible pair.

(d) If $\{x_n\} \in X$, such that $\beta(x_n, x_{n+1}) \geq 1$ and $x_n \rightarrow x$, then $\beta(x_n, x) \geq 1$.

Then there exists $z \in X$ such that $z \in [Sz]_{\alpha_L(z)} \cap [Tz]_{\alpha_L(z)}$.

Proof. We will prove the above result by considering the following three cases:

- (1) $a_1 + a_3 + a_5 = 0$,
- (2) $a_2 + a_4 + a_5 = 0$,
- (3) $a_1 + a_3 + a_5 \neq 0$ and $a_2 + a_4 + a_5 \neq 0$.

Case 1. For $x_0 \in X$ in condition (a), there exist $\alpha_L(x_0) \in L \setminus \{0_L\}$ and $x_1 \in [Sx_0]_{\alpha_L(x_0)}$ such that $\beta(x_0, x_1) \geq 1$ and also there exists $\alpha_L(x_1) \in L \setminus \{0_L\}$ such that $[Sx_0]_{\alpha_L(x_0)}$ and $[Tx_1]_{\alpha_L(x_1)}$ are nonempty closed bounded subsets of X . From Lemma 1, we obtain that

$$\begin{aligned} & d(x_1, [Tx_1]_{\alpha_L(x_1)}) \\ & \leq H([Sx_0]_{\alpha_L(x_0)}, [Tx_1]_{\alpha_L(x_1)}) \\ & \leq \beta(x_0, x_1) [H([Sx_0]_{\alpha_L(x_0)}, [Tx_1]_{\alpha_L(x_1)})] \quad (10) \\ & \leq \max \{ \beta(x_0, x_1), \beta(x_1, x_0) \} \\ & \quad \times [H([Sx_0]_{\alpha_L(x_0)}, [Tx_1]_{\alpha_L(x_1)})]. \end{aligned}$$

Now, inequality (9) implies that

$$\begin{aligned} & d(x_1, [Tx_1]_{\alpha_L(x_1)}) \\ & \leq a_1d(x_0, [Sx_0]_{\alpha_L(x_0)}) + a_2d(x_1, [Tx_1]_{\alpha_L(x_1)}) \\ & \quad + a_3d(x_0, [Tx_1]_{\alpha_L(x_1)}) \\ & \quad + a_4d(x_1, [Sx_0]_{\alpha_L(x_0)}) + a_5d(x_0, x_1). \end{aligned} \quad (11)$$

Using $a_1 + a_3 + a_5 = 0$ together with the fact that $d(x_1, [Sx_0]_{\alpha_L(x_0)}) = 0$, we get

$$d(x_1, [Tx_1]_{\alpha_L(x_1)}) \leq a_2d(x_1, [Tx_1]_{\alpha_L(x_1)}). \quad (12)$$

It follows that $x_1 \in [Tx_1]_{\alpha_L(x_1)}$, which further implies that

$$d(x_1, [Sx_1]_{\alpha_L(x_1)}) \leq H([Tx_1]_{\alpha_L(x_1)}, [Sx_1]_{\alpha_L(x_1)}). \quad (13)$$

By condition (c), for $x_0 \in X$ and $x_1 \in [Sx_0]_{\alpha_L(x_0)}$ such that $\beta(x_0, x_1) \geq 1$, we have $\beta(x_1, z) \geq 1$ for all $z \in [Tx_1]_{\alpha_L(x_1)}$. Since $x_1 \in [Tx_1]_{\alpha_L(x_1)}$, therefore $\beta(x_1, x_1) \geq 1$ and hence

$$\begin{aligned} & d(x_1, [Sx_1]_{\alpha_L(x_1)}) \\ & \leq \beta(x_1, x_1) [H([Sx_1]_{\alpha_L(x_1)}, [Tx_1]_{\alpha_L(x_1)})]. \end{aligned} \quad (14)$$

Again, inequality (9) implies that

$$\begin{aligned} & d(x_1, [Sx_1]_{\alpha_L(x_1)}) \\ & \leq a_1d(x_1, [Sx_1]_{\alpha_L(x_1)}) + a_2d(x_1, [Tx_1]_{\alpha_L(x_1)}) \\ & \quad + a_3d(x_1, [Tx_1]_{\alpha_L(x_1)}) \\ & \quad + a_4d(x_1, [Sx_1]_{\alpha_L(x_1)}) + a_5d(x_1, x_1). \end{aligned} \quad (15)$$

Since $a_1 + a_3 + a_5 = 0$ and $d(x_1, [Tx_1]_{\alpha_L(x_1)}) = 0$, we get

$$d(x_1, [Sx_1]_{\alpha_L(x_1)}) \leq a_4d(x_1, [Sx_1]_{\alpha_L(x_1)}), \quad (16)$$

which implies that $x_1 \in [Sx_1]_{\alpha_L(x_1)}$ and hence

$$x_1 \in [Sx_1]_{\alpha_L(x_1)} \cap [Tx_1]_{\alpha_L(x_1)}. \quad (17)$$

Case 2. For $x_0 \in X$ in condition (a), there exist $\alpha_L(x_0) \in L \setminus \{0_L\}$ and $x_1 \in [Sx_0]_{\alpha_L(x_0)}$ such that $\beta(x_0, x_1) \geq 1$ and also there exists $\alpha_L(x_1) \in L \setminus \{0_L\}$ such that $[Sx_0]_{\alpha_L(x_0)}$ and $[Tx_1]_{\alpha_L(x_1)}$ are nonempty closed bounded subsets of X . By condition (c), we have $\beta(x_1, x_2) \geq 1$ for all $x_2 \in [Tx_1]_{\alpha_L(x_1)}$. From Lemma 1, we obtain that

$$\begin{aligned} & d(x_2, [Sx_2]_{\alpha_L(x_2)}) \\ & \leq H([Tx_1]_{\alpha_L(x_1)}, [Sx_2]_{\alpha_L(x_2)}) \\ & \leq \beta(x_1, x_2) [H([Sx_2]_{\alpha_L(x_2)}, [Tx_1]_{\alpha_L(x_1)})] \\ & \leq \max \{ \beta(x_1, x_2), \beta(x_2, x_1) \} \\ & \quad \times [H([Sx_2]_{\alpha_L(x_2)}, [Tx_1]_{\alpha_L(x_1)})] \quad (18) \\ & \leq a_1d(x_2, [Sx_2]_{\alpha_L(x_2)}) + a_2d(x_1, [Tx_1]_{\alpha_L(x_1)}) \\ & \quad + a_3d(x_2, [Tx_1]_{\alpha_L(x_1)}) + a_4d(x_1, [Sx_2]_{\alpha_L(x_2)}) \\ & \quad + a_5d(x_2, x_1). \end{aligned}$$

Using $a_2 + a_4 + a_5 = 0$ together with the fact that $d(x_2, [Tx_1]_{\alpha_L(x_1)}) = 0$, we get

$$d(x_2, [Sx_2]_{\alpha_L(x_2)}) \leq a_1d(x_2, [Sx_2]_{\alpha_L(x_2)}). \quad (19)$$

It follows that $x_2 \in [Sx_2]_{\alpha_L(x_2)}$, which further implies that

$$d(x_2, [Tx_2]_{\alpha_L(x_2)}) \leq H([Sx_2]_{\alpha_L(x_2)}, [Tx_2]_{\alpha_L(x_2)}). \quad (20)$$

By condition (c), we have $\beta(x_2, x_2) \geq 1$, and hence

$$\begin{aligned} d(x_2, [Tx_2]_{\alpha_L(x_2)}) \\ \leq \beta(x_2, x_2) [H([Sx_2]_{\alpha_L(x_2)}, [Tx_2]_{\alpha_L(x_2)})]. \end{aligned} \quad (21)$$

Again, inequality (9) implies that

$$\begin{aligned} d(x_2, [Tx_2]_{\alpha_L(x_2)}) \\ \leq a_1 d(x_2, [Sx_2]_{\alpha_L(x_2)}) + a_2 d(x_2, [Tx_2]_{\alpha_L(x_2)}) \\ + a_3 d(x_2, [Tx_2]_{\alpha_L(x_2)}) \\ + a_4 d(x_2, [Sx_2]_{\alpha_L(x_2)}) + a_5 d(x_2, x_2). \end{aligned} \quad (22)$$

Since $a_2 + a_4 + a_5 = 0$ and $d(x_2, [Sx_2]_{\alpha_L(x_2)}) = 0$, we get

$$d(x_2, [Tx_2]_{\alpha_L(x_2)}) \leq a_3 d(x_2, [Tx_2]_{\alpha_L(x_2)}), \quad (23)$$

which implies that $x_2 \in [Tx_2]_{\alpha_L(x_2)}$ and hence

$$x_2 \in [Sx_2]_{\alpha_L(x_2)} \cap [Tx_2]_{\alpha_L(x_2)}. \quad (24)$$

Case 3. Let $\lambda = ((a_1 + a_3 + a_5)/(1 - a_2 - a_3))$ and $\mu = ((a_2 + a_4 + a_5)/(1 - a_1 - a_4))$. Next, we show that if $a_1 = a_2$ or $a_3 = a_4$, then $0 < \lambda\mu < 1$.

If $a_3 = a_4$, then $\lambda, \mu < 1$ and so $0 < \lambda\mu < 1$. Now if $a_1 = a_2$, then

$$\begin{aligned} 0 < \lambda\mu &= \left(\frac{a_1 + a_3 + a_5}{1 - a_2 - a_3} \right) \left(\frac{a_2 + a_4 + a_5}{1 - a_1 - a_4} \right) \\ &= \left(\frac{a_1 + a_3 + a_5}{1 - a_1 - a_3} \right) \left(\frac{a_1 + a_4 + a_5}{1 - a_1 - a_4} \right) \\ &= \left(\frac{a_1 + a_3 + a_5}{1 - a_1 - a_4} \right) \left(\frac{a_1 + a_4 + a_5}{1 - a_1 - a_3} \right) < 1. \end{aligned} \quad (25)$$

By condition (a), for $x_1 \in X$, there exists $\alpha_L(x_1) \in L \setminus \{0_L\}$ such that $[Tx_1]_{\alpha_L(x_1)}$ is a nonempty closed bounded subset of X . Since $a_1 + a_3 + a_5 > 0$, by Lemma 2, there exists $x_2 \in [Tx_1]_{\alpha_L(x_1)}$ such that

$$\begin{aligned} d(x_1, x_2) \\ \leq H([Sx_0]_{\alpha_L(x_0)}, [Tx_1]_{\alpha_L(x_1)}) + a_1 + a_3 + a_5 \\ \leq \beta(x_0, x_1) [H([Sx_0]_{\alpha_L(x_0)}, [Tx_1]_{\alpha_L(x_1)})] \\ + a_1 + a_3 + a_5 \\ \leq \max\{\beta(x_0, x_1), \beta(x_1, x_0)\} \\ \times [H([Sx_0]_{\alpha_L(x_0)}, [Tx_1]_{\alpha_L(x_1)})] + a_1 + a_3 + a_5 \end{aligned}$$

$$\begin{aligned} &\leq a_1 d(x_0, [Sx_0]_{\alpha_L(x_0)}) + a_2 d(x_1, [Tx_1]_{\alpha_L(x_1)}) \\ &\quad + a_3 d(x_0, [Tx_1]_{\alpha_L(x_1)}) + a_4 d(x_1, [Sx_0]_{\alpha_L(x_0)}) \\ &\quad + a_5 d(x_0, x_1) + a_1 + a_3 + a_5 \\ &\leq (a_1 + a_5) d(x_0, x_1) + a_2 d(x_1, x_2) \\ &\quad + a_3 d(x_0, x_2) + a_1 + a_3 + a_5 \\ &\leq (a_1 + a_3 + a_5) d(x_0, x_1) + (a_2 + a_3) d(x_1, x_2) \\ &\quad + a_1 + a_3 + a_5. \end{aligned} \quad (26)$$

This implies that

$$d(x_1, x_2) \leq \lambda d(x_0, x_1) + \lambda. \quad (27)$$

By the same argument, for $x_2 \in X$, there exists $\alpha_L(x_2) \in L \setminus \{0_L\}$ such that $[Sx_2]_{\alpha_L(x_2)}$ is a nonempty closed bounded subset of X . Since $a_2 + a_4 + a_5 > 0$, by Lemma 2, there exists $x_3 \in [Sx_2]_{\alpha_L(x_2)}$ such that

$$d(x_2, x_3) \leq H([Tx_1]_{\alpha_L(x_1)}, [Sx_2]_{\alpha_L(x_2)}) + \lambda(a_2 + a_4 + a_5). \quad (28)$$

By condition (c), for $x_0 \in X$ and $x_1 \in [Sx_0]_{\alpha_L(x_0)}$ such that $\beta(x_0, x_1) \geq 1$, we have $\beta(x_1, x_2) \geq 1$ for $x_2 \in [Tx_1]_{\alpha_L(x_1)}$. So we have

$$\begin{aligned} d(x_2, x_3) \\ \leq H([Tx_1]_{\alpha_L(x_1)}, [Sx_2]_{\alpha_L(x_2)}) + \lambda(a_2 + a_4 + a_5) \\ = H([Sx_2]_{\alpha_L(x_2)}, [Tx_1]_{\alpha_L(x_1)}) + \lambda(a_2 + a_4 + a_5) \\ \leq \beta(x_1, x_2) [H([Sx_2]_{\alpha_L(x_2)}, [Tx_1]_{\alpha_L(x_1)})] \\ + \lambda(a_2 + a_4 + a_5) \\ \leq \max\{\beta(x_1, x_2), \beta(x_2, x_1)\} \\ \times [H([Sx_2]_{\alpha_L(x_2)}, [Tx_1]_{\alpha_L(x_1)})] + \lambda(a_2 + a_4 + a_5) \\ \leq a_1 d(x_2, [Sx_2]_{\alpha_L(x_2)}) + a_2 d(x_1, [Tx_1]_{\alpha_L(x_1)}) \\ + a_3 d(x_2, [Tx_1]_{\alpha_L(x_1)}) + a_4 d(x_1, [Sx_2]_{\alpha_L(x_2)}) \\ + a_5 d(x_2, x_1) + \lambda(a_2 + a_4 + a_5) \\ \leq a_1 d(x_2, x_3) + (a_2 + a_5) d(x_1, x_2) \\ + a_4 d(x_1, x_3) + \lambda(a_2 + a_4 + a_5) \\ \leq (a_2 + a_4 + a_5) d(x_1, x_2) \\ + (a_1 + a_4) d(x_2, x_3) + \lambda(a_2 + a_4 + a_5). \end{aligned} \quad (29)$$

This implies that

$$d(x_2, x_3) \leq \mu d(x_1, x_2) + \lambda\mu. \quad (30)$$

By repeating the above process, for $x_3 \in X$, there exists $\alpha_L(x_3) \in L \setminus \{0_L\}$ such that $[Tx_3]_{\alpha_L(x_3)}$ is a nonempty closed bounded subset of X . From Lemma 2, there exists $x_4 \in [Tx_3]_{\alpha_L(x_3)}$ such that

$$d(x_3, x_4) \leq H([Sx_2]_{\alpha_L(x_2)}, [Tx_3]_{\alpha_L(x_3)}) + \lambda\mu(a_1 + a_3 + a_5). \quad (31)$$

By condition (c), for $x_1 \in X$ and $x_2 \in [Tx_1]_{\alpha_L(x_1)}$ such that $\beta(x_1, x_2) \geq 1$, we have $\beta(x_2, x_3) \geq 1$ for $x_3 \in [Sx_2]_{\alpha_L(x_2)}$. So we have

$$\begin{aligned} d(x_3, x_4) &\leq H([Sx_2]_{\alpha_L(x_2)}, [Tx_3]_{\alpha_L(x_3)}) \\ &\quad + \lambda\mu(a_1 + a_3 + a_5) \\ &\leq \beta(x_2, x_3) [H([Sx_2]_{\alpha_L(x_2)}, [Tx_3]_{\alpha_L(x_3)})] \\ &\quad + \lambda\mu(a_1 + a_3 + a_5) \\ &\leq \max\{\beta(x_2, x_3), \beta(x_3, x_2)\} \\ &\quad \times [H([Sx_2]_{\alpha_L(x_2)}, [Tx_3]_{\alpha_L(x_3)})] \\ &\quad + \lambda\mu(a_1 + a_3 + a_5) \\ &\leq a_1 d(x_2, [Sx_2]_{\alpha_L(x_2)}) + a_2 d(x_3, [Tx_3]_{\alpha_L(x_3)}) \\ &\quad + a_3 d(x_2, [Tx_3]_{\alpha_L(x_3)}) + a_4 d(x_3, [Sx_2]_{\alpha_L(x_2)}) \\ &\quad + a_5 d(x_2, x_3) + \lambda\mu(a_1 + a_3 + a_5) \\ &\leq (a_1 + a_5) d(x_2, x_3) + a_2 d(x_3, x_4) \\ &\quad + a_3 d(x_2, x_4) + \lambda\mu(a_1 + a_3 + a_5) \\ &\leq (a_1 + a_3 + a_5) d(x_2, x_3) \\ &\quad + (a_2 + a_3) d(x_3, x_4) + \lambda\mu(a_1 + a_3 + a_5). \end{aligned} \quad (32)$$

This implies that

$$d(x_3, x_4) \leq \lambda d(x_2, x_3) + \lambda(\lambda\mu). \quad (33)$$

By induction, we produce a sequence $\{x_n\}$ in X such that

$$\begin{aligned} x_{2k+1} &\in [Sx_{2k}]_{\alpha_L(x_{2k})}, \\ x_{2k+2} &\in [Tx_{2k+1}]_{\alpha_L(x_{2k+1})}, \quad k = 0, 1, 2, \dots, \\ \beta(x_{n-1}, x_n) &\geq 1, \quad \forall n \in \mathbb{N}. \end{aligned} \quad (34)$$

Now, we have

$$\begin{aligned} d(x_{2k+1}, x_{2k+2}) &\leq H([Sx_{2k}]_{\alpha_L(x_{2k})}, [Tx_{2k+1}]_{\alpha_L(x_{2k+1})}) \\ &\quad + (\lambda\mu)^k (a_1 + a_3 + a_5) \end{aligned}$$

$$\begin{aligned} &\leq \beta(x_{2k}, x_{2k+1}) \\ &\quad \times [H([Sx_{2k}]_{\alpha_L(x_{2k})}, [Tx_{2k+1}]_{\alpha_L(x_{2k+1})})] \\ &\quad + (\lambda\mu)^k (a_1 + a_3 + a_5) \\ &\leq \max\{\beta(x_{2k}, x_{2k+1}), \beta(x_{2k+1}, x_{2k})\} \\ &\quad \times [H([Sx_{2k}]_{\alpha_L(x_{2k})}, [Tx_{2k+1}]_{\alpha_L(x_{2k+1})})] \\ &\quad + (\lambda\mu)^k (a_1 + a_3 + a_5) \\ &\leq a_1 d(x_{2k}, [Sx_{2k}]_{\alpha_L(x_{2k})}) \\ &\quad + a_2 d(x_{2k+1}, [Tx_{2k+1}]_{\alpha_L(x_{2k+1})}) \\ &\quad + a_3 d(x_{2k}, [Tx_{2k+1}]_{\alpha_L(x_{2k+1})}) \\ &\quad + a_4 d(x_{2k+1}, [Sx_{2k}]_{\alpha_L(x_{2k})}) + a_5 d(x_{2k}, x_{2k+1}) \\ &\quad + (\lambda\mu)^k (a_1 + a_3 + a_5) \\ &\leq (a_1 + a_5) d(x_{2k}, x_{2k+1}) \\ &\quad + a_2 d(x_{2k+1}, x_{2k+2}) + a_3 d(x_{2k}, x_{2k+2}) \\ &\quad + (\lambda\mu)^k (a_1 + a_3 + a_5) \\ &\leq (a_1 + a_3 + a_5) d(x_{2k}, x_{2k+1}) \\ &\quad + (a_2 + a_3) d(x_{2k+1}, x_{2k+2}) + (\lambda\mu)^k (a_1 + a_3 + a_5). \end{aligned} \quad (35)$$

This implies that

$$d(x_{2k+1}, x_{2k+2}) \leq \lambda d(x_{2k}, x_{2k+1}) + \lambda(\lambda\mu)^k. \quad (36)$$

Similarly,

$$\begin{aligned} d(x_{2k+2}, x_{2k+3}) &\leq H([Sx_{2k+2}]_{\alpha_L(x_{2k+2})}, [Tx_{2k+1}]_{\alpha_L(x_{2k+1})}) \\ &\quad + (\lambda\mu)^k \lambda (a_2 + a_4 + a_5) \\ &\leq \beta(x_{2k+1}, x_{2k+2}) \\ &\quad \times [H([Sx_{2k+2}]_{\alpha_L(x_{2k+2})}, [Tx_{2k+1}]_{\alpha_L(x_{2k+1})})] \\ &\quad + (\lambda\mu)^k \lambda (a_2 + a_4 + a_5) \\ &\leq \max\{\beta(x_{2k+1}, x_{2k+2}), \beta(x_{2k+2}, x_{2k+1})\} \\ &\quad \times [H([Sx_{2k+2}]_{\alpha_L(x_{2k+2})}, [Tx_{2k+1}]_{\alpha_L(x_{2k+1})})] \\ &\quad + (\lambda\mu)^k \lambda (a_2 + a_4 + a_5) \\ &\leq a_1 d(x_{2k+2}, [Sx_{2k+2}]_{\alpha_L(x_{2k+2})}) \\ &\quad + a_2 d(x_{2k+1}, [Tx_{2k+1}]_{\alpha_L(x_{2k+1})}) \\ &\quad + a_3 d(x_{2k+2}, [Tx_{2k+1}]_{\alpha_L(x_{2k+1})}) \\ &\quad + a_4 d(x_{2k+1}, [Sx_{2k+2}]_{\alpha_L(x_{2k+2})}) \\ &\quad + a_5 d(x_{2k+2}, x_{2k+1}) + (\lambda\mu)^k \lambda (a_2 + a_4 + a_5) \end{aligned}$$

$$\begin{aligned}
&\leq a_1 d(x_{2k+2}, x_{2k+3}) \\
&\quad + (a_2 + a_5) d(x_{2k+1}, x_{2k+2}) \\
&\quad + a_4 d(x_{2k+1}, x_{2k+3}) + (\lambda\mu)^k \lambda (a_2 + a_4 + a_5) \\
&\leq (a_2 + a_4 + a_5) d(x_{2k+1}, x_{2k+2}) \\
&\quad + (a_1 + a_4) d(x_{2k+2}, x_{2k+3}) \\
&\quad + (\lambda\mu)^k \lambda (a_2 + a_4 + a_5).
\end{aligned} \tag{37}$$

This implies that

$$d(x_{2k+2}, x_{2k+3}) \leq \mu d(x_{2k+1}, x_{2k+2}) + (\lambda\mu)^{k+1}. \tag{38}$$

From (36) and (38), it follows that, for each $k = 0, 1, 2, \dots$,

$$\begin{aligned}
&d(x_{2k+1}, x_{2k+2}) \\
&\leq \lambda d(x_{2k}, x_{2k+1}) + \lambda(\lambda\mu)^k \\
&\leq \lambda [\mu d(x_{2k-1}, x_{2k}) + (\lambda\mu)^k] + \lambda(\lambda\mu)^k \\
&= (\lambda\mu) d(x_{2k-1}, x_{2k}) + 2\lambda(\lambda\mu)^k \\
&\leq (\lambda\mu) [\lambda d(x_{2k-2}, x_{2k-1}) + \lambda(\lambda\mu)^{k-1}] + 2\lambda(\lambda\mu)^k \\
&= (\lambda\mu) \lambda d(x_{2k-2}, x_{2k-1}) + 3\lambda(\lambda\mu)^k \\
&\vdots \\
&\leq \lambda(\lambda\mu)^k d(x_0, x_1) + (2k+1) \lambda(\lambda\mu)^k, \\
&d(x_{2k+2}, x_{2k+3}) \\
&\leq \mu d(x_{2k+1}, x_{2k+2}) + (\lambda\mu)^{k+1} \\
&\vdots \\
&\leq (\lambda\mu)^{k+1} d(x_0, x_1) + (2k+2) (\lambda\mu)^{k+1}.
\end{aligned} \tag{39}$$

Then for $m < n$, we have

$$\begin{aligned}
d(x_{2m+1}, x_{2n+1}) &\leq d(x_{2m+1}, x_{2m+2}) + d(x_{2m+2}, x_{2m+3}) \\
&\quad + d(x_{2m+3}, x_{2m+4}) + \dots + d(x_{2n}, x_{2n+1}) \\
&\leq \left[\lambda \sum_{i=m}^{n-1} (\lambda\mu)^i + \sum_{i=m+1}^n (\lambda\mu)^i \right] d(x_0, x_1) \\
&\quad + \lambda \sum_{i=m}^{n-1} (2i+1) (\lambda\mu)^i + \sum_{i=m+1}^n 2i(\lambda\mu)^i.
\end{aligned} \tag{40}$$

Similarly, we obtain that

$$\begin{aligned}
d(x_{2m}, x_{2n+1}) &\leq \left[\sum_{i=m}^n (\lambda\mu)^i + \lambda \sum_{i=m}^{n-1} (\lambda\mu)^i \right] d(x_0, x_1) \\
&\quad + \sum_{i=m}^n 2i(\lambda\mu)^i + \lambda \sum_{i=m}^{n-1} (2i+1) (\lambda\mu)^i, \\
d(x_{2m}, x_{2n}) &\leq \left[\sum_{i=m}^{n-1} (\lambda\mu)^i + \lambda \sum_{i=m}^{n-1} (\lambda\mu)^i \right] d(x_0, x_1) \\
&\quad + \sum_{i=m}^{n-1} 2i(\lambda\mu)^i + \lambda \sum_{i=m}^{n-1} (2i+1) (\lambda\mu)^i, \\
d(x_{2m+1}, x_{2n}) &\leq \left[\lambda \sum_{i=m}^{n-1} (\lambda\mu)^i + \sum_{i=m+1}^{n-1} (\lambda\mu)^i \right] d(x_0, x_1) \\
&\quad + \lambda \sum_{i=m}^{n-1} (2i+1) (\lambda\mu)^i + \sum_{i=m+1}^{n-1} 2i(\lambda\mu)^i.
\end{aligned} \tag{41}$$

Since $0 < \lambda\mu < 1$, so by Cauchy's root test, we get $\sum (2i+1)(\lambda\mu)^i$ and $\sum 2i(\lambda\mu)^i$ are convergent series. Therefore, $\{x_n\}$ is a Cauchy sequence in X . Now, from the completeness of X , there exists $z \in X$ such that $x_n \rightarrow z$ as $n \rightarrow \infty$. By condition (d), we have $\beta(x_{n-1}, z) \geq 1$ for all $n \in \mathbb{N}$. Now, we have

$$\begin{aligned}
&d(x_{2n}, [Sz]_{\alpha(z)}) \\
&\leq H([Tx_{2n-1}]_{\alpha_L(x_{2n-1})}, [Sz]_{\alpha(z)}) \\
&= H([Sz]_{\alpha(z)}, [Tx_{2n-1}]_{\alpha_L(x_{2n-1})}) \\
&\leq \beta(x_{2n-1}, z) H([Sz]_{\alpha(z)}, [Tx_{2n-1}]_{\alpha_L(x_{2n-1})}) \\
&\leq \max\{\beta(x_{2n-1}, z), \beta(z, x_{2n-1})\} \\
&\quad \times H([Sz]_{\alpha(z)}, [Tx_{2n-1}]_{\alpha_L(x_{2n-1})}) \\
&\leq a_1 d(z, [Sz]_{\alpha_L(z)}) \\
&\quad + a_2 d(x_{2n-1}, [Tx_{2n-1}]_{\alpha_L(x_{2n-1})}) \\
&\quad + a_3 d(z, [Tx_{2n-1}]_{\alpha_L(x_{2n-1})}) \\
&\quad + a_4 d(x_{2n-1}, [Sz]_{\alpha_L(z)}) + a_5 d(z, x_{2n-1}) \\
&\leq a_1 d(z, [Sz]_{\alpha_L(z)}) + a_2 d(x_{2n-1}, x_{2n}) + a_3 d(z, x_{2n}) \\
&\quad + a_4 d(x_{2n-1}, [Sz]_{\alpha_L(z)}) + a_5 d(z, x_{2n-1}).
\end{aligned} \tag{42}$$

Since

$$d(z, [Sz]_{\alpha_L(z)}) \leq d(z, x_{2n}) + d(x_{2n}, [Sz]_{\alpha_L(z)}), \tag{43}$$

so we get

$$\begin{aligned}
 d(z, [Sz]_{\alpha_L(z)}) &\leq d(z, x_{2n}) + a_1 d(z, [Sz]_{\alpha_L(z)}) \\
 &\quad + a_2 d(x_{2n-1}, x_{2n}) + a_3 d(z, x_{2n}) \\
 &\quad + a_4 d(x_{2n-1}, [Sz]_{\alpha_L(z)}) + a_5 d(z, x_{2n-1}) \quad (44) \\
 &\leq (1 + a_3) d(z, x_{2n}) \\
 &\quad + (a_4 + a_5) d(z, x_{2n-1}) + a_2 d(x_{2n-1}, x_{2n}) \\
 &\quad + (a_1 + a_4) d(z, [Sz]_{\alpha_L(z)}).
 \end{aligned}$$

This implies that

$$\begin{aligned}
 d(z, [Sz]_{\alpha_L(z)}) &\leq \left(\frac{1 + a_3}{1 - a_1 - a_4} \right) d(z, x_{2n}) \\
 &\quad + \left(\frac{a_4 + a_5}{1 - a_1 - a_4} \right) d(z, x_{2n-1}) \quad (45) \\
 &\quad + \left(\frac{a_2}{1 - a_1 - a_4} \right) d(x_{2n-1}, x_{2n}).
 \end{aligned}$$

Letting $n \rightarrow \infty$, we have $d(z, [Sz]_{\alpha_L(z)}) = 0$. It implies that $z \in [Sz]_{\alpha_L(z)}$. Similarly, by using

$$d(z, [Tz]_{\alpha_L(z)}) \leq d(z, x_{2n+1}) + d(x_{2n+1}, [Tz]_{\alpha_L(z)}), \quad (46)$$

we can show that $z \in [Tz]_{\alpha_L(z)}$. Therefore, $z \in [Sz]_{\alpha_L(z)} \cap [Tz]_{\alpha_L(z)}$. This completes the proof. $\square$

Next, we give an example to support the validity of our result.

Example 15. Let $X = [0, 1]$, $d(x, y) = |x - y|$, whenever $x, y \in X$; then (X, d) is a complete metric space. Let $L = \{\delta, \omega, \tau, \kappa\}$ with $\delta \preceq_L \omega \preceq_L \kappa$, $\delta \preceq_L \tau \preceq_L \kappa$, ω and τ are not comparable; then $(L, \preceq_L)$ is a complete distributive lattice. Define a pair of mappings $S, T : X \rightarrow \mathcal{F}_L(X)$ as follows:

$$\begin{aligned}
 S(x)(t) &= \begin{cases} \kappa, & \text{if } 0 \leq t \leq \frac{x}{6}, \\ \omega, & \text{if } \frac{x}{6} < t \leq \frac{x}{3}, \\ \tau, & \text{if } \frac{x}{3} < t \leq \frac{x}{2}, \\ \delta, & \text{if } \frac{x}{2} < t \leq 1, \end{cases} \\
 T(x)(t) &= \begin{cases} \kappa, & \text{if } 0 \leq t \leq \frac{x}{12}, \\ \delta, & \text{if } \frac{x}{12} < t \leq \frac{x}{8}, \\ \omega, & \text{if } \frac{x}{8} < t \leq \frac{x}{4}, \\ \tau, & \text{if } \frac{x}{4} < t \leq 1. \end{cases} \quad (47)
 \end{aligned}$$

Define $\beta : X \times X \rightarrow [0, \infty)$ as follows:

$$\beta(x, y) = \begin{cases} \frac{1}{|x - y|}, & x \neq y, \\ 1, & x = y. \end{cases} \quad (48)$$

For all $x \in X$, there exists $\alpha_L(x) = \kappa$, such that

$$[Sx]_\kappa = \left[0, \frac{x}{6}\right], \quad [Tx]_\kappa = \left[0, \frac{x}{12}\right], \quad (49)$$

and all conditions of the above theorem are satisfied. Hence, there exists $0 \in X$, such that $0 \in [S0]_{\alpha_L(0)} \cap [T0]_{\alpha_L(0)}$.

Corollary 16. Let (X, d) be a complete metric space, $\beta : X \times X \rightarrow [0, \infty)$, and S, T fuzzy mappings from X into $\mathcal{F}(X)$ satisfying the following conditions.

(a) For each $x \in X$, there exists $\alpha(x) \in (0, 1]$ such that $[Sx]_{\alpha(x)}, [Tx]_{\alpha(x)}$ are nonempty closed bounded subsets of X and for $x_0 \in X$, there exists $x_1 \in [Sx_0]_{\alpha(x_0)}$ with $\beta(x_0, x_1) \geq 1$.

(b) For all $x, y \in X$, we have

$$\begin{aligned}
 &\max\{\beta(x, y), \beta(y, x)\} [H([Sx]_{\alpha(x)}, [Ty]_{\alpha(y)})] \\
 &\leq a_1 (dx, [Sx]_{\alpha(x)}) + a_2 (dy, [Ty]_{\alpha(y)}) \\
 &\quad + a_3 (dx, [Ty]_{\alpha(y)}) \\
 &\quad + a_4 (dy, [Sx]_{\alpha(x)}) + a_5 d(x, y), \quad (50)
 \end{aligned}$$

where a_1, a_2, a_3, a_4 , and a_5 are nonnegative real numbers and $\sum_{i=1}^5 a_i < 1$ and either $a_1 = a_2$ or $a_3 = a_4$.

(c) (S, T) is $\beta_{\mathcal{F}}$ -admissible pair.

(d) If $\{x_n\} \in X$, such that $\beta(x_n, x_{n+1}) \geq 1$ and $x_n \rightarrow x$ then $\beta(x_n, x) \geq 1$.

Then there exists $z \in X$ such that $z \in [Sz]_{\alpha(z)} \cap [Tz]_{\alpha(z)}$.

Proof. Consider an L -fuzzy mapping $A : X \rightarrow \mathcal{F}_L(X)$ defined by

$$Ax = \chi_{L_{Tx}}. \quad (51)$$

Then for $\alpha_L \in L \setminus \{0_L\}$, we have

$$[Ax]_{\alpha_L} = Tx. \quad (52)$$

Hence by Theorem 14, we follow the result.

If we set $\beta(x, y) = 1$ for all $x, y \in X$ in Corollary 16, we get the following result. $\square$

Corollary 17 (see [26]). Let (X, d) be a complete metric space and S, T fuzzy mappings from X into $\mathcal{F}(X)$ satisfying the following conditions:

(a) for each $x \in X$, there exists $\alpha(x) \in (0, 1]$ such that $[Sx]_{\alpha(x)}, [Tx]_{\alpha(x)}$ are nonempty closed bounded subsets of X ;

(b) for all $x, y \in X$, we have

$$\begin{aligned} & H([Sx]_{\alpha(x)}, [Ty]_{\alpha(y)}) \\ & \leq a_1 d(x, [Sx]_{\alpha(x)}) + a_2 d(y, [Ty]_{\alpha(y)}) \\ & \quad + a_3 d(x, [Ty]_{\alpha(y)}) + a_4 d(y, [Sx]_{\alpha(x)}) \\ & \quad + a_5 d(x, y), \end{aligned} \quad (53)$$

where a_1, a_2, a_3, a_4 , and a_5 are nonnegative real numbers and $\sum_{i=1}^5 a_i < 1$ and either $a_1 = a_2$ or $a_3 = a_4$.

Then there exists $z \in X$ such that $z \in [Sz]_{\alpha(z)} \cap [Tz]_{\alpha(z)}$.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- [1] S. Heilpern, "Fuzzy mappings and fixed point theorem," *Journal of Mathematical Analysis and Applications*, vol. 83, no. 2, pp. 566–569, 1981.
- [2] B. Nadler, "Multivalued contraction mappings," *Pacific Journal of Mathematics*, vol. 30, pp. 475–488, 1969.
- [3] A. Azam, M. Waseem, and M. Rashid, "Fixed point theorems for fuzzy contractive mappings in quasi-pseudo-metric spaces," *Fixed Point Theory and Applications*, vol. 2013, article 27, 2013.
- [4] A. Azam, M. Arshad, and I. Beg, "Fixed points of fuzzy contractive and fuzzy locally contractive maps," *Chaos, Solitons and Fractals*, vol. 42, no. 5, pp. 2836–2841, 2009.
- [5] A. Azam and M. Arshad, "A note on "Fixed point theorems for fuzzy mappings" by P. Vijayaraju and M. Marudai," *Fuzzy Sets and Systems*, vol. 161, no. 8, pp. 1145–1149, 2010.
- [6] A. Azam, M. Arshad, and P. Vetro, "On a pair of fuzzy φ -contractive mappings," *Mathematical and Computer Modelling*, vol. 52, no. 1-2, pp. 207–214, 2010.
- [7] A. Azam and I. Beg, "Common fixed points of fuzzy maps," *Mathematical and Computer Modelling*, vol. 49, no. 7-8, pp. 1331–1336, 2009.
- [8] R. K. Bose and D. Sahani, "Fuzzy mappings and fixed point theorems," *Fuzzy Sets and Systems*, vol. 21, no. 1, pp. 53–58, 1987.
- [9] H. Román-Flores, A. Flores-Franulic, M. Rojas-Medar, and R. C. Bassanezi, "Stability of fixed points set of fuzzy contractions," *Applied Mathematics Letters*, vol. 11, no. 4, pp. 33–37, 1998.
- [10] B. Soo Lee and S. Jin Cho, "A fixed point theorem for contractive-type fuzzy mappings," *Fuzzy Sets and Systems*, vol. 61, no. 3, pp. 309–312, 1994.
- [11] B. S. Lee, G. M. Lee, S. J. Cho, and D. S. Kim, "Generalized common fixed point theorems for a sequence of fuzzy mappings," *International Journal of Mathematics and Mathematical Sciences*, vol. 17, no. 3, pp. 437–440, 1994.
- [12] J. Y. Park and J. U. Jeong, "Fixed point theorems for fuzzy mappings," *Fuzzy Sets and Systems*, vol. 87, no. 1, pp. 111–116, 1997.
- [13] R. A. Rashwan and M. A. Ahmad, "Common fixed point theorems for fuzzy mappings," *Archivum Mathematicum*, vol. 38, pp. 219–226, 2002.
- [14] B. E. Rhoades, "A common fixed point theorem for sequence of fuzzy mappings," *International Journal of Mathematics and Mathematical Sciences*, vol. 8, pp. 447–450, 1995.
- [15] T. Som and R. N. Mukherjee, "Some fixed point theorems for fuzzy mappings," *Fuzzy Sets and Systems*, vol. 33, no. 2, pp. 213–219, 1989.
- [16] P. Vijayaraju and M. Marudai, "Fixed point theorems for fuzzy mappings," *Fuzzy Sets and Systems*, vol. 135, no. 3, pp. 401–408, 2003.
- [17] P. Vijayaraju and R. Mohanraj, "Fixed point theorems for sequence of fuzzy mappings," *Southeast Asian Bulletin of Mathematics*, vol. 28, pp. 735–740, 2004.
- [18] L. A. Zadeh, "Fuzzy sets," *Information and Control*, vol. 8, no. 3, pp. 338–353, 1965.
- [19] J. A. Goguen, "L-fuzzy sets," *Journal of Mathematical Analysis and Applications*, vol. 18, pp. 145–174, 1967.
- [20] G. J. Wang, *Theory of L-Fuzzy Topological Spaces*, Shaanxi Normal University Press, Xi'an, China, 1988, (Chinese).
- [21] G.-J. Wang and Y.-Y. He, "Intuitionistic," *Fuzzy Sets and Systems*, vol. 110, no. 2, pp. 271–274, 2000.
- [22] D. S. Zhao, "The N-compactness in L-fuzzy topological spaces," *Journal of Mathematical Analysis and Applications*, vol. 128, no. 1, pp. 64–79, 1987.
- [23] B. Samet, C. Vetro, and P. Vetro, "Fixed point theorems for α - ψ -contractive type mappings," *Nonlinear Analysis, Theory, Methods and Applications*, vol. 75, no. 4, pp. 2154–2165, 2012.
- [24] J. H. Asl, S. Rezapour, and N. Shahzad, "On fixed points of α - ψ -contractive multifunctions," *Fixed Point Theory and Applications*, vol. 2012, article 212, 2012.
- [25] B. Mohammadi, S. Rezapour, and N. Shahzad, "Some results on fixed points of α - ψ -Ciric generalized multifunctions," *Fixed Point Theory and Applications*, vol. 2013, article 24, 2013.
- [26] A. Azam and I. Beg, "Common fuzzy fixed points for fuzzy mappings," *Fixed Point Theory and Applications*, vol. 2013, article 14, 2013.