Research Article

Quorum Sensing Activity of *Mesorhizobium* sp. F7 Isolated from Potable Water

Pei-Ling Yong and Kok-Gan Chan

Division of Genetics and Molecular Biology, Institute of Biological Sciences, Faculty of Science, University of Malaya,
50603 Kuala Lumpur, Malaysia

Correspondence should be addressed to Kok-Gan Chan; kokgan@um.edu.my

Received 19 April 2014; Revised 11 July 2014; Accepted 21 July 2014; Published 6 August 2014

Academic Editor: Christian Staehelin

Copyright © 2014 P.-L. Yong and K.-G. Chan. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We isolated a bacterial isolate (F7) from potable water. The strain was identified as *Mesorhizobium* sp. by 16S rDNA gene phylogenetic analysis and screened for *N*-acyl homoserine lactone (AHL) production by an AHL biosensor. The AHL profile of the isolate was further analyzed using high resolution triple quadrupole liquid chromatography mass spectrometry (LC/MS) which confirmed the production of multiple AHLs, namely, *N*-3-oxo-octanoyl-L-homoserine lactone (3-oxo-C8-HSL) and *N*-3-oxo-decanoyl-L-homoserine lactone (3-oxo-C10-HSL). These findings will open the perspective to study the function of these AHLs in plant-microbe interactions.

1. Introduction

Bacterial cell-cell communication, also termed quorum sensing (QS), occurs when the cell density has reached a threshold level that could regulate gene expressions via signalling molecules [1, 2]. Acyl-homoserine lactones (AHLs) molecules which are responsible for QS activity are synthesized by AHL synthase (LuxI homologue) whereby AHL will bind to its cognate receptor (LuxR homologue) to form a AHL-LuxR complex which in turn modulates a battery of QS-mediated gene expression [3, 4]. Upon sensing the AHLs, the Proteobacteria population works in synchrony to generate a significant impact on the bacterial physiological activities including bioluminescence, antibiotic production, plasmid conjugal transfer, and synthesis of exoenzyme virulence factors in plant and animal pathogens [5, 6]. Structurally, AHL consists of a conserved homoserine lactone moiety and the acyl chain length that ranges from 4 to 18 carbons [7].

Bacteria are commonly known to perform intra- or interspecies communication with the help of AHL. For example, a study on the interaction between *Serratia liquefaciens* MG1 and *Solanum lycopersicum* (tomato) gave first indications that AHL molecules of rhizosphere bacteria stimulate plant defense responses [8].

Rhizobia which include the genera *Rhizobium*, *Sinorhizobium*, *Mesorhizobium*, *Azorhizobium*, and *Bradyrhizobium* exhibit nitrogen fixing properties in root nodules of legumes which can only be achieved when bacteria carry large self-transmissible genetic elements, either plasmids or integrating conjugative elements that include symbiotic genes [9]. Plant growth is often limited by the availability of nitrogen, and hence, plant-rhizobia interactions are particularly important for sustainable and environment-friendly crop production without the use of nitrogen fertilizers [10].

A key mechanism in the symbiotic process is the involvement of QS [11]. Signaling and communication between the host and rhizobia are required for successful symbiotic interactions [12]. The various functions regulated by AHLs of *Rhizobium* and *Sinorhizobium* range from exopolysaccharide production [13], rhizosphere-related gene expression [14], and the conjugal transfer of pSym plasmids [15]. QS is also an important mechanism in other plant growth-promoting rhizobacteria [16].

AHLs have been identified in various rhizobial strains (Table 1). For example, a marine *Mesorhizobium* sp. was isolated that produced novel long-chain *N*-acyl-L-homoserine lactones, namely, 5-*cis*-3-oxo-C12-HSL and 5-*cis*-C12-HSL. It was shown that both these compounds were able to restore

TABLE 1: AHL production in rhizobia.

AHL	Representative organisms
3-oxo-C8-HSL	<i>Rhizobium</i> sp. strain NGR234, <i>Rhizobium leguminosarum</i> , <i>Sinorhizobium meliloti</i> , and <i>Rhizobium etli</i> CFN42 [15, 17–21]
3-OH-C8-HSL	<i>R. leguminosarum</i> , <i>S. meliloti</i> , and <i>R. etli</i> CFN42 [17, 19, 22]
C8-HSL	<i>R. leguminosarum</i> and <i>S. meliloti</i> [14, 17, 21–24]
3-OH-C14-HSL	<i>R. leguminosarum</i> [25]
C16-HSL	<i>S. meliloti</i> [19]

protease and pyoverdine production of an AHL-deficient *Pseudomonas aeruginosa* PAO1 *lasI rhII* double mutant, suggesting the AHLs could be used for intergenus signaling [26]. In other previous studies, when C4-HSL, C6-HSL, 3-oxo-C6-HSL, and 3-oxo-C8-HSL are applied to the plant, these AHLs promoted the growth of *Arabidopsis* [27–29] whereas 3-oxo-C10-HSL induced the formation of adventitious roots in mung beans [30]. On the other hand, 3-oxo-C14-HSL and 3-OH-C14-HSL induced resistance in *Arabidopsis* and barley plants towards biotrophic and hemibiotrophic pathogens [31].

In another study on *Rhizobium* sp. strain NGR234, it was found that the production of 3-oxo-C8-HSL led to activation of the transcriptional regulator TraR which significantly decreased the growth rate of NGR234 [15]. Moreover, it was shown that the regulatory gene TraI of *Rhizobium etli* CFN42 controls synthesis of 3-oxo-C8-HSL and conjugative plasmid transfer [17].

Here, we have identified the AHLs of *Mesorhizobium* sp. F7 which was isolated from potable water. We showed that this strain produces two AHLs, 3-oxo-C8-HSL and 3-oxo-C10-HSL.

2. Experimental Section

2.1. Sample Collection and Processing. A water sample was collected from domestic filtered water with nonwoven fabric and powdered activated carbon features installed in Petaling Jaya, Selangor (Malaysia). The tap was left open to flow for a few minutes before the sample was collected in a sterile plastic tube. The water sample was processed within an hour of sample collection. An aliquot of the water sample (100 μ L) was plated on Difco Reasoner's 2A agar (0.5 g/L proteose; 0.5 g/L casamino acids; 0.5 g/L yeast extract; 0.5 g/L dextrose; 0.5 g/L soluble starch; 0.3 g/L dipotassium phosphate; 0.05 g/L magnesium sulfate; 0.3 g/L sodium pyruvate) and incubated at 37°C under aerobic growth condition for 3 days. The observable different morphologies of the bacteria colonies were isolated and screened for AHL production using cross-streaking with *Chromobacterium violaceum* CV026 [32].

2.2. Isolation and Characterization of Isolate F7. The bacterial isolate F7 was found to exhibit QS properties among all isolates through an AHL biosensor screen (*Chromobacterium violaceum* CV026). The genomic DNA was extracted using a MasterPure™ DNA Purification Kit (EPICENTRE Inc., Madison, WI, USA). The isolate F7 was later characterized

by analyzing its 16S rRNA gene. The 16S rRNA was PCR amplified using 27F forward primer (5'-AGAGTTTGA-TCMTGGCTCAG-3'), 515F forward primer (5'-GTGCCA-GCMGCCGCGGTAA-3'), and 1525R reverse primer (5'-AAGGAGGTGWTCCARCC-3') using a PCR mix (Promega Kit, Madison, WI, USA). The PCR amplification that was carried out consists of an initial denaturation at 94°C for 3 min, followed by 30 repeated cycles at 94°C for 30 s of denaturing, 60°C for 30 s of annealing, and 72°C for 1 min 30 s of extension, and a final extension of 72°C for 7 min. Product sequence alignment was done using GenBank Blastn database and phylogenetic analysis was done using molecular evolutionary genetic analysis (MEGA) version 5.2 [33].

2.3. AHL Extraction. A single pure colony of isolate F7 was cultured overnight in R2 broth [34] buffered to pH 6.5, with 3-(*N*-morpholino) propanesulfonic acid (MOPS, 50 mM, pH 6.5) at 37°C with shaking (220 rpm). The supernatant was extracted twice with 100 mL of acidified (0.1% v/v glacial acetic acid) ethyl acetate [35]. The AHL extract was dried and stored at –20°C prior to further analysis.

2.4. Measurement of Bioluminescence. Preliminary screening of AHLs production by isolate F7 was done using an AHL biosensor (*Escherichia coli* [pSB401]). *E. coli* [pSB401] harbors *lux* from the pSB401 plasmid which will produce bioluminescence activity when exogenous short chain AHLs are supplied [36]. Cell density bioluminescence measurements were done using an Infinite M200 luminometer-spectrophotometer (Tecan, Männedorf, Switzerland). To every well of a 96-well optical bottom microtitre plate, an aliquot of 200 μ L diluted (1 : 100) *E. coli* [pSB401] overnight culture in LB broth supplemented with tetracycline (20 μ g/mL) and 1 μ L extracted AHL were added [37]. Acetonitrile and synthetic 3-oxo-C6-HSL (250 pg/ μ L) were used as the negative and positive standards, respectively. Results were indicated as relative light units/optical density (RLU/OD_{495 nm}) against incubation time (hour) [38].

2.5. Identification of AHL by High Resolution Tandem Liquid Chromatography Quadrupole Mass Spectrometry (LC-MS/MS). LC-MS/MS was performed using Agilent 6490 Triple Quadrupole LC/MS system (Agilent Technologies Inc., Santa Clara, CA, USA) and all experimental conditions were performed essentially as reported previously. The settings for the Agilent Mass Hunter software for MS spectra analysis were applied as described [39, 40].

FIGURE 1: Phylogenetic tree of isolate F7.

FIGURE 2: Detection of short chain AHLs production by *Mesorhizobium* sp.

3. Results and Discussion

3.1. Identification of a Bacterial Isolate from Potable Water. Isolate F7 was identified by analyzing its 16S rRNA gene nucleotide sequence [41–43]. A phylogenetic tree was constructed using Mega 5.2 software to align it with other rRNA sequences obtained from GenBank. A total of 988 unambiguously aligned nucleotides were analysed using the Neighbor-Joining method. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test is shown next to the branches [42]. The bar represents evolutionary distance as 0.002 change per nucleotide. Based on the phylogenetic tree obtained from the 16S rRNA sequencing (Figure 1), F7 was identified as *Mesorhizobium* sp.

3.2. Production of AHL by *Mesorhizobium* sp. F7. Isolate F7 was screened for its production of short chain AHL molecules by using luminometer-spectrophotometer, where the activation of bioluminescence of the biosensor *E. coli* [pSB401] was observed (Figure 2). Bioluminescence measurement was done for 24 h, 37°C. Cells were grown in the presence of AHL extracted from culture supernatant of *Mesorhizobium* sp. and synthetic 3-oxo-C6-HSL and acetonitrile were used as positive and negative controls, respectively.

Data are presented as means of $\pm$ SEM values of triplicate experiments.

To identify the AHLs, triple quadrupole LC/MS analysis was used. The MS results of the material from culture supernatants of *Mesorhizobium* sp. F7 are presented in Figure 3.

FIGURE 3: Mass spectrometry analysis of 3-oxo-C8-HSL: m/z value 242.000; retention time: 2.702 min; abundance: 8394.42 and abundance %: 100, and (b) 3-oxo-C10-HSL: m/z value: 270.200; retention time: 5.369 min; abundance: 3689.5 and abundance %: 100.

The data provide evidence for the presence of both short and long chain AHL molecules, namely, 3-oxo-C8-HSL (m/z 242.0000) and 3-oxo-C10-HSL (m/z 270.0000).

It was shown in other reports that the biological activity of AHLs applied to plants and the plant response is dependent on the length of lipid chains of AHLs. For instance, 3-oxo-C10-HSL was shown to be capable of inducing adventitious roots in mung bean [30] whereas 3-oxo-C8-HSL was active on *Arabidopsis* seedlings where it affected expression of 53 proteins related to plant primary metabolism, energy status, cytoskeleton, and defense [44]. Hence, we speculate that the AHLs produced by *Mesorhizobium* sp. F7 possess similar functions. Future experiments are required to test whether rhizobial strains producing specific AHLs can be used as bio-control bacteria to stimulate plant defense reactions against plant pathogens.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Authors' Contribution

Kok-Gan Chan conceived the idea and secured funding for the work. Pei-Ling Yong carried out the research, and both

authors were involved in data analysis and writing of the paper. All authors read and approved the paper.

Acknowledgment

This work was supported by the University of Malaya for High Impact Research Grant (UM.C/625/1/HIR/MOHE/CHAN/01, Grant no. A-000001-50001) awarded to Kok-Gan Chan.

References

- [1] C. Fuqua, S. C. Winans, and E. P. Greenberg, "Census and consensus in bacterial ecosystems: the LuxR-LuxI family of quorum-sensing transcriptional regulators," *Annual Review of Microbiology*, vol. 50, pp. 727–751, 1996.
- [2] N. D. Robson, A. R. J. Cox, S. J. McGowan, B. W. Bycroft, and G. P. C. Salmond, "Bacterial N-acyl-homoserine-lactone-dependent signalling and its potential biotechnological applications," *Trends in Biotechnology*, vol. 15, no. 11, pp. 458–464, 1997.
- [3] S. Swift, J. P. Throup, P. Williams, G. P. C. Salmond, and G. S. A. B. Stewart, "Quorum sensing: a population-density component in the determination of bacterial phenotype," *Trends in Biochemical Sciences*, vol. 21, no. 6, pp. 214–219, 1996.

- [4] S. Swift, J. Allan Downie, N. A. Whitehead, A. M. L. Barnard, G. P. C. Salmond, and P. Williams, "Quorum sensing as a population-density-dependent determinant of bacterial physiology," *Advances in Microbial Physiology*, vol. 45, pp. 199–270, 2001.
- [5] G. P. C. Salmond, B. W. Bycroft, G. S. A. B. Stewart, and P. Williams, "The bacterial 'enigma': cracking the code of cell-cell communication," *Molecular Microbiology*, vol. 16, no. 4, pp. 615–624, 1995.
- [6] A. M. Hardman, G. S. A. B. Stewart, and P. Williams, "Quorum sensing and the cell-cell communication dependent regulation of gene expression in pathogenic and non-pathogenic bacteria," *Antonie van Leeuwenhoek*, vol. 74, no. 4, pp. 199–210, 1998.
- [7] Y.-H. Dong, J.-L. Xu, X.-C. Li, and L.-H. Zhang, "AiiA, an enzyme that inactivates the acylhomoserine lactone quorum-sensing signal and attenuates the virulence of *Erwinia carotovora*," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 97, no. 7, pp. 3526–3531, 2000.
- [8] R. Schuëgger, A. Ihring, S. Gantner et al., "Induction of systemic resistance in tomato by N-acyl-L-homoserine lactone-producing rhizosphere bacteria," *Plant, Cell & Environment*, vol. 29, no. 5, pp. 909–918, 2006.
- [9] L. Passador, J. M. Cook, M. J. Gambello, L. Rust, and B. H. Iglewski, "Expression of *Pseudomonas aeruginosa* virulence genes requires cell-to-cell communication," *Science*, vol. 260, no. 5111, pp. 1127–1130, 1993.
- [10] C. Freiberg, R. Fellay, A. Bairoch, W. J. Broughton, A. Rosenthal, and X. Perret, "Molecular basis of symbiosis between *Rhizobium* and legumes," *Nature*, vol. 387, no. 6631, pp. 394–401, 1997.
- [11] J. E. González and M. M. Marketon, "Quorum sensing in nitrogen-fixing rhizobia," *Microbiology and Molecular Biology Reviews*, vol. 67, no. 4, pp. 574–592, 2003.
- [12] M. Schultze and A. Kondorosi, "Regulation of symbiotic root nodule development," *Annual Review of Genetics*, vol. 32, pp. 33–57, 1998.
- [13] M. M. Marketon, S. A. Glenn, A. Eberhard, and J. E. González, "Quorum sensing controls exopolysaccharide production in *Sinorhizobium meliloti*," *Journal of Bacteriology*, vol. 185, no. 1, pp. 325–331, 2003.
- [14] B. Rodelas, J. K. Lithgow, F. Wisniewski-Dye et al., "Analysis of quorum-sensing-dependent control of rhizosphere-expressed (*rhi*) genes in *Rhizobium leguminosarum* bv. *viciae*," *Journal of Bacteriology*, vol. 181, no. 12, pp. 3816–3823, 1999.
- [15] X. He, W. Chang, D. L. Pierce, L. O. Seib, J. Wagner, and C. Fuqua, "Quorum sensing in *Rhizobium* sp. strain NGR234 regulates conjugal transfer (*tra*) gene expression and influences growth rate," *Journal of Bacteriology*, vol. 185, no. 3, pp. 809–822, 2003.
- [16] P. N. Bhattacharyya and D. K. Jha, "Plant growth-promoting rhizobacteria (PGPR): emergence in agriculture," *World Journal of Microbiology and Biotechnology*, vol. 28, no. 4, pp. 1327–1350, 2012.
- [17] C. Tun-Garrido, P. Bustos, V. González, and S. Brom, "Conjugative transfer of p42a from *Rhizobium etli* CFN42, which is required for mobilization of the symbiotic plasmid, is regulated by quorum sensing," *Journal of Bacteriology*, vol. 185, no. 5, pp. 1681–1692, 2003.
- [18] I. Hwang, P. Li, L. Zhang et al., "TraI, a luxI homologue, is responsible for production of conjugation factor, the Ti plasmid N-acylhomoserine lactone autoinducer," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 91, no. 11, pp. 4639–4643, 1994.
- [19] M. M. Marketon and J. E. González, "Identification of two quorum-sensing systems in *Sinorhizobium meliloti*," *Journal of Bacteriology*, vol. 184, no. 13, pp. 3466–3475, 2002.
- [20] K. R. Piper, S. Beck Von Bodman, and S. K. Farrand, "Conjugation factor of *Agrobacterium tumefaciens* regulates Ti plasmid transfer by autoinduction," *Nature*, vol. 362, no. 6419, pp. 448–450, 1993.
- [21] A. Wilkinson, V. Danino, F. Wisniewski-Dyé, J. K. Lithgow, and J. A. Downie, "N-acyl-homoserine lactone inhibition of rhizobial growth is mediated by two quorum-sensing genes that regulate plasmid transfer," *Journal of Bacteriology*, vol. 184, no. 16, pp. 4510–4519, 2002.
- [22] F. Wisniewski-Dyé, J. Jones, S. R. Chhabra, and J. A. Downie, "*raiIR* genes are part of a quorum-sensing network controlled by *cinI* and *cinR* in *Rhizobium leguminosarum*," *Journal of Bacteriology*, vol. 184, no. 6, pp. 1597–1606, 2002.
- [23] M. T. Cubo, A. Economou, G. Murphy, A. W. B. Johnston, and J. A. Downie, "Molecular characterization and regulation of the rhizosphere-expressed genes *rhiABCR* that can influence nodulation by *Rhizobium leguminosarum* biovar *viciae*," *Journal of Bacteriology*, vol. 174, no. 12, pp. 4026–4035, 1992.
- [24] M. M. Marketon, M. R. Gronquist, A. Eberhard, and J. E. González, "Characterization of the *Sinorhizobium meliloti* *sinR/sinI* locus and the production of novel N-acyl homoserine lactones," *Journal of Bacteriology*, vol. 184, no. 20, pp. 5686–5695, 2002.
- [25] J. K. Lithgow, A. Wilkinson, A. Hardman et al., "The regulatory locus *cinRI* in *Rhizobium leguminosarum* controls a network of quorum-sensing loci," *Molecular Microbiology*, vol. 37, no. 1, pp. 81–97, 2000.
- [26] A. Krick, S. Kehraus, L. Eberl et al., "A marine *Mesorhizobium* sp. produces structurally novel long-chain N-acyl-L-homoserine lactones," *Applied and Environmental Microbiology*, vol. 73, no. 11, pp. 3587–3594, 2007.
- [27] U. Von Rad, I. Klein, P. I. Dobrev et al., "Response of *Arabidopsis thaliana* to N-hexanoyl-DL-homoserine-lactone, a bacterial quorum sensing molecule produced in the rhizosphere," *Planta*, vol. 229, no. 1, pp. 73–85, 2008.
- [28] F. Liu, Z. Bian, Z. Jia, Q. Zhao, and S. Song, "The GCR1 and GPA1 participate in promotion of *Arabidopsis* primary root elongation induced by N-Acyl-homoserine lactones, the bacterial quorum-sensing signals," *Molecular Plant-Microbe Interactions*, vol. 25, no. 5, pp. 677–683, 2012.
- [29] S. T. Schenk, E. Stein, K. Kogel, and A. Schikora, "Arabidopsis growth and defense are modulated by bacterial quorum sensing molecules," *Plant Signaling & Behavior*, vol. 7, no. 2, pp. 178–181, 2012.
- [30] X. Bai, C. D. Todd, R. Desikan, Y. Yang, and X. Hu, "N-3-oxodecanoyl-L-homoserine-lactone activates auxin-induced adventitious root formation via hydrogen peroxide- and nitric oxide-dependent cyclic GMP signaling in mung bean," *Plant Physiology*, vol. 158, no. 2, pp. 725–736, 2012.
- [31] A. Schikora, S. T. Schenk, E. Stein, A. Molitor, A. Zuccaro, and K. Kogel, "N-acyl-homoserine lactone confers resistance toward biotrophic and hemibiotrophic pathogens via altered activation of AtMPK6," *Plant Physiology*, vol. 157, no. 3, pp. 1407–1418, 2011.
- [32] K. H. McClean, M. K. Winson, L. Fish et al., "Quorum sensing and *Chromobacterium violaceum*: exploitation of violacein production and inhibition for the detection of N-acylhomoserine lactones," *Microbiology*, vol. 143, no. 12, pp. 3703–3711, 1997.
- [33] C. Wong, W. Yin, Y. Choo, C. Sam, C. Koh, and K. Chan, "Coexistence of quorum-quenching and quorum-sensing in

- tropical marine *Pseudomonas aeruginosa* strain MW3A,” *World Journal of Microbiology and Biotechnology*, vol. 28, no. 2, pp. 453–461, 2012.
- [34] N. Y. M. Yunos, W. S. Tan, C. L. Koh et al., “*Pseudomonas cremoricolorata* strain ND07 produces *N*-acyl homoserine lactones as quorum sensing molecules,” *Sensors*, vol. 14, no. 7, pp. 11595–11604, 2014.
- [35] C. A. Ortori, J. Dubern, S. R. Chhabra et al., “Simultaneous quantitative profiling of *N*-acyl-L-homoserine lactone and 2-alkyl-4(1*H*)-quinolone families of quorum-sensing signaling molecules using LC-MS/MS,” *Analytical and Bioanalytical Chemistry*, vol. 399, no. 2, pp. 839–850, 2011.
- [36] M. K. Winson, S. Swift, L. Fish et al., “Construction and analysis of luxCDABE-based plasmid sensors for investigating *N*-acyl homoserine lactone-mediated quorum sensing,” *FEMS Microbiology Letters*, vol. 163, no. 2, pp. 185–192, 1998.
- [37] W. Yin, K. Purmal, S. Chin et al., “*N*-Acyl homoserine lactone production by *Klebsiella pneumonia* isolated from human tongue surface,” *Sensors*, vol. 12, no. 3, pp. 3472–3483, 2012.
- [38] L. Y. Tan, W. Yin, and K. Chan, “Silencing quorum sensing through extracts of melicope lunu-ankenda,” *Sensors*, vol. 12, no. 4, pp. 4339–4351, 2012.
- [39] R. E. Han-Jen, W. F. Yin, and K. G. Chan, “*Pandoraea* sp. RB-44, a novel quorum sensing soil bacterium,” *Sensors*, vol. 13, no. 10, pp. 14121–14132, 2013.
- [40] J. W. Chen, C. L. Koh, C. K. Sam, W. F. Yin, and K. G. Chan, “Short chain *N*-acyl homoserine lactone production by soil isolate *Burkholderia* sp. strain A9,” *Sensors*, vol. 13, no. 10, pp. 13217–13227, 2013.
- [41] N. Saitou and M. Nei, “The neighbor-joining method: a new method for reconstructing phylogenetic trees,” *Molecular Biology and Evolution*, vol. 4, no. 4, pp. 406–425, 1987.
- [42] J. Felsenstein, “Confidence limits on phylogenies: an approach using the bootstrap,” *Evolution*, vol. 39, no. 4, pp. 783–791, 1985.
- [43] K. Tamura, M. Nei, and S. Kumar, “Prospects for inferring very large phylogenies by using the neighbor-joining method,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 101, no. 30, pp. 11030–11035, 2004.
- [44] C. Miao, F. Liu, Q. Zhao, Z. Jia, and S. Song, “A proteomic analysis of *Arabidopsis thaliana* seedling responses to 3-oxo-octanoyl-homoserine lactone, a bacterial quorum-sensing signal,” *Biochemical and Biophysical Research Communications*, vol. 427, no. 2, pp. 293–298, 2012.