Research Article

Hydrochemistry of the Hot Springs in Western Sichuan Province Related to the Wenchuan M_s 8.0 Earthquake

Zhi Chen,^{1,2} Jianguo Du,² Xiaocheng Zhou,² Li Yi,² Lei Liu,² Chao Xie,²
Yueju Cui,² and Ying Li²

¹ School of Earth and Space Sciences, University of Science and Technology of China, Hefei 230026, China

² CEA Key Laboratory of Earthquake Prediction (Institute of Earthquake Science), China Earthquake Administration, Beijing 100036, China

Correspondence should be addressed to Jianguo Du; jianguodu@hotmail.com

Received 1 January 2014; Accepted 5 February 2014; Published 5 May 2014

Academic Editors: J. Glodny, P. C. McGuire, and L. Tosi

Copyright © 2014 Zhi Chen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Hydrogeochemistry of 32 hot springs in the western Sichuan Province after the Wenchuan M_s 8.0 earthquake was investigated by analyzing the concentrations of cation and anion and the isotopic compositions of hydrogen and oxygen. The water samples of the hot springs were collected four times from June 2008 to April 2010. Hydrogeochemical data indicated the water samples can be classified into 9 chemical types. Values of δD and $\delta^{18}O$ indicated that the spring waters were mainly derived from meteoric precipitation and affected by water-rock interaction and mixture of deep fluids. Concentrations of K^+ and SO_4^{2-} of the samples from the Kangding district exhibited evident increases before the Wenchuan earthquake, indicating more supplement of deep fluids under the increase of tectonic stress. The chemical and isotopic variations of the water samples from the area closer to the epicenter area can be attributed to variation of regional stress field when the aftershock activities became weak.

1. Introduction

Great earthquakes usually associate with the physical-chemical variations of groundwater and hot springs. The short-term hydrogeochemical precursors for earthquake [1–4], co-seismic response of hydrochemistry [5, 6] and post-seismic geochemical and isotopic changes of hot springs have been reported [7–12].

The observed geochemical anomalies related to earthquakes are usually attributed to the alteration of groundwater in the circulating system under the action of increasing crustal stress before and after earthquakes [6, 7, 13, 14]. For understanding the hydrogeochemical anomalies related to earthquakes, some genetic mechanism models have been proposed, such as increased solubility of rocks under increased pressure and release of ions from rocks into water [15], pore collapse with fluids expulsion [8], the water-rock interactions at the enhanced reactive surfaces [16], aquifers rupture with mixing of different fluids [6] and expulsion of deep fluids by tectonic pumping [1].

Hydrogeochemical survey of hot springs in western Sichuan before the M_s 8.0 Wenchuan earthquake mainly focused on the origin of waters, the pollutions, the chemical classifications and the heat reservoirs in the sites surrounding Kangding county [17–25].

The M_s 8.0 earthquake occurred in Wenchuan county, Sichuan province, Southwest China on 12 May, 2008, following by hundreds of aftershocks with magnitudes higher than 3.0.

In order to investigate the hydrogeochemical characteristics of the springs related to the seismic activities, the hydrogeochemical survey of 32 hot springs in the western Sichuan were performed four times from June 2008 to April 2010.

2. Geological Setting

The investigated area is located at the eastern margin of the Tibetan Plateau, in which there are four major fault zones

FIGURE 1: Schematic map of geology in western Sichuan Province, showing the sample locations, and earthquake epicenters, the insert figure is time-magnitude plot for aftershocks with $M_s \geq 4$, red lines are active faults.

named as Minjiang fault (MJF), Longmenshan fault (LMSF), Xianshuihe fault (XSHF) and Anninghe fault (ANHF) zones (Figure 1) where earthquakes frequently occurred [26]. Triassic littoral-neritic clastic rocks interbedded with carbonate and intrusive granite are exposed in the west of the LMSF and ANHF zones, but the strata from Late Paleozoic to Cenozoic are widespread exposed in the LMSF and ANHF zones and on the east of the fault zones, and granite is found in the intersection of the LMSF, XSHF and ANHF zones [27–30]. The fault zones act as the important passage for upward migration of thermal fluids from the deep earth, which is indicated by distribution of many hot springs in the MJF, LMSF, XSHF and ANHF zones (Figure 1). Historically, a number of great earthquakes ($M_S > 7.0$) have occurred in the investigated area since 1800. For instance, the 1850 M_S 7.5 Xichang earthquake occurred in the ANHF zone, two M_S 7.2 earthquakes of 16 and 23 August 1976 in the MJF zone, the M_S 7.5 one of 25 August 1933 in the LMSF zone, and the M_S 7.5 one of 1955 and the M_S 7.9 one of 1973 in the XSHF zone [12]. The M_S 8.0 Wenchuan earthquake resulted in a 285 km surface rupture zone along the pre-existing Yingxiu-Beichuan, Guanxian-Anxian and Qingchuan faults, with the maximum vertical surface offset of about 6.2 m [31].

3. Methods

Water samples were repeatedly collected four times (in June and October 2008, June 2009, and April 2010) at 32 sites of spas, wells, and springs (Figure 1) in the MJF, LMSF, XSHF, and ANHF zones. The samples were sealed and stored in 500 mL glass bottles. The values of temperature were measured with a digital thermometer with an error of $\pm 1\%$ in the field. Isotopic compositions of δD and $\delta^{18}O$ were measured with a Picarro LI102 mass spectrometer in the Laboratory of Gas Geochemistry, Institute of Geology and Geophysics, Chinese Academy of Sciences in Lanzhou, and the errors are 0.5‰ for δD and 0.1‰ for $\delta^{18}O$, respectively. The concentrations of cations (K^+ , Na^+ , Mg^{2+} , and Ca^{2+}) and anions (F^- , Cl^- , Br^- , NO_3^- , and SO_4^{2-}) were determined with the Dionex ICS-900 ion chromatography (reproducibility within $\pm 2\%$) that is installed in the Institute of Earthquake Science, China Earthquake Administration. The CO_3^{2-} and HCO_3^- concentrations were measured by the standard titration procedures with a ZDJ-100 potentiometric titrator (reproducibility within $\pm 2\%$). The ion balance (ib) was calculated according to the following equation [11]:

$$ib (\%) = \frac{\sum \text{cations} - \sum \text{anions}}{(\sum \text{cations} + \sum \text{anions}) \times 0.5} \times 100. \quad (1)$$

4. Results

The physicochemical parameters of the 32 springs were listed in Table 1. No water samples were collected from the springs of nos. 2 and 24–26 in June 2008 because the springs were damaged by the Wenchuan earthquake. No samples were collected from the springs nos. 8 and 29 in April 2010 and the spring no. 28 in June 2009 and April 2010 because of the postearthquake reconstruction. The ion balance values of

measured chemical data were calculated to be less than 5% (Table 1).

4.1. Physicochemical Parameters of the Springs in the LMSF Zone. The water samples from in the LMSF zone have $\delta^{18}O$ and δD values between -16.34‰ and -6.18‰ , -116.65‰ and -61.66‰ , respectively. The temperatures and TDS ranged from 10.5 to 58.0°C and 151.69 to 1569.81 mg/L. The concentrations of Na^+ , Ca^{2+} and Mg^{2+} varied from 29.82 to 202.49 mg/L, 4.74 to 252.78 mg/L, and 0.80 to 128.73 mg/L, respectively. The concentrations of Cl^- , SO_4^{2-} , and HCO_3^- varied from 2.31 to 64.74 mg/L, 21.33 to 878.08 mg/L, and 14.63 to 310.70 mg/L respectively (Table 1).

4.2. Physicochemical Parameters of the Springs in the MJF Zone. Fifteen water samples were collected from the sites nos. 6–9 in the MJF zone, of which the $\delta^{18}O$ and δD were between -15.34‰ and -13.20‰ , -112.23‰ and -102.20‰ , the values of temperatures and TDS ranged from 8.8 to 21.7°C and 505.34 to 1657.61 mg/L. The concentrations of Na^+ , Ca^{2+} , and Mg^{2+} ranged from 8.75 to 25.23 mg/L, 49.54 to 183.57 mg/L, and 23.23 to 183.57 mg/L, and the concentrations of Cl^- , SO_4^{2-} , and HCO_3^- from 0.93 to 10.78 mg/L, 3.56 to 17.78 mg/L, and 313.61 to 1324.27 mg/L separately (Table 1).

4.3. Physicochemical Parameters of the Springs in the XSHF Zone. Sixty-two water samples were collected from the sites nos. 10–25 in the XSHF zone, of which $\delta^{18}O$ and δD varied from -18.84‰ to -10.04‰ and -141.98‰ to -78.77‰ . The values of temperatures and TDS ranged from 27.8 to 83.0°C and 427.53 to 2159.43 mg/L, respectively. The concentrations of Na^+ , Ca^{2+} , and Mg^{2+} ranged from 35.14 to 679.23 mg/L, 4.38 to 50.10 mg/L, and 0.12 to 45.71 mg/L, respectively. The concentrations of Cl^- , SO_4^{2-} , and HCO_3^- ranged from 2.35 to 336.16 mg/L, 6.56 to 161.87 mg/L, and 62.48 to 1894.64 mg/L, respectively (Table 1).

4.4. Physicochemical Parameters of the Springs in the ANHF Zone. Twenty-four water samples were collected from the sites nos. 26–32 in the ANHF zone, of which the $\delta^{18}O$ and δD were between -11.64‰ and -96.10‰ , -110.47‰ and -102.20‰ . The temperatures of the spring waters were between 21.1 and 56.9°C, the TDS were between 206.70 and 1212.86 mg/L. The concentrations of Na^+ , Ca^{2+} , and Mg^{2+} ranged from 50.17 to 213.50 mg/L, 2.00 to 81.48 mg/L, and 0.00 to 29.43 mg/L, and the concentrations of Cl^- , SO_4^{2-} , and HCO_3^- from 1.91 to 136.74 mg/L, 10.94 to 257.26 mg/L, and 81.95 to 449.06 mg/L, respectively (Table 1).

5. Discussion

5.1. Chemical Types of the Waters and Their Origins

5.1.1. The Chemical Types. The triangular diagrams in Figure 2 illustrated the proportions of the anionic (Cl^- , SO_4^{2-} , and HCO_3^-) and cationic ($K^+ + Na^+$, Mg^{2+} , and Ca^{2+}) concentrations of the water samples from the 32

TABLE 1: Physicochemical parameters of the 32 hot spring waters.

Number	Site	L (E)	B (N)	Date	T °C	TDS mg/L	K ⁺ mg/L	Na ⁺ mg/L	Ca ²⁺ mg/L	Mg ²⁺ mg/L	Cl ⁻ mg/L	SO ₄ ²⁻ mg/L	CO ₃ ²⁻ mg/L	HCO ₃ ⁻ mg/L	ib %	δ ¹⁸ O ‰	δD ‰	Chemical type
The LMSF zone																		
1	Sangzao Spring	104.35	31.61	11.06.2008	19.0	557.43	2.67	64.78	73.27	12.27	64.74	108.24	0.00	229.38	-0.01	-6.18	-61.66	CaNa- HCO ₃ SO ₄
				01.11.2008	19.1	394.07	2.68	41.92	55.57	12.90	30.39	72.65	0.00	175.98	0.02	-9.79	-69.63	CaNa- HCO ₃ SO ₄
				01.07.2009	19.7	367.75	2.60	40.26	50.64	12.04	30.57	71.52	0.00	158.27	0.02	-9.87	-68.40	CaNa- HCO ₃ SO ₄
				22.04.2010	19.5	361.94	2.61	37.63	45.01	12.88	30.45	65.72	0.00	166.74	0.00	-9.96	-70.34	CaNa- HCO ₃ SO ₄
2	Wenchuan Well	103.77	31.63	10.06.2008	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
				21.10.2008	15.4	1569.81	4.21	71.65	182.01	128.73	47.39	878.08	0.00	248.12	-0.01	-8.37	-72.01	MgCa-SO ₄
				29.06.2009	15.5	1128.39	4.35	61.72	151.53	80.38	29.51	720.73	0.00	70.05	0.00	-10.14	-73.78	MgCa-SO ₄
				20.04.2010	10.8	1033.14	4.55	54.99	137.22	70.43	30.47	687.83	0.00	46.65	-0.01	-10.02	-73.11	MgCa-SO ₄
3	Lixian Well	103.59	31.48	09.06.2008	32.9	1579.54	5.64	35.95	252.78	103.76	8.15	862.32	0.00	310.70	0.00	-9.53	-80.88	CaMg-SO ₄
				22.10.2008	29.5	1195.08	5.00	37.29	182.66	78.59	7.82	812.50	0.00	68.47	-0.01	-11.75	-81.76	CaMg-SO ₄
				29.06.2009	27.1	1105.85	5.44	37.91	161.15	76.82	7.90	780.09	0.00	34.00	-0.01	-11.01	-82.13	CaMg-SO ₄
				20.04.2010	29.5	1025.28	5.72	36.15	146.64	69.97	7.48	741.96	0.00	14.63	-0.02	-11.62	-80.88	CaMg-SO ₄
4	Guergou Spring	103.17	31.46	11.06.2008	n.d.	349.33	2.33	91.11	5.24	0.80	2.58	21.76	0.00	223.52	0.00	-11.74	-116.81	Na-HCO ₃
				21.10.2008	58.0	246.71	2.23	60.76	4.74	0.83	2.45	21.33	0.00	150.62	-0.01	-14.42	-116.49	Na-HCO ₃
				28.06.2009	56.0	240.49	2.18	54.36	5.22	0.83	2.61	22.63	0.00	150.62	-0.03	-16.27	-116.99	Na-HCO ₃
				19.04.2010	55.8	151.69	2.19	29.82	4.76	0.82	2.31	22.16	0.00	88.13	-0.05	-16.34	-116.65	Na-HCO ₃
5	Jiuzhaigou Spring	103.00	31.52	11.06.2008	30.5	837.83	9.06	202.49	26.48	20.48	60.83	237.79	0.00	272.78	0.01	-14.01	-107.26	Na- HCO ₃ SO ₄
				21.10.2008	32.1	813.62	9.15	189.00	26.01	20.66	64.16	225.51	0.00	272.34	0.01	-13.59	-106.10	Na- HCO ₃ SO ₄
				28.06.2009	32.0	739.07	8.86	163.55	28.71	20.55	10.28	232.20	0.00	270.10	0.02	-14.57	-106.62	Na- HCO ₃ SO ₄
				19.04.2010	27.3	728.04	8.75	150.98	26.73	20.41	9.65	241.18	0.00	267.84	0.00	-13.95	-106.54	Na- HCO ₃ SO ₄
The MJF zone																		
6	Erdaohai Spring	103.32	32.73	08.06.2008	21.6	505.34	1.34	8.75	85.17	25.73	0.93	3.80	0.00	379.30	0.02	-13.72	-102.42	CaMg- HCO ₃
				22.10.2008	20.7	511.71	1.32	8.80	84.02	25.05	0.98	4.08	0.00	386.99	0.01	-14.39	-102.46	CaMg- HCO ₃
				29.06.2009	20.0	464.18	1.22	8.57	77.24	23.35	1.02	3.78	0.00	348.94	0.02	-14.57	-103.88	CaMg- HCO ₃
				21.04.2010	19.8	428.70	1.27	8.21	76.73	23.23	0.97	3.56	0.00	313.61	0.04	-14.39	-103.29	CaMg- HCO ₃

TABLE I: Continued.

Number	Site	L (E)	B (N)	Date	T °C	TDS mg/L	K ⁺ mg/L	Na ⁺ mg/L	Ca ²⁺ mg/L	Mg ²⁺ mg/L	Cl ⁻ mg/L	SO ₄ ²⁻ mg/L	CO ₃ ²⁻ mg/L	HCO ₃ ⁻ mg/L	ib %	δ ¹⁸ O ‰	δD ‰	Chemical type
7	Kakagou Spring	103.68	32.98	08.06.2008	18.7	914.21	5.11	25.23	160.21	31.45	2.05	10.98	0.00	678.66	0.01	-13.56	-101.05	Ca-HCO ₃
				23.10.2008	19.7	876.14	4.92	24.08	154.12	30.74	1.93	10.86	0.00	648.69	0.01	-13.37	-101.92	Ca-HCO ₃
				30.06.2009	21.7	873.30	4.84	23.93	142.67	28.67	2.07	10.83	0.00	660.16	-0.01	-14.04	-100.46	Ca-HCO ₃
8	Chuanpanqiao Spring	103.64	32.85	21.04.2010	19.7	815.42	4.79	23.55	134.00	31.65	1.84	10.33	0.00	608.89	0.01	-14.12	-101.80	Ca-HCO ₃
				08.06.2008	8.8	1657.61	6.36	21.82	97.82	182.62	10.78	12.54	0.00	1324.27	-0.01	-14.20	-112.00	Mg-HCO ₃
				23.10.2008	9.6	1598.09	6.52	21.88	103.64	183.57	10.56	12.72	0.00	1258.08	0.00	-14.30	-112.48	Mg-HCO ₃
				30.06.2009	10.7	1557.18	5.89	21.68	103.79	175.29	10.55	11.54	0.00	1228.08	0.00	-15.34	-112.23	Mg-HCO ₃
				21.04.2010	10.7	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	-14.95	-113.33	n.d.
9	Songpan Well	103.60	32.65	08.06.2008	9.0	647.31	1.64	14.05	55.71	65.01	2.34	17.78	0.00	490.23	0.01	-13.25	-101.54	MgCa-HCO ₃
				23.10.2008	9.9	648.33	1.60	14.12	52.12	65.27	2.17	16.32	0.00	496.23	0.00	-13.43	-102.82	MgCa-HCO ₃
				30.06.2009	9.3	610.38	1.57	13.93	49.54	65.43	2.25	16.24	0.00	461.42	0.02	-13.20	-102.20	MgCa-HCO ₃
10	Ganzi Well	100.00	31.62	21.04.2010	9.9	603.35	1.60	13.81	50.12	62.90	2.48	15.96	0.00	456.23	0.02	-13.77	-102.09	MgCa-HCO ₃
				The XSHF zone														
				12.06.2008	27.8	441.46	3.42	49.07	38.84	19.07	4.04	44.27	0.00	281.36	0.00	-15.93	-125.50	NaCa-HCO ₃
				24.10.2008	29.8	447.22	3.53	47.97	37.51	19.97	3.94	43.44	0.00	289.54	-0.01	-16.48	-125.21	NaCa-HCO ₃
				27.06.2009	29.6	416.83	3.43	47.91	37.39	19.13	4.06	43.34	0.00	260.25	0.01	-16.17	-125.79	NaCa-HCO ₃
11	Ganzi Spring	100.00	31.61	18.04.2010	29.2	472.24	3.33	50.76	38.72	19.97	3.80	40.17	0.00	314.28	-0.01	-16.15	-125.75	NaCa-HCO ₃
				12.06.2008	43.1	735.04	8.95	154.67	23.95	16.42	6.38	62.15	0.00	459.17	0.01	-16.33	-124.72	Na-HCO ₃
				24.10.2008	41.5	731.75	9.08	150.20	23.13	16.34	6.19	60.61	0.00	461.22	0.00	-16.28	-124.30	Na-HCO ₃
				27.06.2009	43.0	695.04	9.17	146.66	22.34	16.44	6.00	61.79	0.00	427.90	0.01	-16.45	-126.00	Na-HCO ₃
				18.04.2010	41.4	706.99	9.29	138.30	21.87	16.42	6.17	63.67	0.00	447.27	-0.01	-16.29	-124.46	Na-HCO ₃
12	Guyi Spring	100.80	31.27	13.06.2008	35.5	2218.75	12.68	60707	11.30	21.81	6.09	12.25	142.80	1399.60	0.01	-16.38	-130.98	Na-HCO ₃
				25.10.2008	31.3	2353.19	13.13	619.72	11.36	22.19	5.73	11.62	142.80	1524.39	0.00	-16.16	-131.17	Na-HCO ₃
				26.06.2009	33.2	2239.88	12.63	593.59	10.33	21.82	6.11	11.60	157.21	1424.29	-0.01	-16.97	-132.52	Na-HCO ₃
				18.04.2010	28.2	2366.42	12.63	619.91	10.01	21.31	5.95	12.35	108.56	1573.76	0.00	-16.56	-131.90	Na-HCO ₃

TABLE 1: Continued.

Number	Site	L (E)	B (N)	Date	T °C	TDS mg/L	K ⁺ mg/L	Na ⁺ mg/L	Ca ²⁺ mg/L	Mg ²⁺ mg/L	Cl ⁻ mg/L	SO ₄ ²⁻ mg/L	CO ₃ ²⁻ mg/L	HCO ₃ ⁻ mg/L	ib %	δ ¹⁸ O ‰	δD ‰	Chemical type
13	Longpugou Spring 1	101.24	30.98	13.06.2008	37.8	1180.32	716	310.06	11.00	8.23	13.83	6.83	0.00	820.21	0.01	-16.74	-137.69	Na-HCO ₃
				25.10.2008	37.2	1199.61	6.96	288.86	11.69	7.90	13.02	6.81	0.00	860.75	-0.01	-17.02	-136.73	Na-HCO ₃
				26.06.2009	39.2	1164.18	6.90	280.27	11.28	7.80	13.27	6.61	0.00	836.33	-0.01	-17.41	-137.71	Na-HCO ₃
				18.04.2010	36.5	1145.60	6.83	281.70	11.16	7.85	12.98	6.56	0.00	815.75	-0.01	-17.47	-138.90	Na-HCO ₃
14	Longpugou Spring 2	101.24	30.95	13.06.2008	44.5	1070.89	6.47	221.80	18.30	26.65	8.85	10.09	57.10	719.87	-0.02	-16.85	-134.42	Na-HCO ₃
				25.10.2008	42.5	927.46	6.93	205.03	18.90	26.11	8.28	9.78	0.00	650.41	0.02	-17.52	-134.80	Na-HCO ₃
				26.06.2009	42.5	900.81	6.67	198.93	17.43	25.73	8.39	9.33	0.00	634.33	0.02	-16.63	-133.90	Na-HCO ₃
				18.04.2010	43.2	919.25	6.48	176.77	18.84	25.93	8.53	9.63	0.00	671.47	-0.01	-17.61	-133.70	Na-HCO ₃
15	Bamei Spring	101.62	30.54	14.06.2008	55.2	1966.85	76.60	443.00	15.17	29.06	23.77	6.78	0.00	1368.29	0.01	-17.34	-135.61	Na-HCO ₃
				25.10.2008	55.6	1913.45	75.22	417.08	14.76	28.42	20.55	7.00	0.00	1346.62	0.00	-17.57	-134.51	Na-HCO ₃
				26.06.2009	56.0	2068.93	77.29	463.31	14.32	26.66	20.58	6.81	0.00	1456.10	0.00	-17.54	-135.81	Na-HCO ₃
				17.04.2010	55.6	2724.13	74.40	679.23	14.49	29.42	20.55	7.12	0.00	1894.64	0.02	-17.89	-134.41	Na-HCO ₃
16	Guanding Spring	101.96	29.98	15.06.2008	80.5	1306.05	41.23	396.79	13.16	13.21	336.16	51.96	11.34	440.26	0.02	-14.24	-117.01	Na- ClHCO ₃
				26.10.2008	80.0	1395.37	40.57	367.22	13.41	13.64	328.54	50.02	92.82	485.48	-0.03	-15.26	-118.02	Na- ClHCO ₃
				25.06.2009	83.0	1353.81	42.72	367.40	13.67	13.50	320.93	49.72	99.82	443.09	-0.02	-17.02	-127.74	Na- ClHCO ₃
				17.04.2010	83.0	1273.60	41.52	349.71	13.80	13.81	300.83	50.63	56.00	447.24	-0.01	-16.13	-123.72	Na- ClHCO ₃
17	Longtougou Spring	101.96	29.98	15.06.2008	70.8	2044.87	53.20	510.82	16.02	31.42	220.68	8.33	0.00	1200.36	0.01	-15.99	-128.69	Na-HCO ₃
				26.10.2008	70.2	2106.82	52.72	518.38	17.41	30.95	203.66	8.59	0.00	1270.92	0.00	-16.26	-127.78	Na-HCO ₃
				25.06.2009	73.1	2025.87	52.27	516.68	16.71	30.84	226.15	8.30	0.00	1171.25	0.01	-16.61	-128.57	Na-HCO ₃
				17.04.2010	70.2	2096.92	51.62	507.94	16.33	31.09	207.55	8.30	0.00	1270.92	0.00	-17.04	-130.94	Na-HCO ₃
18	Zheduotang Spring	101.86	30.01	15.06.2008	54.5	681.59	3.13	208.79	4.87	0.12	10.55	17.57	67.83	342.06	-0.01	-18.06	-138.80	Na-HCO ₃
				26.10.2008	53.8	705.52	3.14	219.83	4.38	0.12	10.57	17.31	85.68	338.08	-0.01	-18.21	-137.20	Na-HCO ₃
				25.06.2009	53.4	650.33	3.04	207.09	4.44	0.12	9.59	16.57	35.89	346.59	0.01	-18.64	-141.98	Na-HCO ₃
				17.04.2010	53.8	663.17	2.87	208.84	4.72	0.12	9.76	16.42	88.25	305.09	-0.02	-18.84	-139.88	Na-HCO ₃
19	Erdaociao Spring	101.95	30.09	15.06.2008	39.1	843.39	23.54	150.53	22.10	44.05	43.88	114.59	0.00	443.52	0.02	-14.50	-112.90	NaMg- HCO ₃
				26.10.2008	39.4	919.66	22.47	150.67	22.08	44.68	40.36	107.06	0.00	530.93	0.00	-14.54	-112.16	NaMg- HCO ₃
				26.06.2009	40.6	926.35	22.36	146.31	20.20	44.48	43.92	109.47	0.00	538.41	-0.02	-15.57	-115.14	NaMg- HCO ₃
				16.04.2010	40.6	935.65	23.67	157.07	20.93	43.76	40.97	111.59	0.00	535.44	0.00	-15.01	-115.35	NaMg- HCO ₃

TABLE I: Continued.

Number	Site	L (E)	B (N)	Date	T °C	TDS mg/L	K ⁺ mg/L	Na ⁺ mg/L	Ca ²⁺ mg/L	Mg ²⁺ mg/L	Cl ⁻ mg/L	SO ₄ ²⁻ mg/L	CO ₃ ²⁻ mg/L	HCO ₃ ⁻ mg/L	ib %	δ ¹⁸ O ‰	δD ‰	Chemical type	
20	Xinxing Spring	102.06	29.75	16.06.2008	45.6	1287.50	9.41	422.66	12.53	9.69	148.03	28.85	106.33	545.15	0.03	-15.90	-115.30	Na-HCO ₃	
				27.10.2008	44.1	1167.77	9.68	365.04	12.75	9.43	144.03	24.65	0.00	597.19	0.05	-15.53	-115.83	Na-HCO ₃	
				24.06.2009	42.8	1289.88	9.23	360.63	11.97	9.36	148.41	24.07	124.73	597.17	-0.02	-15.63	-114.96	Na-HCO ₃	
				23.04.2010	48.2	1038.46	9.19	303.10	12.04	8.98	156.23	21.31	0.00	523.35	0.02	-15.21	-115.86	Na-HCO ₃	
21	Gonghe Spring	102.11	30.62	16.06.2008	48.8	578.15	9.94	108.43	19.30	22.06	18.44	60.27	0.00	337.54	0.01	-11.91	-88.74	Na-HCO ₃	
				27.10.2008	47.6	598.18	10.03	106.53	20.12	22.07	17.83	60.07	0.00	359.48	0.00	-12.40	-90.07	Na-HCO ₃	
				24.06.2009	50.0	610.23	9.57	107.62	18.49	21.27	17.36	65.08	0.00	368.95	-0.01	-12.49	-89.29	Na-HCO ₃	
				23.04.2010	44.8	585.91	9.41	111.76	19.32	21.47	17.85	64.11	0.00	339.32	0.01	-12.33	-89.00	Na-HCO ₃	
22	Erhaoying Spring	102.03	29.59	17.06.2008	65.5	597.31	18.34	154.11	13.45	4.62	37.14	59.30	0.00	307.61	0.02	-14.22	-105.92	Na-HCO ₃	
				27.10.2008	67.4	645.11	18.25	156.50	13.48	4.64	36.80	62.21	0.00	349.86	0.00	-14.31	-109.95	Na-HCO ₃	
				25.06.2009	63.0	585.59	18.53	153.02	13.20	4.72	35.31	62.43	0.00	295.11	0.03	-15.13	-108.27	Na-HCO ₃	
				24.04.2010	67.4	631.67	18.29	150.44	13.01	4.75	35.97	59.85	0.00	348.99	0.00	-15.27	-108.87	Na-HCO ₃	
23	Tianwanhe Spring	102.14	29.49	17.06.2008	53.0	613.88	17.60	76.93	13.10	45.32	24.61	152.25	0.00	281.49	-0.01	-10.31	-79.22	NaMg-HCO ₃ SO ₄	
				25.10.2009	n.d.	632.23	16.92	73.37	12.74	45.71	23.42	139.62	0.00	317.73	-0.02	-10.04	-78.77	NaMg-HCO ₃ SO ₄	
				24.06.2009	54.0	611.73	17.58	79.12	12.81	43.62	24.91	145.24	0.00	286.45	-0.01	-10.50	-83.14	NaMg-HCO ₃ SO ₄	
				24.04.2010	53.2	644.79	17.44	79.11	12.71	44.44	25.65	145.44	0.00	317.73	-0.02	-10.59	-80.38	NaMg-HCO ₃ SO ₄	
24	Caoke Spring	102.10	29.39	18.06.2008	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	
				28.10.2008	41.4	407.78	3.00	48.74	50.10	6.19	3.97	155.60	0.00	139.19	-0.02	-12.38	-88.30	CaNa-SO ₄ HCO ₃	
				24.06.2009	42.9	363.08	2.97	46.19	44.43	6.11	3.94	160.52	0.00	98.05	-0.01	-12.50	-91.09	CaNa-SO ₄ HCO ₃	
				24.04.2010	41.4	369.36	3.03	46.29	45.63	6.17	3.57	161.87	0.00	101.89	-0.02	-12.59	-90.94	CaNa-SO ₄ HCO ₃	
25	Shimian Well	102.22	29.44	18.06.2008	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	
				28.10.2008	30.4	150.26	2.35	38.60	6.99	1.20	2.58	17.82	0.00	69.62	0.00	-11.08	-87.15	Na-HCO ₃	
				23.06.2009	31.3	149.10	2.39	37.61	6.46	1.23	2.44	17.29	0.00	70.79	0.00	-11.28	-90.21	Na-HCO ₃	
				24.04.2010	31.2	137.77	2.34	35.14	6.86	1.20	2.35	17.16	0.00	62.48	0.01	-10.51	-88.94	Na-HCO ₃	
26	Gongyihai Spring	102.39	29.02	The ANHF zone															
				18.06.2008	56.9	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	-14.23	-108.54	n.d.
				28.10.2008	54.1	320.27	3.34	109.14	2.37	1.00	1.91	46.17	0.00	82.11	0.00	-14.19	-108.67	Na-HCO ₃ SO ₄	
				23.06.2009	55.4	306.82	3.37	100.47	2.40	1.01	1.99	48.14	0.00	81.95	-0.01	-14.83	-110.47	Na-HCO ₃ SO ₄	
24.04.2010	56.4	323.92	3.23	113.91	2.44	1.02	3.80	47.98	0.00	88.81	0.02	-14.82	-110.20	Na-HCO ₃ SO ₄					

TABLE I: Continued.

Number	Site	L (E)	B (N)	Date	T °C	TDS mg/L	K ⁺ mg/L	Na ⁺ mg/L	Ca ²⁺ mg/L	Mg ²⁺ mg/L	Cl ⁻ mg/L	SO ₄ ²⁻ mg/L	CO ₃ ²⁻ mg/L	HCO ₃ ⁻ mg/L	ib %	δ ¹⁸ O ‰	δD ‰	Chemical type
27	Chuanxin Spring	102.36	28.26	19.06.2008	n.d.	462.71	4.80	86.08	15.24	11.55	5.52	12.67	0.00	325.96	-0.01	-13.46	-106.54	Na-HCO ₃
				28.10.2008	n.d.	449.48	4.25	86.72	15.29	11.45	5.18	12.44	0.00	311.73	0.45	-13.32	-106.98	Na-HCO ₃
				23.06.2009	49.0	483.93	4.23	85.56	13.75	10.18	4.66	12.00	0.00	351.73	-0.04	-14.51	-108.29	Na-HCO ₃
				24.04.2010	45.1	456.34	4.23	85.46	13.21	9.52	4.72	12.74	0.00	325.14	-0.02	-13.41	-107.37	Na-HCO ₃
28	Taihe Spring	102.14	27.91	19.06.2008	21.1	226.41	0.69	58.65	4.03	0.76	1.99	12.76	0.00	146.70	0.01	-13.37	-106.65	Na-HCO ₃
				29.10.2008	21.1	206.70	0.66	53.56	4.02	0.00	1.91	12.72	0.00	132.20	0.00	-13.31	-106.32	Na-HCO ₃
				n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
				n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
29	Xichang Well	102.15	27.91	19.06.2008	22.3	261.87	0.31	78.93	2.08	0.81	21.78	15.22	0.00	137.09	0.01	-13.78	-106.89	Na-HCO ₃
				29.10.2008	n.d.	229.28	0.32	71.76	2.15	0.85	21.32	15.22	0.00	112.05	0.02	-13.23	-106.64	Na-HCO ₃
				22.06.2009	24.2	238.13	0.33	69.68	2.00	0.81	21.38	15.65	0.00	123.05	0.00	-13.24	-106.07	Na-HCO ₃
				n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
30	Zhuhexiang Spring	102.93	28.07	19.06.2008	49.8	256.45	4.65	54.62	13.18	2.42	3.99	11.97	0.00	161.91	0.02	-12.62	-100.01	Na-HCO ₃
				29.10.2008	49.7	248.22	4.57	51.93	13.39	2.35	3.91	12.22	0.00	154.30	0.00	-12.67	-100.50	Na-HCO ₃
				22.06.2009	49.8	244.72	4.27	50.23	13.75	2.36	3.96	12.05	0.00	152.89	0.00	-13.53	-99.12	Na-HCO ₃
				26.04.2010	48.0	255.93	4.27	50.17	13.26	2.44	3.80	12.31	0.00	167.97	0.00	-12.86	-100.54	Na-HCO ₃
31	Lizhouzhen Spring	102.24	28.09	19.06.2008	n.d.	331.37	14.12	54.16	30.00	0.00	5.52	11.33	0.00	212.12	0.01	-14.48	-110.35	NaCa-HCO ₃
				29.10.2008	48.1	326.67	14.66	52.79	30.77	0.00	5.38	11.34	0.00	210.92	0.02	-14.71	-110.23	NaCa-HCO ₃
				22.06.2009	49.0	323.38	13.37	48.45	29.36	0.00	6.14	10.93	0.00	210.00	-0.01	-14.11	-110.74	NaCa-HCO ₃
				25.04.2010	48.6	321.73	14.08	49.49	29.49	0.00	5.45	11.06	0.00	209.92	0.00	-14.63	-110.06	NaCa-HCO ₃
32	Luojishan Spring	102.54	27.40	20.06.2008	n.d.	1212.86	49.31	208.79	81.48	29.43	136.74	257.26	0.00	449.06	0.00	-11.76	-96.43	Na-HCO ₃ SO ₄
				29.10.2008	43.7	1172.03	46.80	213.50	81.42	28.43	135.47	255.58	0.00	407.06	0.01	-11.64	-96.10	Na-HCO ₃ SO ₄
				22.06.2009	43.8	1150.63	46.67	213.09	71.12	26.07	135.76	244.07	0.00	412.82	0.01	-13.41	-96.20	Na-HCO ₃ SO ₄
				25.04.2010	43.8	1127.84	43.86	200.41	73.26	26.02	135.37	252.58	0.00	395.58	0.00	-13.18	-96.77	Na-HCO ₃ SO ₄

The "n.d." represents samples not analysed.

FIGURE 2: Ternary plots of the cation and anion for the waters.

FIGURE 3: Diagram of $\delta^{18}\text{O}$ versus δD of the waters. LMWL stands for the Local Meteoric Water Line: $\delta\text{D} = 7.06 \delta^{18}\text{O} - 4.06$.

sites. The anions and cations of the water samples were mainly distributed in blocks i, ii, and iii, indicating that Na^+ , Ca^{2+} , SO_4^{2-} , and HCO_3^- were the main chemical composition for most of the samples. Based on the Shoka Lev's classification method, the water samples were classified into 9 chemical types according to the main chemical composition, which are $\text{Na}(\text{Ca})\text{-HCO}_3(\text{SO}_4)$, $\text{Na}(\text{Mg})\text{-HCO}_3(\text{SO}_4)$, $\text{Ca}(\text{Na})\text{-HCO}_3(\text{SO}_4)$, $\text{Mg}(\text{Ca})\text{-SO}_4$, $\text{Ca}(\text{Mg})\text{-SO}_4$, $\text{Ca}(\text{Mg})\text{-HCO}_3$, $\text{Mg}(\text{Ca})\text{-HCO}_3$, $\text{Na-Cl}(\text{HCO}_3)$, and $\text{Ca}(\text{Na})\text{-SO}_4(\text{HCO}_3)$ (Table 1). The cations of the samples of the springs nos. 1–5 occur in the LMSF zone were mainly distributed in the blocks 1, 2, and 4, and the anions were

distributed in the blocks i and ii, which together formed 4 chemical types of $\text{Na}(\text{Ca})\text{-HCO}_3(\text{SO}_4)$, $\text{Ca}(\text{Na})\text{-HCO}_3(\text{SO}_4)$, $\text{Mg}(\text{Ca})\text{-SO}_4$, and $\text{Ca}(\text{Mg})\text{-SO}_4$. The cations of the samples of the springs nos. 6–9 occur in the MJF zone were mainly distributed in the blocks 3 and 4, the anions were distributed in the block i, and the chemical types for the samples were $\text{Ca}(\text{Mg})\text{-HCO}_3$ and $\text{Mg}(\text{Ca})\text{-HCO}_3$. Both of the cations and anions of the samples of the springs nos. 26–32 occur in the ANHF zone were distributed in the blocks i–ii and 1–2, these samples have the similar chemical type of $\text{Na}(\text{Ca})\text{-HCO}_3(\text{SO}_4)$. Half of the samples including the samples of the springs nos. 10–25 occur in the XSHF zone, the cations of these samples were distributed in the blocks 1, 2, and 5, with anions in the blocks i, ii, iii, and vi, and the chemical types were $\text{Na}(\text{Ca})\text{-HCO}_3(\text{SO}_4)$, $\text{Na}(\text{Mg})\text{-HCO}_3(\text{SO}_4)$, $\text{Na-Cl}(\text{HCO}_3)$, and $\text{Ca}(\text{Na})\text{-SO}_4(\text{HCO}_3)$.

5.1.2. The Origins

(1) δD and $\delta^{18}\text{O}$. The values of $\delta^{18}\text{O}$ and δD of the 32 springs were plotted along the local meteoric water line (LMWL: $\delta\text{D} = 7.06 \delta^{18}\text{O} + 4.06$) with different extent shifts from the LMWL (Figure 3), which indicated that the spring waters were mainly originated from meteoric water and with different extent alteration. The values of $\delta^{18}\text{O}$ and δD were plotted in three regions, the region I includes the samples of the springs no.1–3, with values of δD and $\delta^{18}\text{O}$ ranged from -82.13‰ to -61.66‰ and -11.75‰ to -6.18‰ , the region III includes all the samples from the XSHF zone (springs nos. 10–25), with values of δD and $\delta^{18}\text{O}$ between -141.98‰ and -78.77‰ , -18.84‰ and -10.04‰ , respectively, and the others in the region II, with the values of δD and $\delta^{18}\text{O}$ ranging from -116.81‰ to -96.10‰ and -16.34‰ to -11.64‰ (Table 1). The values of δD and $\delta^{18}\text{O}$ of the springs nos.10–25 from

the higher mountain area were more negative, while those of the springs nos. 1–3 collected in the lower altitude region were less negative, which is constant with the previous results [32].

(2) *Na(Ca)-HCO₃(SO₄), Na(Mg)-HCO₃(SO₄), and Na-Cl(HCO₃) Waters.* The water samples of the springs nos. 4–5, 10–23 and 25–32 were belong to these types (Table 1). The springs nos. 4–5, 13–23 and 25 occur in granite, while the springs nos. 10–11 in Late Triassic clastic rock interbedded with volcanic rocks. The spring no. 12 is found in Late Permian volcanic tuff, while the spring no. 26 in Sinian pyroclastic rock. The spring nos. 27–29 and 31 are found in Cretaceous clastic rock, the spring no. 30 in Early Triassic sandstone, and the spring no. 32 in Paleozoic mixed layer (Figure 1). The springs nos. 4–5, 10–23 and 25–32 occur in the rocks enriched in feldspar resulting in similar higher concentrations of Na⁺ and HCO₃[−] because of rock-water interaction. In addition, the samples from the springs nos. 5, 19, 23, and 32 were characterized by the similar concentrations of Ca²⁺, Mg²⁺, and SO₄^{2−} (Table 1), which should be attributed to the water-rock interactions between groundwater and Devonian carbonate (Figure 1). Cl[−] is known to be conservative and derive from the deep earth mainly [33, 34]. The chemical type for the samples from the spring no. 16 was Na-Cl (HCO₃), with the Cl[−] concentration as 328.54 mg/l (Table 1), which suggested the input of deep fluid. Meanwhile, the high ³He/⁴He ratio (between 1.43 and 3.73RA, RA = 1.39 × 10^{−6}) [30] together with the high temperatures (between 80.0°C and 70.2°C) also are the evidences for the upwelling of the deep-earth fluids into the spring (Table 1).

(3) *Ca(Na)-HCO₃(SO₄), Ca(Mg)-HCO₃, Mg(Ca)-HCO₃, and Ca(Na)-SO₄(HCO₃) Waters.* These included the samples from the springs nos. 1, 6–9, and 24 (Table 1). The spring no. 1 occurs in Jurassic carbonate interbedded with sandstone. The springs nos. 6 and 9 occur in Late Triassic clastic rock interbedded with carbonate. The springs nos. 7 and 8 occur in Middle Triassic clastic rock interbedded with carbonate, and the spring no. 24 is found in Carboniferous carbonate (Figure 1). The springs nos. 1, 6–9, and 24 occur in strata enriched in carbonate, so that the main components of the waters were Ca²⁺, Mg²⁺ and HCO₃[−] because of the interaction between the groundwater and carbonate. The samples of the springs nos. 8 and 9 have the same chemical type of Mg (Ca)-HCO₃, and similar Ca/Mg ratios (0.54 and 0.86, Table 1), which can be attributed to the dissolution of calcite combined with the low temperatures of the springs nos. 8 and 9 (8.8°C and 9.0°C respectively) (Table 1).

(4) *Mg(Ca)-SO₄ and Ca(Mg)-SO₄ Waters.* The springs nos. 2 and 3 occur in Devonian to Silurian pyrite-bearing shale and marlite (Figure 1), from which the samples had higher SO₄^{2−} concentration (878.08 mg/L and 812.50 mg/L, resp.) and TDS (1571.81 mg/L and 1195.08 mg/L, resp.), with the chemical types of Mg (Ca)-SO₄ and Ca (Mg)-SO₄ (Table 1), indicating interaction between carbonate and groundwater enhanced by the sulfate from the oxidation of pyrite [35].

5.2. Chemical Variations of the Spring Waters Associated with Seismic Activities

5.2.1. *Temporal Variations.* The hydrochemical data of the water samples showed evident trend of temporal variations.

The geochemical parameters of the samples nos. 1, 3–5 from the LMSF zone and the samples nos. 15 and 20 from the XSHF zone varied by more than 20% one month after the Wenchuan earthquake, and similar variations happened after October 2008. The concentrations of Na⁺, Ca²⁺, Cl[−], SO₄^{2−}, and TDS of the sample of the spring no. 1 decreased by 42%, 39%, 53%, 31%, and 35%, respectively; for the sample number 2, the decreasing amplitudes for the concentrations of Na⁺, Mg²⁺, Ca²⁺, Cl[−], SO₄^{2−}, and TDS were 23%, 45%, 25%, 38%, 22%, and 33%, separately. The concentrations of Mg²⁺, Ca²⁺, HCO₃[−], and TDS of the sample from the spring no. 3 decreased by 33%, 42%, 95%, and 35%, respectively. The concentrations of Na⁺, HCO₃[−], and TDS of the sample of the spring no. 4 decreased by 67%, 33% and 57%, respectively. For the sample of the spring no. 5, the concentrations of Na⁺, HCO₃[−] and TDS decreased by 25%, 41%, and 26%, respectively. The concentrations of Na⁺ and SO₄^{2−} of sample of the spring no. 20 decreased by 28% and 26%, respectively. For the sample of the spring no. 15, however the concentrations of Na⁺, HCO₃[−] and TDS increased in 17 April 2010, with the amplitudes of 53%, 38%, and 38%, respectively. Meanwhile, the δ¹⁸O-δD plots of the samples from the springs nos. 1, 3, and 4 approached the LMWL (Local Meteoric Line) after sizeable shift in June 2008, and the similar variation happened for the spring no. 2 after October 2008 (Figure 3), which indicated gradual input decrease of deep-earth fluids that formed by water-rock reaction after the Wenchuan earthquake.

The supplement of deep water characterized by higher mineralization and enriched in δ¹⁸O is considered as a main factor for controlling pre- and co-seismic geochemical variations of the groundwater and springs [2, 6, 7, 11, 13, 14, 16, 36, 37]. The Coulomb stress in the middle-north segment of the LMSF zone, the southeast segment of the XSHF zone, and the south segment of the MJF zone had been enhanced, which resulted in the Wenchuan M_s 8.0 earthquake [38, 39], and the geochemical variations of the samples from the springs nos. 1–5 and 20. Meanwhile, the input decrease of deep-earth water during the sampling period in which seismic activity decreased gradually. 747 aftershocks with M_s higher than 4.0 occurred in the three fault zones, and most of which (including all the 12 events with amplitude ranged from 6.0 to 6.4) occurred before September 10, 2008 (China Earthquake Network Center, Figure 1). As the aftershock activities became weak, the TDS values of the samples from the springs nos. 1–5 and 20 decreased gradually to the normal value (Figure 4), indicating the supplement decrease of the deep-earth fluid as the Coulomb stress being released after the events as indicated by the data of oxygen and hydrogen isotope compositions (Figure 3).

Notably, the concentrations of Na⁺, HCO₃[−], and TDS of the sample of the spring no. 15 increased by 53%, 38%, and 38% on April 17, 2010 (Figure 4), which was closely related to

FIGURE 4: Temporal variations of the geochemical parameters of the 32 hot spring waters after the Wenchuan earthquake. The abscissas indicate the numbers of the samples and the ordinates indicate the ratios of each batch of samples with the first batch of samples.

the 28 April 2010 an M_s 5.4 aftershock 15 kilometers epicenter distance to the spring no. 15 (Figure 1). Therefore, it can be inferred that the hydrogeochemical variations of the spring no. 15 might be the short-term hydrogeochemical precursors for the M_s 5.4 event.

The geochemical parameters of the samples from the springs nos. 16-17 and 19 about 200 kilometers showed evident increase trend before the Wenchuan earthquake [20, 24, 25]. The K^+ concentrations of the samples of the springs nos. 16-17 and 19 increased by 19.3% to 54.4% before the main shock, and rapidly decreased to the normal values after the main shock. The concentrations of SO_4^{2-} for the samples from the springs nos. 16 and 19 increased by 32.0% and 59.6% respectively before the earthquake, and then dropped

FIGURE 5: Temporal variations of concentration of K^+ and SO_4^{2-} . Black squares: waters collected from the Guanding spring; black circles: waters collected from the Longtougou spring; black triangles: waters collected from the Erdaoqiao spring; the dashed vertical line: the time of the occurrence of the Wenchuan earthquake.

to the normal value after the main shock (Figure 5). The springs nos. 16-17 and 19 occur in the intersection of the LMSF, XSHF and ANHF zones where the extrusion stress increased yearly from 2004 to 2007 before the Wenchuan earthquake [40, 41]. Therefore, the increase of K^+ and SO_4^{2-} concentrations for the samples of the springs nos. 16-17 and 19 could be attributed to the more supplement of the deep fluids with higher concentrations of K^+ and SO_4^{2-} [42]. The $^3\text{He}/^4\text{He}$ ratios of the samples from the springs nos. 16-17 and 19 increased before the Wenchuan earthquake, but showed a remarkable drop after the shock, and then became normal indicating the more supplement of mantle-derived fluids before the main shock [30].

5.2.2. Spatial Variations. The geochemical and isotopic compositions of the samples from the springs nos. 1-5 occur in the LMSF zone performed evident variation after the Wenchuan earthquake, with the highest amplitude as 95% for HCO_3^- of the sample from the spring no. 5. However, for the other samples from the other fault zones, just the samples from the springs nos. 15 and 20 presented obvious geochemical variations, with the highest amplitude as 53% for Na^+ of the sample from the spring no. 15 (Figure 4). The Wenchuan earthquake happened in the LMSF zone, which enhanced the Coulomb stress of the zone [38, 39], and with most of the aftershocks occurred there (Figure 1). For the springs nos. 1-5 occurred in the LMSF zone, the epicenter distances are between 50 and 110 kilometers, which are smaller than those of the hot springs nos. 6-32 (between 190 and 410 km). Therefore, the variation of the tectonic stress induced by the Wenchuan earthquake and the aftershocks may resulted in,

the geochemical and isotopic variations for the springs nos. 1–5 after the Wenchuan earthquake.

6. Conclusions

120 water samples of the 32 hot springs in the western Sichuan have been analyzed after the Wenchuan earthquake. The following conclusions can be drawn.

- (1) The waters of the 32 springs were mainly derived from meteoric precipitation. Because of the isotopic elevation effect, the δD and $\delta^{18}O$ of the samples from the higher mountain area were lower, but those of samples from the lower altitude region were higher. All of the analyzed water samples from the 32 springs can be classified into nine chemical types: Na(Ca)-HCO₃(SO₄), Na(Mg)-HCO₃(SO₄), Ca(Na)-HCO₃(SO₄), Mg(Ca)-SO₄, Ca(Mg)-SO₄, Ca(Mg)-HCO₃, Mg(Ca)-HCO₃, Na-Cl(HCO₃), and Ca(Na)-SO₄(HCO₃), which were mainly controlled by water-rock interaction and the input of deep fluid.
- (2) The concentrations of K⁺ and SO₄²⁻ of the samples of the springs nos. 16–17 and 19 in the intersection of LMSE, XSHF, and ANHF varied evidently, with the amplitude ranging from 19.3% to 59.6% before and after the Wenchuan earthquake which may be attributed to the interfusion of the deep fluids with high K⁺ and SO₄²⁻ induced by the increased tectonic stress.
- (3) The hydrogeochemical variations of the springs closer to the epicenter performed more obviously after the Wenchuan earthquake. As the aftershock activities became weak, the geochemical parameters of the samples from the springs nos. 1–5, 15, and 20 located in the regions where the tectonic stress was enhanced before the Wenchuan M_s 8.0 earthquake decreased by more than 20%, and with the $\delta^{18}O$ the LMWL, which may be related to the change of tectonic stress when the aftershock activities became weak.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The authors are grateful to the Seismological Bureau of Sichuan province and Professor Fang Du for helping in the field work. The authors sincerely thank the editor and reviewers for their valuable comments and grammar edits in improving the paper. This research was supported by the National Science and Technology Support Program (the characteristics of gas geochemistry of Longmenshan faults, WFS-10), the Basic Science Research Plan of the Institute of Earthquake Science, China Earthquake Administration (The fluid geochemical survey of the mud volcanoes in the North of the Tianshan Mountain, Xinjiang Uygur Autonomous

Region 02132417), and the Basic Science Research Plan of the Institute of Earthquake Science, China Earthquake Administration (02132417).

References

- [1] A. Dadomo, M. Lemmi, G. Martinelli, M. Menichetti, and L. Telesca, "Springwater continuous monitoring in the L'Aquila area in concomitance with the April 2009 seismic swarm in central Italy: constraining factors to possible deep-seated fluid emissions," *Chemical Geology*, vol. 339, pp. 169–176.
- [2] R. Favara, F. Italiano, and G. Martinelli, "Earthquake-induced chemical changes in the thermal waters of the Umbria region during the 1997–1998 seismic swarm," *Terra Nova*, vol. 13, no. 3, pp. 227–233, 2001.
- [3] C.-Y. King, N. Koizumi, and Y. Kitagawa, "Hydrogeochemical anomalies and the 1995 Kobe earthquake," *Science*, vol. 269, no. 5220, pp. 38–39, 1995.
- [4] F. Quattrocchi, R. Favara, G. Capasso et al., "Thermal anomalies and fluid geochemistry framework in occurrence of the 2000–2001 Nizza Monferrate seismic sequence (northern Italy): episodic changes in the fault zone heat flow or chemical mixing phenomena?" *Natural Hazards and Earth System Science*, vol. 3, no. 3–4, pp. 269–277, 2003.
- [5] D. V. Reddy, P. Nagabhushanam, and B. S. Sukhija, "Earthquake (M 5.1) induced hydrogeochemical and $\delta^{18}O$ changes: Validation of aquifer breaching-mixing model in Koyana, India," *Geophysical Journal International*, vol. 184, no. 1, pp. 359–370, 2011.
- [6] S.-R. Song, Y.-L. Chen, C.-M. Liu et al., "Hydrochemical changes in spring waters in Taiwan: implications for evaluating sites for earthquake precursory monitoring," *Terrestrial, Atmospheric and Oceanic Sciences*, vol. 16, no. 4, pp. 745–762, 2005.
- [7] F. Italiano, P. Bonfanti, L. Pizzino, and F. Quattrocchi, "Geochemistry of fluids discharged over the seismic area of the Southern Apennines (Calabria region, Southern Italy): implications for Fluid-Fault relationships," *Applied Geochemistry*, vol. 25, no. 4, pp. 540–554, 2010.
- [8] S. İnan, W. P. Balderer, F. Leuenberger-West, H. Yakan, A. Özvan, and F. T. Freund, "Springwater chemical anomalies prior to the M_w = 7.2 Van earthquake (Turkey)," *Geochemical Journal*, vol. 46, no. 1, pp. e11–e16, 2012.
- [9] M. Malakootian and J. Nouri, "Chemical variations of ground water affected by the earthquake in Bam region," *International Journal of Environmental Research*, vol. 4, no. 3, pp. 443–454, 2010.
- [10] D. Thomas, "Geochemical precursors to seismic activity," *Pure and Applied Geophysics*, vol. 126, no. 2–4, pp. 241–266, 1988.
- [11] H. Woith, R. Wang, U. Maiwald, A. Pekdeger, and J. Zschau, "On the origin of geochemical anomalies in groundwaters induced by the Adana 1998 earthquake," *Chemical Geology*, vol. 339, pp. 177–186, 2013.
- [12] X. Zhou, J. Du, Z. Chen et al., "Geochemistry of soil gas in the seismic fault zone produced by the Wenchuan M_s 8.0 earthquake, southwestern China," *Geochemical Transactions*, vol. 11, article 5, 2010.
- [13] M. S. Miller, A. Allam, T. W. Becker, J. F. Dileo, and J. Wookey, "Constraints on the geodynamic evolution of the westernmost Mediterranean and northwestern Africa from shear wave splitting analysis," *Earth and Planetary Science Letters*, vol. 375, pp. 234–243, 2013.

- [14] C.-Y. Wang, M. Manga, C.-H. Wang, and C.-H. Chen, "Transient change in groundwater temperature after earthquakes," *Geology*, vol. 40, no. 2, pp. 119–122, 2012.
- [15] G. Etiope, M. Calcara, and F. Quattrocchi, "Seismogeochemical algorithms for earthquake prediction: an overview," *Annali di Geofisica*, vol. 40, no. 6, pp. 1483–1492, 1997.
- [16] J. Du, K. Amita, S. Ohsawa, Y. Zhang, C. Kang, and M. Yamada, "Experimental evidence on formation of imminent and short-term hydrochemical precursors for earthquakes," *Applied Geochemistry*, vol. 25, no. 4, pp. 586–592, 2010.
- [17] Y. Cao, H. Li, Z. Liu, D. Yuan, and L. Shen, "Comparison of geochemical features of warm springs between Chongqing and Kangding," *Carsologica Sinica*, vol. 25, no. 2, pp. 113–121, 2006.
- [18] S. Jia, X. Li, and Y. Hu, "The hydrogeochemistry problems of the shallow heat reservoir in Zhuhai thermal spring," *Journal of Chengdu University of Technology*, vol. 24, no. 1, pp. 78–84, 1997.
- [19] L. L. Luo, "The primary study of distribution and genesis of the springs in western Sichuan province," *Journal of Chongqing Teachers College*, vol. 11, no. 2, pp. 39–47, 1994.
- [20] L. Zaihua, Y. Daoxian, H. Shiyi, Z. Meiliang, and Z. Jiagui, "Geochemical features of the geothermal CO₂-water-carbonate rock system and analysis on its CO₂ sources," *Science in China D: Earth Sciences*, vol. 43, no. 6, pp. 569–576, 2000.
- [21] H. Li, G. Yin, J. Yang, and X. Fan, "Isotopic hydrogeochemical characteristics of water bodies in Munigou, Sichuan," *Acta Geoscientica Sinica*, vol. 26, no. 3, pp. 255–258, 2005.
- [22] M. Liu, "A study on genesis of Hongge hot spring in Yanbian, Sichuan Province," *Coal Geology of China*, vol. 20, no. 3, pp. 45–49, 2008.
- [23] R. Q. Lu, D. Y. Wang, Y. H. Zuo, Q. L. Wu, Y. W. Liu, and X. Y. Li, "Remote-sensing geological features of thermal spring in the Kaka Gulley, Songpan, Sichuan," *Acta Geologica Sinica*, vol. 28, no. 1, pp. 73–75, 2008.
- [24] L. C. Shen, *The study of deep source CO₂ degasification and carbon cycle in the southwest of China [Ph.D. dissertation]*, Southwest University, 2007.
- [25] J. G. Zhang and H. T. Hu, "Quantification research of CO₂ outgassing from hot springs along Xianshuihe fault zone, Sichuan," *Hydrogeology and Engineering Geology*, vol. 27, no. 3, pp. 19–21, 2000.
- [26] B. C. Burchfiel, L. H. Royden, R. D. van der Hilst et al., "A geological and geophysical context for the Wenchuan earthquake of 12 May 2008, Sichuan, People's Republic of China," *GSA Today*, vol. 18, no. 7, pp. 4–11, 2008.
- [27] Z. L. Li, Z. H. Luo, Y. L. Chen, and J. X. Zhao, "Geochronological and geochemical characteristics of Kangding metamorphosed intrusions," *Acta Geoscientica Sinica*, vol. 26, no. 9, pp. 87–89, 2005.
- [28] S. Wu, "The discovery and significance of ductile shear zones in pengguan complex granite mass of Longmenshan mountains," *Journal Mineral Petrology*, vol. 13, no. 2, pp. 71–77, 1993.
- [29] Y. J. Zhao, C. Yuan, M. F. Zhou, D. P. Yan, X. P. Long, and K. D. Cai, "Post-orogenic extension of Songpan-Garz(e) orogen in Early Jurassic: constraints from Niuxingou monzodiorite and Siguniangshan A-type granite of western Sichuan, China," *Geochimica*, vol. 36, no. 2, pp. 139–152, 2007.
- [30] X. C. Zhou, *Gas geochemistry in western Sichuan related to 12 May 2008 Wenchuan M_s 8.0 earthquake [Ph.D. dissertation]*, University of Science and Technology of China, 2011.
- [31] X.-W. Xu, X.-Z. Wen, J.-Q. Ye et al., "The M_s 8.0 Wenchuan earthquake surface ruptures and its seismogenic structure," *Seismology and Geology*, vol. 30, no. 3, pp. 597–629, 2008.
- [32] C. P. Chamberlain and M. A. Poage, "Reconstructing the paleotopography of mountain belts from the isotopic composition of authigenic minerals," *Geology*, vol. 28, no. 2, pp. 115–118, 2000.
- [33] J. Du, C. Liu, B. Fu et al., "Variations of geothermometry and chemical-isotopic compositions of hot spring fluids in the Rehai geothermal field, southwestern China," *Journal of Volcanology and Geothermal Research*, vol. 142, no. 3–4, pp. 243–261, 2005.
- [34] L. Shevenell and F. Goff, "Evolution of hydrothermal waters at Mount St. Helens, Washington, USA," *Journal of Volcanology and Geothermal Research*, vol. 69, no. 1–2, pp. 73–94, 1995.
- [35] M. Floyd, M. A. Czerewko, J. C. Cripps, and D. A. Spears, "Pyrite oxidation in Lower Lias Clay at concrete highway structures affected by thaumasite, Gloucestershire, UK," *Cement and Concrete Composites*, vol. 25, no. 8, pp. 1015–1024, 2003.
- [36] A. Charmoille, O. Fabbri, J. Mudry, Y. Guglielmi, and C. Bertrand, "Post-seismic permeability change in a shallow fractured aquifer following a ML 5.1 earthquake (Fourbanne karst aquifer, Jura outermost thrust unit, eastern France)," *Geophysical Research Letters*, vol. 32, no. 18, Article ID L18406, pp. 1–5, 2005.
- [37] C.-Y. King, W. Zhang, and Z. Zhang, "Earthquake-induced groundwater and gas changes," *Pure and Applied Geophysics*, vol. 163, no. 4, pp. 633–645, 2006.
- [38] B. Shan, X. Xiong, Y. Zheng, and F. Diao, "Stress changes on major faults caused by Mw 7.9 Wenchuan earthquake, May 12, 2008," *Science in China D: Earth Sciences*, vol. 52, no. 5, pp. 593–601, 2009.
- [39] C. D. Xie, Y. Q. Zhu, X. L. Lei, H. Y. Yu, and X. L. Hu, "Pattern of stress change and its effect on seismicity rate caused by M_s 8.0 Wenchuan earthquake," *Science China Earth Sciences*, vol. 53, no. 9, pp. 1260–1270, 2010.
- [40] G. Luo and M. Liu, "Stress evolution and fault interactions before and after the 2008 Great Wenchuan earthquake," *Tectonophysics*, vol. 491, no. 1–4, pp. 127–140, 2010.
- [41] X. Zhang, D. Cui, and F. Jing, "Inversion of seismic parameters of the Wenchuan earthquake and analysis on variation of its Coulomb stress based on GPS observation," *Journal of Seismological Research*, vol. 32, no. 4, pp. 351–356, 2009.
- [42] Y. Fang, Z. S. Jiang, Y. L. Yang, G. H. Gu, and Y. Q. Wu, "Deformation characteristics of Southern segment of Xianshuihe fault before and after Wenchuan M_s 8.0 earthquake," *Journal of Geodesy and Geodynamics*, vol. 30, no. 3, pp. 22–27, 2010.