Research Article

Silicon Improves Maize Photosynthesis in Saline-Alkaline Soils

Zhiming Xie,^{1,2} Ri Song,³ Hongbo Shao,⁴ Fengbin Song,² Hongwen Xu,⁵ and Yan Lu⁵

¹College of Life Sciences, Baicheng Normal University, Baicheng 137000, China

²Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences, Changchun 130102, China

³College of Agronomy, Jilin Agricultural University, Changchun 130118, China

⁴Key Laboratory of Coastal Biology & Bioresources Utilization, Yantai Institute of Coastal Zone Research, Chinese Academy of Sciences (CAS), Yantai 264003, China

⁵School of Urban and Environmental Science, Huaiyin Normal University, Huai'an 223300, China

Correspondence should be addressed to Hongbo Shao; shaohongbochu@126.com and Fengbin Song; fb_song@163.com

Received 10 June 2014; Accepted 26 June 2014

Academic Editor: Marian Brestic

Copyright © 2015 Zhiming Xie et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The research aimed to determine the effects of Si application on photosynthetic characteristics of maize on saline-alkaline soil, including photosynthetic rate (P_n), stomatal conductance (g_s), transpiration rate (E), and intercellular CO_2 concentration (C_i) of maize in the field with five levels (0, 45, 90, 150, and 225 $\text{kg}\cdot\text{ha}^{-1}$) of Si supplying. Experimental results showed that the values of P_n , g_s , and C_i of maize were significantly enhanced while the values of E of maize were dramatically decreased by certain doses of silicon fertilizers, which meant that Si application with proper doses significantly increased photosynthetic efficiency of maize in different growth stages under stressing environment of saline-alkaline soil. The optimal dose of Si application in this experiment was 150 $\text{kg}\cdot\text{ha}^{-1}$ Si. It indicated that increase in maize photosynthesis under saline-alkaline stress took place by Si application with proper doses, which is helpful to improve growth and yield of maize.

1. Introduction

Salinity and sodicity toxicities are worldwide agricultural and environmental problems; it is estimated that there is approximately 27 million hectares of salinised soils in China's coastal and inland areas [1]. The saline and sodic soils, among which about 23% of the cultivated lands are saline and 37% are sodic, cover about 10% of total arable lands worldwide [2]. This kind of soil is widespread in arid and semiarid regions of the world and causes severe environmental and agricultural problems [3]. Soil salinity and sodicity, which seriously affect the stages of germination, seedling growth and vigour, vegetative growth, flowering, and fruit set of crops [4], adversely affect crop production in different regions, especially in arid and semiarid regions of the world [5, 6].

Supplement silicon, which is much cheaper than other methods to minimize salinity and sodicity, such as reclamation, water, and drainage, is an alternative way for overcoming the negative effects of salinity and sodicity on the plant

growth and yield [7]. The beneficial effect of silicon is more evident under stress conditions; this is because silicon is able to protect plants from multiple abiotic and biotic stresses [8]. Silicon can alleviate the adverse effects of salt stress on plants by increasing cell membrane integrity and stability through its ability to stimulate the plants' antioxidant system [9]. Silicon application can moderate the salinity and sodicity stress in plants and plays a multitude of roles in plant existence and crop performance, and silicon is deposited in leaves leading towards decreased transpiration and hence dilutes salts accumulated in saline environment [10]. In general, graminaceous plants accumulate much more silicon in their tissues than other species [11].

Maize (*Zea mays* L.) is reported as salt susceptible [12]. This study aimed to investigate the effects of silicon fertilizer, which was conducted in field tests, on photosynthetic characteristics of maize under the condition of saline-alkaline soil in Northeast China.

2. Materials and Methods

2.1. Experimental Site and Soil Condition. During May to October, 2012, the field experiments were carried out in Agricultural Research Center in Gaoping Village, Baicheng City, Jilin Province, China (45.38°N latitude, 122.50°E longitude), on an saline-alkaline soil, where the climatic conditions are as follows: continental monsoon climate, lying in midtemperate zone, a mean annual temperature being 4.9°C, the annual sunshine time being 2, 919.4 hours, an average frost-free period being 157 days, and the annual average precipitation being 399.9 mm with 88% distributed in May–September. The basic properties of the soil (0–20 cm), which is saline-alkaline soil, are that it contains organic matter 19.6 g·kg⁻¹, available nitrogen 112.46, available phosphorus 11.38 mg·kg⁻¹, available potassium 162.39 mg·kg⁻¹, available silicon 238.60 mg·kg⁻¹, ESP 11.36%, EC 9.88 dS·m⁻¹, K⁺ 83.62 mg·kg⁻¹, Na⁺ 405.76 mg·kg⁻¹, Ca²⁺ 43.26 mg·kg⁻¹, Mg²⁺ 65.68 mg·kg⁻¹, Cl⁻ 69.18 mg·kg⁻¹, HCO₃⁻ 896 mg·kg⁻¹, SO₄²⁻ 40.75, and pH 8.00.

2.2. Experimental Design. The experiment was conducted in the form of randomized complete block design (RCBD) with three replications in a 5 × 10 m² net plot size. Maize (Zhengdan 958) was sown on May 6 with a density of 65 000 plants·ha⁻¹. A uniform dose of basal fertilizer was applied to all experimental plots prior to seed sowing with N 200 kg·ha⁻¹ as urea, P₂O₅ 100 kg·ha⁻¹ as single super phosphate, and K₂O 80 kg·ha⁻¹ as potassium sulphate. Treatments were five levels of SiO₂: T1 with SiO₂ 0 kg·ha⁻¹ (control), T2 with SiO₂ 45 kg·ha⁻¹, T3 with SiO₂ 90 kg·ha⁻¹, T4 with SiO₂ 150 kg·ha⁻¹, and T5 with SiO₂ 225 kg·ha⁻¹. The silicon fertilizer used, which was in the form of a sodium metasilicate (Na₂SiO₃·H₂O) with the content of soluble SiO₂ 30%, was produced by Yubei Fertilizer Company Limited, Xinxiang City, Henan Province, China. All fertilizers were applied as basal applications. In this experiment, maize was evaluated for its physiological parameters of net photosynthetic rate (P_n), transpiration rate (E), stomatal conductance (g_s), and intercellular CO₂ concentration (C_i). Observations were recorded at four key growth stages: big trumpet stage (or the 12-leaf stage), silking stage, grain filling stage, and milk stage. The dates of growth stages are shown in Table 1.

2.3. Measurement of Gas Exchange Parameters. The photosynthetic characteristics of gas exchange parameters, P_n , E , g_s , and C_i of the top second fully expanded leaf at the four growth stages in the field, were measured with a portable open flow gas exchange system LI-6400 (LI-COR Inc., USA) between 9:00 am and 11:00 am. The photosynthetically active radiation was 2000 μmol·m⁻²·s⁻¹, CO₂ concentration was 350 μmol·mol⁻¹, and leaf temperature was 25°C [13–15].

2.4. Statistical Analysis. All the data were analyzed with one-way analysis of variance (ANOVA) procedures using SPSS Version 17.0 for Windows. The differences between means were compared by Duncan's test at 0.05 significance level.

TABLE 1: Dates of growth stages of maize on saline-alkaline soil in 2012.

Sowing stage	Emergence stage	Big trumpet stage	Silking stage	Grain filling stage	Milk stage
4/26	5/20	7/12	8/2	8/16	9/16

3. Results

Observed from Tables 2, 3, and 4, changes in parameters of net photosynthetic rate (P_n), transpiration rate (E), and stomatal conductance (g_s) show that the values measured of these parameters from big trumpet stage to milk stage increased first from the beginning stage, big trumpet stage, and reached the peak values at silking stage, after which these values decreased as maize growing. According to the results of Table 5, there is a similar changing pattern among these Si application treatments T1, T2, T3, T4, and T5, where the values of intercellular CO₂ concentration (C_i) under the five Si treatments continued decreasing from big trumpet stage, got the lowest values of C_i at grain filling stage, and then slowly increased at milk stage.

3.1. Net Photosynthetic Rate (P_n). Results showed that the parameter photosynthetic rate (P_n) in leaves of maize was affected significantly by Si application treatments and P_n was significantly decreased in later growth stages (Table 2). From big trumpet stage to milk stage, the values of P_n under the treatments of T3, T4, and T5 (90 kg·ha⁻¹ Si, 150 kg·ha⁻¹ Si, and 225 kg·ha⁻¹ Si) in the same growth stage were significantly ($P \leq 0.05$) higher than those under the treatment of T1 and T2 (0 kg·ha⁻¹ and 45 kg·ha⁻¹ Si). When the content of Si application reached 90 kg·ha⁻¹ Si, the value of P_n increased significantly ($P \leq 0.05$) with the increased dose of Si application. Low level of Si application (T2, 45 kg·ha⁻¹ Si) did not change P_n significantly and high levels of Si application (T4, 150 kg·ha⁻¹ Si, and T5, 225 kg·ha⁻¹ Si) increased P_n significantly ($P \leq 0.05$), but there were no significant differences between the treatments of 150 kg·ha⁻¹ Si and 225 kg·ha⁻¹ Si dose.

3.2. Transpiration Rate (E). During each stage from big trumpet stage to milk stage, transpiration rate (E) (Table 3) got higher values under the treatments of T1 (without Si application) and T2 (45 kg·ha⁻¹ Si) than those under the treatments of T3, T4, and T5. There were no significant differences between T1 and T2. The values of E significantly ($P \leq 0.05$) decreased with the increase of Si applying dose from 90 kg·ha⁻¹ Si in each growth stage. Comparing the values of E by Si application of T3, T4, and T5 with those by T1, it is shown that, during big trumpet stage, the former decreased 16.28%, 20.32%, and 20.79%, respectively, compared to those of the latter; during silking stage, the former decreased 9.44%, 15.63%, and 16.75%, respectively, compared to those of the latter; during grain filling stage, the former decreased 11.38%, 21.12%, and 22.89%, respectively,

TABLE 2: Effects of Si application on net photosynthetic rate (P_n) in leaves of maize.

Growth stages	Photosynthetic rate (P_n) ($\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$)				
	T1	T2	T3	T4	T5
Big trumpet stage	19.15 $\pm$ 0.25 ^c	18.86 $\pm$ 0.69 ^c	23.02 $\pm$ 0.73 ^b	24.83 $\pm$ 0.75 ^a	24.95 $\pm$ 0.42 ^a
Silking stage	33.62 $\pm$ 0.86 ^c	34.11 $\pm$ 0.62 ^c	37.86 $\pm$ 0.75 ^b	39.30 $\pm$ 0.55 ^a	39.52 $\pm$ 0.39 ^a
Grain filling stage	24.11 $\pm$ 0.33 ^c	23.93 $\pm$ 0.50 ^c	26.10 $\pm$ 0.28 ^b	28.35 $\pm$ 0.37 ^a	28.67 $\pm$ 0.47 ^a
Milk stage	12.70 $\pm$ 0.59 ^c	12.91 $\pm$ 0.36 ^c	14.32 $\pm$ 0.53 ^b	16.85 $\pm$ 0.40 ^a	16.89 $\pm$ 0.33 ^a

Means ($\pm$ SD) labeled with different letters within each column are significantly different ($P < 0.05$) by Duncan's test, $n = 10$.

TABLE 3: Effects of Si application on transpiration rate (E) in leaves of maize.

Growth stages	Transpiration rate (E) ($\text{mmol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$)				
	T1	T2	T3	T4	T5
Big trumpet stage	8.66 $\pm$ 0.21 ^a	8.23 $\pm$ 0.16 ^a	7.25 $\pm$ 0.10 ^b	6.90 $\pm$ 0.12 ^c	6.86 $\pm$ 0.09 ^c
Silking stage	9.85 $\pm$ 0.08 ^a	9.67 $\pm$ 0.18 ^a	8.92 $\pm$ 0.15 ^b	8.31 $\pm$ 0.13 ^c	8.20 $\pm$ 0.16 ^c
Grain filling stage	7.82 $\pm$ 0.15 ^a	7.97 $\pm$ 0.11 ^a	6.93 $\pm$ 0.13 ^b	6.09 $\pm$ 0.12 ^c	6.03 $\pm$ 0.09 ^c
Milk stage	4.31 $\pm$ 0.08 ^a	4.15 $\pm$ 0.10 ^a	3.52 $\pm$ 0.07 ^b	3.03 $\pm$ 0.08 ^c	2.96 $\pm$ 0.05 ^c

Means ($\pm$ SD) labeled with different letters within each column are significantly different ($P < 0.05$) by Duncan's test, $n = 10$.

TABLE 4: Effects of silicon on stomatal conductance (g_s) in leaves of maize.

Growth stages	Stomatal conductance (g_s) ($\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$)				
	T1	T2	T3	T4	T5
Big trumpet stage	0.39 $\pm$ 0.06 ^b	0.37 $\pm$ 0.08 ^b	0.40 $\pm$ 0.02 ^b	0.57 $\pm$ 0.09 ^a	0.56 $\pm$ 0.07 ^a
Silking stage	0.66 $\pm$ 0.03 ^b	0.63 $\pm$ 0.05 ^b	0.65 $\pm$ 0.06 ^b	0.75 $\pm$ 0.08 ^a	0.78 $\pm$ 0.09 ^a
Grain filling stage	0.47 $\pm$ 0.08 ^c	0.45 $\pm$ 0.03 ^c	0.58 $\pm$ 0.07 ^b	0.66 $\pm$ 0.05 ^a	0.69 $\pm$ 0.06 ^a
Milk stage	0.23 $\pm$ 0.02 ^c	0.25 $\pm$ 0.03 ^c	0.32 $\pm$ 0.02 ^b	0.39 $\pm$ 0.03 ^a	0.38 $\pm$ 0.01 ^a

Means ($\pm$ SD) labeled with different letters within each column are significantly different ($P < 0.05$) by Duncan's test, $n = 10$.

TABLE 5: Effects of silicon on intercellular CO_2 concentration (C_i) in leaves of maize.

Growth stages	Intercellular CO_2 concentration (C_i) ($\mu\text{mol}\cdot\text{mol}^{-1}$)				
	T1	T2	T3	T4	T5
Big trumpet stage	139.10 $\pm$ 1.73 ^d	138.23 $\pm$ 2.02 ^d	157.32 $\pm$ 1.35 ^c	162.95 $\pm$ 2.13 ^b	175.68 $\pm$ 2.23 ^a
Silking stage	116.82 $\pm$ 1.69 ^d	120.36 $\pm$ 1.86 ^d	136.20 $\pm$ 1.57 ^c	140.85 $\pm$ 1.86 ^b	152.61 $\pm$ 1.75 ^a
Grain filling stage	103.73 $\pm$ 1.93 ^d	108.92 $\pm$ 1.70 ^d	118.36 $\pm$ 2.33 ^c	131.25 $\pm$ 1.78 ^b	140.62 $\pm$ 1.66 ^a
Milk stage	106.90 $\pm$ 1.28 ^d	112.78 $\pm$ 1.66 ^d	121.32 $\pm$ 1.51 ^c	134.25 $\pm$ 1.36 ^b	142.83 $\pm$ 1.92 ^a

Means ($\pm$ SD) labeled with different letters within each column are significantly different ($P < 0.05$) by Duncan's test, $n = 10$.

compared to those of the latter; during milk stage, the former decreased 18.33%, 29.70%, and 31.32%, compared to those of the latter. There were no significant differences between the treatments of T1 and T2. So during the four studied stages, the value of E of maize began to decreased significantly ($P < 0.05$) when the dose of Si application got to the amount of $90 \text{ kg}\cdot\text{ha}^{-1}$, after which the values of E decreased dramatically with the increased dose of Si application; there was no significant difference between the treatments of doses $150 \text{ kg}\cdot\text{ha}^{-1}$ and $225 \text{ kg}\cdot\text{ha}^{-1}$.

3.3. Stomatal Conductance (g_s). According to the data (Table 4) of big trumpet stage and silking stage, it is shown

that the values of stomatal conductance (g_s) of maize were increased significantly ($P \leq 0.05$) at higher Si levels of T4 and T5 ($150 \text{ kg}\cdot\text{ha}^{-1}$ and $225 \text{ kg}\cdot\text{ha}^{-1}$) as compared to lower levels of Si application of T1, T2, and T3 ($0 \text{ kg}\cdot\text{ha}^{-1}$, $45 \text{ kg}\cdot\text{ha}^{-1}$, and $90 \text{ kg}\cdot\text{ha}^{-1}$). There were no significant differences between the treatments of T4 and T5 as well as among the treatments of T1, T2, and T3. The data (Table 4) showed that, at grain filling stage and milking stage, the values of g_s of maize significantly ($P < 0.05$) increased by Si application at levels of $90 \text{ kg}\cdot\text{ha}^{-1}$ (T3), $150 \text{ kg}\cdot\text{ha}^{-1}$ (T4), and $225 \text{ kg}\cdot\text{ha}^{-1}$ (T5); the values of g_s of maize were significantly enhanced with the increased dose of Si application of $90 \text{ kg}\cdot\text{ha}^{-1}$; there were significant differences between T3 and T4, and there was no significant

difference between T4 and T5. So at big trumpet stage and silking stage, the value of g_s of maize began to increase significantly ($P < 0.05$) when the dose of Si application got to the amount of $150 \text{ kg} \cdot \text{ha}^{-1}$; there was no significant difference in the values of g_s of maize between the treatments of doses $150 \text{ kg} \cdot \text{ha}^{-1}$ and $225 \text{ kg} \cdot \text{ha}^{-1}$; at grain filling stage and milk stage, the value of g_s of maize began to increase significantly ($P < 0.05$) when the dose of Si application got to the amount of $90 \text{ kg} \cdot \text{ha}^{-1}$, after which the values of g_s of maize were enhanced significantly with the increased dose of Si application; there was no significant difference between the treatments of doses $150 \text{ kg} \cdot \text{ha}^{-1}$ and $225 \text{ kg} \cdot \text{ha}^{-1}$.

3.4. Intercellular CO_2 Concentration (C_i). From big trumpet stage to milk stage, the data (Table 5) showed that the values of intercellular CO_2 concentration (C_i) of maize with the Si treatments of T3, T4, and T5 were significantly ($P \leq 0.05$) higher than those with T1 and T2; there was no significant difference between the treatments of T1 and T2. The values of C_i began to increase significantly ($P \leq 0.05$) when the dose of Si application got to the amount of $90 \text{ kg} \cdot \text{ha}^{-1}$, after which the values of C_i were dramatically enhanced with the increased dose of Si application; there were significant ($P \leq 0.05$) differences among the treatments of T3, T4, and T5. During big trumpet stage, comparing the values of C_i by Si application of T3, T4, and T5 with those by T1, it is shown that the former increased 13.1%, 17.1%, and 26.3%, respectively, compared to those of the latter; during silking stage, the former increased 16.6%, 20.6%, and 30.6%, respectively, compared to those of the latter; during grain filling stage, the former increased 14.1%, 26.5%, and 35.6%, respectively, compared to those of the latter; during milk stage, the former increased 13.5%, 25.5%, and 33.6% compared to those of the latter. There was no significant difference between the treatments of T1 and T2. So in these four studied stages, the value of C_i of maize began to increase significantly ($P < 0.05$) when the dose of Si application got to the amount of $90 \text{ kg} \cdot \text{ha}^{-1}$, after which the values of C_i increased dramatically with the increase of dose of Si applying.

4. Discussion

Under abiotic stresses such as toxicity, salinity, and lodging, silicon is reported to improve the growth of many kinds of higher plants [16]. Silicon can improve the growth of plants under salinity stress [17]. Exogenously applied Si significantly increased photosynthetic efficiency (A), stomatal conductance (g_s) and increased internal CO_2 concentration (C_i) in maize under saline conditions [18, 19]. Our results showed that, under the condition of saline-alkaline soil, the values of P_n , C_i , and g_s of maize leaves were significantly enhanced by Si application and that of E was decreased with the increase of Si supplied; similar improvement was reported in crops of strawberry, maize, Chinese cabbage, and rice [20–23]. The main mechanisms of improving crops growth by silicon lie in its functions of stimulation of photosynthesis, reduction of plant transpiration rate, and enhancement of tissue strength [24]. Our researches showed that, in the four

studied growth stages, the values of P_n by Si application were significantly enhanced compared with those with no Si application. Similar results were reported that addition of Si can enhance the photochemical efficiency of plants under salt stress [25]. Photosynthetic capacities of crops treated by Si application can be improved because the size of chloroplasts is enlarged and the number of grana in leaves is increased [26]. Researches on crops of barely (*Hordeum vulgare* L.), rice (*Oryza sativa* L.), sugarcane (*Saccharum officinarum* L.), and wheat (*Triticum aestivum* L.) showed that silicon deposited in leaves is helpful to improve the potential and efficiency of photosynthesis by opening angle of leaves, decreasing self-shading, and keeping the leaf erect [27], which play important roles in increasing P_n of crops.

According to our research, during the studied growth stages, the value of E of maize began to decrease significantly ($P < 0.05$) at the dose of $90 \text{ kg} \cdot \text{ha}^{-1}$ Si application, and above this dose of Si application the values of E decreased dramatically with the increased dose of Si application. By Si application, plant's internal water stress can be reduced; therefore, salt stress can be withstood as the rate of transpiration being influenced by the amount of Si gel associated with the cellulose in the cell walls of epidermal cells [28]. Similar results showed that water loss in maize can be reduced by Si application because Si can change the morphological structures of leaf epidermal cell [29]. Si is deposited in leaves leading towards decreased transpiration and hence dilutes salts accumulated in saline environment [30]. Unnecessary water loss can be limited through the epidermis, which is double layer, and silica combines with cellulose in the epidermal cells of leaf blade [31]. Another reason to explain the decrease of E is that the stomata opening can be influenced by Si [32]. Salt-stressed plants supplied with Si showed values of WUE 17% greater than those of salinized plants which were not supplied with Si by reducing the transpiration [30].

In our research, the value of g_s of maize began to increase significantly when the dose of Si application got to the amount of $150 \text{ kg} \cdot \text{ha}^{-1}$ at big trumpet stage and silking stage; the value of g_s of maize began to increase significantly when the dose of Si application got to the amount of $90 \text{ kg} \cdot \text{ha}^{-1}$, after which the values of g_s of maize were enhanced significantly with the increased dose of Si application at grain filling stage and milk stage. Similar reports on rice (*Oryza sativa* L.) are that Si application can enhance the stomatal conductance of rice plants subjected to salt stress, which shows silicate can reduce Na uptake via decline in the transpiration rate [30]. Silicon added to the saline growth medium improved photosynthetic activity [18, 19]. Si amendment enhanced the stomatal conductance of rice plants subjected to salt stress showing that silicate can reduce Na uptake via decline in the transpiration rate, which ultimately results into the reduction in growth and net photosynthesis [10]. It was also reported that g_s can be increased by Si fertilizer, because the increased g_s can regulate gas exchange, increase CO_2 uptake, and subsequently improve the capacity and efficiency of photosynthesis [18, 19, 33–39].

In the four studied growth stages, the value of C_i of maize began to increase significantly at the dose of $90 \text{ kg} \cdot \text{ha}^{-1}$ Si

application, after which the values of C_i increased dramatically with the increase of dose of Si application, which showed that photosynthetic efficiency of leaves can be enhanced by Si application [18]. Si applying can prevent the activities of photosynthetic enzymes in mesophyll cells from decreasing [34]. Si application reduces the transpiration rate to restrict the Na uptake; as a result, CO_2 intake is enhanced [10]. Under salt stress condition, the double membranes of chloroplasts disappeared, but membrane integrity was markedly improved in the salt treatment supplemented with Si [35]. The values of C_i can be increased dramatically by Si application under salt stress, which is stressful environment with the accumulation of ROS, such as superoxide radicals ($\text{O}_2^{\bullet-}$), hydroxyl radicals ($\text{OH}^\bullet$), and hydrogen peroxide (H_2O_2), and the activity of defense system affected by salinity stress may be enhanced by Si application [10, 19, 36–43].

5. Conclusion

Silicon plays an important role to enhance photosynthesis ability and efficiency of plants under salinity stress [10]. The field research showed that values of photosynthetic rate (P_n), stomatal conductance (g_s), transpiration rate (E), and intercellular CO_2 concentration (C_i) of maize plants in saline-alkaline soil were affected by Si application and salinity stress can be alleviated by Si application. In our research, the optimal dose of Si application on saline-alkaline soil was $150 \text{ kg} \cdot \text{ha}^{-1}$, under which photosynthetic ability of maize was greatly increased at studied growth stages. So in this research, increase in maize photosynthesis under saline-alkaline stress took place by Si application with proper doses, with which it is helpful to improve growth and yield of cereal crops [29].

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

This research was supported by the National Natural Science Foundation of China (31271672; 41301314; 41201559), the National Science and Technology Support Program of China (2012BAD42B01), the Natural Science Foundation of Jiangsu Province (BK2011412), and the CAS/SAFEA International Partnership Program for Creative Research Teams.

References

- [1] Y. Liang, Q. Chen, Q. Liu, W. Zhang, and R. Ding, "Exogenous silicon (Si) increases antioxidant enzyme activity and reduces lipid peroxidation in roots of salt-stressed barley (*Hordeum vulgare* L.)," *Journal of Plant Physiology*, vol. 160, no. 10, pp. 1157–1164, 2003.
- [2] C. N. Deng, G. X. Zhang, X. L. Pan, and K. Y. Zhao, "Chlorophyll fluorescence and gas exchange responses of maize seedlings to saline-alkaline stress," *Bulgarian Journal of Agricultural Science*, vol. 16, no. 1, pp. 49–58, 2010.
- [3] T. A. Ishida, K. Nara, S. Ma, T. Takano, and S. Liu, "Ectomycorrhizal fungal community in alkaline-saline soil in northeastern China," *Mycorrhiza*, vol. 19, no. 5, pp. 329–335, 2009.
- [4] I. Afzal, S. M. A. Basra, T. E. Lodhi, and S. J. Butt, "Improving germination and seedling vigour in wheat by halopriming under saline conditions," *Pakistan Journal of Agricultural Sciences*, vol. 44, no. 1, pp. 40–49, 2007.
- [5] R. Anjum, A. Ahmed, R. Ullah, M. Jahangir, and M. Yousaf, "Effect of soil salinity/sodicity on the growth and yield of different varieties of cotton," *International Journal of Agriculture and Biology*, vol. 7, no. 4, pp. 606–608, 2005.
- [6] Z. Khodarahmpour, M. Ifar, and M. Motamedi, "Effects of NaCl salinity on maize (*Zea mays* L.) at germination and early seedling stage," *African Journal of Biotechnology*, vol. 11, no. 2, pp. 298–304, 2012.
- [7] A. L. Tuna, C. Kaya, D. Higgs, B. Murillo-Amador, S. Aydemir, and A. R. Girgin, "Silicon improves salinity tolerance in wheat plants," *Environmental and Experimental Botany*, vol. 62, no. 1, pp. 10–16, 2008.
- [8] J. F. Ma and N. Yamaji, "Silicon uptake and accumulation in higher plants," *Trends in Plant Science*, vol. 11, no. 8, pp. 392–397, 2006.
- [9] A. C. Marafon and L. Endres, "Silicon: fertilization and nutrition in higher plants," *Amazonian Journal of Agricultural and Environmental Sciences*, vol. 56, no. 4, pp. 380–388, 2013.
- [10] A. Ali, S. M. A. Basra, S. Hussain, J. Iqbal, M. A. A. H. A. Bukhsh, and M. Sarwar, "Salt stress alleviation in field crops through nutritional supplementation of silicon," *Pakistan Journal of Nutrition*, vol. 11, no. 8, pp. 735–735, 2012.
- [11] Y. Liang, J. Si, and V. Römhild, "Silicon uptake and transport is an active process in *Cucumis sativus*," *New Phytologist*, vol. 167, no. 3, pp. 797–804, 2005.
- [12] S. Cha-Um and C. Kirdmanee, "Effect of salt stress on proline accumulation, photosynthetic ability and growth characters in two maize cultivars," *Pakistan Journal of Botany*, vol. 41, no. 1, pp. 87–98, 2009.
- [13] S. Covshoff and J. M. Hibberd, "Integrating C_4 photosynthesis into C_3 crops to increase yield potential," *Current Opinion in Biotechnology*, vol. 23, no. 2, pp. 209–214, 2012.
- [14] S. Q. Zhang, *Plant Physiology Experimental Techniques Tutorial*, Science Press, Beijing, China, 2011, (Chinese).
- [15] H.-W. Xu, F.-B. Song, X.-C. Zhu, and S.-Y. Tong, "Photosynthesis, chlorophyll fluorescence and nonstructural carbohydrates changes in husk leaves of maize in black soils region of Northeast China," *African Journal of Agricultural Research*, vol. 5, no. 9, pp. 785–791, 2010.
- [16] E. Epstein, "Silicon," *Annual Review of Plant Biology*, vol. 50, pp. 641–664, 1999.
- [17] Y. Liang, Q. Shen, Z. Shen, and T. Ma, "Effects of silicon on salinity tolerance of two barley cultivars," *Journal of Plant Nutrition*, vol. 19, no. 1, pp. 173–183, 1996.
- [18] N. Parveen and M. Ashraf, "Role of silicon in mitigating the adverse effects of salt stress on growth and photosynthetic attributes of two maize (*Zea mays* L.) cultivars grown hydroponically," *Pakistan Journal of Botany*, vol. 42, no. 3, pp. 1675–1684, 2010.
- [19] K. Yan, P. Chen, H. Shao, C. Shao, S. Zhao, and M. Brestic, "Dissection of photosynthetic electron transport process in sweet sorghum under heat stress," *PLoS ONE*, vol. 8, no. 5, Article ID e62100, 2013.

- [20] C. Gao, J. B. Liu, H. B. Chang, X. B. Yu, and H. S. Xu, "Effects of silicon on rice leaf photosynthesis and ultrastructure," *Journal of Jilin Agricultural University*, vol. 33, no. 1, pp. 3049–3057, 2011 (Chinese).
- [21] X. D. Yang, *Effect of N Si fertilizer on the growth and yield of Chinese cabbage [M.S. thesis]*, Shandong Agricultural University, 2010 (Chinese).
- [22] L. Z. Shu and H. Y. Liu, "Effects of silicon on growth of maize seedlings under salt stress," *Agro-Environmental Protection*, vol. 20, no. 1, pp. 38–40, 2001.
- [23] J. Y. Wang, M. D. Liu, and D. Li, "Effects of silicon enrichment on photosynthetic characteristics and yield of strawberry," *Northern Horticulture*, no. 12, pp. 90–92, 2009 (Chinese).
- [24] J. F. Ma and E. Takahashi, *Soil, Fertilizer, and Plant Silicon Research in Japan*, Elsevier Science, Amsterdam, The Netherlands, 2002.
- [25] A. Ali, S. M. A. Basra, J. Iqbal et al., "Silicon mediated biochemical changes in wheat under salinized and non-salinized solution cultures," *African Journal of Biotechnology*, vol. 11, no. 3, pp. 606–615, 2012.
- [26] X. D. Yang, *Effect of N Si fertilizer on the growth and yield of Chinese cabbage [M.S. thesis]*, Shandong Agricultural University, 2010, (Chinese).
- [27] R. P. Soratto, C. A. C. Crusciol, G. S. A. Castro, C. H. M. da Costa, and J. F. Neto, "Leaf application of silicic acid to white oat and wheat," *Revista Brasileira de Ciencia do Solo*, vol. 36, no. 5, pp. 1538–1544, 2012.
- [28] T. Aziz, M. A. Gill, and R. Rahmathallah, "Silicon nutrition and crop production: a review," *Pakistan Journal of Agricultural Sciences*, vol. 39, no. 3, pp. 181–187, 2002.
- [29] J. Ren, J. R. Guo, X. Q. Xing, G. Qi, and Z. L. Yuan, "Preliminary exploration into yield increase effects and yield increase mechanism of silicate fertilizer on maize," *Journal of Maize Sciences*, vol. 10, no. 2, pp. 84–86, 2002 (Chinese).
- [30] A. Ali, S. M. A. Basra, S. Hussain, J. Iqbal, M. A. A. H. A. Bukhsh, and M. Sarwar, "Salt stress alleviation in field crops through nutritional supplementation of silicon," *Pakistan Journal of Nutrition*, vol. 11, no. 8, pp. 637–655, 2012.
- [31] J. Lewin and B. E. F. Reimann, "Silicon and plant growth," *Annual Review of Plant Physiology*, vol. 20, pp. 289–304, 1969.
- [32] S. Agarie, H. Uchida, W. Agata, F. Kubota, and P. B. Kaufman, "Effects of silicon on transpiration and leaf conductance in rice plants (*Oryza sativa* L.)," *Plant Production Science*, vol. 1, no. 2, pp. 89–95, 1998.
- [33] K. Kusumi, S. Hirotaka, T. Kumamaru, and K. Iba, "Increased leaf photosynthesis caused by elevated stomatal conductance in a rice mutant deficient in SLAC1, a guard cell anion channel protein," *Journal of Experimental Botany*, vol. 63, no. 15, pp. 5635–5644, 2012.
- [34] B. L. Cao, K. Xu, J. Shi, G. F. Xin, C. Y. Liu, and X. Li, "Effects of silicon on growth, photosynthesis and transpiration of tomato," *Plant Nutrition and Fertilizer Science*, vol. 19, no. 2, pp. 354–360, 2013 (Chinese).
- [35] L. Yongchao, "Effect of silicon on leaf ultrastructure, chlorophyll content and photosynthetic activity of barley under salt stress," *Pedosphere*, vol. 8, no. 4, pp. 289–296, 1998.
- [36] Y. P. Song, *Effects of exogenous Si on growth and physiological characteristics of cucumber seedlings under NO_3^- stress [M.S. thesis]*, Shandong Agricultural University, 2011, (Chinese).
- [37] W. Song, A. Zheng, H. Shao, L. Chu, M. Brestic, and Z. Zhang, "The alleviative effect of salicylic acid on the physiological indices of the seedling leaves in six different wheat genotypes under lead stress," *Plant OMICS*, vol. 5, no. 5, pp. 486–493, 2012.
- [38] M. Brestic, M. Zivcak, H. M. Kalaji, R. Carpentier, and S. I. Allakhverdiev, "Photosystem II thermostability *in situ*: environmentally induced acclimation and genotype-specific reactions in *Triticum aestivum* L.," *Plant Physiology and Biochemistry*, vol. 57, pp. 93–105, 2012.
- [39] W. Tan, Q. W. Meng, M. Brestic, K. Olsovska, and X. Yang, "Photosynthesis is improved by exogenous calcium in heat-stressed tobacco plants," *Journal of Plant Physiology*, vol. 168, no. 17, pp. 2063–2071, 2011.
- [40] S. Hong-Bo, L. Zeng-Hui, Z. Zheng-Bin, C. Qi-Jie, C. Li-Ye, and M. Brestic, "Biological roles of crop NADP-malic enzymes and molecular mechanisms involved in abiotic stress," *African Journal of Biotechnology*, vol. 10, no. 25, pp. 4947–4953, 2011.
- [41] M. Zivcak, H. M. Kalaji, H.-B. Shao, K. Olsovska, and M. Brestic, "Photosynthetic proton and electron transport in wheat leaves under prolonged moderate drought stress," *Journal of Photochemistry and Photobiology B: Biology*, vol. 137, pp. 107–115, 2014.
- [42] H. M. Kalaji, K. Bosa, J. Kościelniak, and K. Żuk-Golaszewska, "Effects of salt stress on photosystem II efficiency and CO_2 assimilation of two Syrian barley landraces," *Environmental and Experimental Botany*, vol. 73, no. 1, pp. 64–72, 2011.
- [43] J. Kościelniak, A. Ostrowska, J. Biesaga-Kościelniak et al., "The effect of zearalenone on PSII photochemical activity and growth in wheat and soybean under salt (NaCl) stress," *Acta Physiologiae Plantarum*, vol. 33, no. 6, pp. 2329–2338, 2011.