Research Article

The Diophantine Equation $8^x + p^y = z^2$

Lan Qi¹ and Xiaoxue Li²

¹College of Mathematics and Statistics, Yulin University, Yulin, Shaanxi 719000, China

²School of Mathematics, Northwest University, Xi'an, Shaanxi 710127, China

Correspondence should be addressed to Xiaoxue Li; lx20072012@163.com

Received 7 July 2014; Accepted 14 December 2014

Academic Editor: Abdelghani Bellouquid

Copyright © 2015 L. Qi and X. Li. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let p be a fixed odd prime. Using certain results of exponential Diophantine equations, we prove that (i) if $p \equiv \pm 3 \pmod{8}$, then the equation $8^x + p^y = z^2$ has no positive integer solutions (x, y, z) ; (ii) if $p \equiv 7 \pmod{8}$, then the equation has only the solutions $(p, x, y, z) = (2^q - 1, (1/3)(q + 2), 2, 2^q + 1)$, where q is an odd prime with $q \equiv 1 \pmod{3}$; (iii) if $p \equiv 1 \pmod{8}$ and $p \neq 17$, then the equation has at most two positive integer solutions (x, y, z) .

1. Introduction

Let $\mathbb{Z}, \mathbb{N}$ be the sets of all integers and positive integers, respectively. Let p be a fixed odd prime. Recently, the solutions (x, y, z) of the equation

$$8^x + p^y = z^2, \quad x, y, z \in \mathbb{N} \quad (1)$$

were determined in the following cases:

- (1) (Sroysang [1]) if $p = 19$, then (1) has no solutions;
- (2) (Sroysang [2]) if $p = 13$, then (1) has no solutions;
- (3) (Rabago [3]) if $p = 17$, then (1) has only the solutions $(x, y, z) = (1, 1, 5), (2, 1, 9)$, and $(3, 1, 23)$.

In this paper, using certain results of exponential Diophantine equations, we prove a general result as follows.

Theorem 1. *If $p \equiv \pm 3 \pmod{8}$, then (1) has no solutions (x, y, z) . If $p \equiv 7 \pmod{8}$, then (1) has only the solutions*

$$(p, x, y, z) = \left(2^q - 1, \frac{q+2}{3}, 2, 2^q + 1\right), \quad (2)$$

where q is an odd prime with $q \equiv 1 \pmod{3}$.

If $p \equiv 1 \pmod{8}$ and $p \neq 17$, then (1) has at most two solutions (x, y, z) .

Obviously, the above theorem contains the results of [1, 2]. Finally, we propose the following conjecture.

Conjecture 2. *If $p \neq 17$, then (1) has at most one solution (x, y, z) .*

2. Preliminaries

Lemma 3. *If $2^n - 1$ is a prime, where n is a positive integer, then n must be a prime.*

Proof. See Theorem 1.10.1 of [4]. □

Lemma 4. *If p is an odd prime with $p \equiv 1 \pmod{4}$, then the equation*

$$u^2 - pv^2 = -1, \quad u, v \in \mathbb{N} \quad (3)$$

has solutions (u, v) .

Proof. See Section 8.1 of [5]. □

Lemma 5. *The equation*

$$X^2 - 2^m = Y^n, \quad X, Y, m, n \in \mathbb{N}, \quad \gcd(X, Y) = 1, \quad Y > 1, \quad m > 1, \quad n > 2 \quad (4)$$

has only the solution $(X, Y, m, n) = (71, 17, 7, 3)$.

Proof. See Theorem 8.4 of [6]. □

Lemma 6. Let D be a fixed odd positive integer. If the equation

$$u^2 - Dv^2 = -1, \quad u, v \in \mathbb{N} \quad (5)$$

has solutions (u, v) , then the equation

$$X^2 - D = 2^n, \quad X, n \in \mathbb{N}, \quad n > 2 \quad (6)$$

has at most two solutions (X, n) , except the following cases:

- (i) $D = 2^{2r} - 3 \cdot 2^{r+1} + 1$, $(X, n) = (2^r - 3, 3)$, $(2^r - 1, r + 2)$, $(2^r + 1, r + 3)$, and $(3 \cdot 2^r - 1, 2r + 3)$, where r is a positive integer with $r \geq 3$;
- (ii) $D = ((1/3)(2^{2r+1} - 17))^2 - 32$, $(X, n) = ((1/3)(2^{2r+1} - 17), 5)$, $(1/3)(2^{2r+1} + 1, 2r + 3)$, and $((1/3)(17 \cdot 2^{2r+1} - 1), 4r + 7)$, where r is a positive integer with $r \geq 3$;
- (iii) $D = 2^{2r_1} + 2^{2r_2} - 2^{r_1+r_2+1} - 2^{r_1+1} - 2^{r_2+1} + 1$, $(X, n) = (2^{r_2} - 2^{r_1} - 1, r_1 + 2)$, $(2^{r_2} - 2^{r_1} + 1, r_2 + 2)$, and $(2^{r_2} + 2^{r_1} - 1, r_1 + r_2 + 2)$, where r_1, r_2 are positive integers with $r_2 > r_1 + 1 > 2$.

Proof. See [7]. $\square$

Lemma 7. If D is an odd prime and D belongs to the exceptional case (i) of Lemma 6, then $D = 17$.

Proof. We now assume that D is an odd prime with $D = 2^{2r} - 3 \cdot 2^{r+1} + 1$. Then we have

$$(2^r - 1)^2 - 2^{r+2} = D, \quad (7)$$

$$(2^r + 1)^2 - 2^{r+3} = D. \quad (8)$$

If $2 \mid r$, since $r \geq 3$, then $r \geq 4$, and by (7), we have

$$(2^r - 1) + 2^{r/2+1} = D, \quad (2^r - 1) - 2^{r/2+1} = 1. \quad (9)$$

But, by the second equality of (9), we get $1 \equiv (2^r - 1) - 2^{r/2+1} \equiv -1 \pmod{8}$, a contradiction.

If $2 \nmid r$, then from (8) we get

$$(2^r + 1) + 2^{(r+3)/2} = D, \quad (2^r + 1) - 2^{(r+3)/2} = 1. \quad (10)$$

Further, by the second equality of (10), we have $2^r = 2^{(r+3)/2}$, $r = 3$, and $D = 17$. Thus, the lemma is proved. $\square$

Lemma 8. If D is an odd prime and D belongs to the exceptional case (iii) of Lemma 6, then $D = 17$.

Proof. Using the same method as in the proof of Lemma 7, we can obtain this lemma without any difficulty. $\square$

Lemma 9. If D belongs to the exceptional case (ii), then (6) has at most one solution (X, n) with $3 \mid n$.

Proof. Notice that, for any positive integer r , there exists at most one number of 5 , $2r + 3$, and $4r + 7$ which is a multiple of 3 . Thus, by Lemma 6, the lemma is proved. $\square$

Lemma 10. The equation

$$X^m - Y^n = 1, \quad X, Y, m, n \in \mathbb{N}, \quad \min\{X, Y, m, n\} > 1 \quad (11)$$

has only the solution $(X, Y, m, n) = (3, 2, 2, 3)$.

Proof. See [8]. $\square$

3. Proof of Theorem

We now assume that (x, y, z) is a solution of (1). Then we have $\gcd(2p, z) = 1$.

If $2 \mid y$, since $\gcd(z + p^{y/2}, z - p^{y/2}) = 2$, then from (1) we get

$$z + p^{y/2} = 2^{3x-1}, \quad z - p^{y/2} = 2, \quad (12)$$

where we obtain

$$z = 2^{3x-2} + 1, \quad (13)$$

$$p^{y/2} = 2^{3x-2} - 1. \quad (14)$$

Since $p > 1$, applying Lemma 10 to (14), we get

$$y = 2, \quad p = 2^{3x-2} - 1. \quad (15)$$

Further, by Lemma 3, we see from the second equality of (15) that

$$p = 2^q - 1, \quad q = 3x - 2 \quad (16)$$

is an odd prime with $q \equiv 1 \pmod{3}$.

Therefore, by (13), (15), and (16), we obtain the solutions given in (2).

Obviously, if p satisfies (2), then $p \equiv 7 \pmod{8}$. Otherwise, since $2 \nmid y$, we see from (1) that $p \equiv p^y \equiv z^2 - 8^x \equiv 1 \pmod{8}$. It implies that if $p \equiv \pm 3 \pmod{8}$, then (1) has no solutions (x, y, z) . If $p \equiv 7 \pmod{8}$, then (1) has only the solutions (2).

Here and below, we consider the remaining cases that $p \equiv 1 \pmod{8}$. By the above analysis, we have $2 \nmid y$. If $y > 1$, then $y \geq 3$ and (4) has the solution $(X, Y, m, n) = (z, p, 3x, y)$ with $3 \mid m$. But, by Lemma 5, it is impossible. Therefore, we have

$$y = 1. \quad (17)$$

Substituting (17) into (1), the equation

$$X^2 - p = 2^n, \quad X, n \in \mathbb{N}, \quad n > 2 \quad (18)$$

has the solution $(X, n) = (z, 3x)$ with $3 \mid n$. Since $p \equiv 1 \pmod{8}$, by Lemma 4, (3) has solutions (u, v) . Therefore, by Lemmas 6–9, (1) has at most two solutions (x, y, z) . Thus, the theorem is proved.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgment

This work is supported by the National Natural Science Foundation of China (11371291).

References

- [1] B. Sroysang, "More on the diophantine equation $8^x + 19^y = z^2$," *International Journal of Pure and Applied Mathematics*, vol. 81, no. 4, pp. 601–604, 2012.
- [2] B. Sroysang, "On the Diophantine equation $8^x + 13^y = z^2$," *International Journal of Pure and Applied Mathematics*, vol. 90, no. 1, pp. 69–72, 2014.
- [3] J. F. T. Rabago, "On an open problem by B. Sroysang," *Konuralp Journal of Mathematics*, vol. 1, no. 2, pp. 30–32, 2013.
- [4] L. G. Hua, *An Introduction to Number Theory*, Science Press, Beijing, China, 1979.
- [5] L. J. Mordell, *Diophantine Equations*, Academic Press, London, UK, 1969.
- [6] M. A. Bennett and C. M. Skinner, "Ternary Diophantine equations via Galois representations and modular forms," *Canadian Journal of Mathematics*, vol. 56, no. 1, pp. 23–54, 2004.
- [7] M.-H. Le, "On the number of solutions of the generalized Ramanujan-Nagell equation $x^2 - D = 2^{n+2}$," *Acta Arithmetica*, vol. 60, no. 2, pp. 149–167, 1991.
- [8] P. Mihăilescu, "Primary cyclotomic units and a proof of Catalan's conjecture," *Journal für die Reine und Angewandte Mathematik*, vol. 2004, no. 572, pp. 167–195, 2004.