Research Article

Effects of a Supraseasonal Drought on the Ecological Attributes of *Plagioscion squamosissimus* (Heckel, 1840) (Pisces, Sciaenidae) in a Brazilian Reservoir

Antonia E. F. Souza,¹ Jônata F. Oliveira,¹ Danielle Peretti,² Rodrigo Fernandes,³ Rodrigo S. Costa,¹ and José Luis Costa Novaes¹

¹Departamento de Ciências Animais, Laboratório de Ecologia de Peixes e Pesca Continental (UFERSA), BR 110, Km 47, Bairro Costa e Silva, 59625-900 Mossoró, RN, Brazil

²Laboratório de Ictiologia, Universidade do Estado do Rio Grande do Norte (UERN), Av. Prof. Antônio Campos s/n, Bairro Costa e Silva, 59625-620 Mossoró, RN, Brazil

³Departamento de Ciências Animais, Laboratório de Ecologia Quantitativa (UFERSA), BR 110, Km 47, Bairro Costa e Silva, 59625-900 Mossoró, RN, Brazil

Correspondence should be addressed to José Luis Costa Novaes; novaes@ufersa.edu.br

Received 19 October 2016; Revised 7 January 2017; Accepted 29 January 2017; Published 23 February 2017

Academic Editor: Joao P. Barreiros

Copyright © 2017 Antonia E. F. Souza et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The aim of this study was to evaluate the effect of a supraseasonal drought on the ecological attributes of *Plagioscion squamosissimus*. The fish were caught quarterly from February 2010 to November 2014 using gill nets in the reservoir of Santa Cruz, Rio Grande do Norte, Brazil. The abundance of the species was evaluated with the catch per unit effort (CPUE) metric and then correlated with the accumulated rainfall and water volume of the reservoir. The diet of the fish was evaluated using the feeding index (IAi). The proportional similarity index (PS_i) was used to evaluate the variation in the niches of the fish. The body condition was inferred through the relative condition factor, and its variation was assessed with ANOVA. A reduction in the abundance of the species that were positively correlated with the reservoir water volume was observed. The diet of the fish comprised shrimp, gastropods, fish, insects, shrimp larvae, and vegetable matter, with shrimp being the major component. PS_i showed the occurrence of individual specialization during November 2013 and November 2014. The relative condition factor was not correlated with a reduction in the water volume of the reservoir. The supraseasonal drought did not affect the relative condition factor, diet, and the trophic niche, but it did affect the species abundance.

1. Introduction

Disturbances are often abrupt and sometimes random and are responsible for major damage to ecosystems, causing changes in the availability of resources and an impact on biological communities, such as the mortality of individuals [1, 2]. Disturbances can be of natural or anthropogenic origin, and their effects have repercussions from the individual to the ecosystem level [3]. Disturbances such as droughts are increasingly common in several regions of the world, particularly in arid and semiarid regions [4]. Among the effects of droughts on aquatic organisms are changes in the

body condition of individuals, a decrease in the abundance and biomass of species, a decrease in the availability of resources, and intensification of competition and predation [5–7].

Directly or indirectly, disturbances influence the niches of species [8]. Despite the niche historically being considered a species or population attribute, it was noted that several species, rather than being categorized as generalists, in fact comprise individuals that only use the niche of a population subgroup [9]. One of the many factors influencing the intrapopulation variation of niches is individual specialization. Such specialization is found in a range of species,

including both vertebrates and invertebrates [10]. An individual specialist is an individual that has adopted a narrower niche compared with the general population [10]. Among the factors driving niche creation is the variation in the diet among individuals, which highlights the importance of ecological interactions (e.g., intra- and interspecific competition, predation) to the development of niches [11].

The South American silver croaker, *Plagioscion squamosissimus* (Heckel, 1840), is a carnivorous Amazonian species of the Sciaenidae family (Owen, 1846) [12], whose diet consists primarily of fish and crustaceans [13]. The distribution of this species has been increasing because of its introduction into many Brazilian reservoirs, mainly to promote fisheries [14]. This species is one of the most abundant in semiarid reservoirs [15] and of the main fishery resources in the region, representing an important source of income and animal protein for human populations living close to the reservoirs [16].

The aim of this study was to evaluate the ecological responses of *P. squamosissimus* in a reservoir in the Brazilian semiarid region to drought conditions. The aspects analyzed were as follows: (1) the catch per unit effort (CPUE) in number and biomass; (2) the correlations among accumulated rainfall, reservoir water volume, and CPUE; (3) the occurrence of individual specialization; and (4) the relative condition factor. The hypotheses tested were as follows: (1) the species abundance and relative condition factor decrease with drought conditions and (2) the amplitude of the niche diversity is lower during the drought period.

2. Materials and Methods

2.1. Study Area. The hydrographic basin of the Oriental Northeast Atlantic occupies an area of 287,348 km² and is characterized by caatinga vegetation and a semiarid climate; the basin has average annual rainfall of 750 mm, predominantly concentrated between February and May [17]. However, during the last 6 years, a decrease in the annual rainfall has been observed, with the exception of the years 2009 and 2011 (Figure 1). Among the small hydrographic basins located in the hydrographic basin of the Oriental Northeast Atlantic is the Apodi/Mossoró basin, with an area of 14,271 km². The basin includes 20 important reservoirs, all with water volumes above 5 m³, constructed for human uses, including agriculture [18]. The Santa Cruz of Apodi Reservoir (05°45'45" S and 37°48'00" W; Figure 2), in the Apodi/Mossoró basin, is classified as oligomesotrophic and has an area of 341.3 ha and the capacity to accumulate up to 600 million m³ of water [19].

2.2. Data Collection. Data were collected quarterly between February 2010 and November 2014 at eight sampling points distributed along the reservoir (Figure 2). Fish were caught using gill nets placed at the reservoir bottom covering a total area of 301.8 m²; the nets had a mesh size of between 12 and 70 mm and were 15 m long and between 1.8 and 2.0 m high. At each point, the nets were set at 05:00 p.m. and removed at 05:00 a.m. the following day. For each individual

FIGURE 1: Variation in annual precipitation in the Santa Cruz of Apodi Reservoir, Brazil, between 2009 and 2014 [21].

collected, the total length (cm) (Lt) and weight (g) (Wt) were measured. The specimens were dissected, and only stomachs with food content were removed, fixed in 4% formaldehyde, and preserved in alcohol; the stomach contents were subsequently analyzed with a stereoscopic microscope. For the feeding study, only individuals caught between February 2011 and November 2014 were analyzed. The food items found were identified with the aid of specialized literature up to the lowest possible taxonomic level and then were grouped taxonomically for the analyses. The volume of the food items was measured by liquid displacement in a graduated cylinder [20]. The accumulated rainfall data (mm) and the accumulated water volume of the reservoir (in %) were used to infer the responses of *P. squamosissimus* to drought; data were obtained from the offices of the Agricultural Research Company of Rio Grande do Norte [21] in the city of Apodi and the Secretariat of the State of the Environment and Hydric Resources [17].

2.3. Data Analyses. To estimate the fish abundance, the numeric and biomass data of the fish captured were transformed into CPUE using the following formula: $CPUE = C/f$, where C is biomass (CPUEb) or N number of captured individuals (CPUEn) and f is effort in m² * h, where m² is 301.8 m² and h is 12 h. The differences in the CPUE among months were tested with a one-way analysis of variance (ANOVA) followed by the Tukey test. The Pearson correlation coefficients between the water volume of the reservoir and accumulated rainfall with the CPUEb and CPUEn were calculated. For the study of the stomach contents, 382 specimens that presented food in the stomach were analyzed using the feeding index (IAi) [20], which combines the frequency of occurrence and volumetric methods [22, 23]. To evaluate whether the diet changed between months, the IAi values were first ordered by nonmetric multidimensional scaling (NMDS) [24] and then the differences between months were tested by permutational multivariate analysis of variance (PERMANOVA) [25]. The intrapopulation variation in trophic niche and individual specialization was quantified using the proportional similarity index (PS_i), which calculates the degree of overlap between the diet of individuals and

FIGURE 2: Location of the Santa Cruz of Apodi Reservoir in the hydrographic Apodi/Mossoró River basin, Brazil. The black line indicates the dam, and numbers (1–8) indicate the collection sites.

the population as a whole [9]. This metric is expressed by the value of each individual (PS_i) and the average population (IS) [9]:

$$PS_i = 1 - 0.5 \sum_j |p_{ij} - q_j| = \sum_j \min(p_{ij}, q_j), \quad (1)$$

where p_{ij} is the frequency of prey type j in the diet of an individual and q_j is the frequency of prey type j in the diet of the population in general. If the amplitude of the IS reaches 1, the consumption of resources by individuals is equal to that of the population as a whole and there is an absence of specialization, whereas values close to 0 indicate strong specialization [26]. Values of probability (P) were obtained using the Monte Carlo procedure (999 simulations). The body condition was inferred by the relative condition factor calculated from the length-weight relationship variables. Body condition infers the changes in body weight based on the increase in length, with the relationship determined according to the following equation: $Wt = a \times Lt^b$, where Wt is the total weight (g), Lt is the total length (cm), a is the intercept, and b is the slope [27]. Differences in the relative condition factor among months were evaluated with a one-way ANOVA followed by a Tukey test; Pearson correlation coefficients were calculated to test the relationship between the relative condition factor and water volume of the reservoir. Statistical analyses were conducted using R software (Development Core Team, 20100) at a significance level of $P < 0.05$.

FIGURE 3: The quarterly variation in precipitation and water volume in the Santa Cruz of Apodi Reservoir, Brazil, between February 2010 and November 2014.

3. Results

A considerable decrease in the water level was evident in the Santa Cruz of Apodi Reservoir during the study period, from 91.37% of its capacity in February 2010 to 43.40% of its capacity in November 2014. The precipitation ranged between 468 mm (May 2011) and 0 mm (November 2010, 2011, 2013, and 2014), with the highest amounts falling in February and May. The exception was in 2012, when precipitation was much lower than expected in February and May and the greatest reduction in water volume in the reservoir was recorded (Figure 3).

FIGURE 4: (a) Catch per unit effort estimated in biomass (CPUEb) and (b) number (CPUEn) of *Plagioscion squamosissimus* in the Santa Cruz of Apodi Reservoir, Brazil, between February 2010 and November 2014. The median is shown with a horizontal line inside the box. The minimum and maximum values are shown with short horizontal lines ("whiskers"). Different letters indicate statistical differences (Tukey, $P < 0.05$).

FIGURE 5: Quarterly percentages of the feeding index (IAi) for each food item category of the stomach contents of *Plagioscion squamosissimus* in the Santa Cruz of Apodi Reservoir, Brazil, between February 2011 and November 2014.

A total of 1,273 specimens were collected with a TL between 3.80 and 54.90 cm (average: 20.31; SD: 6.75) and a TW between 2.70 and 2,233.52 g (average: 120.31; SD: 141.95). The mean abundance in terms of CPUEb was 6,008.21 ($\text{g}/301.8 \text{ m}^2 \cdot 12 \text{ h}$) and that in terms of CPUEn was 50.02 ($\text{ind.}/301.8 \text{ m}^2 \cdot 12 \text{ h}$). The highest monthly average CPUE (i.e., the highest estimated abundance) based on the highest biomass and highest number of captured individuals was in February 2010: CPUEb = 157.43 ($\text{g}/301.8 \text{ m}^2 \cdot 12 \text{ h}$) and CPUEn = 1.18 ($\text{ind.}/301.8 \text{ m}^2 \cdot 12 \text{ h}$), respectively. The ANOVA was significant for CPUEb ($F = 3.1$; $\text{df} = 51.42$; $P < 0.05$) and CPUEn ($F = 1.86$; $\text{df} = 51.29$; $P = 0.04$) (Figure 4). The correlations between the reservoir water volume and CPUEb and CPUEn were positive and significant ($r = 0.67$;

$P = 0.01$ and $r = 0.50$; $P = 0.02$, resp.). In contrast, there were no significant correlations between the mean annual precipitation and the mean CPUEb and CPUEn ($r = 0.42$; $P = 0.06$ and $r = 0.29$; $P = 0.24$, resp.).

The diet of *P. squamosissimus* was composed of six items: (1) shrimp (*Macrobrachium amazonicum* Heller, 1862) (IAi = 84.6 (17.4), 42.76%–100%), (2) fish (scales, bones, and muscle) (IAi = 5.7 (10.2), 0%–40.72%), (3) insects (IAi = 8.3 (15.2), 0%–50.65%), (4) vegetable matter (IAi = 0.01 (0.03), 0%–0.36%), (5) gastropods (*Melanoides tuberculata* Müller, 1774) (IAi = 0.01 (0.04), 0%–0.21%), and (6) shrimp larvae (IAi = 1.3 (5.2), 0%–20.75%) (mean (standard deviation), minimum–maximum, resp.) (Figure 5). The NMDS ordination separated three groups: (i) months with high consumption of insects, specifically February 2011 (IAi = 33.62%), February 2012 (IAi = 50.65%), and February 2014 (IAi = 28.57%); (ii) months with high consumption of fish, specifically November 2013 (IAi = 40.72) and November 2014 (IAi = 8.54%); and (iii) other months with high consumption of shrimp (IAi < 69.82%) (Figure 6). Significant difference in IAi was found among months ($F = 4.33$; $P = 0.01$).

The individual specialization index (IS) ranged from 0.79 to 0.92. The IS values were close to 1, indicating a high degree of similarity in the diets of individuals. In general, the population comprised generalists, except during November 2013 (IS = 0.79; $P = 0.05$) and November 2014 (IS = 0.89; $P = 0.04$), when the presence of specialists was recorded (Figure 7).

The relative condition factor ranged from 0.009 in February 2014 to 0.039 in May 2012 (Figure 8). The average relative condition factor also differed significantly among years ($F = 278.9$; $\text{df} = 167.5$; $P < 0.05$), and the Tukey test indicated differences among months (Table 1). There was no significant correlation between the water volume of the reservoir and relative condition factor ($r = 0.13$; $P = 0.56$).

FIGURE 6: Nonmetric multidimensional scaling (NMDS – stress = 0.04; Axis 1 = 0.81 and Axis 2 = 0.29) of the monthly feeding index (IAi) of *Plagioscion squamosissimus* in the Santa Cruz of Apodi Reservoir, Rio Grande do Norte, Brazil, between February 2011 and November 2014. Feb: February; Aug: August; Nov: November; ○: 2011; □: 2012; ◇: 2013; ×: 2014.

FIGURE 7: Quarterly values of the individual specialization index (IS) of the diet of *Plagioscion squamosissimus* samples in the Santa Cruz of Apodi Reservoir, Brazil, between February 2011 and November 2014. * indicates $P < 0.05$.

4. Discussion

Although high precipitation occurred in 2009 and 2011, a large drop in the reservoir water volume occurred due to the rainfall scarcity from 2012 to 2015. The relationship of a reduction of the reservoir water volume with a decrease in the abundance of *P. squamosissimus* confirmed the hypothesis that the drought negatively affected the abundance of the species; these two variables were positively correlated. Similar results were described in the basin of the Guadalquivir River, Spain, where the water volume was the main environmental factor responsible for the reduction of the abundance of fish after an extended drought period [5]. A study of fish assemblages in streams in the Mira estuary, Portugal, similarly showed that the reduction in fish abundance was significantly influenced by water level [28]. As drought results in a reduction of the available habitat area, individuals of *P. squamosissimus* were possibly forced into a smaller volume

FIGURE 8: Variation in the relative condition factor of *Plagioscion squamosissimus* in the Santa Cruz of Apodi Reservoir, Rio Grande do Norte, Brazil, between February 2010 and November 2014. The median is shown with a horizontal line inside the box. The minimum and maximum values are shown with short horizontal lines (“whiskers”).

of water, whereby a reduction in the availability of prey resulted in a shift from a generalist to a specialist diet. In agreement with this process, ecological interactions together with disturbances can cause a reduction in the abundance of species and may even result in permanent local effects [29]. Moreover, drought can impair the reproductive success and initial development of fish [7]. Therefore, directly and indirectly, disturbances such as drought can result in changes in patterns of species abundance [6], whereas recovery will depend on the life history characteristics of the affected species [30] as well as the frequency and intensity of the disturbance [31].

The diet composition analysis showed that *P. squamosissimus* had predominantly consumed shrimp (*Macrobrachium amazonicum* Heller, 1862), resulting in a considerable overlap in the NMDS ordination. Similar to the present study, *P. squamosissimus* was found to consume mainly shrimp in the Piató Lagoon, Rio Grande do Norte [32]. The consumption of shrimp has been reported for more than 50 species of Amazonian fish, including *P. squamosissimus* [33, 34]. It is likely that some life history aspects of *P. squamosissimus* in the selection of their food partly explain such results. With respect to the selection of the type and size of prey, several factors are involved, including the limitations imposed by the predator morphology, such as the size of the mouth, and behavioral strategies, such as those involved in the hunting and killing of prey [35]. *Macrobrachium amazonicum* is an Amazonian species that can reach 30 cm in length and was introduced in northeastern reservoirs of Brazil [36, 37]. Its smaller body size compared with many other fish may be a factor affecting the food selection of *P. squamosissimus*. The relationship between the size of the prey and the predator has been noted as a key factor in determining the foraging success of aquatic communities [38]. An additional aspect that may have influenced the food selection of *P. squamosissimus* is the

TABLE 1: Tukey test for differences in the relative condition factor between months for *Plagioscion squamosissimus* in the Santa Cruz de Apodi Reservoir, Brazil, between February 2010 and November 2014. Significant differences are in italics.

	Feb/10	May/10	Aug/10	Nov/10	Feb/11	May/11	Aug/11	Nov/11	Feb/12	May/12	Aug/12	Nov/12	Feb/13	May/13	Aug/13	Nov/13	Feb/14	May/14	Aug/14	Nov/14
Feb/10																				
May/10	4.25E - 05																			
Aug/10	15.71	4.25E - 05																		
Nov/10	14.6	14.84	13.72	11.23	3.607	11.23	3.934	17.66	15.16	12.57	15.06	17.66	15.16	12.57	15.06	17.66	15.16	12.57	15.06	17.66
Feb/11	12.1	3.607	11.23	3.934	17.66	15.16	12.57	15.06	17.66	15.16	12.57	15.06	17.66	15.16	12.57	15.06	17.66	15.16	12.57	15.06
May/11	0.4659	16.18	13.41	15.06	12.57	2.592	0.9856	2.691	0.98409	4.25E - 05	0.9984	0.5716	1	0.557	0.001507	4.25E - 05	1	4.25E - 05	4.25E - 05	1
Nov/11	3.157	18.87	4.032	17.75	15.26	0.09856	2.691	2.086	4.679	38.87	36.28	38.97	4.777	0.009352	0.2482	5.65E - 05	1	4.25E - 05	4.25E - 05	0.9998
Feb/12	1.62	14.09	0.7448	12.98	10.48	4.679	38.87	36.28	38.97	4.777	0.009352	0.2482	5.65E - 05	1	4.25E - 05	4.25E - 05	1	4.25E - 05	4.25E - 05	0.9998
May/12	35.81	20.1	34.94	21.21	23.71	2.042	4.635	3.557	1.22	35.41	5.501	0.0478	1	0.1609	0.00835	4.26E - 05	1	4.25E - 05	4.25E - 05	0.04567
Aug/12	5.101	20.81	5.976	19.7	17.21	2.042	4.635	3.557	1.22	35.41	5.501	0.0478	1	0.1609	0.00835	4.26E - 05	1	4.25E - 05	4.25E - 05	0.04567
Nov/12	4.005	15.31	0.4749	14.2	11.7	3.459	0.8665	3.557	1.22	35.41	5.501	0.0478	1	0.1609	0.00835	4.26E - 05	1	4.25E - 05	4.25E - 05	0.04567
Feb/13	4.05	19.76	4.925	18.65	16.15	0.9912	3.584	0.8927	5.67	39.86	1.051	4.45	4.45	9.869	4.25E - 05	0.08968	0.000319	4.25E - 05	0.2726	0.3363
May/13	9.18	6.531	8.304	5.417	6.285	8.878	6.286	8.977	4.199	29.99	10.92	8.779	13.23	3.36	0.00835	4.25E - 05	1	4.25E - 05	4.25E - 05	0.005293
Aug/13	0.697	15.01	0.1784	13.9	11.41	3.755	1.163	3.854	0.9232	35.11	5.798	0.2965	4.747	5.123	8.483	4.28E - 05	4.25E - 05	4.25E - 05	4.25E - 05	1
Nov/13	10.83	26.54	11.7	25.42	22.93	7.769	10.36	7.671	12.45	46.64	5.727	11.23	6.778	16.65	20.01	11.52	4.25E - 05	4.25E - 05	4.25E - 05	1
Feb/14	10.83	26.54	11.7	25.42	22.93	7.769	10.36	7.671	12.45	46.64	5.727	11.23	6.778	16.65	20.01	11.52	4.25E - 05	4.25E - 05	4.25E - 05	1
May/14	25.42	9.71	24.55	10.82	13.32	28.48	25.89	28.58	23.8	10.39	30.52	25.02	29.47	19.6	16.24	24.72	36.25	4.25E - 05	4.25E - 05	4.25E - 05
Aug/14	8.19	23.9	9.065	22.79	20.29	5.132	7.724	5.033	9.81	44	3.089	8.591	4.14	14.01	17.37	8.887	2.638	33.61	4.25E - 05	4.37E - 05
Nov/14	0.05009	15.76	0.9255	14.65	12.15	3.008	0.4158	3.107	1.67	35.86	5.051	0.4506	4	5.87	9.23	0.7471	10.78	25.47	8.14	

fact that when compared with fish (a main food item for the species in many aquatic environments), shrimp require little investment in terms of time and energy in seeking, capturing, killing, and consuming [39, 40]. The large consumption of shrimp may also be related to the fact that shrimp have lower mobility than fish, a characteristic that affects the chance of an encounter between predator and prey [41, 42]. Thus, considering that the investment in time and energy tends to increase when prey items are large, the consumption of small prey items such as shrimp may reflect efficient foraging [38, 43, 44].

During months when high rainfall was recorded (February 2011 and 2014), food items brought in by the rainfall runoff (e.g., insects) increased the contribution of less common items to the fish diet, as evidenced by ordination analysis. Similarly, the diversity of food resources increases during the rainy season, with insects being of central importance, which is attributed primarily to the rising water level and the succession of aquatic plants [45]. During the drier months, insects also contributed to the diet, as strong environmental limitations on their abundance were absent. Insects tolerate periods of water scarcity [46], including those of the Chironomidae and Culicidae families, which can tolerate anoxic conditions [47, 48]. The NMDS analysis showed that, during November 2013 and November 2014, the diet of *P. squamosissimus* also differed due to higher consumption of fish and shrimp larvae. The efficiency of a species' ability to seek, capture, and manipulate prey is related to the size of the prey; therefore, smaller prey are considered a highly important food resource of piscivorous species in the Upper Paraná River floodplain [49]. Thus, the suprasonal drought did not affect the diet of *P. squamosissimus*. The differences observed in this study were related to cycle rain-dry of the region, even though it was weak during the studied period.

The high degree of similarity in the diets of individuals, mainly during November 2013 and November 2014, may have occurred due to the decrease in habitat area and the consequent ecological interactions such as intraspecific competition once individuals were forced into a restricted space. Ecological interactions can also promote changes in the niches of individuals [50]. Empirical and theoretical studies have shown that there is a positive association between intraspecific competition and individual specialization [11]. At the same time, although the premises postulated by optimal foraging theory assume that this interaction promotes an increase in the range of the niche for the population as whole as well as for individuals, many additional and yet unknown factors may restrict such an increase [51]. However, the present study did not find an association between drought and individual specialization, even though the reduction of the reservoir volume occurred over time and not only in a specific month. Moreover, in reservoirs, the feeding strategies of the predatory species are controlled to a larger degree by the availability and susceptibility of their prey than by their food preferences [52].

A decrease in the value of the fish condition factor is expected with a decrease in the water level that usually occurs during periods of drought [6, 53]; however, the relative

condition factor of *P. squamosissimus* was not correlated with the reduction of the water volume of the reservoir. Previous research in the reservoir showed that a reduction of the water volume did not affect the fish condition factor because of an increase in the availability of food of better nutritional value [7]. In addition, the reduction of population abundance decreases competition for food, preventing the condition factor from decreasing.

5. Conclusion

The first hypothesis of the study was partially accepted; the abundance of *P. squamosissimus* decreased with the reduction of the water volume of the reservoir caused by the drought. However, no decrease in the relative condition factor was observed. The second hypothesis was not supported; the amplitude of the niche was not lower during the drought period.

Competing Interests

The authors declare that they have no competing interests.

References

- [1] R. L. Mackey and D. J. Currie, "A re-examination of the expected effects of disturbance on diversity," *Oikos*, vol. 88, no. 3, pp. 483–493, 2000.
- [2] J. R. Svensson, *Ecological Disturbances: The Good, the Bad and the Ugly*, Printed by G. H. Tryck, Göteborgs Universitet, Göteborg, Sweden, 2010.
- [3] P. S. White and A. Jentsch, "The search for generality in studies of disturbance and ecosystem dynamics," *Progress in Botany*, vol. 62, pp. 399–450, 2001.
- [4] E. Jeppesen, S. Brucet, L. Naselli-Flores et al., "Ecological impacts of global warming and water abstraction on lakes and reservoirs due to changes in water level and related changes in salinity," *Hydrobiologia*, vol. 750, no. 1, pp. 201–227, 2015.
- [5] R. Bravo, M. C. Soriguer, N. Villar, and J. A. Hernando, "The dynamics of fish populations in the Palancar stream, a small tributary of the river Guadalquivir, Spain," *Acta Oecologica*, vol. 22, no. 1, pp. 9–20, 2001.
- [6] P. S. Lake, "Ecological effects of perturbation by drought in flowing waters," *Freshwater Biology*, vol. 48, no. 7, pp. 1161–1172, 2003.
- [7] F. Abujanra, A. A. Agostinho, and N. S. Hahn, "Effects of the flood regime on the body condition of fish of different trophic guilds in the Upper Paraná River floodplain, Brazil," *Brazilian Journal of Biology*, vol. 69, no. 2, pp. 469–479, 2009.
- [8] K. Shea, S. H. Roxburgh, and E. S. J. Rauschert, "Moving from pattern to process: coexistence mechanisms under intermediate disturbance regimes," *Ecology Letters*, vol. 7, no. 6, pp. 491–508, 2004.
- [9] D. I. Bolnick, L. H. Yang, J. A. Fordyce, J. M. Davis, and R. Svanbäck, "Measuring individual-level resource specialization," *Ecology*, vol. 83, no. 10, pp. 2936–2941, 2002.
- [10] D. I. Bolnick, R. Svanbäck, J. A. Fordyce et al., "The ecology of individuals: incidence and implications of individual specialization," *American Naturalist*, vol. 161, no. 1, pp. 1–28, 2003.

- [11] M. S. Araújo, D. I. Bolnick, and C. A. Layman, "The ecological causes of individual specialisation," *Ecology Letters*, vol. 14, no. 9, pp. 948–958, 2011.
- [12] L. Casatti, "Revision of the South American freshwater genus *Plagioscion* (Teleostei, Perciformes, Sciaenidae)," *Zootaxa*, vol. 1080, pp. 39–64, 2005.
- [13] N. S. Hahn, V. E. Loureiro, and R. L. Delariva, "Atividade alimentar da curvina *Plagioscion squamosissimus* (Heckel, 1840) (Perciformes, Sciaenidae) no rio Paraná," *Acta Scientiarum Biological Sciences*, vol. 21, pp. 309–314, 1999.
- [14] F. M. S. Braga, "Biologia reprodutiva de *Plagioscion squamosissimus* (Teleostei, Sciaenidae) na represa de Barra Bonita, Rio Piracicaba (SP)," *Revista UNIMAR*, vol. 19, pp. 447–460, 1997.
- [15] J. L. C. Novaes, S. I. L. Moreira, C. E. C. Freire, M. M. O. Sousa, and R. S. Costa, "Fish assemblage in a semi-arid Neotropical reservoir: composition, structure and patterns of diversity and abundance," *Brazilian Journal of Biology*, vol. 74, no. 2, pp. 290–301, 2014.
- [16] J. L. C. Novaes, A. E. Freire, R. D. R. A. Amorim, and R. D. S. Costa, "Diagnosis of the artisanal fisheries in a Brazilian semiarid reservoir," *Boletim do Instituto de Pesca*, vol. 41, no. 1, pp. 31–42, 2015.
- [17] SEMARH—Secretaria do Meio Ambiente e dos Recursos Hídricos, <http://www.semarh.rn.gov.br/Conteudo.asp?TRAN=ITEM&TARG=17381&ACT=&PAGE=0&PARM=&LBL=Servi%20E7os>.
- [18] S. A. S. Almeida, M. Z. Cuellar, A. M. B. Costa, and R. F. Amorim, "Caracterização das bacias hidrográficas dos Rios Apodi/Mossoró e Piranhas/Assú (RN): mapeamento do uso do solo através das imagens do satélite CBERS 2 E ANÁLISE SÓCIO-ECONÔMICA," *Revista FAPERN*, vol. 1, pp. 5–9, 2006.
- [19] G. H. Silva, R. V. Santos, R. S. Moura, and N. C. Bueno, "Primeiro registro de *Chara indica* e *Chara zeylanica* (Charophyceae, Charales, Characeae) em reservatórios do semiárido do estado do Rio Grande do Norte, Brasil," *Biotemas*, vol. 26, no. 3, 2013.
- [20] E. Kawakami and G. Vazzoler, "Método gráfico e estimativa de índice alimentar aplicado no estudo de alimentação de peixes," *Boletim do Instituto Oceanográfico*, vol. 29, no. 2, pp. 205–207, 1980.
- [21] EMPARN—Empresa de Pesquisa Agropecuária do Rio Grande do Norte, <http://189.124.201.150/monitoramento/monitoramento.php>.
- [22] H. B. N. Hynes, "The food of freshwater sticklebacks (*Gasterosteus aculeatus* and *Pygosteus pungitius*) with a review of methods used in studies of the food of fish," *Journal of Animal Ecology*, vol. 19, pp. 35–38, 1950.
- [23] E. J. Hyslop, "Stomach contents analysis—a review of methods and their application," *Journal of Fish Biology*, vol. 17, no. 4, pp. 411–429, 1980.
- [24] Y.-H. Taguchi and Y. Oono, "Relational patterns of gene expression via non-metric multidimensional scaling analysis," *Bioinformatics*, vol. 21, no. 6, pp. 730–740, 2005.
- [25] M. J. Anderson, "A new method for non-parametric multivariate analysis of variance," *Austral Ecology*, vol. 26, no. 1, pp. 32–46, 2001.
- [26] R. Svanbäck and L. Persson, "Individual diet specialization, niche width and population dynamics: implications for trophic polymorphisms," *Journal of Animal Ecology*, vol. 73, no. 5, pp. 973–982, 2004.
- [27] E. D. Le Cren, "The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*)," *The Journal of Animal Ecology*, vol. 20, no. 2, pp. 201–219, 1951.
- [28] M. F. Magalhães, P. Beja, I. J. Schlosser, and M. J. Collares-Pereira, "Effects of multi-year droughts on fish assemblages of seasonally drying Mediterranean streams," *Freshwater Biology*, vol. 52, no. 8, pp. 1494–1510, 2007.
- [29] W. P. Sousa, "The role of disturbance in natural communities," *Annual Review of Ecology and Systematics*, vol. 15, pp. 353–391, 1984.
- [30] R. J. Hobbs and L. F. Huenneke, "Disturbance, diversity, and invasion: implications for conservation," *Conservation Biology*, vol. 6, no. 3, pp. 324–337, 1992.
- [31] I. R. Noble and R. O. Slatyer, "The use of vital attributes to predict successional changes in plant communities subject to recurrent disturbances," *Vegetatio*, vol. 43, no. 1–2, pp. 5–21, 1980.
- [32] S. A. C. Gavilan-Leandro, D. Peretti, J. E. M. P. Júnior, M. A. Fernandes, and A. M. G. Júnior, "Espectro alimentar e variação sazonal da dieta de *Plagioscion squamosissimus* (Heckel, 1840) (Osteichthyes, Sciaenidae) na lagoa do Piató, Assú, Estado do Rio Grande do Norte, Brasil," *Acta Scientiarum Biological Sciences*, vol. 31, pp. 285–292, 2009.
- [33] M. Goulding and E. J. Ferreira, "Shrimp-eating fishes and a case of prey-switching in Amazon Rivers," *Revista Brasileira de Zoologia*, vol. 2, no. 3, pp. 85–97, 1983.
- [34] F. M. S. Braga, "Alimentação de *Plagioscion squamosissimus* (Osteichthyes, Sciaenidae) no reservatório de Barra Bonita, Estado de São Paulo," *Iheringia Série Zoologia*, vol. 84, pp. 11–19, 1998.
- [35] F. Juanes, J. A. Buckel, and F. S. Scharf, "Feeding ecology of piscivorous fishes," in *Handbook of Fish Biology and Fisheries*, vol. 1 of *Fish Biology*, chapter 12, pp. 267–283, Blackwell Science, 2002.
- [36] O. O. Collart, "Aspectos ecológicos do camarão *Macrobrachium amazonicum* (Heller, 1862) no baixo Tocantins (PA-Brasil)," *Memoria de la Sociedad de Ciencias Naturales la Salle*, vol. 48, supplement, pp. 341–353, 1988.
- [37] M. C. N. Silva, F. L. Frédou, and J. S. Rosa Filho, "Estudo do crescimento do camarão *Macrobrachium amazonicum* (Heller, 1862) da Ilha de Combú, Belém, Estado do Pará," *Amazônia, Ciência & Desenvolvimento*, vol. 2, no. 4, pp. 85–104, 2007.
- [38] F. S. Scharf, F. Juanes, and R. A. Rountree, "Predator size—prey size relationships of marine fish predators: interspecific variation and effects of ontogeny and body size on trophic-niche breadth," *Marine Ecology Progress Series*, vol. 208, pp. 229–248, 2000.
- [39] V. L. L. De Almeida, N. S. Hahn, and A. E. A. D. M. Vazzoler, "Feeding patterns in five predatory fishes of the high Paraná River floodplain (PR, Brazil)," *Ecology of Freshwater Fish*, vol. 6, no. 3, pp. 123–133, 1997.
- [40] C. C. G. F. Pereira, W. S. Smith, E. L. G. Espíndola, and O. Rocha, "Alimentação de *Plagioscion squamosissimus* (HECKEL, 1840), uma espécie introduzida no Médio e Baixo rio Tietê, São Paulo, Brasil," in *Espécies Invasoras em águas Doces: Estudos de Casos e Propostas de Manejo*, O. Rocha, E. L. G. Espíndola, N. F. Verani, F. J. R. Verani, and A. C. Rietzler, Eds., p. 416, Editora EdUFSCar, São Paulo, Brazil, 2005.
- [41] A. M. De Roos, E. McCauley, and W. G. Wilson, "Mobility versus density-limited predator-prey dynamics on different spatial scales," *Proceedings of the Royal Society B: Biological Sciences*, vol. 246, no. 1316, pp. 117–122, 1991.

- [42] P. B. Banks, "Nonlinearity in the predation risk of prey mobility," *Proceedings of the Royal Society of London B: Biological Sciences*, vol. 267, pp. 1621–1625, 2000.
- [43] L. Green, E. B. Fisher, S. Perlow, and L. Sherman, "Preference reversal and self control: choice as a function of reward amount and delay," *Behaviour Analysis Letters*, vol. 1, no. 1, pp. 43–51, 1981.
- [44] H. C. G. Vasconcelos and J. C. Sá-Oliveira, "Alimentação de *Potamotrygon motoro* (CHONDRICHTHYES, POTAMOTRYGONIDAE) na planície de inundação da APA do Rio Curiaú, Macapá-Amapá-Brasil," *Biota Amazônia*, vol. 1, pp. 66–73, 2011.
- [45] H. Vasconcelos and J. Oliveira, "Alimentação de *Potamotrygon motoro* (CHONDRICHTHYES, POTAMOTRYGONIDAE) na Planície de Inundação da APA do Rio Curiaú, Macapá-Amapá-Brasil," *Biota Amazônia*, vol. 1, no. 2, pp. 66–73, 2011.
- [46] J. F. Oliveira, R. S. Costa, J. L. C. Novaes, L. G. F. Reboucas, A. L. N. M. Segundo, and D. Peretti, "Efeito da seca e variação espacial na abundância de indivíduos nas guildas tróficas em um reservatório do semiárido brasileiro," *Boletim do Instituto de Pesca*, vol. 1, pp. 51–64, 2016.
- [47] M. Goulart and M. Callisto, "Bioindicadores de qualidade de água como ferramenta em estudos de impacto ambiental," *Revista da FAPAM*, vol. 2, pp. 156–164, 2003.
- [48] E. B. Beserra, C. R. M. Fernandes, J. T. de Sousa, E. M. de Freitas, and K. D. Santos, "The effect of water quality in the life cycle and in the attraction for the egg oviposition of *Aedes Aegypti* (L.) (Diptera: Culicidae)," *Neotropical Entomology*, vol. 39, no. 6, pp. 1016–1023, 2010.
- [49] A. N. Bozza and N. S. Hahn, "Use of food resources by juveniles and adults of piscivorous fish species in a neotropical floodplain," *Biota Neotropica*, vol. 10, no. 3, pp. 217–226, 2010.
- [50] K. Iguchi, N. Matsubara, T. Yodo, and K. Maekawa, "Individual food niche specialization in stream-dwelling charr," *Ichthyological Research*, vol. 51, no. 4, pp. 321–326, 2004.
- [51] R. E. Ricklefs, *Economia da Natureza*, Guanabara Koogan, Rio de Janeiro, Brazil, 6th edition, 2012.
- [52] G. C. Novakowski, N. S. Hahn, and R. Fugi, "Alimentação de peixes piscívoros antes e após a formação do reservatório de Salto Caxias, Paraná, Brasil," *Biota Neotropica*, vol. 7, no. 2, pp. 0–0, 2007.
- [53] B. G. Blackwell, M. L. Brown, and D. W. Willis, "Relative Weight (Wr) status and current use in fisheries assessment and management," *Reviews in Fisheries Science*, vol. 8, no. 1, pp. 1–44, 2000.