ON THE FIRST-PASSAGE TIME OF INTEGRATED BROWNIAN MOTION

CHRISTIAN H. HESSE

Received 30 January 2004 and in revised form 29 September 2004

Let $(B_t; t \geq 0)$ be a Brownian motion process starting from $B_0 = \nu$ and define $X_\nu(t) = \int_0^t B_s ds$. For $a \geq 0$, set $\tau_{a,\nu} := \inf\{t : X_\nu(t) = a\}$ (with $\inf \emptyset = \infty$). We study the conditional moments of $\tau_{a,\nu}$ given $\tau_{a,\nu} < \infty$. Using martingale methods, stopping-time arguments, as well as the method of dominant balance, we obtain, in particular, an asymptotic expansion for the conditional mean $E(\tau_{a,\nu} \mid \tau_{a,\nu} < \infty)$ as $\nu \rightarrow \infty$. Through a series of simulations, it is shown that a truncation of this expansion after the first few terms provides an accurate approximation to the unknown true conditional mean even for small ν .

1. Introduction

First-passage time problems (FPTs) for integrated Markov processes constitute a set of old and challenging problems of probability. They arise both in theoretical and in applied contexts. Our motivation to study them stems from previous work on a stochastic model for particle sedimentation in fluids where one such problem arises naturally; see Hesse and Tory [4]. In stochastic approaches to sedimentation, particle velocity is often modeled as Brownian motion $(B_t; t \geq 0)$. Particle position then is the integral over this Brownian motion and the question when a particle first travels through a specified distance then leads to the FPT for this integrated Markov process.

FPTs for integrated Markov processes are complicated by the fact that the integral of a Markov process is no longer Markovian. However, the two-dimensional process such as

$$\left(B_t, \int_0^t B_s ds; t \geq 0 \right) \quad (1.1)$$

in our case is Markovian. The process in (1.1) is often called the Kolmogorov diffusion since its study was initiated by Kolmogorov [5].

This lifting of the original process into two dimensions so as not to lose the Markovian property often necessitates additional boundary and initial conditions which are frequently not available. Specifically, for our FPT of integrated Brownian motion, we circumvent this impasse by considering a slightly perturbed process, hence introducing a boundary with a specification for the Laplace transform of the first-passage time density

on this boundary and later letting the boundary go to infinity in a controlled fashion. This is demonstrated in Sections 2 and 3. In Section 2, using martingale methods and stopping-time arguments, a system of partial differential equations for the conditional moments of the FPT distribution of integrated Brownian motion is derived. Here and below “conditional” is meant to indicate “given that the first-passage time is finite.” In Section 3, an asymptotic expansion for the conditional mean first-passage time is derived. Section 4 reports on the results of a series of simulations which show that the truncated asymptotic expansion for the conditional mean turns out to be an accurate approximation to the unknown conditional mean.

We now define our problem rigorously and mention related works. Let $(B_t; t \geq 0)$ be a Brownian motion process with diffusion parameter σ^2 starting from $E(B_t) = B_0 = \nu$ and define the position process $(X_\nu(t); t \geq 0)$ through

$$X_\nu(t) = X_\nu(0) + \int_0^t B_s ds. \quad (1.2)$$

Then

$$\tau_{a,\nu} = \begin{cases} \inf \{t : X_\nu(t) = X_\nu(0) + a\} & \text{if this set is not empty,} \\ \infty & \text{otherwise} \end{cases} \quad (1.3)$$

is the first-passage time of the position process to the state at distance a . We are interested in the conditional moments

$$E(\tau_{a,\nu} \mid \tau_{a,\nu} < \infty) \quad (1.4)$$

as functions of a , ν , and σ^2 .

Early studies of integrated Brownian motion are McKean [10], Goldman [2], and Gor'kov [3]. McKean investigated joint probability densities of certain first-passage times τ and corresponding hitting velocities B_τ . Goldman [2] has derived an expression of the joint probability density function of the so-called half-winding time τ' of the process and the velocity $B_{\tau'}$. In Gor'kov [3], a formula for the density of B_{τ^*} , where $\tau^* = \min\{t : X_\nu(t) = b, B_t \geq 0\}$, $b > 0$, is computed.

More recently, Lefebvre [8] has obtained joint moment-generating functions related to certain first-passage times as well as—in a special case—the probability density function of the first hitting place. His method is based on solving the appropriate Kolmogorov backward equations. Lachal [6] presents an elementary procedure for rederiving some of Lefebvre [8] formulas as well as for deriving explicitly the probability laws of certain first-passage times. Also, Lachal [7] contains results on first-passage times which are obtained by martingale methods. Finally, Lefebvre [9] gives results on certain first-passage problems for degenerate two-dimensional diffusion processes. All of these approaches lead to rather complicated integral expressions which one does not seem to be able to evaluate analytically to obtain moment information.

2. A system of equations for the conditional moments

In this section, we derive a system of partial differential equations for the moments of the conditional first-passage time $\tau_{a,\nu}$. Let $(\Omega, \mathcal{F}, P)$ be a probability space on which all upcoming processes will be defined.

Consider the process

$$(X_\nu(s), B_s; s \geq 0), \quad X_\nu(0) = -x \leq 0, \quad B_0 = \nu, \quad (2.1)$$

with an absorbing boundary at the plane $(0, \mathbb{R})$ and let $p(x_t, \nu_t, t|x, \nu)$ be the probability density associated with

$$X_\nu(0) = -x, \quad B_0 = \nu, \quad X_\nu(t) = x_t, \quad B_t = \nu_t, \quad X_\nu(s) < 0, \quad \forall s \in [0, t]. \quad (2.2)$$

The density p depends on the starting distance to the boundary, of course. Also, let

$$P(y, t|x, \nu) = \int_{-\infty}^y \int_{-\infty}^{+\infty} p(x_t, \nu_t, t|x, \nu) d\nu_t dx_t \quad (2.3)$$

so that $P(0, t|x, \nu)$ is the probability that the boundary has not been reached prior to time t . We also introduce the Laplace transform of the FPT distribution, namely,

$$\Psi_s(x, \nu) := \int_0^\infty e^{-st} \frac{\partial}{\partial t} (1 - P(0, t|x, \nu)) dt. \quad (2.4)$$

A key instrument in our derivation will be the random Laplace transform $\Psi_s(-X_\nu(t), B_t)$. To state our results, we also introduce the notation

$$\begin{aligned} \partial_1 \Psi_s(y, z) &= \frac{\partial}{\partial y} \Psi_s(y, z), \\ \partial_2 \Psi_s(y, z) &= \frac{\partial}{\partial z} \Psi_s(y, z), \\ \partial_{22} \Psi_s(y, z) &= \frac{\partial^2}{\partial z^2} \Psi_s(y, z). \end{aligned} \quad (2.5)$$

An important role in obtaining conditional moments of the FPT distribution will be played by the process

$$(Z(t) := \exp(-st) \Psi_s(-X_\nu(t), B_t); 0 \leq t \leq \tau_{x,\nu}), \quad (2.6)$$

which is defined for all fixed positive s .

THEOREM 2.1. *The stochastic Ito-differential $dZ(t)$ of $Z(t)$ is given by*

$$\begin{aligned} dZ(t) &= \exp(-st) \partial_2 \Psi_s(-x_t, \nu_t) \sigma dW_t \\ &+ \exp(-st) \left[\frac{\sigma^2}{2} \partial_{22} \Psi_s(-x_t, \nu_t) - B_t \partial_1 \Psi_s(-x_t, \nu_t) - s \Psi_s(-x_t, \nu_t) \right] dt, \end{aligned} \quad (2.7)$$

where W_t is standard Brownian motion.

Proof. We compute explicitly the increment of $Z(t)$ over a time interval of length Δt . It is

$$\begin{aligned}\Delta Z &:= Z(t + \Delta t) - Z(t) \\ &= [\exp(-s(t + \Delta t)) - \exp(-st)] [\Psi_s(-x_{t+\Delta t}, \nu_{t+\Delta t}) - \Psi_s(-x_t, \nu_t)] \\ &\quad + [\exp(-s(t + \Delta t)) - \exp(-st)] \Psi_s(-x_t, \nu_t) \\ &\quad + \exp(-st) [\Psi_s(-x_{t+\Delta t}, \nu_{t+\Delta t}) - \Psi_s(-x_t, \nu_t)].\end{aligned}\quad (2.8)$$

Retaining terms of order Δt only, this becomes

$$\begin{aligned}\Delta Z &= -s \exp(-st) \Psi_s(-x_t, \nu_t) \Delta t \\ &\quad + \exp(-st) \left[-\partial_1 \Psi_s(-x_t, \nu_t) B_t \Delta t + \partial_2 \Psi_s(-x_t, \nu_t) \Delta B_t \right. \\ &\quad \left. + \frac{1}{2} \partial_{22} \Psi_s(-x_t, \nu_t) (\Delta B_t)^2 \right] + o((\Delta t)^{3/2}).\end{aligned}\quad (2.9)$$

Noting that $\Delta B_t = \sigma \Delta W_t$, $(\Delta B_t)^2 = \sigma^2 (\Delta W_t)^2 = \sigma^2 \Delta t$, and including terms such as $\Delta t \Delta W_t$ and higher-order terms in the $o((\Delta t)^{3/2})$ summand, we arrive at

$$\begin{aligned}\Delta Z &= \exp(-st) \partial_2 \Psi_s(-x_t, \nu_t) \sigma \Delta W_t \\ &\quad + \exp(-st) \left[\frac{\sigma^2}{2} \partial_{22} \Psi_s(-x_t, \nu_t) - B_t \partial_1 \Psi_s(-x_t, \nu_t) - s \Psi_s(-x_t, \nu_t) \right] \Delta t + o((\Delta t)^{3/2}).\end{aligned}\quad (2.10)$$

The result then follows by replacing infinitesimal increments by differentials. $\square$

THEOREM 2.2. *The process $(Z(t); 0 \leq t \leq \tau_{x,\nu})$ as defined before Theorem 2.1 is a local martingale relative to the family of σ -algebras generated by $(Z(s); s \leq t)$.*

Proof. The proof is based on stopping-time arguments and is simple. The details are omitted. $\square$

From Theorem 2.1, we can deduce the representation

$$\begin{aligned}Z(t) &= \Psi_s(x, \nu) + \int_0^t \exp(-s\tau) \partial_2 \Psi_s \sigma dW_\tau \\ &\quad + \int_0^t \exp(-s\tau) \left[\frac{\sigma^2}{2} \partial_{22} \Psi_s - B_\tau \partial_1 \Psi_s - s \Psi_s \right] d\tau.\end{aligned}\quad (2.11)$$

The local martingale property of $Z(t)$ implies that the third summand on the right side of (2.11) must vanish for all $t \geq 0$. From this, we may deduce that

$$\frac{\sigma^2}{2} \frac{\partial^2 \Psi_s(x, \nu)}{\partial \nu^2} - \nu \frac{\partial \Psi_s(x, \nu)}{\partial x} - s \Psi_s(x, \nu) = 0. \quad (2.12)$$

The absorbing boundary condition gives

$$\Psi_s(0, \nu) = 1, \quad \forall s > 0, \quad \forall \nu > 0. \quad (2.13)$$

As defined, the function Ψ_s possesses the representation

$$\Psi_s(x, \nu) = \sum_{n=0}^{\infty} \frac{(-s)^n}{n!} m_n(x, \nu), \quad (2.14)$$

where $m_n(x, \nu)$ is the n th conditional moment of the FPT distribution, that is, $E(\tau_{x,\nu}^n \mid \tau_{x,\nu} < \infty)$.

Substituting (2.14) into (2.12) and equating coefficients of powers of s , we arrive at the following system of partial differential equations for m_n :

$$\frac{\sigma^2}{2} \frac{\partial^2 m_n(x, \nu)}{\partial \nu^2} - \nu \cdot \frac{\partial m_n(x, \nu)}{\partial x} = -n \cdot m_{n-1}(x, \nu), \quad (2.15)$$

$$\begin{aligned} m_n(0, \nu) &= 0, \quad \forall n \geq 1, \nu > 0, \\ m_0(0, \nu) &= 1. \end{aligned} \quad (2.16)$$

For the conditional mean FPT, we get, suppressing the subscripts of m ,

$$\frac{\sigma^2}{2} \frac{\partial^2 m(x, \nu)}{\partial \nu^2} - \nu \cdot \frac{\partial m(x, \nu)}{\partial x} = -1, \quad (2.17)$$

$$m(0, \nu) = 0, \quad \forall \nu > 0. \quad (2.18)$$

A difficulty with the systems (2.15), (2.16) and (2.17), (2.18) arises from the lack of sufficient boundary or initial information. Although (2.16) is obvious since

$$\lim_{x \rightarrow 0+} \lim_{\nu \rightarrow \nu^*} m_n(x, \nu) = m_n(0, \nu^*), \quad \forall \nu^* > 0, \forall n \geq 1, \quad (2.19)$$

it is unclear how this limit behaves for $\nu^* < 0$. Hence, the boundary value problem is not well-posed. On the other hand, to find $\lim_{x \rightarrow x^*} \lim_{\nu \rightarrow \nu^*} m_n(x, \nu)$ for a given ν^* and all $x^* \geq 0$ is as difficult as the original FPTP itself. Therefore, also the initial boundary value problem is ill-posed and it seems that we have arrived at an impasse. In the following section, we demonstrate how to circumvent this difficulty and derive an asymptotic expansion for $m(x, \nu)$ as $\nu \rightarrow \infty$.

3. An asymptotic expansion for the mean

A way out of the dilemma indicated at the end of Section 2 is based on the following modified process. For some $\nu_0 > \nu$ define

$$\tilde{B}_t := \begin{cases} B_t & \text{if } B_s < \nu_0, \quad \forall s \leq t, \\ \nu_0, & \forall t \geq \tau_0 := \min \{t : B_t = \nu_0\}, \end{cases} \quad (3.1)$$

and let $\tilde{X}_\nu(t) = X_\nu(0) + \int_0^t \tilde{B}_s ds$. The process $(\tilde{X}_\nu(t), \tilde{B}_t; t \geq 0)$ has the same sample paths as $(X_\nu(t), B_t; t \geq 0)$ except when the Brownian motion process hits ν_0 . When this happens, the Brownian motion terminates a.s. at ν_0 . Also, let

$$\tau_{a,\nu,\nu_0} := \begin{cases} \min \{t \geq 0 : \tilde{X}_\nu(t) = X_\nu(0) + a\} & \text{if this set is not empty,} \\ \infty & \text{if } \tilde{X}_\nu(t) \neq X_\nu(0) + a, \quad \forall t \geq 0, \end{cases} \quad (3.2)$$

be the FPT of the modified process on $X_\nu(0) + a$ and

$$\tilde{m}_n(x, \nu) := E(\tau_{x, \nu, \nu_0}^n \mid \tau_{x, \nu, \nu_0} < \infty). \quad (3.3)$$

Then the $\tilde{m}_n(x, \nu)$ also satisfy (2.15), (2.16), as well as

$$\tilde{m}_n(x, \nu_0) = \left(\frac{x}{\nu_0}\right)^n, \quad \forall x \geq 0, \quad \forall n \geq 1. \quad (3.4)$$

Furthermore, both $\tau_{x, \nu}$ and τ_{x, ν_0, ν_0} are stopping times with respect to the family of σ -algebras generated by $(X_\nu(t), B_t; t \geq 0)$ and we have $\tau_{x, \nu, \nu_0} \xrightarrow{\nu_0 \rightarrow \infty} \tau_{x, \nu}$ a.s. From these considerations, the following approach is suggested for the mean FPT. For (2.17), the initial boundary value problem is solved with (2.18) and

$$m(x, \nu_0) = \frac{x}{\nu_0}, \quad \forall x \geq 0, \quad \nu_0 \text{ large}. \quad (3.5)$$

Then the limit as $\nu_0 \rightarrow \infty$ is taken. Writing

$$\hat{m}(c, \nu) = \int_0^\infty \exp(-cx) m(x, \nu) dx, \quad (3.6)$$

one obtains

$$\frac{\sigma^2}{2} \frac{d^2 \hat{m}(c, \nu)}{d\nu^2} - c\nu \hat{m}(c, \nu) = -c^{-1}, \quad (3.7)$$

$$\hat{m}(c, \nu_0) = \nu_0^{-1} c^{-2}. \quad (3.8)$$

The substitution

$$\begin{aligned} \omega &= 2^{1/3} c^{1/3} \sigma^{-2/3} \nu, \\ \omega_0 &= 2^{1/3} c^{1/3} \sigma^{-2/3} \nu_0 \end{aligned} \quad (3.9)$$

transforms (3.7) and (3.8) into

$$\frac{d^2 \hat{m}(c, \omega)}{d\omega^2} - \omega \cdot \hat{m}(c, \omega) = -2^{1/3} \sigma^{-2/3} c^{-5/3}, \quad (3.10)$$

$$\hat{m}(c, \omega_0) = 2^{1/3} \sigma^{-2/3} c^{-5/3} \omega_0^{-1}, \quad (3.11)$$

whose homogeneous part is a one-dimensional Schrödinger equation. We now use methods of global analysis such as WKB analysis and the method of dominant balance on this Schrödinger equation (see, e.g., Bender and Orszag [1]). If the power series $\sum_{n=0}^\infty p(n) \omega^n$ is substituted for $\hat{m}$ into (3.10), the coefficients are seen to satisfy

$$\begin{aligned} p(3n+2) &= 0, \\ p(3n+1) &= \frac{p(1) \cdot \Gamma(4/3)}{9^n n! \Gamma(n+4/3)}, \\ p(3n) &= \frac{p(0) \cdot \Gamma(2/3)}{9^n n! \Gamma(n+2/3)}, \quad \forall n \geq 0. \end{aligned} \quad (3.12)$$

Performing a variation of parameters and writing

$$\begin{aligned} A(\omega) &:= 3^{-2/3} \sum_{n=0}^{\infty} \frac{\omega^{3n}}{9^n n! \Gamma(n+2/3)} - 3^{-4/3} \sum_{n=0}^{\infty} \frac{\omega^{3n+1}}{9^n n! \Gamma(n+4/3)}, \\ B(\omega) &:= 3^{-1/6} \sum_{n=0}^{\infty} \frac{\omega^{3n}}{9^n n! \Gamma(n+2/3)} + 3^{-5/6} \sum_{n=0}^{\infty} \frac{\omega^{3n+1}}{9^n n! \Gamma(n+4/3)}, \end{aligned} \quad (3.13)$$

this leads to

$$\hat{m}(c, \omega) = \pi \cdot 2^{1/3} \sigma^{-2/3} c^{-5/3} \left[B(\omega) \int_{\omega}^{\infty} A(t) dt + A(\omega) \int_0^{\omega} B(t) dt \right] + k \cdot A(\omega), \quad (3.14)$$

where $k = k(c)$ is a constant that needs to be fitted so that (3.14) satisfies the initial condition (3.11).

$A(\omega)$ and $B(\omega)$ in (3.14) are the so-called Airy functions and it is well known that $A(\omega)$ decays exponentially as $\omega \rightarrow \infty$. In fact

$$A(\omega) \sim 2^{-1} \pi^{-1/2} \omega^{-1/4} \exp\left(-\frac{2}{3} \omega^{3/2}\right), \quad \text{as } \omega \rightarrow \infty. \quad (3.15)$$

Hence it can be expected that the asymptotic expansion of (3.14) as $\omega \rightarrow \infty$ equals the asymptotic expansion of merely the first two summands of (3.14). We confirm this with the following argument. The strategy is to first peel off the leading asymptotic behavior, then, after having removed this, to determine the leading behavior of the remainder, and so forth. The first step in this procedure is to assume

$$\frac{d^2 \hat{m}(c, \omega)}{d\omega^2} \sim 0, \quad \text{as } \omega \rightarrow \infty, \quad (3.16)$$

which yields

$$\hat{m}(c, \omega) \sim 2^{1/3} \sigma^{-2/3} c^{-5/3} \omega^{-1}, \quad \text{as } \omega \rightarrow \infty. \quad (3.17)$$

Corrections to this leading term are obtained by writing

$$\hat{m}(c, \omega) = 2^{1/3} \sigma^{-2/3} c^{-5/3} (\omega^{-1} + \varepsilon(\omega)), \quad \text{as } \omega \rightarrow \infty, \quad (3.18)$$

where the correction term is of smaller order than ω^{-1} . It is easily determined that $\varepsilon(\omega)$ satisfies the differential equation

$$\frac{d^2 \varepsilon(\omega)}{d\omega^2} + \frac{2}{\omega^3} = \omega \cdot \varepsilon(\omega). \quad (3.19)$$

Again, setting $d^2 \varepsilon(\omega)/d\omega^2 \sim 0$ (as $\omega \rightarrow \infty$), we see that $\varepsilon(\omega) \sim 2\omega^{-4}$ and continuing with this method, one obtains the full asymptotic power series expansion

$$\hat{m}(c, \omega) \sim 2^{1/3} \sigma^{-2/3} c^{-5/3} \sum_{n=0}^{\infty} \frac{(3n)!}{3^n n!} \omega^{-3n-1}. \quad (3.20)$$

This indeed is the asymptotic expansion of the first two summands on the right-hand side of (3.14).

From (3.20), we arrive at

$$m(x, \nu) \sim \sum_{n=0}^{\infty} \frac{(3n)! \sigma^{2n} x^{n+1}}{n!(n+1)! 6^n \nu^{3n+1}}. \quad (3.21)$$

It is easily checked by differentiating termwise that the asymptotic expansion in (3.21) formally satisfies (2.17) and the boundary condition (2.18). However, the sum in (3.21) does not converge for any nonzero value of $\sigma^2 x \nu^{-3}$. Although this might be surprising at first, it is well known that most problems in perturbation theory or WKB analysis lead to divergent series. It was Poincaré who introduced the concept of divergent asymptotic expansions into mathematics and demonstrated that formal solutions of differential equations are asymptotic expansions of actual solutions. In fact, typically, optimally truncated divergent series constitute accurate approximations to the actual solutions, see again Bender and Orszag [1]. We truncate (3.21) after the third summand and use

$$\bar{m}(x, \nu) = \frac{x}{\nu} + \frac{\sigma^2 x^2}{2 \nu^4} + \frac{5\sigma^4 x^3}{3 \nu^7} \quad (3.22)$$

as our approximation.

4. Simulations

We performed an extensive series of simulations to evaluate the quality of the approximation (3.22) for various distances x to boundary, initial velocities ν , and diffusion parameters σ . The simulations indicate that the approximation is accurate whenever $\sigma^2 x \nu^{-3} < 1/4$.

Summaries of the simulations are reported in Tables 4.1, 4.2, and 4.3. The following notation is used (for convenience, the dependence on x, ν, σ was dropped):

$$A_1 = \frac{x}{\nu}, \quad A_2 = \frac{\sigma^2 x^2}{2 \nu^4}, \quad A_3 = \frac{5\sigma^4 x^3}{3 \nu^7}, \quad (4.1)$$

m_s = sample averages of simulated first-passage times for given x, ν, σ^2 ,

$$M_1 = 10^3 \cdot \frac{m_s - A_1}{A_1}, \quad M_2 = 10^3 \cdot \frac{m_s - (A_1 + A_2)}{A_1}, \quad M_3 = 10^3 \cdot \frac{m_s - (A_1 + A_2 + A_3)}{A_1},$$

$$SD(M_i) = \text{sample standard deviation of } M_i, \quad i = 1, 2, 3, \quad (4.2)$$

$$t_i = \frac{M_i}{SD(M_i)}.$$

At time $t = 0$, a total of 2000 particles were started with initial velocity ν (for $\nu = 2, 4, 6, 8$) and diffusion parameter σ (for $\sigma = 0.1, 0.3, 0.7$). We recorded when these particles first crossed boundaries at distances $x = 2, 4, 8, 12, 20$ and computed $m_s, M_i, SD(M_i)$, and t_i .

The values of t_1 indicate that the first-order approximation A_1 tends to underestimate the mean first-passage time. On the other hand, with $|t_3| \leq 2$ in almost all cases

Table 4.1. M_i, t_i for various values of x, ν and $\sigma = 0.1$.

$\nu = 2$							$\nu = 4$					
x	M_1	M_2	M_3	t_1	t_2	t_3	M_1	M_2	M_3	t_1	t_2	t_3
2	6.1	-6.41	-6.51	0.9	-1.00	-1.02	-0.6	-2.12	-2.12	-0.2	-0.95	-0.95
4	9.4	-15.59	-16.01	1.0	-1.71	-1.75	2.1	-1.07	-1.08	0.6	-0.33	-0.33
8	26.0	-24.02	-25.69	1.9	-1.84	-1.97	8.3	2.09	2.06	1.8	0.45	0.45
12	54.9	-20.13	-23.88	3.3	-1.22	-1.45	12.9	3.49	3.43	2.2	0.61	0.60
20	110.0	-14.97	-25.38	5.0	-0.68	-1.15	19.4	3.78	3.62	2.6	0.51	0.49
$\nu = 6$							$\nu = 8$					
x	M_1	M_2	M_3	t_1	t_2	t_3	M_1	M_2	M_3	t_1	t_2	t_3
2	0.7	0.22	0.22	0.5	0.18	0.18	0.8	0.65	0.65	1.1	0.90	0.90
4	0.1	-0.86	-0.86	0.0	-0.49	-0.49	1.5	1.15	1.15	1.3	1.03	1.03
8	-0.4	-2.23	-2.23	-0.1	-0.89	-0.89	1.3	0.53	0.53	0.8	0.33	0.33
12	-0.1	-2.91	-2.92	-0.0	-0.94	-0.95	1.2	-0.01	-0.01	0.6	0.00	0.00
20	2.2	-2.39	-2.41	0.5	-0.60	-0.60	1.9	-0.03	-0.03	0.7	-0.01	-0.01

Table 4.2. M_i, t_i for various values of x, ν and $\sigma = 0.3$.

$\nu = 2$							$\nu = 4$					
x	M_1	M_2	M_3	t_1	t_2	t_3	M_1	M_2	M_3	t_1	t_2	t_3
2	81.0	-31.52	-39.96	3.8	-1.49	-1.89	16.5	2.44	2.31	2.3	0.35	0.33
4	238.4	13.44	-20.31	6.3	0.36	-0.54	26.5	-1.60	-2.13	2.7	-0.16	-0.22
8	846.9	396.91	261.91	6.1	2.85	1.88	43.3	-12.92	-15.02	3.1	-0.91	-1.06
12	1601.0	926.05	622.29	4.3	2.50	1.68	67.9	-16.46	-21.21	3.8	-0.93	-1.20
20	3063.4	1938.43	1094.68	4.0	2.53	1.43	133.9	-6.69	-19.87	5.6	-0.28	-0.83
$\nu = 6$							$\nu = 8$					
x	M_1	M_2	M_3	t_1	t_2	t_3	M_1	M_2	M_3	t_1	t_2	t_3
2	10.1	5.90	5.89	2.7	1.57	1.57	0.1	-1.68	-1.68	0.0	-0.70	-0.70
4	16.4	8.07	8.02	3.1	1.53	1.52	-1.3	-4.85	-4.86	-0.4	-1.43	-1.43
8	25.8	9.15	8.96	3.4	1.22	1.19	0.9	-6.11	-6.14	0.2	-1.28	-1.28
12	31.9	6.87	6.45	3.5	0.74	0.70	3.4	-7.13	-7.21	0.6	-1.22	-1.24
20	52.4	10.76	9.60	4.4	0.90	0.80	10.2	-7.37	-7.58	1.4	-0.99	-1.01

Table 4.3. M_i, t_i for various values of x, ν and $\sigma = 0.7$.

$\nu = 4$							$\nu = 6$					
x	M_1	M_2	M_3	t_1	t_2	t_3	M_1	M_2	M_3	t_1	t_2	t_3
2	52.8	-23.76	-27.61	3.1	-1.41	-1.64	13.6	-9.11	-9.45	1.5	-1.04	-1.08
4	107.6	-45.49	-61.13	4.4	-1.87	-2.51	34.6	-10.75	-12.12	2.7	-0.86	-0.97
8	213.6	-92.68	-155.21	6.0	-2.61	-4.36	88.8	-1.96	-7.45	4.9	-0.11	-0.41
12	300.4	-158.96	-299.64	6.8	-3.63	-6.85	138.4	2.28	-10.07	6.0	0.10	-0.44
20	3195.5	2429.93	2039.14	1.1	0.87	0.74	262.7	35.90	1.59	7.5	1.04	0.05
$\nu = 8$							$\nu = 10$					
x	M_1	M_2	M_3	t_1	t_2	t_3	M_1	M_2	M_3	t_1	t_2	t_3
2	-4.4	-13.97	-14.03	-0.8	-2.52	-2.53	10.4	3.66	3.63	2.1	0.76	0.75
4	-7.0	-26.18	-26.42	-0.8	-3.25	-3.28	18.0	4.52	4.40	2.6	0.66	0.65
8	4.4	-33.89	-34.87	0.3	-2.93	-3.02	39.3	12.40	11.92	4.0	1.27	1.22
12	23.9	-33.54	-35.74	1.6	-2.33	-2.49	55.4	15.03	13.95	4.5	1.23	1.15
20	64.0	-31.74	-37.84	3.4	-1.69	-2.01	80.8	13.56	10.55	5.1	0.86	0.63

considered, the third-order approximation $A_1 + A_2 + A_3$ seems to approximate the mean first-passage time very well. The difference between the second-order and the third-order approximation is rather small in most cases.

References

- [1] C. M. Bender and S. A. Orszag, *Advanced Mathematical Methods for Scientists and Engineers*, International Series in Pure and Applied Mathematics, McGraw-Hill Book, New York, 1978.
- [2] M. Goldman, *On the first passage of the integrated Wiener process*, Ann. Math. Statist. **42** (1971), 2150–2155.
- [3] J. P. Gor'kov, *A formula for the solution of a certain boundary value problem for the stationary equation of Brownian motion*, Dokl. Akad. Nauk SSSR **223** (1975), no. 3, 525–528 (Russian).
- [4] C. H. Hesse and E. M. Tory, *The stochastics of sedimentation*, Adv. Fluid Mech. **7** (1996), 199–240.
- [5] A. Kolmogorov, *Zufällige Bewegungen (zur Theorie der Brownschen Bewegung)*, Ann. of Math. (2) **35** (1934), no. 1, 116–117 (German).
- [6] A. Lachal, *L'intégrale du mouvement brownien*, J. Appl. Probab. **30** (1993), no. 1, 17–27 (French).
- [7] ———, *Some martingales related to the integral of Brownian motion. Applications to the passage times and transience*, J. Theoret. Probab. **11** (1998), no. 1, 127–156.
- [8] M. Lefebvre, *First-passage densities of a two-dimensional process*, SIAM J. Appl. Math. **49** (1989), no. 5, 1514–1523.
- [9] ———, *First-passage problems for degenerate two-dimensional diffusion processes*, Test **12** (2003), no. 1, 125–139.
- [10] H. P. McKean Jr., *A winding problem for a resonator driven by a white noise*, J. Math. Kyoto Univ. **2** (1963), 227–235.
- [11] E. M. Tory and C. H. Hesse, *Theoretical and experimental evidence for a Markov model for sedimentation*, Adv. Fluid Mech. **7** (1996), 241–280.

Christian H. Hesse: Institut für Stochastik und Anwendungen, Fachbereich Mathematik, Universität Stuttgart, 70569 Stuttgart, Germany

E-mail address: hesse@mathematik.uni-stuttgart.de