Research Article

On the Lower Bound for the Number of Real Roots of a Random Algebraic Equation

Takashi Uno

Received 27 February 2007; Accepted 30 October 2007

We estimate a lower bound for the number of real roots of a random algebraic equation whose random coefficients are dependent normal random variables.

Copyright © 2007 Takashi Uno. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let $N_n(\mathbf{R}, \omega)$ be the number of real roots of the random algebraic equation

$$F_n(x, \omega) = \sum_{\nu=0}^n a_\nu(\omega) x^\nu = 0, \quad (1.1)$$

where the $a_\nu(\omega)$, $\nu = 0, 1, \dots, n$, are random variables defined on a fixed probability space $(\Omega, \mathcal{A}, \Pr)$ assuming real values only.

During the past 40–50 years, the majority of published researches on random algebraic polynomials has concerned the estimation of $N_n(R, \omega)$. Works by Littlewood and Offord [1], Samal [2], Evans [3], and Samal and Mishra [4–6] in the main concerned cases in which the random coefficients $a_\nu(\omega)$ are independent and identically distributed.

For dependent coefficients, Sambandham [7] considered the upper bound for $N_n(\mathbf{R}, \omega)$ in the case when the $a_\nu(\omega)$, $\nu = 0, 1, \dots, n$, are normally distributed with mean zero and joint density function

$$|M|^{1/2} (2\pi)^{-(n+1)/2} \exp \left(- (1/2) \mathbf{a}' M \mathbf{a} \right), \quad (1.2)$$

where M^{-1} is the moment matrix with $\sigma_i = 1$, $\rho_{ij} = \rho$, $0 < \rho < 1$, $(i \neq j)$, $i, j = 0, 1, \dots, n$ and $\mathbf{a}'$ is the transpose of the column vector $\mathbf{a}$. Also, Uno and Negishi [8] obtained the same result as Sambandham in the case of the moment matrix with $\sigma_i = 1$, $\rho_{ij} = \rho_{|i-j|}$,

$(i \neq j)$, $i, j = 0, 1, \dots, n$, where ρ_j is a nonnegative decreasing sequence satisfying $\rho_1 < 1/2$ and $\sum_{j=1}^{\infty} \rho_j < \infty$ in (1.2).

The lower bound for $N_n(\mathbf{R}, \omega)$ in the case of dependent normally distributed coefficients was estimated by Renganathan and Sambandham [9] and Nayak and Mohanty [10] under the same condition of Sambandham [7]. Uno [11] pointed out the defect in the proofs of the above papers and obtained the result for the lower bound. Additionally, Uno [12] estimated the *strong* result for this particular problem in the sense of Evans [3]. The term *strong* indicates that the estimation for the exceptional set is independent of the degree n .

The object of this paper is to find the lower bound for $N_n(\mathbf{R}, \omega)$ when the coefficients are nonidentically distributed dependent normal random variables. We remark that this result is the general form of Uno [11] and that the exceptional set is dependent on the degree n . In this paper, we suppose that the $a_\nu(\omega)$, $\nu = 0, 1, \dots, n$, have mean zero, and the moment with

$$\rho_{ij} = \begin{cases} 1 & (i = j), \\ \rho_{|i-j|} & (1 \leq |i-j| \leq m), \\ 0 & (|i-j| > m), \end{cases} \quad i, j = 0, 1, \dots, n, \quad (1.3)$$

for a positive integer m , where $0 \leq \rho_j < 1$, $j = 1, 2, \dots, m$ in (1.2). That is to say we assume the $a_\nu(\omega)$'s to be m -dependent stationary Gaussian random variables. With Yoshihara ([13, page 29]), we see that this assumption is equivalent to the following two statements for a stationary Gaussian sequence:

- (i) $\{a_\nu\}$ is $*$ -mixing;
- (ii) $\{a_\nu\}$ is ϕ -mixing.

Throughout the paper, we suppose n is sufficiently large. We will follow the line of proof of Samal and Mishra [5].

THEOREM 1.1. *Let*

$$f_n(x, \omega) = \sum_{\nu=0}^n a_\nu(\omega) b_\nu x^\nu = 0 \quad (1.4)$$

be a random algebraic equation of degree n , where the $a_\nu(\omega)$'s are dependent normally distributed with mean zero, and the moment matrix given by (1.3) and the b_ν , $\nu = 0, 1, \dots, n$, be positive numbers such that $\lim_{n \rightarrow \infty} (k_n/t_n)$ is finite, where $k_n = \max_{0 \leq \nu \leq n} b_\nu$ and $t_n = \min_{0 \leq \nu \leq n} b_\nu$.

Then for $n > n_0$, the number of real roots of most of the equations $f_n(x, \omega) = 0$ is at least $\varepsilon_n \log n$ outside a set of measure at most

$$\frac{\mu}{\varepsilon_n \log n} + \left(\frac{k_n}{t_n} \right)^\beta \exp \left(-\frac{\mu' \beta}{\varepsilon_n} \right), \quad \beta > 0, \quad (1.5)$$

provided ε_n tends to zero, but $\varepsilon_n \log n$ tends to infinity as n tends to infinity, and μ and μ' are positive constants.

2. Proof of theorem

Let $\{\lambda_n\}$ be any sequence tending to infinity as n tends to infinity and M is the integer defined by

$$M = \left[\alpha^2 \lambda_n^2 \left(\frac{k_n}{t_n} \right)^2 \right] + 1, \quad (2.1)$$

where α is a positive constant and $[x]$ denotes the greatest integer not exceeding x . Let k be the integer determined by

$$M^{2k} \leq n < M^{2k+2}. \quad (2.2)$$

We will consider $f_n(x, \omega)$ at the points

$$x_l = \left(1 - \frac{1}{M^{2l}} \right)^{1/2} \quad (2.3)$$

for $l = [k/2] + 1, [k/2] + 2, \dots, k$.

Let

$$f_n(x_l, \omega) = \sum_1 a_\nu(\omega) b_\nu x_l^\nu + \left(\sum_2 + \sum_3 \right) a_\nu(\omega) b_\nu x_l^\nu = U_l(\omega) + R_l(\omega), \quad (\text{say}), \quad (2.4)$$

where ν ranges from $M^{2l-1} + 1$ to M^{2l+1} in $\sum_1$, from 0 to M^{2l-1} in $\sum_2$ and from $M^{2l+1} + 1$ to n in $\sum_3$.

The following lemmas are necessary for the proof of the theorem. We will use the fact that each $a_\nu(\omega)$ has marginal frequency function $(2\pi)^{-1/2} \exp(-u^2/2)$.

LEMMA 2.1. For $\alpha_1 > 0$,

$$\sigma_l > \alpha_1 t_n M^l, \quad (2.5)$$

where

$$\sigma_l^2 = \sum_{i=M^{2l-1}+1}^{M^{2l+1}} b_i^2 x_l^{2i} + 2 \sum_{i=M^{2l-1}+1}^{M^{2l+1}-1} \sum_{j=i+1}^{M^{2l+1}} b_i b_j x_l^{i+j} \rho_{j-i}. \quad (2.6)$$

Proof. First, we have

$$\sum_{i=M^{2l-1}+1}^{M^{2l+1}} b_i^2 x_l^{2i} > t_n^2 \sum_{i=M^{2l-1}+1}^{M^{2l}} x_l^{2i} > \left(\frac{B}{A} \right) t_n^2 M^{2l}, \quad (2.7)$$

where A and B are positive constants such that $A > 1$ and $0 < B < 1$.

Second, we get

$$\begin{aligned} \sum_{i=M^{2l-1}+1}^{M^{2l+1}-1} \sum_{j=i+1}^{M^{2l+1}} b_i b_j x_l^{i+j} \rho_{j-i} &> t_n^2 \sum_{i=M^{2l-1}+1}^{M^{2l}-1} \sum_{j=i+1}^{M^{2l}} x_l^{i+j} \rho_{j-i} \\ &= t_n^2 \frac{x_l^{2(M^{2l-1}+1)}}{1-x_l^2} \left\{ \sum_{i=1}^m \rho_i x_l^i - \sum_{i=1}^m \rho_i x_l^{2(M^{2l}-M^{2l-1})-i} \right\} \geq \left(\frac{B'}{A'} \right) \rho_0 t_n^2 M^{2l}, \end{aligned} \quad (2.8)$$

where $\rho_0 = \sum_{j=1}^m \rho_j$ and A' and B' are positive constants satisfying $A' > 1$ and $0 < B' < 1$. So we get

$$\sigma_l^2 \geq \alpha_1^2 t_n^2 M^{2l}, \quad (2.9)$$

where α_1 is a positive constant, as required. $\square$

LEMMA 2.2. *Let*

$$\Pr \left(\left\{ \omega; \left| \sum_2 a_v(\omega) b_v x_l^v \right| > \lambda_n \tilde{\sigma}_l \right\} \right) < \sqrt{\frac{2}{\pi}} \frac{e^{-\lambda_n^2/2}}{\lambda_n}, \quad (2.10)$$

where

$$\tilde{\sigma}_l^2 = \sum_{i=0}^{M^{2l-1}} b_i^2 x_l^{2i} + 2 \sum_{i=0}^{M^{2l-1}-1} \sum_{j=i+1}^{M^{2l-1}} b_i b_j x_l^{i+j} \rho_{j-i}. \quad (2.11)$$

Proof. We get

$$\Pr \left(\left\{ \omega; \left| \sum_2 a_v(\omega) b_v x_l^v \right| > \lambda_n \tilde{\sigma}_l \right\} \right) = \sqrt{\frac{2}{\pi}} \int_{\lambda_n}^{\infty} e^{-u^2/2} du < \sqrt{\frac{2}{\pi}} \frac{e^{-\lambda_n^2/2}}{\lambda_n}. \quad (2.12)$$

$\square$

LEMMA 2.3. *Let*

$$\Pr \left(\left\{ \omega; \left| \sum_3 a_v(\omega) b_v x_l^v \right| > \lambda_n \tilde{\sigma}_l \right\} \right) < \sqrt{\frac{2}{\pi}} \frac{e^{-\lambda_n^2/2}}{\lambda_n}, \quad (2.13)$$

where

$$\tilde{\sigma}_l^2 = \sum_{i=M^{2l+1}+1}^n b_i^2 x_l^{2i} + 2 \sum_{i=M^{2l+1}+1}^{n-1} \sum_{j=i+1}^n b_i b_j x_l^{i+j} \rho_{j-i}. \quad (2.14)$$

The proof is similar to that of Lemma 2.2.

LEMMA 2.4. *For a fixed l ,*

$$\Pr \left(\left\{ \omega; |R_l(\omega)| < \sigma_l \right\} \right) > 1 - 2\sqrt{\frac{2}{\pi}} \frac{1}{\lambda_n} e^{-\lambda_n^2/2}. \quad (2.15)$$

Proof. By Lemmas 2.2 and 2.3, we get, for a given l ,

$$|R_l(\omega)| < \lambda_n(\tilde{\sigma}_l + \tilde{\tilde{\sigma}}_l) \quad (2.16)$$

outside a set of measure at most $2(2/\pi)^{1/2} \lambda_n^{-1} \exp(-\lambda_n^2/2)$. Again, we have

$$\begin{aligned} \sum_{i=0}^{M^{2l-1}} b_i^2 x_l^{2i} &\leq 2k_n^2 M^{2l-1}, \\ \sum_{i=0}^{M^{2l-1}-1} \sum_{j=i+1}^{M^{2l-1}} b_i b_j x_l^{i+j} \rho_{j-i} &\leq k_n^2 \sum_{i=1}^m \rho_i \sum_{j=1}^{M^{2l-1}-(i-1)} x_l^{2j+i-2} \leq \rho_0 k_n^2 M^{2l-1}. \end{aligned} \quad (2.17)$$

Hence we get, for a positive constant α_2 ,

$$\tilde{\sigma}_l^2 \leq \alpha_2^2 k_n^2 M^{2l-1}. \quad (2.18)$$

Similarly, we have

$$\tilde{\tilde{\sigma}}_l^2 \leq \alpha_3^2 k_n^2 M^{2l-1} \quad (2.19)$$

for a positive constant α_3 . Therefore, we obtain, outside the exceptional set,

$$|R_l(\omega)| < \lambda_n(\alpha_2 + \alpha_3) k_n M^{l-(1/2)} < \left(\frac{\alpha_2 + \alpha_3}{\alpha_1} \frac{k_n}{t_n} \lambda_n \sigma_l \right) / M^{1/2} < \sigma_l, \quad (2.20)$$

by Lemma 2.1 and (2.1). $\square$

Let us define random events E_p, F_p by

$$\begin{aligned} E_p &= \{\omega; U_{3p}(\omega) \geq \sigma_{3p}, U_{3p+1}(\omega) < -\sigma_{3p+1}\}, \\ F_p &= \{\omega; U_{3p}(\omega) < -\sigma_{3p}, U_{3p+1}(\omega) \geq \sigma_{3p+1}\}. \end{aligned} \quad (2.21)$$

It can be easily seen that

$$\Pr(E_p \cup F_p) = \delta_p \quad (\text{say}) > \delta, \quad (2.22)$$

where $\delta > 0$ is a certain constant. Let η_p be a random variable such that

$$\eta_p = \begin{cases} 1 & \text{on } E_p \cup F_p, \\ 0 & \text{elsewhere.} \end{cases} \quad (2.23)$$

Then we get

$$E(\eta_p) = \delta_p, \quad V(\eta_p) = \delta_p - \delta_p^2. \quad (2.24)$$

Let q be the total number of pairs (U_{3p}, U_{3p+1}) for which

$$\left\lceil \frac{k}{2} \right\rceil + 1 \leq 3p < 3p+1 \leq k, \quad (2.25)$$

q must be at least equal to $[k/3] - [(k/2) + 1]/3 - 1$. Take

$$\eta = \sum \eta_p, \quad (2.26)$$

where the summation is taken over all the q pairs. Applying Tschebyscheff inequality, we have, for $0 < \varepsilon < \delta$,

$$\Pr(\{|\eta - E(\eta)| \geq q\varepsilon\}) \leq \frac{V(\eta)}{q^2\varepsilon^2} \leq \frac{\sum \delta_p}{q^2\varepsilon^2} \leq \frac{1}{q\varepsilon^2}, \quad (2.27)$$

since for n sufficiently large, $\text{Cov}(\eta_i, \eta_j) = 0$ ($i \neq j$). But

$$q \geq \left[\frac{k}{3}\right] - \left[\frac{[k/2] + 1}{3}\right] - 1 \geq \frac{k}{3} - 1 - \left(\frac{(k/2) + 1}{3}\right) - 1 = \frac{1}{6}(k - 14) \geq \mu_1 k, \quad (2.28)$$

where μ_1 is a positive constant. Therefore, outside a set of measure at most μ_2/k ,

$$|\eta - E(\eta)| < q\varepsilon, \quad (2.29)$$

that is,

$$\eta - E(\eta) > -q\varepsilon \quad (2.30)$$

or

$$\eta > E(\eta) - q\varepsilon = \sum \delta_p - q\varepsilon > q(\delta - \varepsilon) \geq \mu_3 k, \quad (2.31)$$

where μ_2 and μ_3 are positive constants. Thus we have proved that outside a set of measure at most μ_2/k , either $U_{3p} \geq \sigma_{3p}$ and $U_{3p+1} < -\sigma_{3p+1}$, or $U_{3p} < -\sigma_{3p}$ and $U_{3p+1} \geq \sigma_{3p+1}$ for at least $\mu_3 k$ values of l .

Define

$$\zeta_p = \begin{cases} 0 & \text{if } |R_{3p}| < \sigma_{3p}, \quad |R_{3p+1}| < \sigma_{3p+1}, \\ 1 & \text{elsewhere.} \end{cases} \quad (2.32)$$

Let $\xi_p = \eta_p - \eta_p \zeta_p$. If $\xi_p = 1$, there is a root of the polynomial in the interval (x_{3p}, x_{3p+1}) . Hence the number of real roots in the interval $(x_{[k/2]+1}, x_k)$ must exceed $\sum \xi_p$, where the summation is taken over all the q pairs. Now, by using Lemma 2.4, we have

$$\begin{aligned} E\left(\sum \eta_p \zeta_p\right) &= \sum E(\eta_p \zeta_p) \leq \sum E(\zeta_p) = \sum \Pr(\zeta_p = 1) \\ &\leq \sum \{\Pr(|R_{3p}| \geq \sigma_{3p}) + \Pr(|R_{3p+1}| \geq \sigma_{3p+1})\} \\ &< \mu_4(k+1) \frac{1}{\lambda_n} e^{-\lambda_n^2/2}, \end{aligned} \quad (2.33)$$

where μ_4 is a constant. Hence we have, for $\beta > 0$,

$$\Pr\left(\left\{\sum \eta_p \zeta_p > \mu_4(k+1) \lambda_n^\beta \frac{1}{\lambda_n} e^{-\lambda_n^2/2}\right\}\right) < \frac{E(\sum \eta_p \zeta_p)}{\mu_4(k+1) \lambda_n^{\beta-1} e^{-\lambda_n^2/2}} < \frac{1}{\lambda_n^\beta}. \quad (2.34)$$

So we get

$$\sum \eta_p \zeta_p \leq \mu_4(k+1)\lambda_n^{\beta-1} e^{-\lambda_n^2/2}, \quad (2.35)$$

except for a set of measure at most $1/\lambda_n^\beta$. Therefore, we have, outside a set of measure at most $\mu_2/k + 1/\lambda_n^\beta$,

$$N_n > \sum \xi_p > \mu_3 k - \mu_4(k+1)\lambda_n^{\beta-1} e^{-\lambda_n^2/2} \geq k(\mu_3 - \varepsilon_1), \quad (2.36)$$

where $0 < \varepsilon_1 < \mu_3$ (since $\mu_4 \lambda_n^{\beta-1} \exp(-\lambda_n^2/2)$ tends to zero as n tends to infinity). But it follows from (2.1) and (2.2) that

$$\begin{aligned} \mu_5 \left(\frac{k_n}{t_n} \right)^2 \lambda_n^2 &\leq M \leq \mu_6 \left(\frac{k_n}{t_n} \right)^2 \lambda_n^2, \\ \frac{\mu_7 \log n}{\log((k_n/t_n)\lambda_n)} &\leq k \leq \frac{\mu_8 \log n}{\log((k_n/t_n)\lambda_n)}, \end{aligned} \quad (2.37)$$

where μ_i , $i = 5, 6, 7, 8$, are constants. Hence we get outside the exceptional set

$$N_n > \frac{\mu_9 \log n}{\log((k_n/t_n)\lambda_n)}, \quad (2.38)$$

where μ_9 is a constant.

Taking $\lambda_n = (t_n/k_n) \exp(\mu_9/\varepsilon_n)$, we obtain

$$N_n > \varepsilon_n \log n \quad (2.39)$$

outside a set of measure at most

$$\frac{\mu}{\varepsilon_n \log n} + \left(\frac{k_n}{t_n} \right)^\beta \exp\left(-\frac{\mu' \beta}{\varepsilon_n}\right), \quad (2.40)$$

where μ and μ' are constants. This completes the proof of the theorem.

Acknowledgment

The author wishes to thank the referee for his/her valuable comments.

References

- [1] J. E. Littlewood and A. C. Offord, "On the number of real roots of a random algebraic equation II," *Proceedings of the Cambridge Philosophical Society*, vol. 35, pp. 133–148, 1939.
- [2] G. Samal, "On the number of real roots of a random algebraic equation," *Proceedings of the Cambridge Philosophical Society*, vol. 58, pp. 433–442, 1962.
- [3] E. A. Evans, "On the number of real roots of a random algebraic equation," *Proceedings of the London Mathematical Society. Third Series*, vol. 15, no. 3, pp. 731–749, 1965.
- [4] G. Samal and M. N. Mishra, "On the lower bound of the number of real roots of a random algebraic equation with infinite variance," *Proceedings of the American Mathematical Society*, vol. 33, pp. 523–528, 1972.

- [5] G. Samal and M. N. Mishra, "On the lower bound of the number of real roots of a random algebraic equation with infinite variance. II," *Proceedings of the American Mathematical Society*, vol. 36, pp. 557–563, 1972.
- [6] G. Samal and M. N. Mishra, "On the lower bound of the number of real roots of a random algebraic equation with infinite variance. III," *Proceedings of the American Mathematical Society*, vol. 39, no. 1, pp. 184–189, 1973.
- [7] M. Sambandham, "On the upper bound of the number of real roots of a random algebraic equation," *The Journal of the Indian Mathematical Society. New Series*, vol. 42, no. 1–4, pp. 15–26, 1978.
- [8] T. Uno and H. Negishi, "On the upper bound of the number of real roots of a random algebraic equation," *The Journal of the Indian Mathematical Society. New Series*, vol. 62, no. 1–4, pp. 215–224, 1996.
- [9] N. Renganathan and M. Sambandham, "On the lower bounds of the number of real roots of a random algebraic equation," *Indian Journal of Pure and Applied Mathematics*, vol. 13, no. 2, pp. 148–157, 1982.
- [10] N. N. Nayak and S. P. Mohanty, "On the lower bound of the number of real zeros of a random algebraic polynomial," *The Journal of the Indian Mathematical Society. New Series*, vol. 49, no. 1–2, pp. 7–15, 1985.
- [11] T. Uno, "On the lower bound of the number of real roots of a random algebraic equation," *Statistics & Probability Letters*, vol. 30, no. 2, pp. 157–163, 1996.
- [12] T. Uno, "Strong result for real zeros of random algebraic polynomials," *Journal of Applied Mathematics and Stochastic Analysis*, vol. 14, no. 4, pp. 351–359, 2001.
- [13] K. Yoshihara, *Weakly dependent stochastic sequences and their applications. Vol. I. Summation theory for weakly dependent sequences*, Sanseido, Tokyo, Japan, 1992.

Takashi Uno: Faculty of Urban Science, Meijo University, 4-3-3 Nijigaoka, Kani,
Gifu 509-0261, Japan
Email address: uno@urban.meijo-u.ac.jp