

Research Article

On the Lower Classes of Some Mixed Fractional Gaussian Processes with Two Logarithmic Factors

Charles El-Nouty

U.F.R. de Mathématiques, Université Pierre et Marie Curie, 175 rue du Chevaleret, 75013 Paris, France

Correspondence should be addressed to Charles El-Nouty, elnouty@ccr.jussieu.fr

Received 5 February 2007; Revised 10 July 2007; Accepted 12 September 2007

Recommended by Hideo Nagai

We introduce the fractional mixed fractional Brownian sheet and investigate the small ball behavior of its sup-norm statistic by establishing a general result on the small ball probability of the sum of two not necessarily independent joint Gaussian random vectors. Then, we state general conditions and characterize the sufficiency part of the lower classes of some statistics of the above process by an integral test. Finally, when we consider the sup-norm statistic, the necessity part is given by a second integral test.

Copyright © 2008 Charles El-Nouty. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let $\{B_{H_i}(s), s \geq 0\}$ be a fractional Brownian motion (FBM) with index $0 < H_i < 1$, $i \in \mathbb{N}^*$, and $\{B_{H_j, H_k}(s, s'), s \geq 0, s' \geq 0\}$ a fractional Brownian sheet (FBS) with index $0 < H_j, H_k < 1$, $j \in \mathbb{N}^*$, $k \in \mathbb{N}^*$. We refer to [1] for further information about the FBM and the FBS. Denote by λ_1 and λ_2 two real numbers such that $\lambda_1 \lambda_2 \neq 0$.

Define a fractional mixed fractional Gaussian process by a suitable combination of some appropriate fractional Gaussian processes. In the sequel, we consider the following three examples.

Example 1.1. The fractional mixed fractional Brownian motion (FMFBM) is defined by

$$X(\omega_1, \omega_2, s) = \lambda_1 s^{H_2} B_{H_1}(\omega_1) + \lambda_2 s^{H_1} B_{H_2}(\omega_2), \quad (1.1)$$

where B_{H_1} and B_{H_2} are independent FBM with $H_1 \neq H_2$.

Example 1.2. The fractional mixed fractional Brownian motion and fractional Brownian sheet (FMFBMFBS) are defined by

$$X(w_1, w_2, w_3, s) = \lambda_1 s^{H_2+H_3} B_{H_1}(w_1) + \lambda_2 s^{H_1} B_{H_2, H_3}(w_2, w_3), \quad (1.2)$$

where the FBM B_{H_1} and the FBS B_{H_2, H_3} are independent.

Example 1.3. The fractional mixed fractional Brownian sheet (FMFBS) is defined by

$$X(w_1, w_2, w_3, w_4, s) = \lambda_1 s^{H_3+H_4} B_{H_1, H_2}(w_1, w_2) + \lambda_2 s^{H_1+H_2} B_{H_3, H_4}(w_3, w_4), \quad (1.3)$$

where B_{H_1, H_2} and B_{H_3, H_4} are independent FBS with $(H_1, H_2) \neq (H_3, H_4)$.

The motivation supporting this paper is threefold.

- (i) The first goal of the FMFBS deals with the potential applications. Since the FMFBM, the FMFBMFBS, and the FMFBS can be analyzed based on the large bodies of knowledge on FBM and FBS, it can be used in the same fields, that is, natural time series in economics, fluctuations in solid, hydrology, and, more recently, by new problems in mathematical finance, telecommunication networks, and the environment (see [2–4]).
- (ii) A second application deals with the small ball probability problem of the sum of two not necessarily joint centered Gaussian random vectors X and Y in a separable Banach space E with norm $\|\cdot\|$ (see [5]). The small ball behavior of the FMFBS under the uniform norm can be investigated as a special case of the small ball probability problem of the sum of two centered Gaussian random vectors, having a log-type small ball factor (see [6]).
- (iii) Last but not least, this article extends El-Nouty's results [6–9] and consequently answers some new questions. Recall first two definitions of the Lévy classes, stated in [10]. Let $\{Z(t), t \geq 0\}$ be a stochastic process defined on the basic probability space (Ω, \mathcal{A}) .

Definition 1.4. The function $f(t), t \geq 0$, belongs to the lower-lower class of the process Z , ($f \in \text{LLC}(Z)$), if, for almost all $\omega \in \Omega$, there exists $t_0 = t_0(\omega)$ such that $Z(t) \geq f(t)$ for every $t > t_0$.

Definition 1.5. The function $f(t), t \geq 0$, belongs to the lower-upper class of the process Z , ($f \in \text{LUC}(Z)$), if, for almost all $\omega \in \Omega$, there exists a sequence $0 < t_1 = t_1(\omega) < t_2 = t_2(\omega) < \dots$ with $t_n \rightarrow +\infty$, as $n \rightarrow +\infty$, such that $Z(t_n) \leq f(t_n)$, $n \in \mathbb{N}^*$.

In the spirit of [6–9, 11], the main aim of this paper is to characterize the lower classes of the uniform norm of the FMFBS for any $0 < H_1, H_2, H_3, H_4 < 1$. More precisely, we want to compare the influence of two FBSs and to measure the weight of a log-type small ball factor versus another one.

2. Main results

Our first result is given in the following theorem.

Theorem 2.1. *Let X and Y be any two joint Gaussian random vectors in a separable Banach space with norm $\|\cdot\|$. Assume that there exist $C_X \geq 1$ and $C_Y \geq 1$ such that one has, for any $\epsilon > 0$ small enough,*

$$\begin{aligned} -C_X &\leq \frac{e^{1/\alpha}}{(\log(1/\epsilon))^\beta} \log \mathbb{P}(\|X\| \leq \epsilon) \leq -\frac{1}{C_X}, \\ -C_Y &\leq \frac{e^{1/\tilde{\alpha}}}{(\log(1/\epsilon))^{\tilde{\beta}}} \log \mathbb{P}(\|Y\| \leq \epsilon) \leq -\frac{1}{C_Y}, \end{aligned} \quad (2.1)$$

with $0 < \alpha, \tilde{\alpha} < +\infty, 0 \leq \beta, \tilde{\beta} < +\infty$ and $(\alpha, \beta) \neq (\tilde{\alpha}, \tilde{\beta})$.

If $(\alpha < \tilde{\alpha})$ or $(\alpha = \tilde{\alpha}$ and $\beta > \tilde{\beta})$, then there exists $K_X \geq C_X$ depending on C_X only such that one has, for any $\epsilon > 0$ small enough,

$$-K_X \leq \frac{e^{1/\alpha}}{(\log(1/\epsilon))^\beta} \log \mathbb{P}(\|X + Y\| \leq \epsilon) \leq -\frac{1}{K_X}. \quad (2.2)$$

Since the study of the lower classes of the FMFBM (resp., FMFBMFBS) under the uniform norm was investigated in [8] (resp., [9]), we focus our attention to the FMFBS. Set

$$Y(t) = \sup_{0 \leq s \leq t} \sup_{0 \leq w_1, w_2, w_3, w_4 \leq s} |X(w_1, w_2, w_3, w_4, s)|, \quad t \geq 0. \quad (2.3)$$

Note first that, by the scaling property, we have, for any $\epsilon > 0$,

$$\begin{aligned} \mathbb{P}(Y(t) \leq \epsilon t^{H_1+H_2+H_3+H_4}) &= \mathbb{P}\left(\sup_{0 \leq s \leq 1} \sup_{0 \leq w_1, w_2, w_3, w_4 \leq s} |X(w_1, w_2, w_3, w_4, s)| \leq \epsilon\right) \\ &= \mathbb{P}(Y(1) \leq \epsilon) := \phi(\epsilon), \end{aligned} \quad (2.4)$$

where ϕ is named the small ball function and $\gamma := H_1 + H_2 + H_3 + H_4$ the scaling factor.

Recall that the small ball behavior of the FBS under the uniform norm was studied in [12, 13].

Set $\alpha = \min(H_1, H_2, H_3, H_4)$, which is in $]0, 1[$. We introduce the number β taking its values in $\{0, 1 + 1/\alpha\}$. As a direct consequence of Theorem 2.1, we have the following corollary.

Corollary 2.2. *There is a constant $K_0, 0 < K_0 \leq 1$, depending on $H_1, H_2, H_3, H_4, \lambda_1$ and λ_2 only, such that one has, for any $\epsilon > 0$ small enough,*

$$\exp\left(-\frac{(\log(1/\epsilon))^\beta}{K_0 e^{1/\alpha}}\right) \leq \phi(\epsilon) \leq \exp\left(-\frac{K_0 (\log(1/\epsilon))^\beta}{e^{1/\alpha}}\right). \quad (2.5)$$

Recall that we suppose $(H_1, H_2) \neq (H_3, H_4)$. In the sequel, there is no loss of generality to assume also that

$$H_1 \leq H_2, \quad H_3 \leq H_4. \quad (2.6)$$

Thus when $(H_1 = H_2, H_3 < H_4 \text{ and } H_1 \leq H_3)$, $(H_1 = H_2, H_3 = H_4 \text{ and } H_1 < H_3)$, or $(H_1 < H_2, H_3 = H_4 \text{ and } H_3 \leq H_1)$, we emphasize that $\beta = 1 + 1/\alpha$, that is, we have a log-type small ball factor.

Note first that the minimum α plays a key role. This is not really surprising. Indeed, this phenomenon was already observed in [8, 9].

It appears that the sufficiency part of the lower classes of Y can be stated in a general framework. Roughly speaking, we follow the same lines as those of [6, 7].

Let $\{Y_0(t), t \geq 0\}$ be a real-valued statistic of the two independent FBSs, B_{H_1, H_2} and B_{H_3, H_4} , such that $Y_0(t)$ is a nondecreasing function of $t \geq 0$.

The following notation is needed. If \mathbb{K} is a Hausdorff compact space, we denote, by $C(\mathbb{K})$, the space of all continuous functions from \mathbb{K} to \mathbb{R} equipped with the classical sup-norm. Let $\mathbb{X} = C([0, 1]^2) \times C([0, 1]^2)$ be the product space equipped with the product topology. Denote, by $\mathbb{L}(B_{H_1, H_2}, B_{H_3, H_4})$, the Gaussian measure associated to B_{H_1, H_2} and B_{H_3, H_4} and defined on \mathbb{B} , the Borel σ -field of \mathbb{X} .

We assume that Y_0 satisfies the following three conditions:

(i) *The scaling condition.* There exists $\gamma_0 > 0$ such that

$$\mathbb{P}(Y_0(t) \leq \epsilon t^{\gamma_0}) = \mathbb{P}(Y_0(1) \leq \epsilon) := \phi(\epsilon). \quad (2.7)$$

(ii) *The convexity condition.* There exists a convex and \mathbb{B} -measurable function $g: (\mathbb{X}, \mathbb{L} \times (B_{H_1, H_2}, B_{H_3, H_4})) \rightarrow \mathbb{R}$ such that, for any $t \geq 0$, $Y_0(t) = g(B_{H_1, H_2}(s_1 t, s_2 t), B_{H_3, H_4}(s_3 t, s_4 t))$; $0 \leq s_1, s_2, s_3, s_4 \leq 1$, and $Y_0(t) < +\infty$, with probability 1.

(iii) *The log-type small ball condition.* There exist $\alpha_0 \in]0, \gamma_0]$, $\beta_0 \in \mathbb{R}$ and a constant K , $0 < K \leq 1$, depending on $H_1, H_2, H_3, H_4, \gamma_0, \alpha_0$ and β_0 only such that we have, for any $\epsilon > 0$ small enough,

$$\exp\left(-\frac{(\log(1/\epsilon))^{\beta_0}}{K\epsilon^{1/\alpha_0}}\right) \leq \phi(\epsilon) \leq \exp\left(-\frac{K(\log(1/\epsilon))^{\beta_0}}{\epsilon^{1/\alpha_0}}\right). \quad (2.8)$$

Note that these conditions generalize those of [6, 7]. The small ball function still plays a key role. The convexity of the function ψ defined by $\psi(\epsilon) = -\log \phi(\epsilon)$, $0 < \epsilon < 1$, is ensured by (ii) (see [14, 15]).

Our second result is given in the following theorem.

Theorem 2.3. *Let $f(t)$ be a positive nondecreasing function of $t \geq 0$. Assume that there exists $m > 0$ such that $(f(t)/t^{\gamma_0 - \alpha_0})(\log t^{\gamma_0}/f(t))^{-\beta_0 \alpha_0} \geq m$.*

If

$$\frac{f(t)}{t^{\gamma_0}} \text{ is bounded and } \int_0^{+\infty} f(t)^{-1/\alpha_0} t^{(\gamma_0/\alpha_0)-1} \left(\log \frac{t^{\gamma_0}}{f(t)}\right)^{\beta_0} \phi\left(\frac{f(t)}{t^{\gamma_0}}\right) dt < +\infty, \quad (2.9)$$

then one has

$$f \in LLC(Y_0). \quad (2.10)$$

The sup-norm statistic Y clearly satisfies the three above conditions with $\gamma_0 = \gamma = H_1 + H_2 + H_3 + H_4$, $\alpha_0 = \alpha = \min(H_1, H_2, H_3, H_4) = \min(H_1, H_3)$, $\beta_0 = \beta \in \{0, 1 + 1/\alpha\}$, and $K = K_0$. Now, we characterize the necessity part of the lower classes of the FMFBMFBS. Our main result is stated in the following theorem.

Theorem 2.4. *Let $f(t)$ be a positive nondecreasing function of $t \geq 0$ such that $f(t)/t^\gamma$ is a nonincreasing function of $t > 0$.*

If

$$f \in LLC(Y), \quad (2.11)$$

then one has

$$\lim_{t \rightarrow +\infty} \frac{f(t)}{t^\gamma} = 0, \quad \int_0^{+\infty} f(t)^{-1/\gamma} \phi\left(\frac{f(t)}{t^\gamma}\right) dt < +\infty. \quad (2.12)$$

First, we can notice that Theorem 2.3 involves γ_0 , α_0 and β_0 . If $\beta_0 = 0$, Theorem 2.3 looks like [7, Theorem 1] or else like [6, Theorem 1.1]. Theorem 2.4 has the same form as the necessity part of [11, Theorem 1.2] or as the theorems obtained in [6–9]. In his previous works on the study of the lower classes, the author showed that the methodology in [11] led to two integral tests, these tests are actually identical when $\gamma = \alpha$ and $\beta = 0$. Here, $\alpha = \min(H_1, H_3) < \gamma = H_1 + H_2 + H_3 + H_4$. This is why the integral tests of Theorems 2.3 and 2.4 have different forms. Moreover, since $\alpha < \gamma$, we must assume, as in [6–9], that $f(t)/t^\gamma$ is not only bounded, but also a nonincreasing function of $t > 0$. This last assumption will play a key role in some proofs. Finally, although they are two different integral tests, Theorems 2.3 and 2.4 are sharp. Indeed, set, if $\beta = 0$,

$$f(t) = \frac{\lambda t^\gamma}{(\log \log t)^\alpha}, \quad t \geq 3, \lambda > 0, \quad (2.13)$$

or else (i.e., $\beta = 1 + (1/\alpha)$)

$$f(t) = t^\gamma \frac{(\lambda \log \log \log t)^{1+\alpha}}{(\log \log t)^\alpha}, \quad t \geq 16, \lambda > 0. \quad (2.14)$$

If λ is small enough, then Theorem 2.3 yields $f \in LLC(Y)$, or else if λ is large enough, then $f \in LUC(Y)$ by applying Theorem 2.4.

In Section 3, we prove Theorem 2.1. The proof of Theorem 2.4 is postponed to Sections 4 and 5. In the latter, we establish some key small ball estimates. Note also that these estimates can be of independent interest. The proofs, which are modifications of those of [6, 7], will be consequently omitted, in particular, the proof of Theorem 2.3.

3. Proof of Theorem 2.1

Recall, first, a Gaussian correlation inequality stated in [5].

Theorem A. Let μ be a centered Gaussian measure on a separable Banach space E . Then for any $0 < \lambda < 1$, and any symmetric convex sets A and B in E ,

$$\mu(A \cap B) \mu(\lambda^2 A + (1 - \lambda^2) B) \geq \mu(\lambda A) \mu((1 - \lambda^2)^{1/2} B). \quad (3.1)$$

In particular,

$$\mu(A \cap B) \geq \mu(\lambda A) \mu((1 - \lambda^2)^{1/2} B). \quad (3.2)$$

Roughly speaking, the proof follows the same lines as those in [15] and will be split into two parts: the lower bound and the upper one.

Part I. The lower bound

Theorem A implies, for any $0 < \delta < 1$ and $0 < \lambda < 1$,

$$\begin{aligned} \mathbb{P}(\|X + Y\| \leq \epsilon) &\geq \mathbb{P}(\|X\| \leq (1 - \delta)\epsilon, \|Y\| \leq \delta\epsilon) \\ &\geq \mathbb{P}(\|X\| \leq \lambda(1 - \delta)\epsilon) \mathbb{P}(\|Y\| \leq (1 - \lambda^2)^{1/2} \delta\epsilon). \end{aligned} \quad (3.3)$$

Then we get, by using (2.1),

$$\log \mathbb{P}(\|X + Y\| \leq \epsilon) \geq -C_X \frac{(\log(1/\lambda(1 - \delta)\epsilon))^\beta}{(\lambda(1 - \delta)\epsilon)^{1/\alpha}} - C_Y \frac{(\log(1/(1 - \lambda^2)^{1/2} \delta\epsilon))^{\tilde{\beta}}}{((1 - \lambda^2)^{1/2} \delta\epsilon)^{1/\tilde{\alpha}}}. \quad (3.4)$$

Hence, since $(\alpha < \tilde{\alpha})$ or $(\alpha = \tilde{\alpha} \text{ and } \beta > \tilde{\beta})$, there exists $C'_X \geq C_X$ depending on C_X only such that we have, for any $\epsilon > 0$ small enough,

$$\frac{\epsilon^{1/\alpha}}{(\log(1/\epsilon))^\beta} \log \mathbb{P}(\|X + Y\| \leq \epsilon) \geq -\frac{C'_X}{(\lambda(1 - \delta))^{1/\alpha}}, \quad (3.5)$$

and the lower bound follows by taking $\delta \rightarrow 0$ and $\lambda \rightarrow 1$.

Part II. The upper bound

A new use of Theorem A implies, for any $0 < \delta < 1$ and $0 < \lambda < 1$,

$$\begin{aligned} \mathbb{P}\left(\|X\| \leq \frac{\epsilon}{(1 - \delta)\lambda}\right) &\geq \mathbb{P}\left(\|X + Y\| \leq \frac{\epsilon}{\lambda}, \|Y\| \leq \frac{\delta\epsilon}{(1 - \delta)\lambda}\right) \\ &\geq \mathbb{P}(\|X + Y\| \leq \epsilon) \mathbb{P}\left(\|Y\| \leq (1 - \lambda^2)^{1/2} \frac{\delta\epsilon}{(1 - \delta)\lambda}\right). \end{aligned} \quad (3.6)$$

Then combining (2.1) with the fact that $(\alpha < \tilde{\alpha})$ or $(\alpha = \tilde{\alpha} \text{ and } \beta > \tilde{\beta})$, there exists $C''_X \geq C_X$ depending on C_X only such that we have, for any $\epsilon > 0$ small enough,

$$\frac{\epsilon^{1/\alpha}}{(\log(1/\epsilon))^\beta} \log \mathbb{P}(\|X + Y\| \leq \epsilon) \leq -\frac{(\lambda(1 - \delta))^{1/\alpha}}{C''_X}, \quad (3.7)$$

and the upper bound follows by taking $\delta \rightarrow 0$ and $\lambda \rightarrow 1$.

By choosing $K_X = \max(C'_X, C''_X)$, we complete the proof of Theorem 2.1.

Remark 3.1. When X and Y are independent, there is a simple proof without using the correlation inequality in the spirit of [5]. A direct proof of Corollary 2.2 can also be done as in [9].

4. Proof of Theorem 2.4, Part I

To simplify the reading of our paper, we introduce the following notation. Set $a_t = f(t)/t^\gamma$ and $b_t = \phi(a_t)$.

Suppose here that, with probability 1, $f(t) \leq Y(t)$ for all t large enough. We want to prove that $\lim_{t \rightarrow +\infty} a_t = 0$ and $\int_0^\infty a_t^{-1/\gamma} b_t (dt/t) < +\infty$.

In the sequel, there is no loss of generality to assume that $f(t)$ is a continuous function of $t \geq 0$.

Lemma 4.1. *One has*

$$\lim_{t \rightarrow +\infty} a_t = 0. \quad (4.1)$$

To prove Theorem 2.4, we will show that $f \in \text{LUC}(Y)$ when $\int_0^\infty a_t^{-1/\gamma} b_t (dt/t) = +\infty$ and $\lim_{t \rightarrow +\infty} a_t = 0$.

Following the same lines as those in [11], our aim is to construct a suitable subset J of \mathbb{N} such that we have the following property for an appropriate family of sets $(E_i)_{i \in J}$ in a basic probability space: given $\epsilon > 0$, there exist a number K and an integer p such that

$$\forall n \in J, \quad n \geq p \implies \sum_{m \in J, m > n} \mathbb{P}(E_n \cap E_m) \leq \mathbb{P}(E_n) \left(K + (1 + \epsilon) \sum_{m \in J, m > n} \mathbb{P}(E_m) \right). \quad (4.2)$$

Lemma 4.2. *When $\int_0^\infty a_t^{-1/\gamma} b_t (dt/t) = +\infty$ and $\lim_{t \rightarrow +\infty} a_t = 0$, one can find a sequence $\{t_n, n \geq 1\}$ with the two following properties:*

$$t_{n+1} \geq t_n(1 + a_{t_n}^{1/\gamma}), \quad \sum_{n=1}^{\infty} b_{t_n} = +\infty. \quad (4.3)$$

Remark 4.3. The condition “ $f(t)/t^\gamma$ is a nonincreasing function of $t > 0$ ” is essential to prove Lemma 4.2 (see [7, page 373]).

To continue the construction of the set J , we need the following definition and notation.

Definition 4.4. Consider the interval $A_k = [2^k, 2^{k+1}[$, $k \in \mathbb{N}$. If $a_{t_i}^{-1/\gamma} \in A_k$, $i \in \mathbb{N}^*$, then one notes $u(i) = k$.

Next, set $I_k = \{i \in \mathbb{N}^*, u(i) = k \in \mathbb{N}\}$ which is finite by Lemma 4.1 and

$$N_k = \exp(K_0 (\gamma \log 2)^\beta k^\beta 2^{\gamma(k-1)/\alpha}), \quad (4.4)$$

where K_0 was defined in Corollary 2.2.

Notation

- (i) $U_{m,k} = \{i \in \mathbb{N}^*, i \in I_k, i < m, \text{card}(I_k \cap [i, m]) \leq N_k\}$, $m \in \mathbb{N}^*$, $k \in \mathbb{N}$;
- (ii) $k_0 = \inf \{n \in \mathbb{N}, 2^{\gamma n/\alpha} \geq 2^{\gamma/\alpha} / (K_0(2^{\gamma/\alpha} - 1)(\gamma \log 2)^\beta) + 2^{2\gamma/\alpha} / K_0^2(2^{\gamma/\alpha} - 1)\}$, (k_0 depends on γ , α , β and K_0 only);
- (iii) $V_m = \bigcup_{k \in \mathbb{N}} U_{m,k}$, where m is fixed, $u(m) = k_1$, and $k \geq k_1 + k_0$;
- (iv) $W = \bigcup_{m \geq 1} V_m$.

Now, we can define the set J as follows:

$$J = \mathbb{N}^* - W. \quad (4.5)$$

Since it is assumed that $f(t)/t^\gamma$ is a nonincreasing function of $t > 0$ (being a particular case of the condition “ $f(t)/t^\gamma$ is bounded”),

$$i < m \implies k = u(i) \leq u(m) = k_1. \quad (4.6)$$

Moreover, since $k_0 \geq 1$, we get $k < k_1 + k_0$. Hence V_m is always an empty set (by construction). Thus we obtain $J = \mathbb{N}^*$.

We have, by Lemma 4.2,

$$\sum_{n \in J} b_{t_n} = +\infty. \quad (4.7)$$

Lemma 4.5. $n \in J$, $m \in J$, $n < m$, such that $\text{card}(I_{u(m)} \cap [n, m]) > \exp(K_0 2^{u(m)-1})$, one has

$$\frac{t_m}{t_n} \geq \exp\left(\exp\left(\frac{K_0}{4} 2^{\min(u(n), u(m))}\right)\right). \quad (4.8)$$

Proof. Set $k = u(n)$, $k_1 = u(m)$ and $G = I_{k_1} \cap [n, m] = \{i_1, i_2, \dots, i_z\}$, where $n \leq i_1 < i_2 < \dots < i_z \leq m$. We have

$$\frac{t_m}{t_n} = \frac{t_m}{t_{i_z}} \frac{t_{i_z}}{t_{i_{z-1}}} \dots \frac{t_{i_1}}{t_n}. \quad (4.9)$$

Note that, when $i \in I_{k_1}$, we have $t_{i+1} \geq t_i(1 + a_i^{1/\gamma}) \geq t_i(1 + 2^{-k_1-1})$. Moreover, since $\text{card}(G) > \exp(K_0 2^{k_1-1})$ by hypothesis, (4.9) implies

$$\frac{t_m}{t_n} \geq \exp(\exp(K_0 2^{k_1-1}) \log(1 + 2^{-k_1-1})) \geq \exp\left(\exp\left(\frac{K_0}{4} 2^{k_1}\right)\right), \quad (4.10)$$

when n hence k_1 are large enough.

Thus, since $k \leq k_1$, (4.10) implies (4.8).

The proof of Lemma 4.5 is now complete. \square

5. Proof of Theorem 2.4, Part II

Consider, now, the events $E_n = \{Y(t_n) < f(t_n)\}$. We have directly $\mathbb{P}(E_n) = b_{t_n}$, and, by (4.7), $\sum_{n \in J} b_{t_n} = +\infty$. To prove (4.2), we remark that, given $n \in J$, J can be rewritten as follows: $J = J' \cup (\bigcup_{k \in \mathbb{N}} J_k) \cup J''$, where $J' = \{m \in J, t_n \leq t_m \leq 2t_n\}$, $J_k = \{m \in J \cap I_k, t_m > 2t_n, \text{card}(I_k \cap [n, m]) \leq \exp(K_0 2^{k-1})\}$, and $J'' = J - (J' \cup (\bigcup_{k \in \mathbb{N}} J_k))$.

Our first key small ball estimate is given in the following lemma.

Lemma 5.1. *Consider $0 < t < u$, and $\theta, \nu > 0$. Then one has*

$$\mathbb{P}(\{Y(t) \leq \theta t^\gamma\} \cap \{Y(u) \leq \nu\}) \leq \exp(K_5) \mathbb{P}(Y(t) \leq \theta t^\gamma) \exp\left(-\frac{K_5(u-t)}{\nu^{1/\gamma}}\right), \quad (5.1)$$

where K_5 depends on $H_1, H_2, H_3, H_4, \lambda_1$, and λ_2 only.

Proof. Set $F_1 = \{Y(t) \leq \theta t^\gamma\}$ and $F_2 = \{Y(u) \leq \nu\}$. We have

$$\begin{aligned} \mathbb{P}(F_1 \cap F_2) &= \mathbb{P}\left(F_1 \cap \left\{ \sup_{0 \leq s \leq u} \sup_{0 \leq w_1, w_2, w_3, w_4 \leq s} |X(w_1, w_2, w_3, w_4, s)| \leq \nu \right\}\right) \\ &\leq \mathbb{P}\left(F_1 \cap \left\{ \sup_{t \leq s \leq u} \sup_{0 \leq w_1, w_2, w_3, w_4 \leq s} |X(w_1, w_2, w_3, w_4, s)| \leq \nu \right\}\right) \\ &\leq \mathbb{P}\left(F_1 \cap \left\{ \sup_{t \leq s \leq u} \sup_{0 \leq w_1, w_2 \leq s} |\lambda_1 s^{H_3+H_4} B_{H_1, H_2}(w_1, w_2)| \leq \nu \right\}\right). \end{aligned} \quad (5.2)$$

Denote, by $[x]$, the integer part of a real x . Let $\delta > 0$. We consider the sequence $t_k, k \in \{0, \dots, n\}$, where $t_0 = t, t_{k+1} = t_k + \delta$, and $n = [(u-t)/\delta]$. Consider also the rectangles $R_j = [t_j, t_{j+1}] \times [t_j, t_{j+1}]$, where $j \in \{0, \dots, n-1\}$. Their area is $(t_{j+1} - t_j)^2 = \delta^2$. Let G_j be the event defined by

$$G_j = F_1 \cap \left\{ \sup_{t \leq s \leq t_j} \sup_{0 \leq w_1, w_2 \leq s} |\lambda_1 s^{H_3+H_4} B_{H_1, H_2}(w_1, w_2)| \leq \nu \right\}. \quad (5.3)$$

We have $F_1 \cap F_2 \subset G_j$.

Moreover, we have also

$$G_{j+1} \subset G_j \cap \{Z_j \leq 4\nu\}, \quad (5.4)$$

where

$$Z_j = \lambda_1 t_{j+1}^{H_3+H_4} (B_{H_1, H_2}(t_{j+1}, t_{j+1}) - B_{H_1, H_2}(t_j, t_{j+1}) - B_{H_1, H_2}(t_{j+1}, t_j) + B_{H_1, H_2}(t_j, t_j)). \quad (5.5)$$

Before rewriting Z_j , we recall the integral representation of B_{H_k, H_l} , $k \in \mathbb{N}^*, l \in \mathbb{N}^*$, given by

$$B_{H_k, H_l}(s_k, s_l) = \int_{-\infty}^{s_k} \int_{-\infty}^{s_l} g_{H_k}(s_k, u_k) g_{H_l}(s_l, u_l) W(d(u_k, u_l)), \quad (5.6)$$

where $W(u_k, u_l)$, $u_k \in \mathbb{R}$, $u_l \in \mathbb{R}$, is a standard Brownian sheet, $g_H(s, u) = k_{2H}^{-1} (\max(s - u, 0)^{H-1/2} - \max(-u, 0)^{H-1/2})$, and k_{2H} is a normalizing constant.

Hence Z_j can be rewritten by (5.6) as follows: $Z_j = Z_{j,1} + Z_{j,2}$, where

$$Z_{j,1} = \lambda_1 k_{2H_1}^{-1} k_{2H_2}^{-1} t_{j+1}^{H_3+H_4} \int_{t_j}^{t_{j+1}} \int_{t_j}^{t_{j+1}} (t_{j+1} - u_1)^{H_1-1/2} (t_{j+1} - u_2)^{H_2-1/2} W(d(u_1, u_2)). \quad (5.7)$$

Note also that $Z_{j,1}$ and $Z_{j,2}$ are independent.

Since $\mathbb{P}(|Z_{j,1} + x| \leq 4\nu)$ is maximum at $x = 0$ and $Z_{j,1}$ and G_j are independent, we have

$$\mathbb{P}(G_{j+1}) \leq \mathbb{P}(G_j) \mathbb{P}(|Z_{j,1}| \leq 4\nu). \quad (5.8)$$

The integral representation of $Z_{j,1}$ implies that $\mathbb{E}(Z_{j,1}) = 0$ and

$$\text{Var } Z_{j,1} = \frac{\lambda_1^2 k_{2H_1}^{-2} k_{2H_2}^{-2}}{4H_1 H_2} \delta^{2(H_1+H_2)} t_{j+1}^{2(H_3+H_4)} \geq \frac{\lambda_1^2 k_{2H_1}^{-2} k_{2H_2}^{-2}}{4H_1 H_2} \delta^{2\gamma} := L^2 \delta^{2\gamma}. \quad (5.9)$$

Denote, by Φ , the distribution function of the absolute value of a standard Gaussian random variable. Then we obtain

$$\mathbb{P}(Z_{j+1}) \leq \mathbb{P}(Z_j) \Phi\left(\frac{4\nu}{L \delta^\gamma}\right), \quad (5.10)$$

and therefore, $\mathbb{P}(F_1 \cap F_2) \leq \mathbb{P}(F_1) \Phi(4\nu/L \delta^\gamma)^n$.

Choosing $\delta = \nu^{1/\gamma}$, we get $K_5 = -\log \Phi(2/L)$. Lemma 5.1 is proved. \square

Lemma 5.2. $\sum_{m \in J'} \mathbb{P}(E_n \cap E_m) \leq K' b_{t_n}$ and $\sum_{m \in (\cup_k J_k)} \mathbb{P}(E_n \cap E_m) \leq K'' b_{t_n}$, where K' and K'' are numbers.

Proof. Setting $u = t_m$, $t = t_n$, $\theta = a_{t_n}$ and $\nu = f(t_m)$, Lemma 5.1 implies

$$\mathbb{P}(E_n \cap E_m) \leq \exp(K_5) b_{t_n} \exp\left(-\frac{K_5(t_m - t_n)}{f(t_m)^{1/\gamma}}\right). \quad (5.11)$$

Consider, first, the case when $m \in J'$.

Lemma 4.2 implies that, for all $i \geq n$, we have $t_{i+1} - t_i \geq t_i a_i^{1/\gamma} = f(t_i)^{1/\gamma} \geq f(t_n)^{1/\gamma}$. Then we can establish

$$t_m - t_n \geq (m - n) f(t_n)^{1/\gamma}, \quad f(t_m) \leq f(t_n) \left(\frac{t_m}{t_n}\right)^\gamma \leq 2^\gamma f(t_n). \quad (5.12)$$

Combining (5.11) with (5.12), we get

$$\mathbb{P}(E_n \cap E_m) \leq \exp(K_5) b_{t_n} \exp\left(-\frac{K_5(m - n)}{2}\right), \quad (5.13)$$

which is the first part of Lemma 5.2.

Consider, now, the case $m \in J_k$.

Combining (5.11) with the definition of J_k , we have

$$\mathbb{P}(E_n \cap E_m) \leq \exp(K_5) b_{t_n} \exp\left(-\frac{K_5}{2(a_{t_m})^{1/\gamma}}\right). \quad (5.14)$$

Since $u(m) = k$, we get

$$\mathbb{P}(E_n \cap E_m) \leq \exp(K_5) b_{t_n} \exp(-K_5 2^{k-1}), \quad (5.15)$$

and consequently, by noting that $\text{card } J_k \leq \text{card}(I_k \cap [n, m]) \leq \exp(K_0 2^{k-1})$ and by assuming $K_0 < K_5$, we have

$$\sum_{m \in J_k} \mathbb{P}(E_n \cap E_m) \leq \exp(K_5) b_{t_n} \exp((K_0 - K_5) 2^{k-1}). \quad (5.16)$$

Lemma 5.2 is, therefore, proved. \square

To deal with the set J'' , we first state a standard large deviation result and a technical lemma (see [6]).

Lemma A. *Let $X = \{X(s_1, s_2), (s_1, s_2) \in [0, 1]^2\}$ be a separable real-valued centered Gaussian process such that $X(0, 0) = 0$ with probability 1 and satisfying, for any $[s_1, s_1 + h_1] \times [s_2, s_2 + h_2] \subset [0, 1]^2$,*

$$\left(\mathbb{E}X([s_1, s_1 + h_1] \times [s_2, s_2 + h_2])^2\right)^{1/2} \leq \kappa(h_1, h_2) \leq c_\kappa h_1^{\alpha_1} h_2^{\alpha_2}, \quad \alpha_1 > 0, \alpha_2 > 0, \quad (5.17)$$

where

$$X([s_1, t_1] \times [s_2, t_2]) = \int_{[s_1, t_1] \times [s_2, t_2]} X(d(u_1, u_2)). \quad (5.18)$$

Then one has, for $c_\kappa^{-1} \delta > 1$,

$$\mathbb{P}\left(\sup_{(s_1, s_2) \in [0, 1]^2} |X(s_1, s_2)| \geq \delta\right) \leq \frac{1}{C} \exp(-C(c_\kappa^{-1} \delta)^2), \quad (5.19)$$

where C is a positive constant independent of c_κ and δ .

Lemma B. *One has, for $\epsilon_1 > \epsilon/2$, where ϵ is small enough,*

$$\exp\left(-K_3 \frac{|\epsilon_1 - \epsilon| (\log(1/\epsilon))^\beta}{\epsilon^{1+1/\alpha}}\right) \leq \frac{\phi(\epsilon_1)}{\phi(\epsilon)} \leq \exp\left(K_3 \frac{|\epsilon_1 - \epsilon| (\log(1/\epsilon))^\beta}{\epsilon^{1+1/\alpha}}\right), \quad (5.20)$$

where $K_3 > 0$.

Building on Lemmas A and B, we can establish our last key small ball estimate in the following result.

Lemma 5.3. Let λ be a real number such that $1/2 < \lambda < 1$. Set

$$r = \min \left(\frac{1 - \max(H_1, H_2, H_3, H_4)}{3}, \frac{(1 - \lambda)\alpha}{3} \right). \quad (5.21)$$

Then one has, for $u \geq 2t$,

$$\begin{aligned} & \mathbb{P}(Y(t) \leq \theta t^\lambda, Y(u) \leq \nu u^\lambda) \\ & \leq \phi(\theta)\phi(\nu) \exp \left(2 \left(\frac{t}{u} \right)^r K_3 \left(\frac{(\log(1/\theta))^\beta}{\theta^{1+(1/\alpha)}} + \frac{(\log(1/\nu))^\beta}{\nu^{1+(1/\alpha)}} \right) \right) \\ & \quad + 3 \left(\frac{1}{C_{12,2}} \exp \left(- \frac{C_{12,2}}{4\lambda_1^2 K_{H_1,2}^2} \left(\frac{u}{t} \right)^r \right) + \frac{1}{C_{34,2}} \exp \left(- \frac{C_{34,2}}{4\lambda_2^2 K_{H_3,2}^2} \left(\frac{u}{t} \right)^r \right) \right) \\ & \quad + 3 \left(\frac{1}{C_{12,1}} \exp \left(- \frac{C_{12,1}}{4\lambda_1^2 K_{H_1,1}^2} \left(\frac{u}{t} \right)^r \right) + \frac{1}{C_{34,1}} \exp \left(- \frac{C_{34,1}}{4\lambda_2^2 K_{H_3,1}^2} \left(\frac{u}{t} \right)^r \right) \right), \end{aligned} \quad (5.22)$$

where $K_{H_1,1}, K_{H_1,2} > 0$ depend on H_1 ($H_1 \leq H_2$) only, $K_{H_3,1}, K_{H_3,2} > 0$ depend on H_3 ($H_3 \leq H_4$) only, $K_3 > 0$ is defined as in Lemma B, and $C_{12,1}, C_{12,2}, C_{34,1}, C_{34,2} > 0$ are defined as in Lemma A.

Proof. Set $Q = \mathbb{P}(Y(t) \leq \theta t^\lambda, Y(u) \leq \nu u^\lambda)$.

Set $v = \sqrt{ut}$. If $t = o(u)$, then $t = o(v)$ and $v = o(u)$.

Based on (5.6), B_{H_1, H_2} and B_{H_3, H_4} can be split as follows:

$$B_{H_1, H_2} = B_{H_1, H_2, 1} + B_{H_1, H_2, 2}, \quad B_{H_3, H_4} = B_{H_3, H_4, 1} + B_{H_3, H_4, 2}, \quad (5.23)$$

where we have, for $(i, j) \in \{(1, 2), (3, 4)\}$,

$$B_{H_i, H_j, 1}(w_i, w_j) = \int_{|x_i| \leq v} \int_{-\infty}^{w_j} g_{H_i}(w_i, x_i) g_{H_j}(w_j, x_j) W(d(x_i, x_j)). \quad (5.24)$$

Note that $B_{H_1, H_2, 1}$ and $B_{H_1, H_2, 2}$ are independent as $B_{H_3, H_4, 1}$ and $B_{H_3, H_4, 2}$.

Equation (5.23) implies that the FMFBS X can be rewritten as follows: $X = X_1 + X_2$, where

$$X_i(w_1, w_2, w_3, w_4, s) = \lambda_1 s^{H_3 + H_4} B_{H_1, H_2, i}(w_1, w_2) + \lambda_2 s^{H_1 + H_2} B_{H_3, H_4, i}(w_3, w_4). \quad (5.25)$$

Set

$$Y_i(t) = \sup_{0 \leq s \leq t} \sup_{0 \leq w_1, w_2, w_3, w_4 \leq s} |X_i(w_1, w_2, w_3, w_4, s)|, \quad t \geq 0, \quad i \in \{1, 2\}. \quad (5.26)$$

Then, given $\delta > 0$, we have (see [11])

$$Q \leq \phi(\theta + 2\delta) \phi(\nu + 2\delta) + 3 \mathbb{P}(Y_2(t) > \delta t^\lambda) + 3 \mathbb{P}(Y_1(u) > \delta u^\lambda). \quad (5.27)$$

Equation (5.20) implies

$$\phi(\theta + 2\delta) \leq \phi(\theta) \exp \left(2\delta K_3 \left(\frac{(\log(1/\theta))^\beta}{\theta^{1+(1/\alpha)}} \right) \right), \quad (5.28)$$

and, consequently,

$$\phi(\theta + 2\delta)\phi(\nu + 2\delta) \leq \phi(\theta)\phi(\nu) \exp\left(2\delta K_3 \left(\frac{(\log(1/\theta))^\beta}{\theta^{1+(1/\alpha)}} + \frac{(\log(1/\nu))^\beta}{\nu^{1+(1/\alpha)}}\right)\right). \quad (5.29)$$

If we choose $\delta = (t/u)^r$, then we get the first term of the RHS of Lemma 5.3.

Next, we want to obtain an upper bound of

$$\begin{aligned} & \mathbb{P}(Y_2(t) > \delta t^r) \\ &= \mathbb{P}\left(\sup_{0 \leq s \leq 1} \sup_{0 \leq w_1, w_2, w_3, w_4 \leq s} |\lambda_1 s^{H_3+H_4} L_{H_1, H_2, 2}(w_1, w_2) + \lambda_2 s^{H_1+H_2} L_{H_3, H_4, 2}(w_3, w_4)| > \delta\right), \end{aligned} \quad (5.30)$$

where we have, for $(i, j) \in \{(1,2), (3,4)\}$,

$$L_{H_i, H_j, 2}(w_i, w_j) = \int_{|x_i| \geq \nu/t}^{\int_{-\infty}^{w_j}} g_{H_i}(w_i, x_i) g_{H_j}(w_j, x_j) W(d(x_i, x_j)). \quad (5.31)$$

We can show, by standard computations, that

$$\begin{aligned} \mathbb{P}(Y_2(t) > \delta t^r) &\leq \mathbb{P}\left(\sup_{0 \leq w_1, w_2 \leq 1} |L_{H_1, H_2, 2}(w_1, w_2)| > \frac{\delta}{2|\lambda_1|}\right) \\ &\quad + \mathbb{P}\left(\sup_{0 \leq w_3, w_4 \leq 1} |L_{H_3, H_4, 2}(w_3, w_4)| > \frac{\delta}{2|\lambda_2|}\right). \end{aligned} \quad (5.32)$$

Denote, by σ_H , the covariance function of a FBM B_H . Set $\sigma_{H,2}$, the covariance function of the process $\{B_{H,2}(w), 0 \leq w \leq 1\}$, defined by

$$B_{H,2}(w) = \int_{|x| \geq \nu/t} g_H(w, x) \widetilde{W}(dx), \quad (5.33)$$

where $\widetilde{W}(x)$, $x \in \mathbb{R}$, is a Wiener process.

Since

$$\mathbb{E}(L_{H_1, H_2, 2}(w_1, w_2) L_{H_1, H_2, 2}(w'_1, w'_2)) = \sigma_{H_1, 2}(w_1, w'_1) \times \sigma_{H_2}(w_2, w'_2), \quad (5.34)$$

we have, for any $[w_1, w_1 + h_1] \times [w_2, w_2 + h_2] \subset [0, 1]^2$,

$$\begin{aligned} & \mathbb{E}\left(L_{H_1, H_2, 2}([w_1, w_1 + h_1] \times [w_2, w_2 + h_2])^2\right) \\ &= \mathbb{E}\left(\int_{[w_1, w_1 + h_1] \times [w_2, w_2 + h_2]} L_{H_1, H_2, 2}(d(x_1, x_2)) \right. \\ &\quad \times \left. \int_{[w_1, w_1 + h_1] \times [w_2, w_2 + h_2]} L_{H_1, H_2, 2}(d(x'_1, x'_2))\right) \\ &\leq \int_{w_1}^{w_1 + h_1} \int_{w_1}^{w_1 + h_1} |\sigma_{H_1, 2}(x_1, x'_1)| dx_1 dx'_1 \\ &\quad \times \int_{w_2}^{w_2 + h_2} \int_{w_2}^{w_2 + h_2} |\sigma_{H_2}(x_2, x'_2)| dx_2 dx'_2 := I \times II. \end{aligned} \quad (5.35)$$

Consider II first. We get, by the inequality of Cauchy-Schwarz,

$$II \leq \int_{w_2}^{w_2+h_2} \int_{w_2}^{w_2+h_2} x_2^{H_2} x_2'^{H_2} dx_2 dx_2' \leq h_2^2. \quad (5.36)$$

Consider I now.

A straight computation implies that there exists $K_{H_1,2} > 0$ depending on H_1 such that

$$\mathbb{E}(B_{H_1,2}(x_1))^2 \leq K_{H_1,2}^2 x_1^2 (v/t)^{2H_1-2}, \quad (5.37)$$

and, consequently, by the inequality of Cauchy-Schwarz,

$$|\sigma_{H_1,2}(x_1, x_1')| \leq K_{H_1,2}^2 x_1 x_1' (v/t)^{2H_1-2}. \quad (5.38)$$

So we get

$$I \leq K_{H_1,2}^2 (v/t)^{2H_1-2} h_1^2. \quad (5.39)$$

Hence, combining (5.35) with (5.36) and (5.39), we have

$$\mathbb{E}\left(L_{H_1, H_2, 2}([w_1, w_1 + h_1] \times [w_2, w_2 + h_2])^2\right) \leq K_{H_1,2}^2 (v/t)^{2H_1-2} h_1^2 h_2^2. \quad (5.40)$$

An application of Lemma A with $\alpha_1 = \alpha_2 = 1$, $c_\kappa = K_{H_1,2}(v/t)^{H_1-1}$, and $c_\kappa^{-1}\delta > 1$ implies that

$$\mathbb{P}\left(\sup_{0 \leq w_1, w_2 \leq 1} |L_{H_1, H_2, 2}(w_1, w_2)| > \frac{\delta}{2|\lambda_1|}\right) \leq \frac{1}{C_{12,2}} \exp\left(-\frac{C_{12,2}}{K_{H_1,2}^2 (v^2/t^2)^{H_1-1}} \frac{\delta^2}{4\lambda_1^2}\right). \quad (5.41)$$

Similarly, we can establish

$$\mathbb{P}\left(\sup_{0 \leq w_3, w_4 \leq 1} |L_{H_3, H_4, 2}(w_3, w_4)| > \frac{\delta}{2|\lambda_2|}\right) \leq \frac{1}{C_{34,2}} \exp\left(-\frac{C_{34,2}}{K_{H_3,2}^2 (v^2/t^2)^{H_3-1}} \frac{\delta^2}{4\lambda_2^2}\right). \quad (5.42)$$

Set $\delta = (t/u)^r$. Recall that $v^2 = ut$ and $r \leq (1 - \max(H_1, H_2, H_3, H_4))/3$. Combining (5.32) with (5.41) and (5.42), we get

$$\mathbb{P}(Y_2(t) > \delta t^r) \leq \frac{1}{C_{12,2}} \exp\left(-\frac{C_{12,2}}{4\lambda_1^2 K_{H_1,2}^2} \left(\frac{u}{t}\right)^r\right) + \frac{1}{C_{34,2}} \exp\left(-\frac{C_{34,2}}{4\lambda_2^2 K_{H_3,2}^2} \left(\frac{u}{t}\right)^r\right), \quad (5.43)$$

that is the second term of the RHS of Lemma 5.3.

Finally, we can establish a similar result for $\mathbb{P}(Y_1(u) > \delta u^r)$, that is,

$$\mathbb{P}(Y_1(u) > \delta u^r) \leq \frac{1}{C_{12,1}} \exp\left(-\frac{C_{12,1}}{4\lambda_1^2 K_{H_1,1}^2} \left(\frac{u}{t}\right)^r\right) + \frac{1}{C_{34,1}} \exp\left(-\frac{C_{34,1}}{4\lambda_2^2 K_{H_3,1}^2} \left(\frac{u}{t}\right)^r\right), \quad (5.44)$$

which achieves the proof of Lemma 5.3. \square

Finally, we state the last technical lemma.

Lemma 5.4. *There exists an integer p such that if $n > \sup_{s \leq p} (\sup I_s)$, then for $m \in J''$, $m > n$, given $\epsilon > 0$, one has $\mathbb{P}(E_n \cap E_m) \leq (1 + \epsilon) b_{t_n} b_{t_m}$.*

Proof. Let $u(n) = k'$ and $u(m) = k_1$. We have, by Lemma 4.5,

$$\frac{t_m}{t_n} \geq \exp\left(\exp\left(\frac{K_0}{4} 2^{\min(k', k_1)}\right)\right). \quad (5.45)$$

Let $p \in \mathbb{N}$. Then $k' > p$ and $k_1 > p$. Thus we have $\min(k', k_1) > p$.

Set $t = t_n$, $u = t_m$, $\theta = a_{t_n}$ and $\nu = a_{t_m}$. Note that $\log(1/\theta) \leq 1/\theta \leq 2^{(k'+1)\alpha}$, $\log(1/\nu) \leq 1/\nu \leq 2^{(k_1+1)\alpha}$, and $1/b_{t_n} b_{t_m} = \exp(\psi(\theta) + \psi(\nu))$.

By using Lemma 5.3 and letting $p \rightarrow +\infty$, we end the proof of Lemma 5.4. \square

Lemmas 5.2 and 5.4 yield that (4.2) holds. Combining Borel-Cantelli's second lemma with (4.2) and (4.7), we show that, given $\epsilon > 0$,

$$\sum_{n \in J} \mathbb{P}(E_n) \geq \frac{1+2K}{\epsilon} \implies \frac{1}{1+2\epsilon} \leq \mathbb{P}\left(\bigcup_{n \in J} E_n\right) = \mathbb{P}\left(\bigcup_{n \in J} \{Y(t_n) \leq f(t_n)\}\right), \quad (5.46)$$

and, consequently, $f \in \text{LUC}(Y)$. The proof of Theorem 2.4 is now complete.





Acknowledgment

The author thanks the referee for the insightful comments.

References

- [1] W. V. Li and Q.-M. Shao, "Gaussian processes: inequalities, small ball probabilities and applications," in *Stochastic Processes: Theory and Methods*, pp. 533–597, North-Holland, Amsterdam, 2001.
- [2] P. Cheridito, "Mixed fractional Brownian motion," *Bernoulli*, vol. 7, no. 6, pp. 913–934, 2001.
- [3] F. Gassmann and D. Bürki, "Experimental investigation of atmospheric dispersion over the swiss plain—experiment SIESTA," *Boundary-Layer Meteorology*, vol. 41, pp. 295–307, 1987.
- [4] F. Gassmann, P. Gaglione, S. E. Gryning, et al., "Experimental Investigation of Atmospheric Dispersion over Complex Terrain in a Prealpine Region (experiment SIESTA)," Swiss Federal Institute for Reactor Research EIR 604, 1986.
- [5] W. V. Li, "A Gaussian correlation inequality and its applications to small ball probabilities," *Electronic Communications in Probability*, vol. 4, pp. 111–118, 1999.
- [6] C. El-Nouty, "The influence of a log-type small ball factor in the study of the lower classes," *Bulletin des Sciences Mathématiques*, vol. 129, no. 4, pp. 318–338, 2005.
- [7] C. El-Nouty, "On the lower classes of fractional Brownian motion," *Studia Scientiarum Mathematicarum Hungarica*, vol. 37, no. 3-4, pp. 363–390, 2001.
- [8] C. El-Nouty, "The fractional mixed fractional Brownian motion," *Statistics & Probability Letters*, vol. 65, no. 2, pp. 111–120, 2003.

- [9] C. El-Nouty, "The fractional mixed fractional Brownian motion and fractional Brownian sheet," *ESAIM. Probability and Statistics*, vol. 11, pp. 448–465, 2007.
- [10] P Révész, *Random Walk in Random and Nonrandom Environments*, World Scientific, Teaneck, NJ, USA, 1990.
- [11] M. Talagrand, "Lower classes for fractional Brownian motion," *Journal of Theoretical Probability*, vol. 9, no. 1, pp. 191–213, 1996.
- [12] E. Belinsky and W. Linde, "Small ball probabilities of fractional brownian sheets via fractional integration operators," *Journal of Theoretical Probability*, vol. 15, no. 3, pp. 589–612, 2002.
- [13] D. M. Mason and Z. Shi, "Small deviations for some multi-parameter Gaussian processes," *Journal of Theoretical Probability*, vol. 14, no. 1, pp. 213–239, 2001.
- [14] C. Borell, "Convex measures on locally convex spaces," *Arkiv för Matematik*, vol. 12, pp. 239–252, 1974.
- [15] M. A Lifshits, *Gaussian Random Functions*, vol. 322, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1995.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

