

Research Letter

The Effect of Tb and Sm Ions on the Photochromic Behavior of Two Spiropyrans of Benzoxazine Series in Solution

Esam Bakeir,¹ G. M. Attia,¹ Maria Lukyanova,² Boris Lukyanov,² and M. S. A. Abdel-Mottaleb¹

¹Nanophotochemistry Lab, Department of Chemistry, Faculty of Science, Ain Shams University, Abbassia, Cairo 11566, Egypt

²Department of Chemistry, Institute of Physical and Organic Chemistry, Southern Federal University, Rostov-on-Don 344006, Russia

Correspondence should be addressed to M. S. A. Abdel-Mottaleb, phochem08@photoenergy.org

Received 18 November 2007; Accepted 17 January 2008

Recommended by Bern Kohler

The photochromism of [7'-hydroxy-8'-formyl-3-methyl-4-oxospiro[1,3-benzoxazin-2,2'-[2H-1]benzopyran], SP(I), [7'-hydroxy-8'-formyl-3-benzyl-4-oxospiro[1,3-benzoxazin-2,2'-[2H-1]benzopyran] SP(II) and their coordination with Tb³⁺ and Sm³⁺ ions have been studied in DMF. UV/vis induced-color development due to heterolytic bond cleavage of SP(I) and SP(II) is greatly influenced by complexation with the lanthanide ions. The irradiation-induced color enhancement due to ring opening and thermal decoloration of the open forms of SP(I), SP(II) follows first-order kinetics. Physical characteristics of the studied systems such as colorability and relaxation time of thermal bleaching parameters were determined. Moreover, light-energy transfer-induced luminescence of lanthanide ions via coordination with the two spirobenzoxazines was monitored.

Copyright © 2008 Esam Bakeir et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. INTRODUCTION

Recently, many profitable applications of photochromic dyes, particularly spirooxazines, either as passive or active devices, have been proposed [1–13].

In some cases, it has been reported that thermal equilibrium between the closed (colorless form) and opened form (colored merocyanine quinonoid form) is affected by the change in solvent polarity [14, 15], since polar solvents promote the formation of the colored form at room temperature in the absence of light. The equilibrium between both forms is strongly displaced upon irradiation to the side of open-chain colored photomerocyanine and spontaneously converts to the colorless spiro form to reach thermal equilibrium immediately after removing the light [15]. The metal-ion coordination ability of photochromic spirocyclic compounds adequately substituted is of a great interest and is being the topic of several recent studies [12, 13]. Search for molecules possessing better performances is valuable and it is important to continue to explore this subject. Here, we report on the possibility to stabilize the colored open forms of the recently synthesized spirobenzoxazines SP(I) and SP(II) toward thermal bleaching by coordination with Tb³⁺ and Sm³⁺ ions in polar DMF solvent. Moreover, the expected light-energy transfer-induced characteristic lumines-

cence of lanthanide ions via complexation with the merocyanine quinonoid open forms of the two spirobenzoxazines will be explored.

2. EXPERIMENTAL

2.1. Materials

The metal chlorides (Sigma-Aldrich, 99.99%) were used as received. The synthesis of [7'-hydroxy-8'-formyl-3-methyl-4-oxospiro [1,3-benzoxazin-2,2'-[2H-1]benzopyran], SP(I), [7'-hydroxy-8'-formyl-3-benzyl-4-oxospiro [1,3-benzoxazin-2,2'-[2H-1] benzopyran] SP(II) were described previously [16, 17]. Spectroscopic pure grade solvents were used.

2.2. Instruments

UV-visible absorption spectra were recorded on a range (200–650 nm) using λ -Helios SP Pye-Unicam spectrophotometer and/or Ocean Optics US4000 fiber optics spectrophotometer. Continuous irradiation experiments were performed using a 150 W xenon arc lamp (PTI-LPS-220 Photon Technology International, USA) operated at 70 W. Photochemical reactions were carried out in the spectrophotometric quartz cell with a homogeneously spread light on

TABLE 1: The experimental values of absorption maxima of SP and its open form (λ_{SP} , λ_{open}), SP ring opening rate constant (k_1), thermal open form ring closure rate constant (k_{-1}), equilibrium constant of the reversible reaction (K_e), relaxation time of open form ($\tau_{open-SP}$), and colorability in DMF at 295 ± 1 K. Error limits of the kinetics parameters are in the order of about 5%.

Compound	λ_{SP} (nm)	λ_{open} (nm)	$k_{-1} \times 10^3$ (min ⁻¹)	$k_1 \times 10^3$ (min ⁻¹)	$\tau_{open-SP}$ (min)	K_e	Colorability M ⁻¹ cm ⁻¹ × 10 ⁻²
SP(I)	276	409	3.16	192	316	60	7.36
	370						
SP(II)	276	404	7.96	529	125	66	1.30
	370						

the cell window to avoid stirring. Measurements were made on aerated solutions.

Fluorescence spectra were measured in the range (290–750 nm) using Shimadzu RF5301 (PC) spectrofluorophotometer.

2.3. Kinetics measurements

The ring-closure reaction after photocoloration was monitored directly after removal of light at room temperature. First-order rate constants were obtained from the linear $\ln A$ versus time descending curves. By extrapolation of the obtained $\ln A/t$ plots to zero time, the absorbance A_0 of the open form or its complexes at $t = 0$ was related to their “colorabilities” [18–21] using the expression ($A_0/c_{SP}b$), where c_{SP} is the initial concentration of SP and b is the optical path length.

3. RESULTS AND DISCUSSION

3.1. Absorption spectra

The absorption spectrum of SP(I) and SP(II) displayed two bands at 276 and 370 nm, Table 1, which are solvent independent (methanol, ethanol, and DMF solvents were tried). UV light-induced color development of both compounds was only remarkably observed in DMF solution.

Upon irradiation of 60 μ M SP(I) solutions, the absorption spectra at 276 nm and 370 nm decreased and a new band at 409 nm appeared and its intensity increased by increasing the irradiation time. Addition of lanthanide chloride solution in DMF showed no change in the absorption band of SP(I) in the dark. However, irradiation of the (1 : 1 molar ratio) solution of SP(I) in presence of Tb³⁺ and Sm³⁺, respectively, led to a new band at 412 nm and 414 nm, Figure 1. Similar behavior observed for SP(II) in presence of lanthanide ions (Tb³⁺, Sm³⁺) in DMF, Table 1. Thermal bleaching was monitored spectrophotometrically. First-order kinetics rate constants of the reversible close \rightleftharpoons open reactions (see Scheme 2) were determined graphically (Figure 2, as an example) and data are collected in Tables 1 and 2.

Color development rate constant (k_1) of the photomercyanine form and the thermal bleaching rate constant (k_{-1}) are used to estimate the equilibrium constant ($K_e = k_1/k_{-1}$) of the forward photochemical ring opening reaction and the backward thermal bleaching one. The results of K_e are summarized in Table 1 for the SP(I) and SP(II). Data for Ln³⁺-SP (I and II) appear in Table 2. The relaxation time of the open





FIGURE 1: UV/vis absorption spectra of SP(I) and its complexes with Tb³⁺ and Sm³⁺ ions in DMF before and after light irradiation ([SP(I)] = [Ln³⁺] = 60 μ M) (at 295 ± 1 K).



SCHEME 1

form ($\tau_{open-SO}$) given in Table 1 was obtained from the first-order rate constant using the expression $\tau = 1/k_{-1}$ [18–21]. The obtained relaxation time of SP and its complexes with lanthanide ions are relatively high [22], reflecting highly stabilized color form. The colored open MC form of the methyl-substituted SP(I) is more stable than that of the bulky benzyl derivative reflecting the influence of the constituent’s nature and size. Moreover, the presence of the lanthanide ions

3.3. Conclusion

The substituent nature and the presence of lanthanide metal ions in solution of recently synthesized photochromic spirobenzopyrans of benzoxazine series induced significant changes in its photochromic parameters. It could be generally concluded that the presence of lanthanide ions significantly enhances the rate constants of both color development of the light-induced formation of the open merocyanine-like quinonoid species and the rate constant of its thermal bleaching. In most cases, colorability is enhanced. Moreover, strong characteristic sensitized luminescence of the lanthanide ions was observed due to efficient population of its emissive states via energy transfer from the open forms of the spirobenzoxazines.

ACKNOWLEDGEMENT

Russian Foundation for Basic Research supported the Russian team (Grant 07-03-00234).

REFERENCES

- [1] F. Ebisawa, M. Hoshino, and K. Sukegawa, "Self-holding photochromic polymer Mach—Zehnder optical switch," *Applied Physics Letters*, vol. 65, no. 23, pp. 2919–2921, 1994.
- [2] M. Seibold, H. Port, and H. C. Wolf, "Fulgides as light switches for intra-supermolecular energy transfer," *Molecular Crystals and Liquid Crystals*, vol. 283, no. 1, pp. 75–80, 1996.
- [3] M. Hamano and M. Irie, "Rewritable near-field optical recording on photochromic thin films," *Japanese Journal of Applied Physics*, vol. 35, no. 3, pp. 1764–1767, 1996.
- [4] M. Irie and M. Mohri, "Thermally irreversible photochromic systems. Reversible photocyclization of diarylethene derivatives," *Journal of Organic Chemistry*, vol. 53, no. 4, pp. 803–808, 1988.
- [5] R. Wortmann, P. M. Lundquist, R. J. Twieg, et al., "A novel sensitized photochromic organic glass for holographic optical storage," *Applied Physics Letters*, vol. 69, no. 12, pp. 1657–1659, 1996.
- [6] R. M. Tarkka, M. E. Talbot, D. J. Brady, and G. B. Schuster, "Holographic storage in a near-ir sensitive photochromic dye," *Optics Communications*, vol. 109, no. 1-2, pp. 54–58, 1994.
- [7] J. P. Hagen, I. Becerra, D. Drakulich, and R. O. Dillon, "Effect of antenna porphyrins and phthalocyanines on the photochromism of benzospiropyrans in poly(methyl methacrylate) films," *Thin Solid Films*, vol. 398-399, no. 1, pp. 104–109, 2001.
- [8] R. C. Bertelson, "Photochromic processes involving heterolytic cleavage," in *Photochromism*, G. H. Brown, Ed., vol. 3 of *Techniques of Chemistry*, pp. 45–431, John Wiley & Sons, New York, NY, USA, 1971.
- [9] R. Guglielmetti, "Spiroprans have been extensively studied," in *Photochromism: Molecules and Systems*, H. Dürr and H. Bouas-Laurent, Eds., vol. 40 of *Studies in Organic Chemistry*, pp. 314–466, Elsevier, Amsterdam, The Netherlands, 1990.
- [10] J. B. Flannery Jr., "Photo- and thermochromic transients from substituted 1',3',3'-trimethylindolinobenzospiropyrans," *Journal of the American Chemical Society*, vol. 90, no. 21, pp. 5660–5671, 1968.
- [11] T. Bercovici, R. Heiligman-Rim, and E. Fischer, "Photochromism in spiropyrans. Part VI trimethylindolino-benzospiropyran and its derivatives," *Molecular Photochemistry*, vol. 1, pp. 23–55, 1969.
- [12] V. I. Minkin, "Photo-, thermo-, solvato-, and electrochromic spiroheterocyclic compounds," *Chemical Reviews*, vol. 104, no. 5, pp. 2751–2776, 2004.
- [13] A. V. Chernyshev, N. A. Voloshin, I. M. Raskita, A. V. Metelitsa, and V. I. Minkin, "Photo- and ionochromism of 5'-(4,5-diphenyl-1,3-oxazol-2-yl) substituted spiro[indoline-naphthopyrans]," *Journal of Photochemistry and Photobiology A*, vol. 184, no. 3, pp. 289–297, 2006.
- [14] T. Deligeorgiev, S. Minkovska, B. Jeliakova, and S. Rakovsky, "Synthesis of photochromic chelating spironaphthoxazines," *Dyes and Pigments*, vol. 53, no. 2, pp. 101–108, 2002.
- [15] S. Minkovska, B. Jeliakova, E. Borisova, L. Avramov, and T. Deligeorgiev, "Substituent and solvent effect on the photochromic properties of a series of spiroindolinonaphthooxazines," *Journal of Photochemistry and Photobiology A*, vol. 163, no. 1-2, pp. 121–126, 2004.
- [16] Yu. S. Alekseenko, B. Lukyanov, A. N. Utenyshev, et al., "Photo- and thermochromic spiranes. 24. Novel photochromic spiropyrans from 2,4-dihydroxyisophthalaldehyde," *Chemistry of Heterocyclic Compounds*, vol. 42, no. 6, pp. 803–812, 2006.
- [17] B. Lukyanov and M. B. Lukyanova, "Spiroprans: synthesis, properties, and application," *Chemistry of Heterocyclic Compounds*, vol. 41, no. 3, pp. 281–311, 2005.
- [18] G. Favaro, V. Malatesta, U. Mazzucato, G. Ottavi, and A. Romani, "Thermally reversible photoconversion of spiroindoline-naphtho-oxazines to photomerocyanines: a photochemical and kinetic study," *Journal of Photochemistry and Photobiology A*, vol. 87, no. 3, pp. 235–241, 1995.
- [19] S. Kawauchi, H. Yoshida, N. Yamashina, M. Ohira, S. Saeda, and M. Irie, "A new photochromic spiro[3H-1,4-oxazine]," *Bulletin of the Chemical Society of Japan*, vol. 63, no. 1, pp. 267–268, 1990.
- [20] J.-L. Pozzo, A. Samat, R. Guglielmetti, and D. de Keukeleire, "Solvatochromic and photochromic characteristics of new 1,3-dihydrospiro[2H-indole-2,2'-[2H]-bipyrido[3,2-f][2,3-h][1,4]benzoxazines]," *Journal of the Chemical Society, Perkin Transactions 2*, no. 7, pp. 1327–1332, 1993.
- [21] V. Pimienta, C. Frouté, M. H. Deniel, D. Lavabre, R. Guglielmetti, and J. C. Micheau, "Kinetic modelling of the photochromism and photodegradation of a spiro[indoline-naphthoxazine]," *Journal of Photochemistry and Photobiology A*, vol. 122, no. 3, pp. 199–204, 1999.
- [22] B. B. Safoklov, B. Lukyanov, A. O. Bulanov, et al., "Photo- and thermochromic spiropyrans. 21. 2,8''-Formyl-3,6''-dimethyl-4-oxospiro(3,4-dihydro-2H-1,3-benzoxazine-2,2''-[2H]chromene) possessing photochromic properties in the solid phase," *Russian Chemical Bulletin*, vol. 51, no. 3, pp. 462–466, 2002.
- [23] K. A. Chibisov and H. Görner, "Complexes of spiropyran-derived merocyanines with metal ions: relaxation kinetics, photochemistry and solvent effects," *Chemical Physics*, vol. 237, no. 3, pp. 425–442, 1998.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

