

Two New Cytotoxic Candidaspongiolides from an Indonesian Sponge

Agus TRIANTO,^{a,b*} Idam HERMAWAN,^a Toshimasa SUZUKA,^a and Junichi
TANAKA^{a,*}


^a Department of Chemistry, Biology and Marine Science, University of the Ryukyus,
Senbaru 1, Nishihara, Okinawa 903-0213, Japan.

^b Department of Marine Sciences, Faculty of Fisheries and Marine Sciences,
University of Diponegoro, Semarang, Central Java 50234, Indonesia.


Agus Trianto: trianto_telawur@yahoo.co.id, TEL/FAX +81-98-895-8560

Supporting Information


Supporting information available: ¹H NMR spectra (for compounds **1**, **2**, **3**, and **4**) and
¹³C NMR spectra (for compounds **1**, **2**, and **3**).




^1H NMR of compounds **2** (upper spectrum) and **1** (lower spectrum) (Jeol 500 MHz, CDCl_3).




¹H NMR of compounds **2** (upper spectrum) and **3** (lower spectrum) (Bruker 500 MHz, CD₃OD).



^{13}C NMR spectrum of compound **1** (Jeol 500 MHz, CDCl_3).



^{13}C NMR spectrum of compound **3** (Jeol 500 MHz, CD_3OD).



¹H NMR spectra of **4** derived from compounds (**1** upper spectrum) and **2** (lower spectrum) (500 MHz, CDCl₃).