Research Article

A Logistic L -Moment-Based Analog for the Tukey g - h , g , h , and h - h System of Distributions

Todd C. Headrick¹ and Mohan D. Pant²

¹ Section on Statistics and Measurement, Department of EPSE, Southern Illinois University Carbondale, 222-J Wham Building, Carbondale, IL 62901-4618, USA

² Department of Curriculum and Instruction, University of Texas at Arlington, 320B Science Hall, Arlington, TX 76019, USA

Correspondence should be addressed to Todd C. Headrick, headrick@siu.edu

Received 19 May 2012; Accepted 20 June 2012

Academic Editors: F. Fagnola, D. Fiems, and C. Proppe

Copyright © 2012 T. C. Headrick and M. D. Pant. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper introduces a standard logistic L -moment-based system of distributions. The proposed system is an analog to the standard normal conventional moment-based Tukey g - h , g , h , and h - h system of distributions. The system also consists of four classes of distributions and is referred to as (i) asymmetric γ - κ , (ii) log-logistic γ , (iii) symmetric κ , and (iv) asymmetric κ_L - κ_R . The system can be used in a variety of settings such as simulation or modeling events—most notably when heavy-tailed distributions are of interest. A procedure is also described for simulating γ - κ , γ , κ , and κ_L - κ_R distributions with specified L -moments and L -correlations. The Monte Carlo results presented in this study indicate that estimates of L -skew, L -kurtosis, and L -correlation associated with the γ - κ , γ , κ , and κ_L - κ_R distributions are substantially superior to their corresponding conventional product-moment estimators in terms of relative bias and relative standard error.

1. Introduction

Simulating or modeling phenomena that involve heavy-tailed distributions have increasingly become of interest in many areas. Some examples include investigations associated with modeling stream flow and flood frequency [1–3], regional frequency analysis [4], aircraft landing processes [5], disaster analysis [6], finance [7], signal and image processing [8], and latent traits in the social and behavioral sciences [9]. Further, it is also common practice for methodologists to investigate the type I error and power properties associated with inferential statistics using heavy-tailed distributions [10–14]. In many cases, these investigations may only require an elementary transformation to produce heavy-tailed distributions with specified values of conventional skew and kurtosis. A system of transformations that is

particularly well suited for this task is the Tukey g - h , g , h , and h - h classes, which can be used for simulating or modeling univariate and multivariate heavy-tailed distributions [14–20].

The quantile function associated with g - h distributions can be succinctly described as

$$q(Z) = g^{-1}(e^{gZ} - 1)e^{hZ^2}, \quad (1.1)$$

where $Z \sim N(0, 1)$, the parameters $g \neq 0$ and $h \geq 0$ control the skew and kurtosis of $q(Z)$, and the quantile functions for the one-parameter log-normal g or symmetric h distributions can be obtained by taking the $\lim_{h \rightarrow 0} q(Z)$ or $\lim_{g \rightarrow 0} q(Z)$, respectively. The class of asymmetric h - h distributions can be obtained by a doubling technique of h distributions that is described in [18, 21]. Hence, an attractive feature of the Tukey system (g - h , g , h , h - h classes) is that it is computationally efficient because each class of distributions only requires the knowledge of one or two parameters and an algorithm that generates standard normal random deviates.

One problem that arises in the context of heavy-tailed distributions is that conventional moment-based estimators of skew and kurtosis have unfavorable attributes insofar as they can be substantially biased, have high variance, or can be influenced by outliers [4, 22–25]. To obviate this problem, L -moment-based estimators such as L -skew and L -kurtosis were introduced to address the limitations associated with conventional estimators [22]. Specifically, some of the advantages that L -moments have over conventional moments are that they (i) exist whenever the mean of the distribution exists, (ii) are nearly unbiased for all sample sizes and distributions, and (iii) are more robust in the presence of outliers [4, 22–25].

Thus, it would be advantageous to have a L -moment-based Tukey system of transformations for the purpose of simulating or modeling univariate and multivariate heavy-tailed distributions. However, because of the complexities associated with (1.1) and the fact that the standard normal distribution function is not available in closed form, the derivation of a L -moment-based Tukey system would be problematic. We would note that the Tukey h and h - h classes of distributions have been characterized in the context of L -moments and the L -correlation (see [21]).

In view of the above, the present aim is to introduce an L -moment Tukey system analog that is based on the standard logistic distribution. More specifically, the system consists of four quantile functions that produce continuous symmetric and asymmetric distributions with specified values of L -skew, L -kurtosis, and L -correlation. The four classes of distributions are referred to as (i) asymmetric γ - κ , (ii) log-logistic γ , (iii) symmetric κ , and (iv) asymmetric κ_L - κ_R . Some of the advantages that the new system has is that the class of log-logistic (γ) distributions has demonstrated to be more efficient than its log-normal (g) counterpart in terms of computing its hazard function or when censored data are encountered in the context of survival analysis [26]. Note that the κ and κ_L - κ_R classes of distributions were recently introduced in an article by Headrick and Pant [27].

The rest of the paper is outlined as follows. In Section 2, the cumulative distribution function and probability density function as well as other properties associated with the four classes of distributions are derived. A summary of univariate L -moment theory is also provided, and the derivations of the systems of equations for specifying values of L -skew and L -kurtosis for the four classes of distributions are subsequently provided. In Section 3, the coefficient of L -correlation is introduced, and the equations are developed for determining intermediate correlations for specified L -correlations associated with the four classes of distributions. In Section 4, the steps for implementing a simulation procedure are described. Numerical examples and the results of a simulation are also provided to confirm

the derivations and compare the new methodology with its conventional moment-based counterpart. In Section 5, the results of the simulation are discussed.

2. Methodology

2.1. Definitions and Properties for the System of γ - κ , γ , κ , and κ_L - κ_R Distributions

The derivation of the cumulative distribution function (cdf), probability density function (pdf), and other properties associated with the system of γ - κ , γ , κ , and κ_L - κ_R distributions begins with the following definitions.

Definition 2.1. Let X be a random variable that has a standard logistic distribution with cdf and pdf expressed as

$$F_X(x) = \Psi(x) = \Pr(X \leq x) = \frac{1}{(1 + e^{-x})}, \quad (2.1)$$

$$f_X(x) = \psi(x) = \frac{e^{-x}}{(1 + e^{-x})^2}, \quad -\infty < x < +\infty. \quad (2.2)$$

Definition 2.2. Let the quantile function $q(X)$ associated with the system of γ - κ , γ , κ , and κ_L - κ_R distributions be defined as

$$q(X) = q_{\gamma,\kappa}(X) = \gamma^{-1} \left(e^{\gamma X} - 1 \right) e^{\kappa|X|}, \quad (2.3)$$

$$q(X) = q_{\gamma,0}(X) = \lim_{\kappa \rightarrow 0} q_{\gamma,\kappa}(X) = \gamma^{-1} \left(e^{\gamma X} - 1 \right), \quad (2.4)$$

$$q(X) = q_{0,\kappa}(X) = \lim_{\gamma \rightarrow 0} q_{\gamma,\kappa}(X) = X e^{\kappa|X|}, \quad (2.5)$$

$$q(X) = q_{\kappa_L, \kappa_R}(X) = \begin{cases} X e^{\kappa_L|X|}, & \text{for } X \leq 0, \\ X e^{\kappa_R|X|}, & \text{for } X \geq 0, \end{cases} \quad (2.6)$$

where $q(X)$ in (2.3)–(2.6) is a strictly monotone increasing function with real-valued parameters γ , κ , κ_L , κ_R . Equations (2.3)–(2.6) produce distributions defined as (i) asymmetric γ - κ ($\gamma \neq 0$, $\kappa \geq 0$), (ii) log-logistic γ ($\gamma \neq 0$), (iii) symmetric κ ($\kappa \geq 0$), and (iv) asymmetric κ_L - κ_R ($\kappa_L \geq 0$, $\kappa_R \geq 0$, $\kappa_L \neq \kappa_R$). The parameter $\pm\gamma$ in (2.3) and (2.4) controls the degree of skew associated with a distribution. Taking the negative of γ will change the direction of the skew but not its magnitude that is, $q_{-\gamma,\kappa}(X) = -q_{\gamma,\kappa}(-X)$. The parameter κ in (2.3) and (2.5) controls the tail weight of a distribution where the function $e^{\kappa|X|}$ (i) preserves symmetry, (ii) is increasing for $X \geq 0$ and $\kappa \geq 0$, and (iii) produces increased tail weight as the value of κ becomes larger. Analogous to κ , the real-valued parameter κ_L (κ_R) in (2.6) controls the left (right) tail weight of a κ_L - κ_R distribution.

Theorem 2.3. The cdf and pdf associated with the γ - κ , γ , κ , and κ_L - κ_R classes of distributions in (2.3)–(2.6) are

$$F_{q(X)}(x) = \Psi(x), \quad (2.7)$$

$$f_{q(X)}(x) = \frac{\psi(x)}{q'(x)} \quad (2.8)$$

with derivatives $q'(x)$ expressed as

$$\begin{aligned} q'(x) &= q'_{\gamma,\kappa}(x) = e^{\gamma x + \kappa|x|} + \gamma^{-1} \left(e^{\kappa|x|} (e^{\gamma x} - 1) \kappa \frac{d|x|}{dx} \right), \\ q'(x) &= q'_{\gamma,0}(x) = e^{\gamma x}, \\ q'(x) &= q'_{0,\kappa}(x) = e^{\kappa|x|} + x\kappa e^{\kappa|x|} \frac{d|x|}{dx}, \\ q'(x) &= q'_{\kappa_L,\kappa_R}(x) = \begin{cases} e^{\kappa_L|x|} - x\kappa_L e^{\kappa_L|x|}, & \text{for } x \leq 0, \\ e^{\kappa_R|x|} + x\kappa_R e^{\kappa_R|x|}, & \text{for } x \geq 0, \end{cases} \end{aligned} \quad (2.9)$$

where $q'(x) > 0$ and $q'(x = 0) = 1$ for (2.9).

Proof. The requirement that $q(X)$ in (2.3)–(2.6) is a strictly monotone increasing function implies that an inverse function (q^{-1}) exists and thus $y = F_{q(X)}(x) = \Psi(x)$. Differentiating both sides with respect to $q(x)$ yields $dy/dq(x) = f_{q(x)}(x)$. Hence, $dy/dq(x) = (dy/dx)/(dq(x)/dx) = \psi(x)/q'(x)$. Whence, the pdf in (2.8) integrates to one because $\psi(x)$ is the logistic pdf in (2.2) and will be nonnegative on its support $q(x)$ for $x \in (-\infty, +\infty)$ given from Definition 2.2 that $\gamma \neq 0$, $\kappa \geq 0$, $\kappa_L \geq 0$, $\kappa_R \geq 0$, and where $\lim_{x \rightarrow \pm\infty} \psi(x)/q'(x) = 0$ as $\lim_{x \rightarrow \pm\infty} \psi(x) = 0$ and $\lim_{x \rightarrow \pm\infty} 1/q'(x) = 0$. $\square$

Definition 2.4. If the monotonicity assumption ($q'(x) > 0$) holds for the pdf in (2.8) for all $x \in (-\infty, +\infty)$, then (2.8) is defined to be a *global pdf*.

Remark 2.5. Inspection of (2.2) and (2.3)–(2.9) indicates that the height of a global pdf in (2.8) for any γ - κ , γ , κ , or κ_L - κ_R distribution at $x = 0$ will be $\psi(x) = 1/4$.

Remark 2.6. The mode associated with a global pdf in (2.8) is located at $q(\tilde{x})$ where $x = \tilde{x}$ is the critical number that solves $d(\psi(x)/q'(x))/dq(x) = 0$ and globally maximizes $\psi(\tilde{x})/q'(\tilde{x})$ at the mode $q(\tilde{x})$. It is noted that a global pdf in (2.8) will have a global maximum because the standard logistic pdf in (2.2) has a global maximum, and the transformations in (2.3)–(2.6) are assumed to be strictly monotone increasing functions.

Remark 2.7. The median associated with a global pdf in (2.8) is located at $q(x = 0) = 0$. This can be shown by letting $\Psi(x) = 0.50$ denote the 50th percentile. In general, we must have $x = 0$ such that $\Psi(0) = \Pr\{X \leq 0\} = 0.50$ holds in (2.7) for the standard logistic distribution. As such, the limit of the quantile function $q(x)$ locates the median at $\lim_{x \rightarrow 0} q(x) = 0$ for the system of γ - κ , γ , κ , and κ_L - κ_R distributions.

Remark 2.8. In terms of the κ_L - κ_R (or κ) class of distributions, the monotonicity assumption ($q'(x) > 0$) holds *locally* in (2.8) for cases where $\kappa_L < 0$ and $(1/\kappa_L < x < 0$ or $0 \leq x < -1/\kappa_R)$. This leads to Definition 2.9.

Definition 2.9. A κ_L - κ_R (or κ) distribution is defined to have a *local* pdf if the monotonicity assumption holds in accordance with Remark 2.8 and $1 - \Pr\{q(x \leq |1/\kappa_L|)\} \leq \varepsilon$, where ε is a user specified threshold probability based on the cdf in (2.7) (e.g., $\varepsilon \leq 0.001$).

Examples of γ - κ , γ , κ , and κ_L - κ_R distributions based on the cdf and pdf in (2.7) and (2.8) are presented in Figure 1 through Figure 4, respectively. The heavy-tailed distributions in Figures 1(a), 2(a), 3(b), and 4(a) are used in the simulation portion of this study in Section 4. In the next section, univariate L -moments are introduced, and the L -moments for the system of γ - κ , γ , κ , and κ_L - κ_R distributions are subsequently derived, and other properties are discussed.

2.2. Preliminaries on Univariate L -Moments

Let $Y_1, \dots, Y_j, \dots, Y_n$ be independent and identically distributed random variables each with continuous pdf $f_Y(y)$, cdf $F_Y(y)$, order statistics denoted as $Y_{1:n} \leq \dots \leq Y_{j:n} \leq \dots \leq Y_{n:n}$, and L -moments defined in terms of either linear combinations of (i) expectations of order statistics or (ii) probability weighted moments (β_i). For the purposes considered herein, the first four L -moments associated with $Y_{j:n}$ are expressed as [4, pages 20–22]

$$\lambda_1 = E[Y_{1:1}] = \beta_0, \quad (2.10)$$

$$\lambda_2 = \frac{1}{2}E[Y_{2:2} - Y_{1:2}] = 2\beta_1 - \beta_0, \quad (2.11)$$

$$\lambda_3 = \frac{1}{3}E[Y_{3:3} - 2Y_{2:3} + Y_{1:3}] = 6\beta_2 - 6\beta_1 + \beta_0, \quad (2.12)$$

$$\lambda_4 = \frac{1}{4}E[Y_{4:4} - 3Y_{3:4} + 3Y_{2:4} - Y_{1:4}] = 20\beta_3 - 30\beta_2 + 12\beta_1 - \beta_0, \quad (2.13)$$

where the β_i are determined from

$$\beta_i = \int y \{F_Y(y)\}^i f_Y(y) dy, \quad (2.14)$$

where $i = 0, \dots, 3$. The coefficients associated with β_i in (2.14) are obtained from shifted orthogonal Legendre polynomials and are computed as shown in [4, page 20] or in [22].

The L -moments λ_1 and λ_2 in (2.10) and (2.11) are measures of location and scale and are the arithmetic mean and one-half the coefficient of mean difference, respectively. Higher order L -moments are transformed to dimensionless quantities referred to as L -moment ratios defined as $\tau_r = \lambda_r/\lambda_2$ for $r \geq 3$, and where τ_3 and τ_4 are the analogs to the conventional measures of skew and kurtosis. In general, L -moment ratios are bounded in the interval $-1 < \tau_r < 1$ as is the index of L -skew (τ_3) where a symmetric distribution implies that all L -moment ratios with odd subscripts are zero. Other smaller boundaries can be found for more specific

Pdf characteristics	Parameters	Percentiles
Mean = 0.3284	Skew: $\alpha_3 = 3.488$	$q(x)_{0.05} = -2.96843$
Median = 0	Kurtosis: $\alpha_4 = 120$	$q(x)_{0.25} = -1.09985$
Mode = 0	L-skew: $\tau_3 = 0.1664$	$q(x)_{0.75} = 1.29689$
Max(height) = 0.25	L-kurtosis: $\tau_4 = 0.2506$	$q(x)_{0.95} = 4.6168$

(a)

Pdf characteristics	Parameters	Percentiles
Mean = -0.492	Skew: $\alpha_3 = -14.68$	$q(x)_{0.05} = -5.84095$
Median = 0	Kurtosis: $\alpha_4 = +\infty$	$q(x)_{0.25} = -1.411107$
Mode = 0	L-skew: $\tau_3 = -0.2296$	$q(x)_{0.75} = 1.07221$
Max(height) = 0.25	L-kurtosis: $\tau_4 = 0.2892$	$q(x)_{0.95} = 2.79766$

(b)

Figure 1: Two asymmetric γ - κ pdfs and their cdfs. The parameters of skew (α_3) and kurtosis (α_4) are based on (A.2) in Appendix A. The parameters of L -skew (τ_3) and L -kurtosis (τ_4) are based on (2.19) and (2.20).

cases. For example, the index of L -kurtosis (τ_4) has the boundary condition for continuous distributions of [28]

$$\frac{5\tau_3^2 - 1}{4} < \tau_4 < 1. \quad (2.15)$$

Pdf characteristics	Parameters	Percentiles
Mean = 0.3448	Skew: $\alpha_3 = 2.485$	$q(x)_{0.05} = -2.225279$
Median = 0	Kurtosis: $\alpha_4 = 26.57$	$q(x)_{0.25} = -0.986291$
Mode = -0.3895	L-skew: $\tau_3 = 0.2$	$q(x)_{0.75} = 1.228655$
Max(height) = 0.2603	L-kurtosis: $\tau_4 = 0.2$	$q(x)_{0.95} = 4.0099$

(a)

Pdf characteristics	Parameters	Percentiles
Mean = 0.8033	Skew: $\alpha_3 = +\infty$	$q(x)_{0.05} = -1.92255$
Median = 0	Kurtosis: $\alpha_4 = +\infty$	$q(x)_{0.25} = -0.889015$
Mode = -0.7187	L-skew: $\tau_3 = 0.4$	$q(x)_{0.75} = 1.37961$
Max(height) = 0.2947	L-kurtosis: $\tau_4 = 0.3$	$q(x)_{0.95} = 5.61784$

(b)

Figure 2: Two log-logistic γ pdfs and their cdfs. The parameters of skew (α_3) and kurtosis (α_4) are based on (A.2) in Appendix A. The parameters of L -skew (τ_3) and L -kurtosis (τ_4) are based on (2.23) and (2.24).

2.3. L -Moments for the System of γ - κ , γ , κ , and κ_L - κ_R Distributions

The derivation of the first four L -moments associated with the γ - κ class of distributions begins by defining the probability weighted moments based on (2.14) in terms of (2.1)–(2.3) as

$$\beta_i = \int_{-\infty}^{+\infty} q(x, \gamma, \kappa) \{\Psi(x)\}^i \psi(x) dx. \quad (2.16)$$

Pdf characteristics	Parameters	Percentiles
Mean = 0	Skew: $\alpha_3 = 0$	$q(x)_{0.05} = -3.41145$
Median = 0	Kurtosis: $\alpha_4 = 3.585$	$q(x)_{0.25} = -1.16065$
Mode = 0	L-skew: $\tau_3 = 0$	$q(x)_{0.75} = 1.16065$
Max(height) = 0.25	L-kurtosis: $\tau_4 = 0.1876$	$q(x)_{0.95} = 3.41145$

(a)

Pdf characteristics	Parameters	Percentiles
Mean = 0	Skew: $\alpha_3 = 0$	$q(x)_{0.05} = -4.57946$
Median = 0	Kurtosis: $\alpha_4 = 39.83$	$q(x)_{0.25} = -1.295424$
Mode = 0	L-skew: $\tau_3 = 0$	$q(x)_{0.75} = 1.295424$
Max(height) = 0.25	L-kurtosis: $\tau_4 = 0.2958$	$q(x)_{0.95} = 4.57946$

(b)

Figure 3: Two symmetric κ pdfs and their cdfs. The parameters of skew (α_3) and kurtosis (α_4) are based on (A.2) in Appendix A. The parameters of L-skew (τ_3) and L-kurtosis (τ_4) are based on (2.27) and (2.28).

In general, to obtain finite values of $\lambda_1, \lambda_2, \tau_3, \tau_4$ based on (2.16), we must have (i) $\gamma + \kappa < 1$, (ii) $\kappa < 1$, and (iii) $1 + \gamma > \kappa$. As such, integrating (2.16) for $i = 0, 1, 2, 3$ and using (2.10)–(2.13) gives $\lambda_1, \lambda_2, \tau_3$, and τ_4 as

$$\lambda_1 = \frac{((- \gamma - \kappa)h_1 + (\gamma - \kappa)h_2 + (-\gamma + \kappa)h_3 + 2\kappa h_4 + (\gamma + \kappa)h_5 - 2\kappa h_6)}{(2\gamma)}, \quad (2.17)$$

$$\lambda_2 = \frac{(2\gamma - (\gamma + \kappa)^2 h_1 + (\gamma - \kappa)^2 (h_2 - h_3) + (\gamma + \kappa)^2 h_5)}{(2\gamma)}, \quad (2.18)$$

Global Pdf characteristics	Parameters	Percentiles
Mean = 0.177	Skew: $\alpha_3 = 1.83$	$q(x)_{0.05} = -3.38644$
Median = 0	Kurtosis: $\alpha_4 = 23.48$	$q(x)_{0.25} = -1.157465$
Mode = 0	L-skew: $\tau_3 = 0.1$	$q(x)_{0.75} = 1.275371$
Max(height) = 0.25	L-kurtosis: $\tau_4 = 0.25$	$q(x)_{0.95} = 4.3919$

(a)

Local Pdf characteristics	Parameters	Percentiles
Mean = 0.331	Skew: $\alpha_3 = 2.47$	$q(x)_{0.05} = -2.57544$
Median = 0	Kurtosis: $\alpha_4 = 24.21$	$q(x)_{0.25} = -1.04507$
Mode = -0.1817	L-skew: $\tau_3 = 0.18$	$q(x)_{0.75} = 1.26292$
Max(height) = 0.2521	L-kurtosis: $\tau_4 = 0.22$	$q(x)_{0.95} = 4.278$

(b)

Figure 4: Two asymmetric κ_L - κ_R distributions pdfs and their cdfs. The parameters of skew (α_3) and kurtosis (α_4) are based on (A.2) in Appendix A. The parameters of L-skew (τ_3) and L-kurtosis (τ_4) are based on (2.32) and (2.33). The distribution in (b) is a local pdf in the range of $q(x) \in (-8.085, +\infty)$ where (2.7) yields $1 - \Pr\{q(x) \leq -8.085\} = 0.0003$.

$$\tau_3 = \frac{(2\gamma^2 - (\gamma + \kappa)^3 h_1 + (\gamma - \kappa)^3 h_2 - (\gamma - \kappa)^3 h_3 + 2\kappa^3 h_4 + (\gamma + \kappa)^3 h_5 - 2\kappa^3 h_6)}{2\gamma\lambda_2}, \quad (2.19)$$

$$\tau_4 = \left(2\gamma(1 + 5\gamma^2 + 5\kappa(1 - 3\kappa)) - (\gamma + \kappa)^2(1 + 5(\gamma + \kappa)^2)h_1\right. \\ \left.+ (1 + 5(\gamma - \kappa)^2)(\gamma - \kappa)^2(h_2 - h_3) + (\gamma + \kappa)^2(1 + 5(\gamma + \kappa)^2)h_5 / (12\gamma\lambda_2)\right) \quad (2.20)$$

with harmonic number functions $H[\cdot]$ of $h_1 = H[(1/2)(-1-\gamma-\kappa)]$, $h_2 = H[(1/2)(-1+\gamma-\kappa)]$, $h_3 = H[(1/2)(\gamma-\kappa)]$, $h_4 = H[(1/2)(-1-\kappa)]$, $h_5 = H[(-1/2)(\gamma+\kappa)]$, and $h_6 = H[-(1/2)\kappa]$. As such, given user specified values of τ_3 and τ_4 , (2.19) and (2.20) can be numerically solved to obtain the parameters for γ and κ .

Taking the limit of (2.17)–(2.20) as $\kappa \rightarrow 0$ gives the class associated with the log-logistic γ distributions in (2.4) as

$$\lambda_1 = \pi \csc(\gamma\pi) - \frac{1}{\gamma}, \quad (2.21)$$

$$\lambda_2 = \gamma\pi \csc(\gamma\pi), \quad (2.22)$$

$$\tau_3 = \gamma, \quad (2.23)$$

$$\tau_4 = \frac{1}{6}(1 + 5\gamma^2). \quad (2.24)$$

Analogously, taking the limit of (2.17)–(2.20) as $\gamma \rightarrow 0$ gives the system associated with the symmetric κ family of distributions in (2.5) as in [27]

$$\lambda_1 = 0, \quad (2.25)$$

$$\lambda_2 = \frac{1}{2}(2 + \kappa(-4p_1 + 4p_2 + \kappa(p_3 - p_4))), \quad (2.26)$$

$$\tau_3 = 0, \quad (2.27)$$

$$\tau_4 = \frac{1}{6} \left(1 + 5\kappa^2 + \frac{10\kappa(1 + 2\kappa + 2\kappa^2(h_6 - h_4))}{2\lambda_2} \right) \quad (2.28)$$

with polygamma functions $P[\cdot]$ of $p_1 = P[0, 1/2 - \kappa/2]$, $p_2 = P[0, 1 - \kappa/2]$, $p_3 = P[1, 1/2 - \kappa/2]$, and $p_4 = P[1, 1 - \kappa/2]$.

The L -moments for the class of asymmetric κ_L - κ_R distributions in (2.6) can be determined by separately evaluating and summing two integrals of the form in (2.16) as

$$\beta_i = \int_{-\infty}^0 q(x, \kappa_L) \{\Psi(x)\}^i \psi(x) dx + \int_0^{+\infty} q(x, \kappa_R) \{\Psi(x)\}^i \psi(x) dx. \quad (2.29)$$

For an asymmetric κ_L - κ_R distribution with a global pdf to have finite values of λ_1 , λ_2 , τ_3 , τ_4 based on (2.29) will require $\kappa_L < 1$ and $\kappa_R < 1$. As such, integrating (2.29) to obtain $\beta_0, \dots, \beta_3$ and subsequently substituting these terms into (2.10)–(2.13) yield as in [27]

$$\lambda_1 = \frac{1}{4}(2p_5 - 2p_6 - 2p_7 + 2p_8 - \kappa_L p_9 + \kappa_L p_{10} + \kappa_R p_{11} - \kappa_R p_{12}), \quad (2.30)$$

$$\lambda_2 = \frac{1}{4}(4 + \kappa_L(-4p_5 + 4p_6 + \kappa_L(p_9 - p_{10})) + 4 + \kappa_R(-4p_7 + 4p_8 + \kappa_R(p_{11} - p_{12}))), \quad (2.31)$$

$$\tau_3 = \frac{\{4\kappa_R - 4\kappa_L + \kappa_L^2(6p_5 - 6p_6 + \kappa_L(p_{10} - p_9)) + \kappa_R^2(-6p_7 + 6p_8 + \kappa_R(p_{11} - p_{12}))\}}{(4\lambda_2)}, \quad (2.32)$$

$$\begin{aligned}
\tau_4 = & \left\{ 2(2 + 5\kappa_{\mathcal{L}}(1 + 3\kappa_{\mathcal{L}}) + 5\kappa_{\mathcal{R}}(1 + 3\kappa_{\mathcal{R}})) - 4(\kappa_{\mathcal{L}} + 10\kappa_{\mathcal{L}}^3)p_5 - 4(\kappa_{\mathcal{R}} + 10\kappa_{\mathcal{R}}^3)p_7 \right. \\
& + 5\kappa_{\mathcal{L}}\left((4 + 40\kappa_{\mathcal{L}}^2)p_6 + \kappa_{\mathcal{L}}(1 + 5\kappa_{\mathcal{L}}^2)(p_9 - p_{10}) \right. \\
& \left. \left. + \kappa_{\mathcal{R}}\left((4 + 40\kappa_{\mathcal{R}}^2)p_8 + \kappa_{\mathcal{R}}(1 + 5\kappa_{\mathcal{R}}^2)(p_{11} - p_{12})\right)\right) \right\} / (24\lambda_2)
\end{aligned} \tag{2.33}$$

with polygamma functions $P[\cdot]$ of $p_5 = P[0, 1/2 - \kappa_{\mathcal{L}}/2]$, $p_6 = P[0, 1 - \kappa_{\mathcal{L}}/2]$, $p_7 = P[0, 1/2 - \kappa_{\mathcal{R}}/2]$, $p_8 = P[0, 1 - \kappa_{\mathcal{R}}/2]$, $p_9 = P[1, 1/2 - \kappa_{\mathcal{L}}/2]$, $p_{10} = P[1, 1 - \kappa_{\mathcal{L}}/2]$, $p_{11} = P[1, 1/2 - \kappa_{\mathcal{R}}/2]$, $p_{12} = P[1, 1 - \kappa_{\mathcal{R}}/2]$.

As with the γ - κ class of distributions, given specified values of τ_3 and τ_4 , (2.32) and (2.33) can be numerically solved to obtain the corresponding values of $\kappa_{\mathcal{L}}$ and $\kappa_{\mathcal{R}}$. Inspection of (2.32) and (2.33) reveals that interchanging values for the parameters of $\kappa_{\mathcal{L}}$ and $\kappa_{\mathcal{R}}$ reverses the direction of τ_3 and has no effect on τ_4 . As such, a graph of the region for feasible combinations of $|\tau_3|$ and τ_4 for (2.32) and (2.33) is provided in Figure 5. Feasible combinations of L -skew and L -kurtosis will lie in the region above the curve graphed in the $|\tau_3|$, τ_4 plane of Figure 5. Note that the curve in Figure 5 was graphed by setting $\kappa_{\mathcal{L}} = 0$ with $\kappa_{\mathcal{R}} \in [0, 1]$ in (2.32) and (2.33).

The conventional moment-based systems for the γ - κ and $\kappa_{\mathcal{L}}$ - $\kappa_{\mathcal{R}}$ classes of distributions are given in Appendices A and B, respectively. These systems were used to determine the values of skew and kurtosis associated with the distributions given in Figures 1–4. It is worthy to point out that the conventional moment-based systems have a disadvantage in terms of moment existence. That is, the integral in (A.1) of Appendix A reveals that for the r th moment to exist we must have in general (i) $\gamma + \kappa < 1/r$, (ii) $\kappa < 1/r$ and (iii) $1/r + \gamma < \kappa$. For example, if the mean, variance, skew, and kurtosis exist, then we must have $1/r = 0.25$. And, analogously for the $\kappa_{\mathcal{L}}$ - $\kappa_{\mathcal{R}}$ class of distributions in Appendix B, the parameters are bounded in the range of $0 \leq \kappa_{\mathcal{L}} < 0.25$ and $0 \leq \kappa_{\mathcal{R}} < 0.25$. The advantage that the L -moment system has in this context is attributed to Hosking's Theorem 1 [22] which states that if the mean (λ_1) exists, then all other L -moments will have finite expectations. We note that the conventional moment-based systems for the log-logistic γ and symmetric κ classes of distributions can be obtained by simplifying the system as described in Appendix A. In the next section we first introduce the topic of the L -correlation and subsequently develop the methodology for simulating γ - κ , γ , κ , and $\kappa_{\mathcal{L}}$ - $\kappa_{\mathcal{R}}$ distributions with specified L -correlations.

3. L -Correlations for the System of γ - κ , γ , κ , and $\kappa_{\mathcal{L}}$ - $\kappa_{\mathcal{R}}$ Distributions

The coefficient of L -correlation [29] is introduced by considering two random variables Y_j and Y_k with distribution functions $F(Y_j)$ and $F(Y_k)$, respectively. The second L -moments of Y_j and Y_k can alternatively be expressed as

$$\lambda_2(Y_j) = 2\text{Cov}(Y_j, F(Y_j)), \tag{3.1}$$

$$\lambda_2(Y_k) = 2\text{Cov}(Y_k, F(Y_k)). \tag{3.2}$$

The second L -comoments of Y_j toward Y_k and Y_k toward Y_j are

$$\lambda_2(Y_j, Y_k) = 2\text{Cov}(Y_j, F(Y_k)), \tag{3.3}$$

$$\lambda_2(Y_k, Y_j) = 2\text{Cov}(Y_k, F(Y_j)). \tag{3.4}$$

Figure 5: Graph of the region for feasible combinations of (absolute value) L -skew $|\tau_3|$ and L -kurtosis τ_4 for global κ_L - κ_R pdfs. Asymmetric κ_L - κ_R distributions will lie in the area above the curve graphed in the $|\tau_3|$ and τ_4 plane.

As such, the L -correlations of Y_j toward Y_k and Y_k toward Y_j are expressed as

$$\eta_{jk} = \frac{\lambda_2(Y_j, Y_k)}{\lambda_2(Y_j)}, \quad (3.5)$$

$$\eta_{kj} = \frac{\lambda_2(Y_k, Y_j)}{\lambda_2(Y_k)}. \quad (3.6)$$

The L -correlation in (3.5) (or (3.6)) is bounded such that $-1 \leq \eta_{jk} \leq 1$ where a value of $\eta_{jk} = 1$ ($\eta_{jk} = -1$) indicates a strictly increasing (decreasing) *monotone* relationship between the two variables. In general, we would also note that $\eta_{jk} \neq \eta_{kj}$.

In the context of the L -moment-based γ - κ , γ , κ , and κ_L - κ_R classes of distributions, suppose that it is desired to simulate a T -variate distribution based on quantile functions of the forms in (2.3)–(2.6) with a specified L -correlation matrix and where each distribution has its own specified values of τ_3 and τ_4 . Let $Z_1, \dots, Z_T$ denote standard normal variables where the distribution functions and bivariate density function associated with Z_j and Z_k are expressed as

$$\Phi(z_j) = \Pr\{Z_j \leq z_j\} = \int_{-\infty}^{z_j} (2\pi)^{-1/2} \exp\left\{-\frac{w_j^2}{2}\right\} dw_j, \quad (3.7)$$

$$\Phi(z_k) = \Pr\{Z_k \leq z_k\} = \int_{-\infty}^{z_k} (2\pi)^{-1/2} \exp\left\{-\frac{w_k^2}{2}\right\} dw_k, \quad (3.8)$$

$$f_{jk} = \left(2\pi(1 - \rho_{jk}^2)^{1/2}\right)^{-1} \exp\left\{-\left(2(1 - \rho_{jk}^2)\right)^{-1}(z_j^2 + z_k^2 - 2\rho_{jk}z_jz_k)\right\}. \quad (3.9)$$

Using (3.7), it follows that the j th distribution associated with (2.3)–(2.6) can be expressed as $q_j(g(\Phi(Z_j)))$, where $g(\Phi(Z_j)) = \ln(\Phi(Z_j)/(1 - \Phi(Z_j)))$ is standard logistic because $\Phi(Z_j) \sim U(0, 1)$. As such, using (3.5), the L -correlation of $q_j(g(\Phi(Z_j)))$ toward $q_k(g(\Phi(Z_k)))$ can

be evaluated using the solved value(s) of the parameter(s) (i.e., γ_j - κ_j ; γ_j ; κ_j ; $\kappa_{\mathcal{L}_j}$ - $\kappa_{\mathcal{R}_j}$) for $q_j(g(\Phi(Z_j)))$, a specified *intermediate correlation* (IC) ρ_{jk} in (3.9), and the following integral generally expressed as

$$\eta_{jk} = 2\sqrt{\pi} \int_{-\infty}^{+\infty} s_j \left(q_j \left(g(\Phi(z_j)), \gamma_j, \kappa_j, \kappa_{\mathcal{L}_j}, \kappa_{\mathcal{R}_j} \right) \right) \Phi(z_k) f_{jk} dz_j dz_k. \quad (3.10)$$

We would point out that the purpose of the IC (ρ_{jk}) in (3.9) and (3.10) is to adjust for the effect of the transformation $q_j(g(\Phi(Z_j)))$, which is induced by the parameters, such that $q_j(g(\Phi(Z_j)))$ has its specified L -correlation (η_{jk}) toward $q_k(g(\Phi(Z_k)))$. Further, to simplify the computation, the quantile function in (3.10) is standardized by a linear transformation such that it has a mean of zero and one-half the coefficient of mean difference equal to that of the unit-normal distribution as

$$s_j \left(q_j \left(g(\Phi(z_j)), \gamma_j, \kappa_j, \kappa_{\mathcal{L}_j}, \kappa_{\mathcal{R}_j} \right) \right) = \delta \times \left(q_j \left(g(\Phi(z_j)), \gamma_j, \kappa_j, \kappa_{\mathcal{L}_j}, \kappa_{\mathcal{R}_j} \right) - \lambda_1 \right), \quad (3.11)$$

where λ_1 is a mean from (2.17), (2.21), (2.25), or (2.30) and δ is a constant that scales λ_2 in (2.18), (2.22), (2.26), or (2.31) and in the denominator of (3.5) to $1/\sqrt{\pi}$ as

$$\begin{aligned} \delta = \delta_{\gamma_j, \kappa_j} &= \frac{2\gamma_j}{\left(\sqrt{\pi} \left(2\gamma_j - h_1(\gamma_j + \kappa_j)^2 + (\gamma_j - \kappa_j)^2 (h_2 - h_3) + h_5(\gamma_j + \kappa_j)^2 \right) \right)'}, \\ \delta = \delta_{\gamma_j} &= \frac{\sin(\gamma_j \pi)}{\gamma_j \pi^{3/2}}, \\ \delta = \delta_{\kappa_j} &= \frac{2}{\left(\sqrt{\pi} (2 + \kappa_j (-4p_1 + 4p_2 + \kappa_j (p_3 - p_4))) \right)'}, \\ \delta = \delta_{\kappa_{\mathcal{L}_j}, \kappa_{\mathcal{R}_j}} &= 4 / \left(\sqrt{\pi} \left(4 - 4p_5 \kappa_{\mathcal{L}_j} + 4p_6 \kappa_{\mathcal{L}_j} + p_9 \kappa_{\mathcal{L}_j}^2 - p_{10} \kappa_{\mathcal{L}_j}^2 - 4p_7 \kappa_{\mathcal{R}_j} + 4p_8 \kappa_{\mathcal{R}_j} \right. \right. \\ &\quad \left. \left. + p_{11} \kappa_{\mathcal{R}_j}^2 - p_{12} \kappa_{\mathcal{R}_j}^2 \right) \right). \end{aligned} \quad (3.12)$$

Analogously, the L -correlation of $q_k(g(\Phi(Z_k)))$ toward $q_j(g(\Phi(Z_j)))$ is expressed as

$$\eta_{kj} = 2\sqrt{\pi} \int_{-\infty}^{+\infty} s_k \left(q_k \left(g(\Phi(z_k)), \gamma_k, \kappa_k, \kappa_{\mathcal{L}_k}, \kappa_{\mathcal{R}_k} \right) \right) \Phi(z_j) f_{jk} dz_k dz_j. \quad (3.13)$$

Note also for the special case that if $q_j(g(\Phi(Z_j)))$ in (3.10) and $q_k(g(\Phi(Z_k)))$ in (3.13) have the same parameters, that is, $\gamma_j = \gamma_k$; $\kappa_j = \kappa_k$; $\kappa_{\mathcal{L}_j} = \kappa_{\mathcal{L}_k}$; $\kappa_{\mathcal{R}_j} = \kappa_{\mathcal{R}_k}$, then $\eta_{jk} = \eta_{kj}$. Source code written in Mathematica [30] that implements the computation of an IC (ρ_{jk}) based on (3.10) is provided in Algorithm 1. The details for simulating γ - κ , γ , κ , and $\kappa_{\mathcal{L}}$ - $\kappa_{\mathcal{R}}$ distributions with specified values of L -skew, L -kurtosis, and L -correlations are described in the next section.

```

(*Intermediate Correlation for Distributions 1 and 2. See Table 3.*)
 $\rho_{jk} = 0.380206;$ 

Needs["MultivariateStatistics'"]
 $f_{jk} = \text{PDF}[\text{MultinormalDistribution}[\{0, 0\}, \{\{1, \rho_{jk}\}, \{\rho_{jk}, 1\}\}], \{Z_j, Z_k\}];$ 
 $F_j = \text{CDF}[\text{NormalDistribution}[0, 1], Z_j];$ 
 $F_k = \text{CDF}[\text{NormalDistribution}[0, 1], Z_k];$ 

(*Parameters for Figure 1(a)*)
 $\gamma_j = 0.15;$ 
 $\kappa_j = 0.075;$ 

 $X_j = \text{Log}[F_j / (1 - F_j)];$ 

(*Quantile function from (2.3)*)
 $q_{\gamma_j, \kappa_j} = \gamma_j^{-1} * (\text{Exp}[\gamma_j X_j] - 1) * \text{Exp}[\kappa_j * \text{Abs}[X_j]];$ 

(*Standardizing constants  $\lambda_1$  from (2.17) and  $\delta_{\gamma_j, \kappa_j}$  from (3.12)*)
 $Sq_{\gamma_j, \kappa_j} = \delta_{\gamma_j, \kappa_j} * (q_{\gamma_j, \kappa_j} - \lambda_1);$ 

(*Compute the specified  $L$ -correlation in Table 1 for Distributions 1 and 2*)
 $\eta_{jk} = 2\sqrt{\pi} * \text{NIntegrate}[Sq_{\gamma_j, \kappa_j} * F_k * f_{jk}, \{Z_j, -10, 10\}, \{Z_k, -10, 10\}$ 
Method  $\rightarrow$  MultiDimensional]

0.40

```

Algorithm 1: Mathematica source code for computing intermediate correlations for specified L -correlations. The example is for distribution $j = 1$ towards distribution $k = 2$ (η_{12}) in Figures 1(a) and 2(a). See Tables 1 and 3.

4. The Procedure for Simulation and Monte Carlo Study

To implement the method for simulating γ - κ , γ , κ , and κ_L - κ_R distributions with specified L -moments and L -correlations we suggest the following six steps.

- (1) Specify the L -moments for T transformations of the forms in (2.3)–(2.6) that is, $q_1(g(\Phi(Z_1))), \dots, q_T(g(\Phi(Z_T)))$ and obtain the solutions for the parameters of γ - κ , γ , κ , or κ_L - κ_R distributions by solving (2.19), (2.20); (2.23), (2.24); (2.27), (2.28); (2.32), (2.33) using the specified values of L -skew (τ_3) and L -kurtosis (τ_4) for each distribution. Specify a $T \times T$ matrix of L -correlations (η_{jk}) for $q_j(g(\Phi(Z_j)))$ toward $q_k(g(\Phi(Z_k)))$, where $j < k \in \{1, 2, \dots, T\}$.
- (2) Compute the (Pearson) intermediate correlations (ICs) ρ_{jk} by substituting the solutions of the parameters from Step (1) into (3.10) and then numerically integrate to solve for ρ_{jk} (see Algorithm 1 for an example). Repeat this step separately for all $T(T-1)/2$ pairwise combinations of correlations.
- (3) Assemble the ICs into a $T \times T$ matrix and decompose this matrix using a Cholesky factorization. Note that this step requires the IC matrix to be positive definite.

```

(*Intermediate Correlation for Distributions 1 and 4. See Table 2.*)
ρjk = 0.835142;

Needs["MultivariateStatistics'"]
fjk = PDF[MultinormalDistribution[{0,0}, {{1, ρjk}, {ρjk, 1}}], {Zj, Zk}]
Fj = CDF[NormalDistribution[0,1], Zj];
Fk = CDF[NormalDistribution[0,1], Zk];

(*Parameters for Figure 1(a) and Figure 4(a) in Figure 4*)
γj = 0.15;
κj = 0.075;
κℓk = 0.0475;
κℓk = 0.1358;

Xj = Log[Fj/(1-Fj)];
Xk = Log[Fk/(1-Fk)];

(*Quantile functions from (2.3) and (2.6)*)
qγj, κj = γj-1 * (Exp[γjXj] - 1) * Exp[κj * Abs[Xj]];
qℓk = Xk * Exp[κℓk * Abs[Xk]];
qℓk = Xk * Exp[κℓk * Abs[Xk]];

(*Standardizing constants λ1j, λ1k ((2.17), (2.30)) and α2j, α2k ((A.2), (B.2)) from
Appendices A and B*)
Sqγj, κj = (qγj, κj - λ1j)/α2j;
Sqℓk = (qℓk - λ1k)/α2k;
Sqℓk = (qℓk - λ1k)/α2k;

(*Compute the specified Pearson correlation in Table 1 for Distributions 1 and 4.*)
ρjk* = NIntegrate[Sqγj, κj * Piecewise[{{Sqℓk, Xk ≤ 0}, {Sqℓk, Xk > 0}}] * fjk, {Zj, -10, 10}, {Zk,
-10, 10}, Method → MultiDimensional]

0.80

```

Algorithm 2: Mathematica source code for computing intermediate correlations for specified conventional Pearson correlations. The example is for distributions $j = 1$ and $k = 4$ (ρ_{14}^*) in Figures 1(a) and 4(a). See Tables 1 and 2.

- (4) Use the results of the Cholesky factorization from Step (3) to generate T standard normal variables $(Z_1, \dots, Z_T)$ correlated at the intermediate levels as follows:

$$\begin{aligned}
 Z_1 &= a_{11}V_1, \\
 Z_2 &= a_{12}V_1 + a_{22}V_2, \\
 &\vdots \\
 Z_j &= a_{1j}V_1 + a_{2j}V_2 + \dots + a_{ij}V_i + \dots + a_{jj}V_j, \\
 &\vdots \\
 Z_T &= a_{1T}V_1 + a_{2T}V_2 + \dots + a_{iT}V_i + \dots + a_{jT}V_j + \dots + a_{TT}V_T,
 \end{aligned} \tag{4.1}$$

Table 1: Specified correlation matrix for the distributions in Figures 1(a), 2(a), 3(b), and 4(a).

	1	2	3	4
1	1			
2	0.40	1		
3	0.60	0.50	1	
4	0.80	0.70	0.60	1

Table 2: Intermediate correlations for the conventional moment procedure.

	1	2	3	4
1	1			
2	0.446226	1		
3	0.671890	0.566802	1	
4	0.835142	0.740484	0.653826	1

where $V_1, \dots, V_T$ are independent standard normal random variables and where a_{ij} represents the element in the i th row and the j th column of the matrix associated with the Cholesky factorization performed in Step (3).

- (5) Substitute $Z_1, \dots, Z_T$ from Step (4) into the following Taylor series-based expansion for the standard normal cdf [30]:

$$\Phi(Z_j) = \left(\frac{1}{2}\right) + \phi(Z_j) \left\{ Z_j + \frac{Z_j^3}{3} + \frac{Z_j^5}{(3 \cdot 5)} + \frac{Z_j^7}{(3 \cdot 5 \cdot 7)} + \dots \right\}, \quad (4.2)$$

where $\phi(Z_j)$ denotes the standard normal pdf and where the absolute error associated with (4.2) is less than 8×10^{-16} .

- (6) Substitute the zero-one uniform deviates, $\Phi(Z_j)$, generated from Step (5) into the T equations of the form of $q_j(g(\Phi(Z_j)))$, where $g(\Phi(Z_j)) = \ln(\Phi(Z_j)/(1 - \Phi(Z_j)))$ is standard logistic to generate the γ - κ , γ , κ , and κ_L - κ_R distributions with the specified L -moments and L -correlations.

To demonstrate the steps above and evaluate the proposed method, a comparison between the proposed L -moment and conventional product-moment-based procedures is subsequently described. Specifically, the heavy-tailed distributions in Figures 1(a), 2(a), 3(b), and 4(a) are used as a basis for a comparison using the specified correlation matrix in Table 1. Tables 2 and 3 give the solved IC matrices for the conventional moment and L -moment-based methods, respectively. See Algorithm 2 for an example of computing ICs for the conventional method. Tables 4 and 5 give the results of the Cholesky decompositions on the IC matrices, which are then used to create $Z_1, \dots, Z_4$ with the specified ICs by making use of the formulae given in (4.1) of Step (4) with $T = 4$. The values of $Z_1, \dots, Z_4$ are subsequently transformed to $\Phi(Z_1), \dots, \Phi(Z_4)$ using (4.2) and then substituted into equations of the forms in (2.3)–(2.6) to produce $q_1(g(\Phi(Z_1))), \dots, q_4(g(\Phi(Z_4)))$ for both methods.

In terms of the simulation, a Fortran algorithm was written for both methods to generate 25,000 independent sample estimates for the specified parameters of (i) conventional skew (α_3), kurtosis (α_4), and Pearson correlation (ρ_{jk}^*) and (ii) L -skew (τ_3), L -kurtosis (τ_4), and L -correlation (η_{jk}). All estimates were based on sample sizes of $n = 25$ and $n = 1000$.

Table 3: Intermediate correlations for the L -moment procedure.

	1	2	3	4
1	1			
2	0.380206	1		
3	0.576862	0.486201	1	
4	0.782049	0.686737	0.568733	1

Table 4: Cholesky decompositions for the conventional moment procedure.

$a_{11} = 1$	$a_{12} = 0.446226$	$a_{13} = 0.671890$	$a_{14} = 0.835142$
0	$a_{22} = 0.894920$	$a_{23} = 0.298336$	$a_{24} = 0.411011$
0	0	$a_{33} = 0.677908$	$a_{34} = -0.044131$
0	0	0	$a_{44} = 0.362849$

The formulae used for computing estimates of $\alpha_{3,4}$ were based on Fisher's k -statistics that is, the formulae currently used by most commercial software packages such as SAS, SPSS, and Minitab, for computing indices of skew and kurtosis (where $\alpha_{3,4} = 0$ for the standard normal distribution). The formulae used for computing estimates of $\tau_{3,4}$ were Headrick's Equations (2.4) and (2.6) [25]. The estimate for ρ_{jk}^* was based on the usual formula for the Pearson product moment of correlation statistic, and the estimate for η_{jk} was computed based on (3.5) using the empirical forms of the cdfs in (3.1) and (3.3). The estimates for ρ_{jk}^* and η_{jk} were both transformed using Fisher's z' transformation. Bias-corrected accelerated bootstrapped average (mean) estimates, confidence intervals (CIs), and standard errors were subsequently obtained for the estimates associated with the parameters $(\alpha_{3,4}, \tau_{3,4}, z'_{\rho_{jk}^*}, z'_{\eta_{jk}})$ using 10,000 resamples via the commercial software package Spotfire S+ [31]. The bootstrap results for the estimates of the means and CIs associated with $z'_{\rho_{jk}^*}$ and $z'_{\eta_{jk}}$ were transformed back to their original metrics (i.e., estimates for ρ_{jk}^* and η_{jk}). Further, if a parameter (P) was outside its associated bootstrap CI, then an index of relative bias (RB) was computed for the estimate (E) as $RB = ((E - P) / P) \times 100$. Note that the small amount of bias associated with any bootstrap CI containing a parameter was considered negligible and thus not reported. The results of the simulation are reported in Tables 6–13 and are discussed in the next section.

5. Discussion and Conclusion

One of the primary advantages that L -moments have over conventional moment-based estimators is that they can be far less biased when sampling is from distributions with more severe departures from normality (e.g., [21, 25]). Inspection of the simulation results in Tables 6 and 7 of this study clearly indicates that this is also the case for the system of γ - κ , γ , κ , and κ_L - κ_R distributions. Specifically, the superiority that estimates of L -moment ratios (τ_3, τ_4) have over their corresponding conventional moment-based counterparts (α_3, α_4) is obvious. For example, with samples of size $n = 25$ the estimates of skew and kurtosis for Distribution 1 (Tables 6 and 8) were, on average, only 26.80% and 2.38% of their associated population parameters whereas the estimates of L -skew and L -kurtosis were 87.80% and 94.81% of their respective parameters. It is also evident from Tables 6 and 8 that L -skew and L -kurtosis are more efficient estimators as their relative standard errors $RSE = (\text{standard error} / \text{estimate}) \times 100$ are substantially smaller than the conventional estimators of skew and kurtosis. For example, in terms of Distribution 1, inspection of Tables 7 and 9 indicates RSE measures of

Table 5: Cholesky decompositions for the L -moment procedure.

$a_{11} = 1$	$a_{12} = 0.380206$	$a_{13} = 0.576862$	$a_{14} = 0.782049$
0	$a_{22} = 0.924902$	$a_{23} = 0.288544$	$a_{24} = 0.421015$
0	0	$a_{33} = 0.764181$	$a_{34} = -0.005081$
0	0	0	$a_{44} = 0.459478$

Table 6: Skew (α_3) and Kurtosis (α_4) results for the conventional moment procedure. Sample size of $n = 25$.

Dist.	Parameter	Estimate	95% bootstrap C.I.	Standard error	Relative bias %
1	$\alpha_3 = 3.488$	0.9348	0.9229, 0.9495	0.00676	-73.2
	$\alpha_4 = 120.0$	2.861	2.8107, 2.9087	0.02499	-97.6
2	$\alpha_3 = 2.485$	1.074	1.0641, 1.0853	0.00541	-56.8
	$\alpha_4 = 26.57$	2.240	2.1963, 2.2856	0.02272	-91.6
3	$\alpha_3 = 0.0$	0.0125	-0.0052, 0.0294	0.00875	—
	$\alpha_4 = 39.83$	3.365	3.3157, 3.4122	0.02464	-91.6
4	$\alpha_3 = 1.83$	0.5777	0.5633, 0.5907	0.00693	-68.4
	$\alpha_4 = 23.48$	2.5360	2.4915, 2.5778	0.02190	-89.2

$\text{RSE}(\hat{\alpha}_3) = 0.436$ and $\text{RSE}(\hat{\alpha}_4) = 1.12$ compared with $\text{RSE}(\hat{\tau}_3) = 0.109$ and $\text{RSE}(\hat{\tau}_4) = 0.048$. This demonstrates that L -skew and L -kurtosis have more precision because they have less variance around their estimates.

The results associated with the conventional Pearson and L -correlations are presented in Tables 10–13. Overall inspection of these tables indicates that the L -correlation is substantially superior to the Pearson correlation in terms of relative bias. For example, in terms of a moderate correlation (Table 10, $n = 25$, $\rho_{12}^* = 0.40$) the relative bias for Distributions 1 and 2 was 9.73% for the Pearson correlation compared to only 1.85% for the L -correlation (Table 12, $n = 25$, $\eta_{12} = 0.40$). Further, for large sample sizes (Table 13, $n = 1000$), the L -correlation bootstrap CIs contained all of the population parameters whereas the Pearson correlation CIs contained none of the parameters (Table 11).

In summary, the proposed L -moment-based system of γ - κ , γ , κ , and κ_L - κ_R distributions is an attractive alternative to the traditional conventional-moment-based system. In particular, the L -moment-based system has distinct advantages when heavy-tailed distributions are of concern. Finally, we would note that Mathematica Version 8.0 [30] source code is available from the authors for implementing the L -moment-based method.

Appendices

A. System of Conventional Moment-Based Equations for γ - κ Distributions

The moments ($\mu_{r=1,\dots,4}$) associated with the class of γ - κ distributions in (2.3) can be determined from

$$\mu_r = \int_{-\infty}^{+\infty} q(x, \gamma, \kappa)^r f_X(x) dx. \quad (\text{A.1})$$

Table 7: Skew (α_3) and Kurtosis (α_4) results for the conventional moment procedure. Sample size of $n = 1000$.

Dist.	Parameter	Estimate	95% bootstrap CI	Standard error	Relative bias %
1	$\alpha_3 = 3.488$	2.562	2.5383, 2.5823	0.01117	-26.5
	$\alpha_4 = 120.0$	22.15	21.6873, 22.6698	0.24850	-81.5
2	$\alpha_3 = 2.485$	2.180	2.1668, 2.1944	0.00697	-12.3
	$\alpha_4 = 26.57$	13.36	13.0936, 13.6467	0.14100	-49.7
3	$\alpha_3 = 0.0$	-0.0051	-0.0265, 0.0163	0.01100	—
	$\alpha_4 = 39.83$	18.57	18.2203, 18.9412	0.18330	-53.4
4	$\alpha_3 = 1.83$	1.54	1.5246, 1.5539	0.00743	-15.8
	$\alpha_4 = 23.48$	12.91	12.6537, 13.1903	0.13610	-45.0

Table 8: L -skew (τ_3) and L -kurtosis (τ_4) results. Sample size of $n = 25$.

Dist.	Parameter	Estimate	95% bootstrap CI	Standard error	Relative bias %
1	$\tau_3 = 0.1664$	0.1461	0.1443, 0.1480	0.00095	-12.2
	$\tau_4 = 0.2506$	0.2376	0.2362, 0.2389	0.00069	-5.19
2	$\tau_3 = 0.20$	0.1833	0.1819, 0.1849	0.00079	-8.35
	$\tau_4 = 0.20$	0.1911	0.1898, 0.1923	0.00064	-3.35
3	$\tau_3 = 0.0$	0.0014	-0.0008, 0.0037	0.00115	—
	$\tau_4 = 0.2958$	0.2809	0.2795, 0.2823	0.00070	-5.04
4	$\tau_3 = 0.10$	0.0879	0.0859, 0.0897	0.00096	-12.1
	$\tau_4 = 0.25$	0.2400	0.2388, 0.2414	0.00065	-4.00

Table 9: L -skew (τ_3) and L -kurtosis (τ_4) results. Sample size of $n = 1000$.

Dist.	Parameter	Estimate	95% bootstrap CI	Standard error	Relative bias %
1	$\tau_3 = 0.1664$	0.1657	0.1653, 0.1660	0.00018	-0.421
	$\tau_4 = 0.2506$	0.2502	0.2499, 0.2504	0.00012	-0.160
2	$\tau_3 = 0.20$	0.1995	0.1992, 0.1997	0.00013	-0.250
	$\tau_4 = 0.20$	0.1997	0.1995, 0.1999	0.00011	-0.150
3	$\tau_3 = 0.0$	-0.0001	-0.0005, 0.0003	0.00022	—
	$\tau_4 = 0.2958$	0.2954	0.2951, 0.2956	0.00012	-0.135
4	$\tau_3 = 0.10$	0.0995	0.0992, 0.0999	0.00017	-0.500
	$\tau_4 = 0.25$	0.2497	0.2495, 0.2499	0.00011	-0.120

Table 10: Correlation results for the conventional moment procedure. Sample size of $n = 25$.

Parameter	Estimate	95% bootstrap CI	Standard error	Relative bias %
$\rho_{12}^* = 0.40$	0.4389	0.4365, 0.4413	0.00153	9.73
$\rho_{13}^* = 0.60$	0.6522	0.6504, 0.6540	0.00153	8.70
$\rho_{14}^* = 0.80$	0.8296	0.8287, 0.8308	0.00165	3.70
$\rho_{23}^* = 0.50$	0.5441	0.5420, 0.5460	0.00142	8.82
$\rho_{24}^* = 0.70$	0.7306	0.7293, 0.7321	0.00157	4.37
$\rho_{34}^* = 0.60$	0.6417	0.6400, 0.6435	0.00154	6.95

Table 11: Correlation results for the conventional moment procedure. Sample size of $n = 1000$.

Parameter	Estimate	95% bootstrap CI	Standard error	Relative bias %
$\rho_{12}^* = 0.40$	0.4040	0.4035, 0.4044	0.00029	1.00
$\rho_{13}^* = 0.60$	0.6067	0.6063, 0.6072	0.00036	1.12
$\rho_{14}^* = 0.80$	0.8041	0.8038, 0.8044	0.00042	0.51
$\rho_{23}^* = 0.50$	0.5037	0.5032, 0.5040	0.00029	0.74
$\rho_{24}^* = 0.70$	0.7027	0.7024, 0.7030	0.00033	0.39
$\rho_{34}^* = 0.60$	0.6041	0.6036, 0.6044	0.00032	0.68

Table 12: Correlation results for the L -moment procedure. Sample size of $n = 25$.

Parameter	Estimate	95% bootstrap CI	Standard error	Relative bias %
$\eta_{12} = 0.40$	0.4074	0.4049, 0.4099	0.00154	1.85
$\eta_{13} = 0.60$	0.6068	0.6049, 0.6087	0.00153	1.13
$\eta_{14} = 0.80$	0.8058	0.8046, 0.8067	0.00152	0.73
$\eta_{23} = 0.50$	0.5078	0.5056, 0.5100	0.00151	1.56
$\eta_{24} = 0.70$	0.7076	0.7061, 0.7091	0.00154	1.09
$\eta_{34} = 0.60$	0.6063	0.6045, 0.6083	0.00155	1.05

In general, to obtain defined values of μ_r based on (A.1) we must have (i) $\gamma + \kappa < 1/r$, (ii) $\kappa < 1/r$, and (iii) $1/r + \gamma < \kappa$.

The mean, variance, skew, and kurtosis are defined in general as in [17]

$$\begin{aligned}
\alpha_1 &= \mu_1, \\
\alpha_2^2 &= \mu_2 - \mu_1^2, \\
\alpha_3 &= \frac{(\mu_3 - 3\mu_2\mu_1 + 2\mu_1^3)}{\alpha_2^3}, \\
\alpha_4 &= \frac{(\mu_4 - 4\mu_3\mu_1 - 3\mu_2^2 + 12\mu_2\mu_1^2 - 6\mu_1^4)}{\alpha_2^4}.
\end{aligned} \tag{A.2}$$

The moments associated with the location and scale parameters in (A.2) are

$$\alpha_1 = \mu_1 = \frac{((- \gamma - \kappa)h_{11} + (\gamma - \kappa)h_{12} + (- \gamma + \kappa)h_{13} + 2\kappa h_{14} + (\gamma + \kappa)h_{15} - 2\kappa h_{16})}{(2\gamma)} \tag{A.3}$$

with harmonic number functions $H[\cdot]$ of $h_{11} = H[(1/2)(-1-\gamma-\kappa)]$, $h_{12} = H[(1/2)(-1+\gamma-\kappa)]$, $h_{13} = H[(1/2)(\gamma - \kappa)]$, $h_{14} = H[(1/2)(-1 - \kappa)]$, $h_{15} = H[(-1/2)(\gamma + \kappa)]$, $h_{16} = H[(-1/2)\kappa]$, and

$$\begin{aligned}
\mu_2 &= (- (\gamma - 2\kappa)h_{21} - 2\kappa h_{22} - (\gamma + \kappa)h_{23} + (\gamma + \kappa)h_{24} + (\gamma + 2\kappa)h_{25} \\
&\quad - (\gamma + 2\kappa)h_{26} + (\gamma - 2\kappa)h_{27} + (\gamma - \kappa)h_{28} + (- \gamma + \kappa)h_{29} + 2\kappa h_{210} / (2\gamma^2))
\end{aligned} \tag{A.4}$$

Table 13: Correlation results for the L -moment procedure. Sample size of $n = 1000$.

Parameter	Estimate	95% bootstrap CI	Standard error	Relative bias %
$\eta_{12} = 0.40$	0.4000	0.3996, 0.4004	0.00023	—
$\eta_{13} = 0.60$	0.6002	0.5999, 0.6005	0.00023	—
$\eta_{14} = 0.80$	0.8001	0.7999, 0.8002	0.00023	—
$\eta_{23} = 0.50$	0.4997	0.4994, 0.5001	0.00023	—
$\eta_{24} = 0.70$	0.7002	0.6999, 0.7004	0.00023	—
$\eta_{34} = 0.60$	0.5999	0.5996, 0.6002	0.00023	—

with harmonic number functions $H[\cdot]$ of $h_{21} = H[(1/2)(-1 + \gamma - 2\kappa)]$, $h_{22} = H[-1/2 - \kappa]$, $h_{23} = H[-1/2 - \gamma - \kappa]$, $h_{24} = H[-\gamma - \kappa]$, $h_{25} = H[-1/2 - \gamma/2 - \kappa]$, $h_{26} = H[-\gamma/2 - \kappa]$, $h_{27} = H[\gamma/2 - \kappa]$, $h_{28} = H[-1/2 + \gamma - \kappa]$, $h_{29} = H[\gamma - \kappa]$, and $h_{210} = H[-\kappa]$.

The moments associated with the shape parameters of skew and kurtosis in (A.2) are

$$\begin{aligned} \mu_3 = & 3(-(\gamma + \kappa)h_{31} - (\gamma + 3\kappa)h_{32} + (\gamma - 3\kappa)h_{33} - (\gamma - 3\kappa)h_{34} + (\gamma - \kappa)h_{35} + 2\kappa h_{36} \\ & + (2\gamma + 3\kappa)h_{37} + (-2\gamma - 3\kappa)h_{38} + (\gamma + 3\kappa)h_{39} + (-2\gamma + 3\kappa)h_{310} \\ & + (2\gamma - 3\kappa)h_{311} + (-\gamma + \kappa)h_{312} - 2\kappa h_{313} + (\gamma + \kappa)h_{314}) / (2\gamma^3) \end{aligned} \quad (\text{A.5})$$

with harmonic number functions $H[\cdot]$ of $h_{31} = H[(1/2)(-1 - 3\gamma - 3\kappa)]$, $h_{32} = H[(1/2)(-1 - \gamma - 3\kappa)]$, $h_{33} = H[(1/2)(-1 + \gamma - 3\kappa)]$, $h_{34} = H[(1/2)(\gamma - 3\kappa)]$, $h_{35} = H[(1/2)(-1 + 3\gamma - 3\kappa)]$, $h_{36} = H[-1/2 - 3\kappa/2]$, $h_{37} = H[-1/2 - \gamma - 3\kappa/2]$, $h_{38} = H[-\gamma - 3\kappa/2]$, $h_{39} = H[-\gamma/2 - 3\kappa/2]$, $h_{310} = H[-1/2 + \gamma - 3\kappa/2]$, $h_{311} = H[\gamma - 3\kappa/2]$, $h_{312} = H[3(\gamma - \kappa)/2]$, $h_{313} = H[-3\kappa/2]$, $h_{314} = H[-3(\gamma + \kappa)/2]$, and

$$\begin{aligned} \mu_4 = & 2(-(\gamma - 4\kappa) - 2\kappa h_{41} - (\gamma + \kappa)h_{42} + (3\gamma + 4\kappa)h_{43} + (-3\gamma - 4\kappa)h_{44} \\ & - 3(\gamma + 2\kappa)h_{45} + 3(\gamma + 2\kappa)h_{46} + (\gamma + 4\kappa)h_{47} - (\gamma + 4\kappa)h_{48} \\ & + (\gamma - 4\kappa)h_{49} + 3(\gamma - 2\kappa)h_{410} + (-3\gamma - 2\kappa)h_{411} + (-3\gamma + 4\kappa)h_{412} \\ & + (3\gamma - 4\kappa)h_{413} + (\gamma - \kappa)h_{414} + (-\gamma + \kappa)h_{415} + 2\kappa h_{416} \\ & + (\gamma + \kappa)h_{417}) / (\gamma^4) \end{aligned} \quad (\text{A.6})$$

with harmonic number functions $H[\cdot]$ of $h_{41} = H[-1/2 - 2\kappa]$, $h_{42} = H[-1/2 - 2\gamma - 2\kappa]$, $h_{43} = H[-1/2 - 3\gamma/2 - 2\kappa]$, $h_{44} = H[-3\gamma/2 - 2\kappa]$, $h_{45} = H[-1/2 - \gamma - 2\kappa]$, $h_{46} = H[-\gamma - 2\kappa]$, $h_{47} = H[-1/2 - \gamma/2 - 2\kappa]$, $h_{48} = H[-\gamma/2 - 2\kappa]$, $h_{49} = H[\gamma/2 - 2\kappa]$, $h_{410} = H[-1/2 + \gamma - 2\kappa]$, $h_{411} = H[\gamma - 2\kappa]$, $h_{412} = H[-1/2 + 3\gamma/2 - 2\kappa]$, $h_{413} = H[3\gamma/2 - 2\kappa]$, $h_{414} = H[-1/2 + 2\gamma - 2\kappa]$, $h_{415} = H[2\gamma - 2\kappa]$, $h_{416} = H[-2\kappa]$, and $h_{417} = H[-2\gamma - 2\kappa]$.

The moments for the asymmetric γ class of distributions in (2.4) can be obtained by taking the limit of (A.3)–(A.6) as $\kappa \rightarrow 0$. Analogously, the moments for the symmetric κ class of distributions in (2.5) can be obtained by taking the limit of (A.3)–(A.6) as $\gamma \rightarrow 0$.

B. System of Conventional Moment-Based Equations for $\kappa_{\mathcal{L}}\text{-}\kappa_{\mathcal{R}}$ Distributions

The moments ($\mu_{r=1,\dots,4}$) associated with the asymmetric $\kappa_{\mathcal{L}}\text{-}\kappa_{\mathcal{R}}$ class of distributions in (2.6) can be determined from

$$\mu_r = \int_{-\infty}^0 q(x, \kappa_{\mathcal{L}})^r f_X(x) dx + \int_0^{+\infty} q(x, \kappa_{\mathcal{R}})^r f_X(x) dx. \quad (\text{B.1})$$

Using the expressions in (A.2), the moments for the location and scale parameters are

$$\alpha_1 = \mu_1 = \frac{1}{4}(2p_{11} - 2p_{12} - 2p_{13} + 2p_{14} - \kappa_{\mathcal{L}}p_{15} + \kappa_{\mathcal{L}}p_{16} + \kappa_{\mathcal{R}}p_{17} - \kappa_{\mathcal{R}}p_{18}), \quad (\text{B.2})$$

$$\begin{aligned} \mu_2 = \frac{1}{2} & \left(\kappa_{\mathcal{L}} \text{HZ} \left[3, \frac{1}{2} - \kappa_{\mathcal{L}} \right] - \kappa_{\mathcal{L}} \text{HZ} [3, 1 - \kappa_{\mathcal{L}}] + p_{15} + p_{16} \right) \\ & + \frac{1}{2} \left(\kappa_{\mathcal{R}} \text{HZ} \left[3, \frac{1}{2} - \kappa_{\mathcal{R}} \right] - \kappa_{\mathcal{R}} \text{HZ} [3, 1 - \kappa_{\mathcal{R}}] + p_{17} - p_{19} \right), \end{aligned} \quad (\text{B.3})$$

where the polygamma functions $P[\cdot]$ are: $p_{11} = P[0, 1/2 - \kappa_{\mathcal{L}}/2]$, $p_{12} = P[0, 1 - \kappa_{\mathcal{L}}/2]$, $p_{13} = P[0, 1/2 - \kappa_{\mathcal{R}}/2]$, $p_{14} = P[0, 1 - \kappa_{\mathcal{R}}/2]$, $p_{15} = P[1, 1/2 - \kappa_{\mathcal{L}}/2]$, $p_{16} = P[1, 1 - \kappa_{\mathcal{L}}/2]$, $p_{17} = P[1, 1/2 - \kappa_{\mathcal{R}}/2]$, $p_{18} = P[1, 1 - \kappa_{\mathcal{R}}/2]$, and $p_{19} = P[1, 1 - \kappa_{\mathcal{R}}]$. The notation $\text{HZ}[\cdot]$ in (B.3) is the Hurwitz zeta function. The moments related to skew and kurtosis are as follows:

$$\begin{aligned} \mu_3 = \frac{3}{16} & \left(2p_{31} - 2p_{32} - \kappa_{\mathcal{L}}p_{33} + \kappa_{\mathcal{L}}p_{34} \right. \\ & \left. + \frac{3}{8} \left(p_{35} + 2Z \left[3, \frac{1}{2}(1 - 3\kappa_{\mathcal{R}}) \right] + 3\kappa_{\mathcal{R}}Z \left[4, \frac{1}{2}(1 - 3\kappa_{\mathcal{R}}) \right] - 3\kappa_{\mathcal{R}}Z \left[4, 1 - \frac{3\kappa_{\mathcal{R}}}{2} \right] \right) \right), \end{aligned} \quad (\text{B.4})$$

$$\mu_4 = \frac{1}{8}(p_{41} - p_{42} - \kappa_{\mathcal{L}}p_{43} + \kappa_{\mathcal{L}}p_{44}) + \frac{1}{8}(2p_{45} - 2p_{46} - \kappa_{\mathcal{R}}p_{47} + \kappa_{\mathcal{R}}p_{48}), \quad (\text{B.5})$$

where the polygamma functions $P[\cdot]$ are $p_{31} = P[2, 1/2 - 3\kappa_{\mathcal{L}}/2]$, $p_{32} = P[2, 1 - 3\kappa_{\mathcal{L}}/2]$, $p_{33} = P[3, 1/2 - 3\kappa_{\mathcal{L}}/2]$, $p_{34} = P[3, 1 - 3\kappa_{\mathcal{L}}/2]$, $p_{35} = P[2, 1 - 3\kappa_{\mathcal{R}}/2]$, $p_{41} = P[3, 1/2 - 2\kappa_{\mathcal{L}}]$, $p_{42} = P[3, 1 - 2\kappa_{\mathcal{L}}]$, $p_{43} = P[4, 1/2 - 2\kappa_{\mathcal{L}}]$, $p_{44} = P[4, 1 - 2\kappa_{\mathcal{L}}]$, $p_{45} = P[3, 1/2 - 2\kappa_{\mathcal{R}}]$, $p_{46} = P[3, 1 - 2\kappa_{\mathcal{R}}]$, $p_{47} = P[4, 1/2 - 2\kappa_{\mathcal{R}}]$, and $p_{48} = P[4, 1 - 2\kappa_{\mathcal{R}}]$. The notation $Z[\cdot]$ in (B.4) is the zeta function.

References

- [1] J. Kysely and J. Picek, "Probability estimates of heavy precipitation events in a flood-prone central-European region with enhanced influence of Mediterranean cyclones," *Advances in Geosciences*, vol. 12, no. 1, pp. 43–50, 2007.
- [2] K. Kochanek, W. Strupczewski, and V. P. Singh, "The PWM large quantile estimates of heavy tailed distributions from samples deprived of their largest element," *Hydrological Sciences Journal*, vol. 53, no. 2, pp. 367–386, 2008.
- [3] W. H. Asquith, *Distributional Analysis with L-Moment Statistics Using the R Environment for Statistical Computing*, Lubbock, Tex, USA, 2011.

- [4] J. R. M. Hosking and J. R. Wallis, *Regional Frequency Analysis: An Approach Based on L-Moments*, Cambridge University Press, Cambridge, UK, 1997.
- [5] B. Ghalebaz-Jeddi, G. L. Donohue, and J. F. Shortle, "A statistical analysis of the aircraft landing process," *Journal of Industrial and Systems Engineering*, vol. 3, no. 3, pp. 152–169, 2009.
- [6] V. Pisarenko and M. Rodkin, *Heavy-Tailed Distributions in Disaster Analysis*, Springer, New York, NY, USA, 2010.
- [7] A. Ravi and F. B. Butar, "An insight into heavy-tailed distribution," *Journal of Mathematical Sciences and Mathematics Education*, vol. 5, no. 1, 15 pages, 2010.
- [8] A. Achim, E. E. Kuruoğlu, and J. Zerubia, "SAR image filtering based on the heavy-tailed rayleigh model," *IEEE Transactions on Image Processing*, vol. 15, no. 9, pp. 2686–2693, 2006.
- [9] K. Yuan and P. M. Bentler, "Structural equation modeling with heavy tailed distributions," *Psychometrika*, vol. 69, no. 3, pp. 421–436, 2004.
- [10] I. Berkovits, G. R. Hancock, and J. Nevitt, "Bootstrap resampling approaches for repeated measure designs: relative robustness to sphericity and normality violations," *Educational and Psychological Measurement*, vol. 60, no. 6, pp. 877–892, 2000.
- [11] C. K. Enders, "The impact of nonnormality on full information maximum-likelihood estimation for structural equation models with missing data," *Psychological Methods*, vol. 6, no. 4, pp. 352–370, 2001.
- [12] T. C. Headrick and S. S. Sawilowsky, "Simulating correlated multivariate nonnormal distributions: extending the fleishman power method," *Psychometrika*, vol. 64, no. 1, pp. 25–35, 1999.
- [13] T. C. Headrick and S. S. Sawilowsky, "Properties of the rank transformation in factorial analysis of covariance," *Communications in Statistics*, vol. 29, no. 4, pp. 1059–1087, 2000.
- [14] T. C. Headrick, *Statistical Simulation: Power Method Polynomials and Other Transformations*, Chapman & Hall, Boca Raton, Fla, USA, 2010.
- [15] J. W. Tukey, "Modern techniques in data analysis," in *Proceedings of the NSF-Sponsored Regional Research Conference*, Southern Massachusetts University, Boston, Mass, USA, 1977.
- [16] J. Martinez and B. Iglewicz, "Some properties of the Tukey g and h family of distributions," *Communications in Statistics A*, vol. 13, no. 3, pp. 353–369, 1984.
- [17] T. C. Headrick, R. K. Kowalchuk, and Y. Sheng, "Parametric probability densities and distribution functions for Tukey g -and- h transformations and their use for fitting data," *Applied Mathematical Sciences*, vol. 2, no. 9, pp. 449–462, 2008.
- [18] S. Morgenthaler and J. W. Tukey, "Fitting quantiles: doubling, HR, HQ, and HHH distributions," *Journal of Computational and Graphical Statistics*, vol. 9, no. 1, pp. 180–195, 2000.
- [19] C. Field and M. G. Genton, "The multivariate g -and- h distribution," *Technometrics*, vol. 48, no. 1, pp. 104–111, 2006.
- [20] R. K. Kowalchuk and T. C. Headrick, "Simulating multivariate g -and- h distributions," *British Journal of Mathematical and Statistical Psychology*, vol. 63, no. 1, pp. 63–74, 2010.
- [21] T. C. Headrick and M. D. Pant, "Characterizing tukey h and hh -distributions through L -moments and the L -correlation," *ISRN Applied Mathematics*, vol. 2012, Article ID 980153, 20 pages, 2012.
- [22] J. R. M. Hosking, "L-moments: analysis and estimation of distributions using linear combinations of order statistics," *Journal of the Royal Statistical Society B*, vol. 52, no. 1, pp. 105–124, 1990.
- [23] R. M. Vogel and N. M. Fennessey, "L-moment diagrams should replace product moment diagrams," *Water Resources Research*, vol. 29, no. 6, pp. 1745–1752, 1993.
- [24] F. A. Hodis, T. C. Headrick, and Y. Sheng, "Power method distributions through conventional moments and L -moments," *Applied Mathematical Sciences*, vol. 6, no. 44, pp. 2159–2193, 2012.
- [25] T. C. Headrick, "A characterization of power method transformations through L -moments," *Journal of Probability and Statistics*, vol. 2011, Article ID 497463, 22 pages, 2011.
- [26] K. P. Singh, C. M. S. Lee, and E. O. George, "On generalized log-logistic model for censored survival data," *Biometrical Journal*, vol. 30, no. 7, pp. 843–850, 1988.
- [27] T. C. Headrick and M. D. Pant, "A method for simulating nonnormal distributions with specified L -Skew, L -Kurtosis, and L -Correlation," *ISRN Applied Mathematics*, vol. 2012, Article ID 980827, 23 pages, 2012.
- [28] M. C. Jones, "On some expressions for variance, covariance, skewness and L -moments," *Journal of Statistical Planning and Inference*, vol. 126, no. 1, pp. 97–106, 2004.
- [29] R. Serfling and P. Xiao, "A contribution to multivariate L -moments: L -comoment matrices," *Journal of Multivariate Analysis*, vol. 98, no. 9, pp. 1765–1781, 2007.
- [30] Wolfram Research Inc., *Mathematica Version 8*, Wolfram Research, Champaign, Ill, USA, 2010.
- [31] G. Marsaglia, "Evaluating the normal distribution function," *Journal of Statistical Software*, vol. 11, no. 5, pp. 1–11, 2004.