Clinical Study

Carbonic Anhydrase I as a New Plasma Biomarker for Prostate Cancer

Michiko Takakura,¹ Akira Yokomizo,² Yoshinori Tanaka,³ Michimoto Kobayashi,³ Giman Jung,³ Miho Banno,⁴ Tomohiro Sakuma,⁴ Kenjiro Imada,^{2, 5} Yoshinao Oda,⁵ Masahiro Kamita,¹ Kazufumi Honda,¹ Tesshi Yamada,¹ Seiji Naito,² and Masaya Ono¹

¹ Division of Chemotherapy and Clinical Research, National Cancer Center Research Institute, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

² Department of Urology, Graduate School of Medical Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, Japan

³ New Frontiers Research Laboratories, Toray Industries, Inc., 10-1 Tebiro, Kanagawa, Kamakura 248-8555, Japan

⁴ Bio Science Department, Research and Development Center, Mitsui Knowledge Industry Co., Ltd., 2-7-14 Higashinakano, Nakano-Ku, Tokyo 164-8555, Japan

⁵ Department of Anatomic Pathology, Graduate School of Medical Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, Japan

Correspondence should be addressed to Masaya Ono, masono@ncc.go.jp

Received 17 September 2012; Accepted 2 October 2012

Academic Editors: A. E. Bilsland, B. Comin-Anduix, G. Ferrandina, and S. Holdenrieder

Copyright © 2012 Michiko Takakura et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Serum prostate-specific antigen (PSA) levels ranging from 4 to 10 ng/mL is considered a diagnostic gray zone for detecting prostate cancer because biopsies reveal no evidence of cancer in 75% of these subjects. Our goal was to discover a new highly specific biomarker for prostate cancer by analyzing plasma proteins using a proteomic technique. Enriched plasma proteins from 25 prostate cancer patients and 15 healthy controls were analyzed using a label-free quantitative shotgun proteomics platform called 2DICAL (2-dimensional image converted analysis of liquid chromatography and mass spectrometry) and candidate biomarkers were searched. Among the 40,678 identified mass spectrum (MS) peaks, 117 peaks significantly differed between prostate cancer patients and healthy controls. Ten peaks matched carbonic anhydrase I (CAI) by tandem MS. Independent immunological assays revealed that plasma CAI levels in 54 prostate cancer patients were significantly higher than those in 60 healthy controls (P = 0.022, Mann-Whitney U test). In the PSA gray-zone group, the discrimination rate of prostate cancer patients increased by considering plasma CAI levels. CAI can potentially serve as a valuable plasma biomarker and the combination of PSA and CAI may have great advantages for diagnosing prostate cancer in patients with gray-zone PSA level.

1. Introduction

Prostate cancer is the most common malignancy in the United State. In 2009, 192,280 men were estimated to have been diagnosed with prostate cancer, and 27,360 of these patients died in the United States [1]. The prostate-specific antigen (PSA) is used for the detection of prostate cancer in daily practice, but its diagnostic reliability is hampered by its low specificity. Thus, serum PSA levels ranging from 4 to 10 ng/mL are called the "gray zone" in which it is

very difficult to discriminate between patients with prostate cancer and those with benign prostatic hyperplasia (BPH), prostatitis, or normal prostate. Furthermore, among the patients with serum PSA levels between 4 to 10 ng/mL, only 25% will be found to have prostate cancer [2]. Serum PSA levels can also increase in prostatitis, [3, 4] and approximately 20%–30% of prostate cancers are missed when the cut-off value is set to 4 ng/mL [5–7]. The false negative rate in the first biopsy is estimated between 12% and 32% [8, 9], and a large population of men with chronically high serum PSA

levels undergo repeated biopsies to eliminate the possibility of prostate cancer [3, 4].

Our quantitative label-free shotgun proteomics analysis system, called 2-dimensional image converted analysis of liquid chromatography and mass spectrometry (2DICAL), can accurately align different liquid chromatography-mass spectrometry (LC-MS) data sets, enabling rapid comparison of a statistically sufficient number of clinical samples [10-16]. 2DICAL has a characteristic of top-down proteomics in shotgun proteomics. It converts the LC-MS spectrum data into peaks on a 2-dimensional plane with axes of mass-tocharge ratio (m/z) and retention time (RT). The peaks with the same m/z and RT are compared across the samples, and statistically significant peaks are selected. Targeted tandem mass spectrometry (MS) is conducted on the selected peaks, and the peaks are annotated by sequence search programs (see Supplemental Materials available online at doi: 10.5402/2012/768190).

Here we describe the discovery of a new candidate biomarker for prostate cancer diagnosis that we uncovered using 2DICAL to compare the plasma proteomes of prostate cancer patients with those of healthy controls.

2. Materials and Methods

2.1. Clinical Samples. Plasma samples were prospectively collected at the Department of Urology and Ophthalmology, Graduate School of Medical Sciences, Kyushu University (Fukuoka, Japan), between October 2000 and January 2008 from 162 individuals, including those suffering from prostate cancer (n = 54), renal cell cancer (RCC; n = 20), prostatitis (n = 6), and BPH (n = 22) and 60 healthy individuals who had no symptom and PSA below 10 ng/mL, and those with PSA over 4 ng/mL were periodically followed in outpatient clinic with no evidence of prostate cancer. Prostate cancer patients were definitively diagnosed by prostate biopsy. Patient characteristics including age, PSA levels, Gleason score, and TNM classification are shown in Table 1. For 2DICAL analysis, we selected prostate cancer patients and age-matched healthy controls. All patients provided written informed consent authorizing the collection and use of their samples for research. The institutional ethics committee boards of the National Cancer Center Research Institute (Tokyo, Japan) and the Kyushu University reviewed and approved our protocol.

2.2. Sample Preparation. To exclude sampling bias, 7 mL of each patient's whole blood was collected in a tube containing ethylenediaminetetraacetic acid-2Na (Venoject II, Terumo, Japan) before surgery or first treatment. Plasma was prepared by centrifuging samples at $1,050 \times g$ for 10 min at 4°C. Aliquots of 1 mL were added to 1.5-mL Eppendorf tubes and frozen at -80° C until analysis. Control samples were collected and stored identically. All samples were subjected to only 1 freeze-thaw cycle. To enrich low molecular weight plasma proteins, $500 \,\mu$ L of plasma was diluted to 4 mL by adding 25 mM ammonium bicarbonate buffer (pH 8.0), and the diluted plasma samples were processed using a hollow fiber membrane-based low molecular weight (LMW) protein

enrichment device as described previously [14, 17]. The device employs multistage filtration and cascaded cross-flow processes, and the proteins smaller than a predetermined molecular weight can be separated in a fully automated operation. The solution enriched for LMW proteins was recovered for 1 h operation and the LMW proteins were digested at 37°C for 18 h with sequencing grade modified trypsin (Promega, Madison, WI).

2.3. LC-MS. Trypsin-digested samples were analyzed in duplicate by nanoflow high-performance liquid chromatography (NanoFrontier nLC, Hitachi High-technologies, Tokyo, Japan) connected to an electrospray ionization quadrupole time-of-flight mass spectrometer (Q-Tof Ultima, Waters, Milford, MA). MS peaks were detected, normalized, and quantified using our 2DICAL software package [10, 12]. A serial identification (ID) number was applied to each of the MS peaks detected. The reproduction of LC-MS was monitored by calculating the correlation coefficient (CC) and coefficient of variance (CV) of every measurement.

2.4. Protein Identification by Tandem Mass Spectrometry (MS/MS). Peak lists were generated using the Mass Navigator software package (Mitsui Knowledge Industry, Tokyo, Japan) and searched against the SwissProt database using the Mascot software package (Matrix Science, London, UK). Search parameters used were as follows: human protein sequences were selected, trypsin was designated as the enzyme, and up to 1 missed cleavage was allowed. Mass tolerances for precursor and fragment ions were ± 0.6 Da and ± 0.2 Da, respectively. The score threshold was set to a Mascot score >30. If a peptide matched to multiple proteins, the protein name with the highest Mascot score was selected.

2.5. Western Blot Analysis. Plasma samples were analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis using 10%–20% (w/v) ready-made gels (Pagel; ATTO, Tokyo, Japan) with the Laemmli buffer and electroblotted to a polyvinylidene difluoride membrane (Millipore, Billerica, MA). Primary antibodies were goat polyclonal carbonic anhydrase I (CAI) antibody (Millipore, Billerica, MA) and mouse monoclonal antibody against human complement C3b- α (PROGEN, Heidelberg, Germany). The membrane was incubated with the primary antibodies and then with the relevant horseradish peroxidase (HRP)-conjugated antigoat or anti-mouse IgG as described previously [18]. Blots were developed using an enhanced chemiluminescence plus detection system (GE Healthcare, Buckinghamshire, UK).

2.6. Enzyme-Linked Immunosorbent Assay (ELISA). To develop sandwich ELISA, the capture antibody (rabbit polyclonal CAI antibody, Abnova, Taipei, Taiwan) was immobilized on a 96-well plate (Thermo Fisher Scientific, MA) at a final concentration of 2.5 μ g/mL and incubated at 4°C overnight. A mouse monoclonal CAI antibody (0.5 μ g/mL; Abnova, Taipei, Taiwan) was used as the detection antibody. After incubation with HRP-conjugated goat anti-mouse IgG (Vector Laboratories, Burlingame, CA) for 1 h and then with

				All cases ((n = 162)			Cases anal	vzed by 2DICAL	n = 40
		$PCa^*(n = 54)$	BPH $(n = 22)$	Prostatitis $(n = 6)$	RCC $(n = 20)$	Healthy $(n = 60)$	P value	PCa* $(n = 25)$	Healthy $(n = 15)$	P value
Age	$(mean \pm SD)$	66.3 ± 7.3	72.3 ± 7.5	65.8 ± 5.9	63.8 ± 8.5	67.2 ± 8.8	0.544^{****}	67.0 ± 7.2	65.1 ± 8.0	0.440^{****}
042.00	I	0						0		
olage	II	50						22		
	III	2						2		
	IV	2						1		
		ц						01		
	TONIOUTO	C.4 11						01 г		
	1 ZAINUMIU	CI								
TNIM classification **	T2bN0M0	2						1		
TINIM CIASSIIICAUOII	T2cN0M0	8						4		
	T3aN0M0	2						2		
	T4N0M1b	1						1		
	T4N1M1b	1						0		
	ng/mL									
*** v UC	<4	2	11	4		51		2	12	
V VCJ	$4\leq,\leq 10$	30	10	2		6	<0.001****	14	ŝ	<0.001****
	10<	22	1	0		0		6	0	
	ſ	ç						-		
	2	1 [+ F		
5	5 1	CI 5								
Gleason score	/	C7						12		
	8	8						4		
	6	2						1		
	10	2						0		
*Abbreviation: PCa: prc	state cancer; ** a *Fisher's evect fee	iccording to TNM	Classification of	Malignant Tumors	(International Unio	n Against Cancer), 6	5th Edition; ***	* measured by Tand	em R kit before the	e first treatment;

TABLE 1: Clinicopathological characteristics of individuals examined in this study.

ISRN Oncology

3



FIGURE 1: (a) Two-dimensional display of all MS peaks. The 117 MS peaks whose mean intensities significantly differed between prostate cancer patients and healthy controls (P < 0.05, Welch's *t*-test) are highlighted in red. RT: retention time. (b) 2DICAL images of peak ID 396 in a representative prostate cancer patient (left) and a healthy control (right).

OPD solution for 10 min, absorbance was measured at 490 nm using EnSpire AlphaPLUS (PerkinElmer, Waltham, MA).

2.7. Cell Lines. The human prostate cancer cell line 22Rv1 was purchased from Riken BRC Cell Bank (Tsukuba, Japan) and cultured in Roswell Park Memorial Institute Medium 1640 supplemented with 10% fetal bovine serum. Normal human prostate epithelial cells (CC-3165 PrEBM) were purchased from Lonza (Basel, Switzerland). The cells were cultured at 37°C under 5% CO₂. The culture medium was

changed every 3 days. After incubation of cell lines for 48 h, the cultured conditioned media were collected, filtered through a sterile 0.22-mm filter unit, and concentrated 100-fold by freeze drying. A 10 μ L sample of concentrated medium was used for Western blot analysis.

2.8. Immunohistochemistry. Cells were seeded on BioCoat collagen I-coated 8-well culture slides (Becton Dickinson Labware, Bedford, MA) and cultured overnight. The cells were washed twice with PBS (pH 7.2) and fixed with 4% paraformaldehyde in PBS for 20 min at room temperature.



FIGURE 2: (a) MS peaks of peak ID 396 in triplicate LC-MS runs (25 with prostate cancer (left) and 15 with healthy controls (right)) aligned along LC RT. Columns represent the mean intensities of triplicate runs. (b) Immunoblotting of CAI and complement C3b- α (loading control) for 7 prostate cancer patients selected from the most intense MS peaks and 7 healthy controls from the least intense peaks.

Fixed cells were permeabilized with 0.1% Triton in PBS for 5 min, washed 3 times with PBS, blocked for 30 min with 5% normal donkey serum (Chemicon International, Inc., Temecula, CA) in PBS, and stained with the indicated primary antibodies diluted in 1% normal donkey serum overnight at room temperature. After washing 3 times with PBS, fluorophore-conjugated secondary antibodies were applied for 1 h at room temperature and washed with PBS. Alexa Fluor 568-phalloidin was used to visualize the actin cytoskeleton. The slides were mounted with Vectashield antifade reagent (Vector Laboratories), covered with glass coverslips, and observed under Zeiss LSM510 fluorescence microscope equipped with 488/514-nm argon and 543-nm helium-neon lasers.

To stain human tissues, 10 radical prostatectomy specimens were selected with five high and five low plasma CAI concentrations. Sections $(4\,\mu\text{m})$ from 10% formalin-fixed paraffin-embedded material were deparaffinized in xylene and rehydrated in ethanol. Endogenous peroxidase activity was blocked by methanol containing 0.3% hydrogen peroxidase for 30 min. Microwave heating was used for antigen retrieval. The sections were incubated at 4°C overnight with a primary antibody and then incubated with a second antibody for 40 min at room temperature. The reaction products were visualized using diaminobenzidine tetrahydrochloride and counterstained with hematoxylin.

2.9. Statistical Analysis. Statistical significance of intergroup differences was assessed by the Welch's *t*-test and Mann-Whitney *U* test. Area under the receiver operating characteristic (ROC) curve (AUC) was calculated for each marker to evaluate its diagnostic significance. A composite index of 2 markers was generated using the results of multivariate logistic regression analysis, which also enabled the calculation of sensitivity, specificity, and ROC curve. Statistical analyses were performed using an open-source statistical language R with the optional module design package.

3. Results

3.1. Plasma Biomarker Discovery by Quantitative MS. To identify a diagnostic biomarker for prostate cancer patients we compared the plasma proteins of 25 prostate cancer patients with those of 15 healthy controls using 2DICAL (Table 1). Among 40,678 independent MS peaks detected within 250–1, 600 *m/z*, and 30–62.5 min, we found 117 peaks showing significant differences between prostate cancer patients and healthy controls (P < 0.05, Welch's t-test). Figure 1(a) shows a representative 2-dimensional view of all MS peaks displayed with m/z along the x-axis and LC retention time along the y-axis. The 117 MS peaks whose expression levels differed significantly between cancer patients and healthy controls are highlighted in red. MS/MS spectra acquired from these 117 MS peaks matched 4 proteins in the database with Mascot scores of >30. Ten peaks matched amino acid sequences of CAI (Table 2). The CAIderived peak (ID 396) that clearly differed between cancer patients and healthy controls is shown as a representative peak (Figure 1(b)). The intensity distribution of peak ID 396 was different between prostate cancer patients (left) and healthy controls (right) (Figure 2(a)). Immunoblotting with a CAI probe confirmed the results of the 2DICAL findings (Figure 2(b)).

3.2. Large-Scale Validation of Plasma CAI by ELISA. To further validate plasma CAI levels in prostate cancer patients determined using 2DICAL, we performed ELISA to quantify plasma CAI levels in numerous plasma samples. The plasma samples were derived from patients suffering from prostate cancer (n = 54), prostatitis (n = 6), BPH (n = 22), and RCC (n = 20) as well as from healthy controls (n = 60). Plasma CAI levels were significantly different between prostate cancer patients and healthy controls (P = 0.022, Mann-Whitney U test). Plasma CAI levels of patients with BPH or RCC did not show significant differences from healthy

D $m/2$ (min) $undigo (mean \pm SD) (mean \pm SD) Values** 3 396 643.7 41.6 3 27.35 \pm 13.42 63.53 \pm 67.39 0.015 310 790.9 47.8 2 24.96 \pm 9.73 62.84 \pm 72.61 0.016 983 753.0 47.5 3 16.02 \pm 4.60 30.40 \pm 27.50 0.017 1463 719.4 51.8 2 25.20 \pm 22.26 11.26 \pm 2.32 0.030 11656 872.4 46.0 2 16.99 \pm 5.22 25.86 \pm 19.63 0.041 311 513.8 37.2 2 37.71 \pm 16.18 86.33 \pm 85.17 0.010 798 585.3 45.4 2 21.08 \pm 6.68 43.99 \pm 34.82 0.013 798 585.3 47.6 2 21.08 \pm 6.68 43.99 \pm 34.82 0.013 798 493.2 41.06 2 21.108 \pm 6.68 43.99 \pm 34.82 0.013 $	L PCa	-	INTRON L	Dantida comanca	Drotain description	I Inimat ID
$\overline{396}$ 643.7 41.6 3 27.35 ± 13.42 63.53 ± 67.39 0.015 310 790.9 47.8 2 24.96 ± 9.73 62.84 ± 72.61 0.016 983 753.0 47.5 3 16.02 ± 4.60 30.40 ± 27.50 0.017 1463 719.4 51.8 2 25.20 ± 22.26 11.26 ± 2.32 0.030 1656 872.4 46.0 2 16.99 ± 5.22 25.86 ± 19.63 0.041 311 513.8 37.2 2 37.71 ± 16.18 86.33 ± 85.17 0.010 798 585.3 45.4 2 21.08 ± 6.68 43.99 ± 34.82 0.004 278 485.8 47.6 2 40.70 ± 20.28 91.91 ± 92.78 0.016 278 485.8 47.6 2 20.19 ± 5.79 47.30 ± 52.18 0.016 278 485.8 47.6 2 20.19 ± 5.79 47.30 ± 52.18 0.016 2742 680.4 47.5 2 20.19 ± 5.79 47.30 ± 52.18 0.016 2429 680.4 47.5 2 27.43 ± 10.68 20.70 ± 3.90 0.032 1369 714.4 48.1 3 21.16 ± 5.02 31.96 ± 19.53 0.014 916 494.3 48.9 2 12.19 ± 4.64 31.68 ± 32.02 0.006 747 920.8 45.3 3 9.93 ± 2.84 19.94 ± 16.30 0.013 9156 55.3 36.6 2 9.93 ± 2.84 19.9	\pm SD) (mean \pm SD)	Values ^{**} s	score	repute sequence	rioteni aescripuon	
310 790.9 47.8 2 24.96 ± 9.73 62.84 ± 72.61 0.016 983 753.0 47.5 3 16.02 ± 4.60 30.40 ± 27.50 0.017 1463 719.4 51.8 2 25.20 ± 22.26 11.26 ± 2.32 0.030 1656 872.4 46.0 2 16.99 ± 5.22 25.86 ± 19.63 0.041 311 513.8 37.2 2 37.71 ± 16.18 86.33 ± 85.17 0.010 798 585.3 45.4 2 21.08 ± 6.68 43.99 ± 34.82 0.004 778 485.8 47.6 2 40.70 ± 20.28 91.91 ± 92.78 0.013 538 493.2 41.0 2 20.19 ± 5.79 47.30 ± 52.18 0.016 278 485.8 47.6 2 20.19 ± 5.79 47.30 ± 52.18 0.016 278 495.2 41.0 2 20.19 ± 5.79 47.30 ± 52.18 0.016 218 493.2 41.0 2 20.19 ± 5.79 47.30 ± 52.18 0.016 2429 680.4 47.5 2 27.43 ± 10.68 20.70 ± 3.90 0.032 1369 714.4 48.1 3 21.16 ± 5.02 31.96 ± 19.53 0.014 916 494.3 48.9 2 12.19 ± 4.64 31.68 ± 32.02 0.006 747 92.08 45.3 3 9.93 ± 2.84 19.94 ± 16.30 0.013 9156 55.3 36.6 2 9.93 ± 2.84 19.94 ± 1	-13.42 63.53 ± 67.39	0.015	95.02	HDTSLKPISVSYNPATAK	Carbonic anhydrase 1	CAH1_HUMAN
983753.0 47.5 3 16.02 ± 4.60 30.40 ± 27.50 0.017 1463 719.4 51.8 2 25.20 ± 22.26 11.26 ± 2.32 0.030 1656 872.4 46.0 2 16.99 ± 5.22 25.86 ± 19.63 0.041 311 513.8 37.2 2 37.71 ± 16.18 86.33 ± 85.17 0.010 798 585.3 45.4 2 37.71 ± 16.18 86.33 ± 85.17 0.010 798 585.3 45.4 2 21.08 ± 6.68 43.99 ± 34.82 0.004 278 485.8 47.6 2 40.70 ± 20.28 91.91 ± 92.78 0.016 278 493.2 41.0 2 20.19 ± 5.79 47.30 ± 52.18 0.016 2429 680.4 47.5 2 20.19 ± 5.79 47.30 ± 52.18 0.016 2429 680.4 47.5 2 27.43 ± 10.68 20.70 ± 3.90 0.032 1369 714.4 48.1 3 21.16 ± 5.02 31.96 ± 19.53 0.014 916 494.3 48.9 2 12.19 ± 4.64 31.68 ± 32.02 0.006 747 920.8 45.3 36.6 2 9.93 ± 2.84 19.94 ± 16.30 0.013 5156 556.3 36.6 2 9.93 ± 2.84 19.94 ± 16.30 0.016	± 9.73 62.84 ± 72.61	0.016	72.71	ESISVSSEQLAQFR	Carbonic anhydrase 1	CAH1_HUMAN
1463 719.4 51.8 2 25.20 ± 22.26 11.26 ± 2.32 0.030 1656 872.4 46.0 2 16.99 ± 5.22 25.86 ± 19.63 0.041 311 513.8 37.2 2 37.71 ± 16.18 86.33 ± 85.17 0.010 798 585.3 45.4 2 21.08 ± 6.68 43.99 ± 34.82 0.004 798 585.3 47.6 2 40.70 ± 20.28 91.91 ± 92.78 0.013 538 493.2 41.0 2 20.19 ± 5.79 47.30 ± 52.18 0.016 2429 680.4 47.5 2 20.19 ± 5.79 47.30 ± 52.18 0.016 2429 680.4 47.5 2 20.19 ± 5.79 47.30 ± 52.18 0.016 3166 714.4 48.1 3 21.16 ± 5.02 31.96 ± 19.53 0.014 916 494.3 48.9 2 12.19 ± 4.64 31.68 ± 32.02 0.006 747 920.8 45.3 36.6 2 9.93 ± 2.84 19.94 ± 16.30 0.013	± 4.60 30.40 ± 27.50	0.017	71.62	EIINVGHSFHVNFEDNDNR	Carbonic anhydrase 1	CAH1_HUMAN
1656 872.4 46.0 2 16.99 ± 5.22 25.86 ± 19.63 0.041 311 513.8 37.2 2 37.71 ± 16.18 86.33 ± 85.17 0.010 798 585.3 45.4 2 21.08 ± 6.68 43.99 ± 34.82 0.004 278 485.8 47.6 2 40.70 ± 20.28 91.91 ± 92.78 0.013 538 493.2 41.0 2 20.19 ± 5.79 47.30 ± 52.18 0.016 2429 680.4 47.5 2 20.19 ± 5.79 47.30 ± 52.18 0.016 1369 714.4 48.1 3 21.16 ± 5.02 31.96 ± 19.53 0.014 916 494.3 48.9 2 12.19 ± 4.64 31.68 ± 32.02 0.006 747920.8 45.3 3 9.93 ± 2.84 19.94 ± 16.30 0.013 1596 656.3 36.6 2 9.93 ± 2.84 19.94 ± 16.30 0.016	± 22.26 11.26 ± 2.32	0.030	66.87	GLEEELQFSLGSK	Complement C4-A	CO4A_HUMAN
311513.8 37.2 2 37.71 ± 16.18 86.33 ± 85.17 0.010 798585.3 45.4 2 21.08 ± 6.68 43.99 ± 34.82 0.004 278 485.8 47.6 2 40.70 ± 20.28 91.91 ± 92.78 0.013 538 493.2 41.0 2 20.19 ± 5.79 47.30 ± 52.18 0.016 2429 680.4 47.5 2 27.43 ± 10.68 20.70 ± 3.90 0.032 1369 714.4 48.1 3 21.16 ± 5.02 31.96 ± 19.53 0.014 916 494.3 48.9 2 12.19 ± 4.64 31.68 ± 32.02 0.006 747 920.8 45.3 3 9.93 ± 2.84 19.94 ± 16.30 0.013 1596 55.3 36.6 2 9.93 ± 2.84 19.94 ± 16.30 0.006	± 5.22 25.86 ± 19.63	0.041	66.24	LYPIANGNNQSPVDIK	Carbonic anhydrase 1	CAH1_HUMAN
798585.345.42 21.08 ± 6.68 43.99 ± 34.82 0.004 278485.847.62 40.70 ± 20.28 91.91 ± 92.78 0.013 538493.241.02 20.19 ± 5.79 47.30 ± 52.18 0.016 2429680.4 47.5 2 27.43 ± 10.68 20.70 ± 3.90 0.032 1369714.448.13 21.16 ± 5.02 31.96 ± 19.53 0.014 916494.348.92 12.19 ± 4.64 31.68 ± 32.02 0.006 747920.8 45.3 3 14.00 ± 4.82 35.85 ± 40.39 0.013 1596 656.3 36.6 2 9.93 ± 2.84 19.94 ± 16.30 0.006	$\pm 16.18 86.33 \pm 85.17$	0.010	59.09	YSSLAEAASK	Carbonic anhydrase 1	CAH1_HUMAN
278 485.8 47.6 2 40.70 ± 20.28 91.91 ± 92.78 0.013 538 493.2 41.0 2 20.19 ± 5.79 47.30 ± 52.18 0.016 2429 680.4 47.5 2 27.43 ± 10.68 20.70 ± 3.90 0.032 1369 714.4 48.1 3 21.16 ± 5.02 31.96 ± 19.53 0.014 916 494.3 48.9 2 12.19 ± 4.64 31.68 ± 32.02 0.006 747 920.8 45.3 3 14.00 ± 4.82 35.85 ± 40.39 0.013 1596 656.3 36.6 2 9.93 ± 2.84 19.94 ± 16.30 0.006	$\pm 6.68 43.99 \pm 34.82$	0.004	56.32	SADFTNFDPR	Carbonic anhydrase 2	CAH2_HUMAN
538493.241.02 20.19 ± 5.79 47.30 ± 52.18 0.016 2429680.4 47.5 2 27.43 ± 10.68 20.70 ± 3.90 0.032 1369714.448.13 21.16 ± 5.02 31.96 ± 19.53 0.014 916494.348.9212.19 \pm 4.64 31.68 ± 32.02 0.006 747920.8 45.3 3 14.00 ± 4.82 35.85 ± 40.39 0.013 1596656.3 36.6 2 9.93 ± 2.84 19.94 ± 16.30 0.006	$\pm 20.28 91.91 \pm 92.78$	0.013	50.23	VLDALQAIK	Carbonic anhydrase 1	CAH1_HUMAN
2429 680.4 47.5 2 27.43 ± 10.68 20.70 ± 3.90 0.032 1369 714.4 48.1 3 21.16 ± 5.02 31.96 ± 19.53 0.014 916 494.3 48.9 2 12.19 ± 4.64 31.68 ± 32.02 0.006 747 920.8 45.3 3 14.00 ± 4.82 35.85 ± 40.39 0.013 1596 656.3 36.6 2 9.93 ± 2.84 19.94 ± 16.30 0.006	$\pm 5.79 47.30 \pm 52.18$	0.016	44.71	GGPFSDSYR	Carbonic anhydrase 1	CAH1_HUMAN
1369 714.4 48.1 3 21.16 ± 5.02 31.96 ± 19.53 0.014 916 494.3 48.9 2 12.19 ± 4.64 31.68 ± 32.02 0.006 747 920.8 45.3 3 14.00 ± 4.82 35.85 ± 40.39 0.013 1596 656.3 36.6 2 9.93 ± 2.84 19.94 ± 16.30 0.006	-10.68 20.70 \pm 3.90	0.032	43.87	LNDLEDALQQAK	Keratin, type II cytoskeletal 1	K2C1_HUMAN
916 494.3 48.9 2 12.19 \pm 4.64 31.68 \pm 32.02 0.006 747 920.8 45.3 3 14.00 \pm 4.82 35.85 \pm 40.39 0.013 1596 656.3 36.6 2 9.93 \pm 2.84 19.94 \pm 16.30 0.006	± 5.02 31.96 ± 19.53	0.014	43.57	YDPSLKPLSVSYDQATSLR	Carbonic anhydrase 2	CAH2_HUMAN
747 920.8 45.3 3 14.00 ± 4.82 35.85 ± 40.39 0.013 1596 656.3 36.6 2 9.93 ± 2.84 19.94 ± 16.30 0.006	± 4.64 31.68 ± 32.02	0.006	39.1	VVDVLDSIK	Carbonic anhydrase 2	CAH2_HUMAN
1596 656.3 36.6 2 9.93 ± 2.84 19.94 ± 16.30 0.006	± 4.82 35.85 ± 40.39	0.013	36.65	SLLSNVEGDNAVPMQHNNRPTQPLK	Carbonic anhydrase 1	CAH1_HUMAN
	$2.2.84$ 19.94 \pm 16.30	0.006	33.83	QSPVDIDTHTAK	Carbonic anhydrase 2	CAH2_HUMAN
$3781 826.1 40.2 3 4.72 \pm 0.94 9.10 \pm 9.25 0.027$	-0.94 9.10 \pm 9.25	0.027	32.29	TSETKHDTSLKPISVSYNPATAK	Carbonic anhydrase 1	CAH1_HUMAN
1199 538.3 40.7 3 19.72 ± 3.65 30.15 ± 23.16 0.036	$\pm 3.65 30.15 \pm 23.16$	0.036	30.33	YSAELHVAHWNSAK	Carbonic anhydrase 1	CAH1_HUMAN

s spectromet	
mas	
v tandem	
Ę,	
luc	
catio	
ntifi	
ide	
rotein	
ā	
of	
Summarv	
::	
TABLE (



FIGURE 3: (a) Plasma CAI levels in patients with prostate cancer (n = 54), BPH (n = 22), prostatitis (n = 6), and RCC (n = 20) and healthy controls (n = 60) were 1.43 ± 0.69, 1.03 ± 0.51, 0.98 ± 0.46, 0.99 ± 0.36, and 1.09 ± 1.82 µg/mL (mean ± SD), respectively. There was a significant difference between prostate cancer patients and healthy controls (P = 0.020, Mann-Whitney U test). Horizontal lines represented the average levels. (b) Scatter plot correlating PSA and CAI levels in prostate cancer patients (red) and healthy controls (black). The dotted lines were added to indicate the PSA gray zone. Serum PSA levels were measured using the Tandem-R kit before the first treatment in each patient. (c) ROC curve of PSA plus CAI (red line) and PSA alone (blue line) confined to the cases with PSA levels in the gray zones. ROC curves were created using a composite index of the 2 markers generated from the results of multivariate logistic regression analysis. As a reference, ROC curves of CAI (black line) for all cases were included.

controls. Plasma CAI levels of prostate cancer patients were clearly higher than those of any other patients (Figure 3(a)).

3.3. Combination of Plasma CAI Levels and PSA Assays. To determine whether plasma CAI levels and PSA assays together would be useful for diagnosing prostate cancer, CAI and PSA levels from the same cases were compared (Figure 3(b)). Pearson's CC between them was -0.106, which meant that these 2 proteins had different vectors in blood concentration. From the distribution data in the plot, CAI levels in prostate cancer patients with PSA levels of >20 ng/mL were low and CAI levels were higher in prostate cancer patients with PSA levels in the gray zone, compared to healthy controls.

To understand the diagnostic significance of CAI levels that coincided with PSA levels in the gray zone, subjects



FIGURE 4: CAI expression in cell lines. (a) Immunofluorescence detection of CAI in normal prostate epithelial cells and prostate cancer cells. The cells were fixed for 1 h, permeabilized, and stained with antibodies to detect CAI. Alexa Fluor 488- or Alexa Fluor 568-phalloidin was used to visualize normal prostate epithelial cells (upper panels) and prostate cancer cells of 22Rv1 (lower panels). (b) Western blot analysis of media conditioned by the normal prostate epithelial cells and prostate cancer cells probed using a CAI antibody. C3b- α was used as a loading control.

whose PSA levels were within this gray zone were investigated further. Selected cases were as follows: prostate cancer (n = 30), BPH (n = 10), prostatitis (n = 2), and healthy controls (n = 9). ROC curves for PSA alone and PSA plus CAI were generated for prostate cancer patients relative to the other cases as described in Materials and Methods (Figure 3(c)). AUCs were 0.763 and 0.694, respectively, for PSA plus CAI and PSA alone in the gray zone. PSA levels had a great discriminatory result with an AUC of 0.939 for all cases. However, PSA levels in the gray zone did not provide sufficient discriminatory power when considered alone. This indicates that CAI levels could improve the PSA assay.

3.4. Subcellular Localization and Secretion of CAI. The subcellular localization of CAI was investigated by immunofluorescence microscopy. CAI was hardly detected in the normal prostate epithelial cells. In comparison to actin as a cytoskeletal marker (red, Figure 4(a), b and e), immunofluorescence staining with anti-CAI (green, Figure 4(a), a and d) was observed in the cytoplasm of prostate cancer cells (merged, Figure 4(a), f). CAI was clearly detected in media harvested from 22Rv1 cells (Figure 4(b)).

3.5. The Staining of Human Prostate Cancer Cells. CAI was expressed in every case of prostate cancer. Cancer cells from patients with high plasma CAI concentrations tended to have stronger CAI staining than normal prostatic glands (Figure 5).

4. Discussion

We took advantage of our originally developed label-free proteomic technique (2DICAL) [16] to discover a better biomarkers for prostate cancer diagnosis. We identified that CAI peptide fragments were detected at higher levels in plasma samples from prostate cancer patients than in plasma



FIGURE 5: Immunohistochemical staining of CAI. CAI was strongly stained at prostate cancer (magnification $\times 100$).

samples from healthy controls. The 2DICAL results were confirmed by Western blot analysis and ELISA using numerous plasma samples including those from patients with urological diseases other than prostate cancer. We found that the combination of CAI and PSA assays has a potential for improving the specificity of PSA assay especially for PSA levels in the gray zone. These initial results suggest that it is reasonable to vigorously pursue CAI as a potentially valuable new biomarker for prostate cancer.

The PSA assay has largely improved the detection of prostate cancer, but only approximately 25% of patients with PSA levels in the gray zone indeed have prostate cancer [19]. Despite the development of several variations of the PSA assay such as free PSA, PSA velocity, or PSA density, these methods do not substantially outperform analysis of total PSA and are not more specific [19, 20]. The reason is because healthy males are known to have PSA levels in the gray zone. Therefore, improved prostate cancer markers such as CAI are eagerly awaited to overcome these problems and enhance diagnostic specificity and sensitivity.

CAI is a zinc metalloenzyme that catalyzes the reversible hydration of carbon dioxide to bicarbonate. Sixteen CA isoforms exist in mammals [21]. Some of them are cytosolic (CAI, CAII, CAIII, CAVII, and CAXIII) catalyzing the hydration of CO_2 to H⁺ and HCO_3^- , and subsequently, exporting them from the cell in exchange for Na⁺ and Cl⁻ ions [22]. CAI is a specific marker for the cytoplasm or apical cell membranes of colonic epithelial cells [22, 23] and is related to enterocyte proliferation [24]. CAI is also involved in electroneutral NaCl reabsorption and short chain fatty acid uptake [22]. The biological functions of CAs are of great interest, but CA family members are also being studied as drug targets for treating several diseases such as glaucoma, cancer, obesity, and infections [25].

To the best of our knowledge this is the first report to consider a correlation between CAI blood levels and prostate cancer. To verify the upregulation of plasma CAI in prostate cancer, we investigated the cell biology of prostate cancer using immunofluorescent staining to elucidate its subcellular localization, and Western blotting of culture media to determine if CAI was secreted. We demonstrated increased CAI production and secretion in prostate cancer cell lines. Stronger staining of CAI was observed in prostate cancer cells from patients with high plasma CAI concentrations compared to that in normal prostate glands. However, to clarify the mechanism of the plasma concentration change of CAI, further investigations will be needed considering the secretion mechanism of CAI from prostate cancer cells.

Our present proteome research is new in that we found increased plasma CAI levels in prostate cancer patients. Furthermore, it indicates the possibility that CAI was produced and secreted by prostate cancer cell lines. This study may lead to clinically using CAI as a new prostate cancer marker and the combination of PSA and CAI may have great advantages for diagnosing prostate cancer in patients with gray-zone PSA levels.

Conflict of Interests

The authors have no conflict of interests.

Authors' Contribution

M. Takakura and A. Yokomizo contributed equally to the study.

Acknowledgments

The authors thank Ms. Ayako Ikarashi, Ms. Tomoko Umaki, and Ms. Yuka Nakamura for their technical assistance. Funding was provided by the Program for Promotion of Fundamental Studies in Health Sciences conducted by the National Institute of Biomedical Innovation of Japan, the Third-Term Comprehensive Control Research for Cancer and Research on Biological Markers for New Drug Development conducted by the Ministry of Health and Labor of Japan. These sponsors had no role in the design of the study, collection of the data, analysis and interpretation of the data, decision to submit the paper for publication, or writing of the paper.

References

- A. Jemal, R. Siegel, E. Ward, Y. Hao, J. Xu, and M. J. Thun, "Cancer statistics, 2009," *CA—A Cancer Journal for Clinicians*, vol. 59, no. 4, pp. 225–249, 2009.
- [2] W. J. Catalona, D. S. Smith, T. L. Ratliff, and J. W. Basler, "Detection of organ-confined prostate cancer is increased through prostate- specific antigen-based screening," *JAMA*, vol. 270, no. 8, pp. 948–954, 1993.
- [3] J. Pannek and A. W. Partin, "Prostate-specific antigen: what's new in 1997," *Oncology*, vol. 11, no. 9, pp. 1273–1278, 1997.
- [4] S. Loeb, S. N. Gashti, and W. J. Catalona, "Exclusion of inflammation in the differential diagnosis of an elevated prostatespecific antigen (PSA)," *Urologic Oncology*, vol. 27, no. 1, pp. 64–66, 2009.
- [5] W. J. Catalona, J. P. Richie, F. R. Ahmann et al., "Comparison of digital rectal examination and serum prostate specific antigen in the early detection of prostate cancer: Results of a

multicenter clinical trial of 6,630 men," *Journal of Urology*, vol. 151, no. 5, pp. 1283–1290, 1994.

- [6] W. J. Catalona, D. S. Smith, and D. K. Ornstein, "Prostate cancer detection in men with serum PSA concentrations of 2.6 to 4.0 ng/mL and benign prostate examination: Enhancement of specificity with free PSA measurements," *JAMA*, vol. 277, no. 18, pp. 1452–1455, 1997.
- [7] A. Magklara, A. Scorilas, W. J. Catalona, and E. P. Diamandis, "The combination of human glandular Kallikrein and free prostrate- specific antigen (PSA) enhances discrimination between prostate cancer and benign prostatic hyperplasia in patients with moderately increased total PSA," *Clinical Chemistry*, vol. 45, no. 11, pp. 1960–1966, 1999.
- [8] J. Cervera Deval, F. J. Morales Olaya, J. Jornet Fayos, and M. González Añón, "Diagnostic value of the second prostate biopsies in males of risk. Study stratified by value of PSA," *Actas Urologicas Espanolas*, vol. 28, no. 9, pp. 666–671, 2004.
- [9] M. Raber, V. Scattoni, A. Salonia, P. Consonni, and P. Rigatti, "Repeated ultrasound-guided transrectal prostate biopsy in patients with negative histologic test," *Archivio Italiano di Urologia, Andrologia*, vol. 72, no. 4, pp. 197–199, 2000.
- [10] M. Ono, M. Shitashige, K. Honda et al., "Label-free quantitative proteomics using large peptide data sets generated by nanoflow liquid chromatography and mass spectromery," *Molecular and Cellular Proteomics*, vol. 5, no. 7, pp. 1338–1347, 2006.
- [11] A. Negishi, M. Ono, Y. Handa et al., "Large-scale quantitative clinical proteomics by label-free liquid chromatography and mass spectrometry," *Cancer Science*, vol. 100, no. 3, pp. 514– 519, 2009.
- [12] M. Ono, J. Matsubara, K. Honda et al., "Prolyl 4-hydroxylation of α-fibrinogen. A novel protein modification revealed by plasma proteomics," *The Journal of Biological Chemistry*, vol. 284, no. 42, pp. 29041–29049, 2009.
- [13] Y. Murakoshi, K. Honda, S. Sasazuki et al., "Plasma biomarker discovery and validation for colorectal cancer by quantitative shotgun mass spectrometry and protein microarray," *Cancer Science*, vol. 102, no. 3, pp. 630–638, 2011.
- [14] J. Matsubara, K. Honda, M. Ono et al., "Reduced plasma level of CXC chemokine ligand 7 in patients with pancreatic cancer," *Cancer Epidemiology Biomarkers and Prevention*, vol. 20, no. 1, pp. 160–171, 2011.
- [15] A. Yokomizo, M. Takakura, Y. Kanai et al., "Use of quantitative shotgun proteomics to identify fibronectin 1 as a potential plasma biomarker for clear cell carcinoma of the kidney," *Cancer Biomarkers*, vol. 10, no. 3-4, pp. 175–183, 2011.
- [16] M. Ono, M. Kamita, Y. Murakoshi et al., "Biomarker discovery of pancreatic and gastrointestinal cancer by 2DICAL: 2dimensional image-converted analysis of liquid chromatography and mass spectrometry," *International Journal of Proteomics*, vol. 2012, Article ID 897412, 10 pages, 2012.
- [17] Y. Tanaka, H. Akiyama, T. Kuroda et al., "A novel approach and protocol for discovering extremely low-abundance proteins in serum," *Proteomics*, vol. 6, no. 17, pp. 4845–4855, 2006.
- [18] M. Idogawa, T. Yamada, K. Honda, S. Sato, K. Imai, and S. Hirohashi, "Poly(ADP-ribose) polymerase-1 is a component of the oncogenic T-cell factor-4/beta;-catenin complex," *Gastroenterology*, vol. 128, no. 7, pp. 1919–1936, 2005.
- [19] H. Lilja, D. Ulmert, and A. J. Vickers, "Prostate-specific antigen and prostate cancer: Prediction, detection and monitoring," *Nature Reviews Cancer*, vol. 8, no. 4, pp. 268–278, 2008.
- [20] S. Loeb and W. J. Catalona, "What to do with an abnormal PSA test," Oncologist, vol. 13, no. 3, pp. 299–305, 2008.

- [21] C. T. Supuran and A. Scozzafava, "Carbonic anhydrases as targets for medicinal chemistry," *Bioorganic and Medicinal Chemistry*, vol. 15, no. 13, pp. 4336–4350, 2007.
- [22] E. R. Swenson, "Distribution and functions of carbonic anhydrase in the gastrointestinal tract," in *Carbonic Anhydrases: Cellular Physiology and Molecular Genetics*, S. J. Dodgson, R. E. Tashian, G. Gros, and N. D. Carter, Eds., pp. 265–287, Plenum Press, New York, NY, USA, 1991.
- [23] G. Lonnerholm and P. Wistrand, "Carbonic anhydrase in the human fetal gastrointestinal tract," *Biology of the Neonate*, vol. 44, no. 3, pp. 166–176, 1983.
- [24] I. B. Renes, M. Verburg, D. J. P. M. Van Nispen et al., "Epithelial proliferation, cell death, and gene expression in experimental colitis: Alterations in carbonic anhydrase I, mucin MUC2, and trefoil factor 3 expression," *International Journal of Colorectal Disease*, vol. 17, no. 5, pp. 317–326, 2002.
- [25] C. T. Supuran, "Carbonic anhydrase inhibition/activation: Trip of a scientist around the world in the search of novel chemotypes and drug targets," *Current Pharmaceutical Design*, vol. 16, no. 29, pp. 3233–3245, 2010.



The Scientific **World Journal**



Gastroenterology Research and Practice





Journal of Diabetes Research



Disease Markers



Immunology Research





Submit your manuscripts at http://www.hindawi.com





BioMed **Research International**



Journal of Ophthalmology

Computational and Mathematical Methods in Medicine













Research and Treatment





Oxidative Medicine and Cellular Longevity



Stem Cells International



Behavioural Neurology