

Supporting Information for

Microwave-assisted, DBU promoted a new route for the synthesis of 3-hydroxy-benzofurans -2-yl)coumarins from 4-bromomethyl coumarins

Kallimeledoddi Boregowda. Puttaraju¹, Kalegowda. Shivashankar^{1,*}, Vijaykumar P. Rasal², Jagadeeshreddy Eluru², Ebenezer Koyye²

¹ *P.G. Department of Chemistry, Central College Campus, Bangalore University, Bangalore 560 001, Karnataka, India.*


² *Department of Pharmacology, K.L.E. University College of Pharmacy, Belgaum-590 010, Karnataka, India*


*Corresponding author. Tel.: +91 80 22961249. E-mail address: shivashankark@gmail.com

Electronic supplementary information (ESI) available: ¹H NMR spectra of all new compounds, ¹³C NMR spectra of **3h** and LC-MS spectra of **3g**.


Table of content

¹ H NMR spectrum of 1g-j, 2b-k, 3a-k	S1-S25
LC-MS spectra of 3g	S26
¹³ C NMR and HSQC spectrum of 3h	S27-S28








S2





S3







7.83
7.81
7.90
7.75
7.64
7.52
7.50
7.48
7.47
7.46
7.44
7.43
7.42
7.41
7.39
7.38
7.37
7.36
7.35
7.34
7.33
7.32
7.31
7.30
7.29





























S13






S14


































LC-Mass spectra of 3g

¹³C NMR spectra of **3h**



HSQC spectra of **3h**