

## Research Article

# Determining the $IC_{50}$ Values for Vorozole and Letrozole, on a Series of Human Liver Cytochrome P450s, to Help Determine the Binding Site of Vorozole in the Liver

Lendelle Raymond, Nikita Rayani, Grace Polson, Kylie Sikorski, Ailin Lian, and Melissa A. VanAlstine-Parris

Chemistry Department, Adelphi University, Garden City, NY 11530, USA

Correspondence should be addressed to Melissa A. VanAlstine-Parris; [mvanalstine@adelphi.edu](mailto:mvanalstine@adelphi.edu)

Received 3 July 2015; Accepted 26 October 2015

Academic Editor: Hartmut Kuhn

Copyright © 2015 Lendelle Raymond et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Vorozole and letrozole are third-generation aromatase (cytochrome P450 19A1) inhibitors. [ $^{11}C$ ]-Vorozole can be used as a radiotracer for aromatase in living animals but when administered by IV, it collects in the liver. Pretreatment with letrozole does not affect the binding of vorozole in the liver. In search of finding the protein responsible for the accumulation of vorozole in the liver, fluorometric high-throughput screening assays were used to test the inhibitory capability of vorozole and letrozole on a series of liver cytochrome P450s (CYP1A1, CYP1A2, CYP2A6, and CYP3A4). It was determined that vorozole is a potent inhibitor of CYP1A1 ( $IC_{50} = 0.469 \mu M$ ) and a moderate inhibitor of CYP2A6 and CYP3A4 ( $IC_{50} = 24.4$  and  $98.1 \mu M$ , resp.). Letrozole is only a moderate inhibitor of CYP1A1 and CYP2A6 ( $IC_{50} = 69.8$  and  $106 \mu M$ ) and a very weak inhibitor of CYP3A4 (<10% inhibition at 1 mM). Since CYP3A4 makes up the majority of the CYP content found in the human liver, and vorozole inhibits it moderately well but letrozole does not, CYP3A4 is a good candidate for the protein that [ $^{11}C$ ]-vorozole is binding to in the liver.

## 1. Introduction

Vorozole and letrozole (Figure 1) are nonsteroidal, triazole-containing compounds that are competitive, reversible, third-generation aromatase (CYP19A1) inhibitors [1, 2]. Both vorozole and letrozole were initially developed and underwent clinical trials as antineoplastic agents [3]. Letrozole (Femara) is currently used in the treatment of breast cancer [4] but vorozole was not pushed forward after phase III clinical trials because it did not show any significant advantage over the current drugs [5–7]. However, vorozole, labeled with  $^{11}C$ , is currently being used as a tracer for positron emission topography (PET) imaging to study CYP19A1 distribution in living animals [8–13].

[ $^{11}C$ ]-Vorozole has been shown to display high and specific binding *in vitro* to CYP19A1-rich human placenta [14], human granulosa cells [15], and rat brains [16]. *In vivo* studies have shown that [ $^{11}C$ ]-vorozole binds regionally specifically to CYP19A1 in rhesus monkey [16], baboon [11, 17], and

human [8] brain. The brain accumulation has been shown to be specific by being blocked by both vorozole and letrozole [8, 16, 17]. However, when [ $^{11}C$ ]-vorozole is administered to rats [10], rhesus monkeys [14], baboons [17], or humans [18] by IV, some of it binds to the liver. It has been shown that this binding in the baboon and human liver is not caused by CYP19A1 because pretreatment with letrozole does not block its binding [18, 19]. While vorozole has been shown to be selective against other cytochrome P450s- (CYP-) dependent reactions in steroid biosynthesis [6], there is limited data available on other CYPs, especially those found in the liver.

Cytochrome P450s are heme containing monooxygenases responsible for oxidative metabolism of more than 95% of pharmaceutical drugs in the human liver. There are a number of xenobiotic metabolizing CYPs that are expressed in a typical human liver and the top CYP isoforms that contribute to the metabolism of small molecule drugs are CYP3A, CYP2C, CYP1A2, CYP2A6, and CYP2E1 [20]. Many imidazole and triazole ring-containing inhibitors of CYPs

TABLE 1: Fluorometric enzyme assay conditions for recombinant cytochrome P450s (CYPs). 3-Cyano-7-ethoxycoumarin (CEC), 3-cyano-7-hydroxycoumarin (CHC), 7-hydroxycoumarin (HC), 7-benzyloxy-4-(trifluoromethyl)coumarin (BFC), 7-hydroxy-4-(trifluoromethyl)coumarin (HFC), 7-methoxy-4-(trifluoromethyl)coumarin (MFC), excitation wavelength ( $\lambda_{Ex}$ ), and emission wavelength ( $\lambda_{Em}$ ).

Enzyme	Enzyme amount (pmol)	Substrate	Substrate concentration ( $\mu$ M)	Product	$\lambda_{Ex}$ (nm)	$\lambda_{Em}$ (nm)
CYP1A1	1	CEC	5	CHC	410	450
CYP1A2	0.5	CEC	5	CHC	410	450
CYP2A6	1	Coumarin	3	HC	330	460
CYP3A4	1	BFC	50	HFC	409	530
CYP19A1	1.5	MFC	100	HFC	409	530




FIGURE 1: Structures of vorozole and letrozole.

form a noncovalent ligand interaction with the ferric ion heme and therefore have the potential to inhibit multiple isoforms [21]. Vorozole and letrozole are both triazole-containing compounds so it is likely that they will bind to other CYPs besides CYP19A1.

Since vorozole has been shown to bind to the liver and pretreatment with letrozole does not block this binding, by determining and comparing the binding affinity of both vorozole and letrozole on a series of liver CYPs, we can potentially identify the protein that is responsible for vorozole binding in the liver. This CYP can be identified by having a high binding affinity to vorozole but not letrozole. Fluorometric high-throughput screening (HTS) assays for CYPs have been developed for 13 recombinant human CYPs [22]. These assays use nonnatural coumarin substrates that are converted into fluorescent products by the CYPs. These fluorometric HTS assays can be used to determine the  $IC_{50}$  values of vorozole and letrozole on human liver CYPs. By comparing the potency of vorozole and letrozole (as a negative control) on human liver CYPs we can have a better idea of the CYP responsible for vorozole's accumulation in the liver.

## 2. Materials and Methods

**2.1. Materials.** Coumarin, glucose-6-phosphate dehydrogenase, 7-methoxy-4-(trifluoromethyl)coumarin (MFC), 7-hydroxy-4-(trifluoromethyl)coumarin (HFC), magnesium chloride ( $MgCl_2$ ), and nicotine adenine dinucleotide phosphate ( $NADP^+$ ) were purchased from Sigma Aldrich (St. Louis, MO). 3-Cyano-7-ethoxycoumarin (CEC), 7-benzyloxy-4-(trifluoromethyl)coumarin (BFC), and all recombinant microsomes from baculovirus-infected insect cells (supersomes) were purchased from BD Bioscience (Woburn, MA). 3-Cyano-7-hydroxycoumarin (CHC) was purchased from Indofine Chemical Company. Potassium phosphate dibasic ( $K_2HPO_4$ ) was obtained from Merck. Potassium phosphate

TABLE 2:  $IC_{50}$  values ( $\pm$ SEM) for vorozole and letrozole on CYP1A1, CYP1A2, CYP2A6, CYP3A4, and CYP19A1 for 3 replicate assays.

	Vorozole ( $\mu$ M)	Letrozole ( $\mu$ M)	Fold difference
CYP1A1	0.469 $\pm$ 0.031	69.8 $\pm$ 5.3	149
CYP1A2	321 $\pm$ 49	332 $\pm$ 195	1.03
CYP2A6	24.4 $\pm$ 3.6	106 $\pm$ 34	4.34
CYP3A4	98.1 $\pm$ 15.6	<10% @ 1mM	>10
CYP19A1	0.00417 $\pm$ 0.00019	0.00727 $\pm$ 0.00070	1.73

monobasic ( $KH_2PO_4$ ), D-glucose 6-phosphate sodium salt, 7-hydroxycoumarin (HC), and dimethyl sulfoxide (DMSO) were purchased from Fisher Scientific Company. Vorozole and letrozole were provided by Brookhaven National Laboratory. All experiments were completed in all black, flat bottom Costar 96-well plates (Corning Incorporated, Corning, NY).

**2.2. Enzyme Assay.**  $IC_{50}$  determinations for human CYPs were similar to HTS methods described by Crespi et al. [23]. Unless otherwise stated, the incubations were carried out in a total volume of 200  $\mu$ L of 50 mM potassium phosphate buffer (pH 7.4) with 1% acetonitrile. The reaction mixtures contained NADPH-regenerating system in potassium phosphate buffer (pH 7.4) and varying amounts of inhibitor (dissolved in 100% acetonitrile) were preincubated at 37°C for 10 minutes (except CYP2A6 which was in 100 mM Tris buffer pH 7.4). The reactions were initiated by the addition of enzyme/substrate mixture (see Table 1), followed by incubation at 37°C for 30 minutes. The fluorescent signal was measured using a SpectraMax GEMINI XPS (Molecular Devices) with excitation and emission wavelengths listed in Table 1. The data were fit to sigmoidal dose-response curves with nonlinear regression and  $IC_{50}$  values calculated using GraphPad Prism 5.  $IC_{50}$  values were converted to  $K_i$  values using the Cheng-Prusoff equation ( $K_i = IC_{50}/(1 + [S]/K_m)$ ) and literature  $K_m$  values when available.

## 3. Results and Discussion

$IC_{50}$  values for vorozole and letrozole on CYP1A1, CYP1A2, CYP2A6, CYP3A4, and CYP19A1 are given in Table 2. Vorozole and letrozole are potent inhibitors of CYP19A1 ( $IC_{50} = 4.17$  and  $7.27$  nM, resp.). Vorozole is a moderate inhibitor of CYP1A1 ( $IC_{50} = 0.469$   $\mu$ M) and a weak inhibitor of CYP1A2, CYP2A6, and CYP3A4 ( $IC_{50} = 321$ ,  $24.4$ , and  $98.1$   $\mu$ M, resp.). Letrozole is a weak inhibitor of CYP1A1, CYP1A2, and

CYP2A6 ( $IC_{50} = 69.8, 332, \text{ and } 106 \mu\text{M}$ , resp.). Letrozole was an extremely weak inhibitor of CYP3A4 (<10% inhibition at 1 mM).

[ $^{11}\text{C}$ ]-Vorzole is currently being used as a radiotracer for CYP19A1 using PET imaging; however when it is administered by IV, some of it binds to the liver but not through CYP19A1. Since vorzole contains a triazole ring and binds to the heme of CYP19A1, it is likely that vorzole can bind to the heme of other CYPs. Since there is a high concentration of CYP protein in the liver, it is likely that vorzole is binding to a CYP other than CYP19A1 in the liver. Letrozole is another potent CYP19A1 inhibitor but it does not block [ $^{11}\text{C}$ ]-vorzole's binding in the liver. Therefore we can use letrozole as our negative control when searching for the protein responsible for vorzole accumulation in the liver.

To show that our assays are valid, we confirmed that vorzole and letrozole are very potent inhibitors of CYP19A1. We used the  $K_m$  value of  $25 \mu\text{M}$  from the BD Bioscience CYP19A1/MFC kit, to convert our  $IC_{50}$  values (4.17 and 7.27 nM) to  $K_i$  values of 0.9 and 1.6 nM for vorzole and letrozole, respectively. While there are no literature values for vorzole and letrozole on CYP19A1 with the synthetic substrate MFC, our  $K_i$  values are in agreement with the literature  $K_i$  values with the natural steroid substrates (testosterone and androstenedione) of around 1 nM for vorzole [24–29] and 2 nM for letrozole [30].

The  $IC_{50}$  values for vorzole and letrozole were then compared on each CYP. It was found that vorzole and letrozole bind equally poor to CYP1A2 (1.03-fold difference) with  $IC_{50}$  values in the hundred micromolar range. While vorzole is slightly more potent than letrozole on CYP2A6 (4.34-fold difference), they are both still weak inhibitors. Therefore both CYP1A2 and CYP2A6 are not likely candidates for the protein that is causing the accumulation of vorzole in the liver.

On CYP1A1, vorzole was almost 150-fold more potent than letrozole with an  $IC_{50}$  of  $0.47 \mu\text{M}$ . However, there is conflicting evidence as to whether CYP1A1 is expressed in the human liver [31]. Even if CYP1A1 is expressed, it is in such small quantities [32] that it is an unlikely candidate.

On CYP3A4, 1 mM letrozole inhibited the reaction by less than 10% so we could not calculate an  $IC_{50}$  value. However, vorzole inhibited CYP3A4 with an  $IC_{50}$  of  $98.1 \mu\text{M}$ . Therefore, there is at least a 10-fold difference in the potency of vorzole and letrozole on CYP3A4. This data, combined with the fact that CYP3A4 makes up the majority of the CYP content found in the liver [20], implies that CYP3A4 is a good candidate for the enzyme that [ $^{11}\text{C}$ ]-vorzole is binding to in the liver but letrozole does not block.

The inhibition potential of vorzole and letrozole on some other liver CYPs has previously been reported in the literature. Letrozole was not a potent inhibitor of CYP1B1 with an  $IC_{50} \geq 100 \mu\text{M}$  for estradiol 4-hydroxylation and 2-hydroxylation. However, vorzole was shown to inhibit CYP1B1 activity with  $IC_{50}$  values of 17 and  $21 \mu\text{M}$  for 4-hydroxy estradiol and 2-hydroxy estradiol [33]. Since there is at least a 5-fold difference in potency between vorzole and letrozole and vorzole inhibits CYP1B1 moderately well, CYP1B1 should not be overlooked.

With this information, our collaborators at Brookhaven National Laboratories used a CYP3A4 inhibitor (ketoconazole) to see if it would block the binding of [ $^{11}\text{C}$ ]-vorzole to the liver. When [ $^{11}\text{C}$ ]-vorzole was given with a pretreatment of ketoconazole, the liver pharmacokinetics of [ $^{11}\text{C}$ ]-vorzole binding was altered [19]. While ketoconazole is typically thought of as a selective CYP3A4 inhibitor [34], it has also been shown to inhibit CYP1A1 [35]. This data further confirms that CYP3A4 may be responsible for [ $^{11}\text{C}$ ]-vorzole accumulation in the liver but CYP1A1 cannot be completely ruled out.

## 4. Conclusion

In conclusion, this data shows that vorzole is at least 10-fold more potent than letrozole on CYP3A4 and vorzole inhibits CYP3A4 with an  $IC_{50}$  value of  $98 \mu\text{M}$ . Therefore CYP3A4 is a likely candidate for the protein responsible for binding vorzole in the liver. Pretreatment with ketoconazole, a CYP3A4 inhibitor, affects liver pharmacokinetics of [ $^{11}\text{C}$ ]-vorzole binding, further confirming that CYP3A4 may be responsible for [ $^{11}\text{C}$ ]-vorzole accumulation in the liver.

## Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.




## Acknowledgments

Vorzole and letrozole were kindly supplied by Dr. Joanna Fowler at Brookhaven National Laboratories. This work was carried out at Adelphi University and was supported by an Adelphi University Faculty Development Grant and, in part, by the Horace McDonell Fellowship for summer research for an undergraduate student.

## References

- [1] A. S. Bhatnagar, A. Häusler, K. Schieweck, M. Lang, and R. Bowman, "Highly selective inhibition of estrogen biosynthesis by CGS 20267, a new non-steroidal aromatase inhibitor," *Journal of Steroid Biochemistry and Molecular Biology*, vol. 37, no. 6, pp. 1021–1027, 1990.
- [2] W. Wouters, R. De Coster, R. W. Tuman et al., "Aromatase inhibition by R 76713: experimental and clinical pharmacology," *Journal of Steroid Biochemistry*, vol. 34, no. 1–6, pp. 427–430, 1989.
- [3] R. De Coster, R. F. Van Ginckel, M. J. Callens, N. K. Goeminne, and B. L. Janssens, "Antitumoral and endocrine effects of (+)-vorzole in rats bearing dimethylbenzanthracene-induced mammary tumors," *Cancer Research*, vol. 52, no. 5, pp. 1240–1244, 1992.
- [4] S. Jeong, M. M. Woo, D. A. Flockhart, and Z. Desta, "Inhibition of drug metabolizing cytochrome P450s by the aromatase inhibitor drug letrozole and its major oxidative metabolite 4,4'-methanol-bisbenzotriazole in vitro," *Cancer Chemotherapy and Pharmacology*, vol. 64, no. 5, pp. 867–875, 2009.

- [5] P. M. Wood, L. W. L. Woo, M. P. Thomas, M. F. Mahon, A. Purohit, and B. V. L. Potter, "Aromatase and dual aromatase-steroid sulfatase inhibitors from the letrozole and vorozole templates," *ChemMedChem*, vol. 6, no. 8, pp. 1423–1438, 2011.
- [6] P. E. Goss, "Pre-clinical and clinical review of vorozole, a new third generation aromatase inhibitor," *Breast Cancer Research and Treatment*, vol. 49, no. 1, pp. S59–S65, 1998.
- [7] A. Hamilton and M. Piccart, "The third-generation non-steroidal aromatase inhibitors: a review of their clinical benefits in the second-line hormonal treatment of advanced breast cancer," *Annals of Oncology*, vol. 10, no. 4, pp. 377–384, 1999.
- [8] A. Biegon, S. W. Kim, D. L. Alexoff et al., "Unique distribution of aromatase in the human brain: in vivo studies with PET and [ $N$ -methyl- $^{11}$ C]vorozole," *Synapse*, vol. 64, no. 11, pp. 801–807, 2010.
- [9] A. Biegon, N. Alia-Klein, and J. S. Fowler, "Potential contribution of aromatase inhibition to the effects of nicotine and related compounds on the brain," *Frontiers in Pharmacology*, vol. 3, article 185, Article ID Article 185, 2012.
- [10] M. Ozawa, K. Takahashi, K.-H. Akazawa et al., "PET of aromatase in gastric parietal cells using  $^{11}$ C-vorozole," *Journal of Nuclear Medicine*, vol. 52, no. 12, pp. 1964–1969, 2011.
- [11] A. Biegon, S.-W. Kim, J. Logan, J. M. Hooker, L. Muench, and J. S. Fowler, "Nicotine blocks brain estrogen synthase (aromatase): in vivo positron emission tomography studies in female baboons," *Biological Psychiatry*, vol. 67, no. 8, pp. 774–777, 2010.
- [12] K. Takahashi, K. Onoe, H. Doi et al., "Increase in hypothalamic aromatase in macaque monkeys treated with anabolic-androgenic steroids: PET study with [ $^{11}$ C]vorozole," *Neuroreport*, vol. 22, no. 7, pp. 326–330, 2011.
- [13] K. Takahashi, G. Yamagishi, T. Hiramatsu et al., "Practical synthesis of precursor of [ $N$ -methyl- $^{11}$ C]vorozole, an efficient PET tracer targeting aromatase in the brain," *Bioorganic and Medicinal Chemistry*, vol. 19, no. 4, pp. 1464–1470, 2011.
- [14] P. Lidström, T. A. Bonasera, D. Kirilovas et al., "Synthesis, *in vivo* rhesus monkey biodistribution and *in vitro* evaluation of a  $^{11}$ C-labelled potent aromatase inhibitor: [ $N$ -methyl- $^{11}$ C]vorozole," *Nuclear Medicine and Biology*, vol. 25, no. 5, pp. 497–501, 1998.
- [15] D. Kirilovas, M. Bergström, T. A. Bonasera et al., "In vitro evaluation of aromatase enzyme in granulosa cells using a [ $^{11}$ C]vorozole binding assay," *Steroids*, vol. 64, no. 4, pp. 266–272, 1999.
- [16] K. Takahashi, M. Bergström, P. Frändberg, E.-L. Vesström, Y. Watanabe, and B. Långström, "Imaging of aromatase distribution in rat and rhesus monkey brains with [ $^{11}$ C]vorozole," *Nuclear Medicine and Biology*, vol. 33, no. 5, pp. 599–605, 2006.
- [17] S. W. Kim, A. Biegon, Z. E. Katsamanis et al., "Reinvestigation of the synthesis and evaluation of [ $N$ -methyl- $^{11}$ C]vorozole, a radiotracer targeting cytochrome P450 aromatase," *Nuclear Medicine and Biology*, vol. 36, no. 3, pp. 323–334, 2009.
- [18] A. Biegon, D. L. Alexoff, S. W. Kim et al., "Aromatase imaging with [ $N$ -methyl- $^{11}$ C]vorozole PET in healthy men and women," *Journal of Nuclear Medicine*, vol. 56, no. 4, pp. 580–585, 2015.
- [19] S. W. Kim, A. Biegon, J. M. Hooker et al., "Saturable, non-aromatase binding of [ $^{11}$ C]vorozole in baboon liver," *Journal of Nuclear Medicine*, vol. 52, article 415, 416 pages, 2011.
- [20] S. C. Khojasteh, S. Prabhu, J. R. Kenny, J. S. Halladay, and A. Y. H. Lu, "Chemical inhibitors of cytochrome P450 isoforms in human liver microsomes: a re-evaluation of P450 isoform selectivity," *European Journal of Drug Metabolism and Pharmacokinetics*, vol. 36, no. 1, pp. 1–16, 2011.
- [21] C. Tang, M. Chiba, J. Nishime et al., "Comparison of imidazole- and 2-methyl imidazole-containing farnesyl-protein transferase inhibitors: interaction with and metabolism by rat hepatic cytochrome P450s," *Drug Metabolism and Disposition*, vol. 28, no. 6, pp. 680–686, 2000.
- [22] A. Ghosal, N. Hapangama, Y. Yuan et al., "Rapid determination of enzyme activities of recombinant human cytochromes P450, human liver microsomes and hepatocytes," *Biopharmaceutics and Drug Disposition*, vol. 24, no. 9, pp. 375–384, 2003.
- [23] C. L. Crespi, V. P. Miller, and B. W. Penman, "Microtiter plate assays for inhibition of human, drug-metabolizing cytochromes P450," *Analytical Biochemistry*, vol. 248, no. 1, pp. 188–190, 1997.
- [24] Y.-C. Kao, K. R. Korzekwa, C. A. Loughton, and S. Chen, "Evaluation of the mechanism of aromatase cytochrome p450," *European Journal of Biochemistry*, vol. 268, no. 2, pp. 243–251, 2001.
- [25] H. Vanden Bossche, G. Willemsens, I. Roels et al., "R 76713, A new specific non-steroidal aromatase inhibitor," *Biochemical Pharmacology*, vol. 40, pp. 1707–1718, 1990.
- [26] M. D. W. G. Krekels, W. Wouters, and R. De Coster, "Aromatase inhibition by R 76 713: a kinetic analysis in rat ovarian homogenates," *Steroids*, vol. 55, no. 2, pp. 69–73, 1990.
- [27] M. D. W. G. Krekels, W. Wouters, R. De Coster, R. Van Ginckel, A. Leonaers, and P. A. J. Janssen, "Aromatase in the human choriocarcinoma JEG-3: inhibition by R 76 713 in cultured cells and in tumors grown in nude mice," *Journal of Steroid Biochemistry and Molecular Biology*, vol. 38, no. 4, pp. 415–422, 1991.
- [28] H. V. Bossche, G. Willemsens, I. Roels et al., "R 76713 and enantiomers: selective, nonsteroidal inhibitors of the cytochrome P450-dependent oestrogen synthesis," *Biochemical Pharmacology*, vol. 40, no. 8, pp. 1707–1718, 1990.
- [29] W. Wouters, R. De Coster, D. Beerens et al., "Potency and selectivity of the aromatase inhibitor R76,713. A study in human ovarian, adipose stromal, testicular and adrenal cells," *Journal of Steroid Biochemistry*, vol. 36, no. 1-2, pp. 57–65, 1990.
- [30] Q. Lu, Y. Liu, B. J. Long, D. Grigoryev, M. Gimbel, and A. Brodic, "The effect of combining aromatase inhibitors with antiestrogens on tumor growth in a nude mouse model for breast cancer," *Breast Cancer Research and Treatment*, vol. 57, no. 2, pp. 183–192, 1999.
- [31] A. T. Drahushuk, B. P. McGarrigle, K. E. Larsen, J. J. Stegeman, and J. R. Olson, "Detection of CYP1A1 protein in human liver and induction by TCDD in precision-cut liver slices incubated in dynamic organ culture," *Carcinogenesis*, vol. 19, no. 8, pp. 1361–1368, 1998.
- [32] W. M. A. Westerink, J. C. R. Stevenson, and W. G. E. J. Schoonen, "Pharmacologic profiling of human and rat cytochrome P450 1A1 and 1A2 induction and competition," *Archives of Toxicology*, vol. 82, no. 12, pp. 909–921, 2008.
- [33] M. Rahman, S. F. Lax, C. H. Sutter et al., "CYP1B1 is not a major determinant of the disposition of aromatase inhibitors in epithelial cells of invasive ductal carcinoma," *Drug Metabolism and Disposition*, vol. 36, no. 5, pp. 963–970, 2008.
- [34] V. A. Eagling, J. F. Tjia, and D. J. Back, "Differential selectivity of cytochrome P450 inhibitors against probe substrates in human and rat liver microsomes," *British Journal of Clinical Pharmacology*, vol. 45, no. 2, pp. 107–114, 1998.
- [35] M. F. Paine, P. Schmiedlin-Ren, and P. B. Watkins, "Cytochrome P-450 1A1 expression in human small bowel: interindividual variation and inhibition by ketoconazole," *Drug Metabolism and Disposition*, vol. 27, no. 3, pp. 360–364, 1999.



**Hindawi**

Submit your manuscripts at  
<http://www.hindawi.com>

