

Research Article

Establishment of MDCK Stable Cell Lines Expressing TMPRSS2 and MSPL and Their Applications in Propagating Influenza Vaccine Viruses in Absence of Exogenous Trypsin

Zhiyuan Wen, Chao Wu, Weiye Chen, Xianying Zeng, Jianzhong Shi, Jinying Ge, Hualan Chen, and Zhigao Bu

State Key Laboratory of Veterinary Biotechnology, Harbin Veterinary Research Institute, Chinese Academy of Agricultural Sciences, 427 Maduan Street, Harbin 150001, China

Correspondence should be addressed to Zhigao Bu; buzhigao@caas.cn

Received 6 January 2015; Revised 19 March 2015; Accepted 20 March 2015

Academic Editor: Manuel Canovas

Copyright © 2015 Zhiyuan Wen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We established two Madin-Darby canine kidney (MDCK) cell lines stably expressing human airway transmembrane protease: transmembrane protease, serine 2 (TMPRSS2) and mosaic serine protease large form (MSPL) which support multicycle growth of two H5 highly pathogenic avian influenza viruses (HPAIV) recombinant vaccines (Re-5 and Re-6) and an H9 avian influenza virus (AIV) recombinant vaccine (Re-9) in the absence of trypsin. Data showed that the cell lines stably expressed TMPRSS2 and MSPL after 20 serial passages. Both MDCK-TMPRSS2 and MDCK-MSPL could proteolytically cleave the HA of Re-5, Re-6, and Re-9 and supported high-titer growth of the vaccine without exogenous trypsin. Re-5, Re-6, and Re-9 efficiently infected and replicated within MDCK-TMPRSS2 and MDCK-MSPL cells and viral titer were comparable to the virus grown in MDCK cells with TPCK-trypsin. Thus, our results indicate a potential application for these cell lines in cell-based influenza vaccine production and may serve as a useful tool for HA proteolytic cleavage-related studies.

1. Introduction

Influenza is a major zoonotic threat to public health, which is caused by 3 types (A, B, and C) of influenza viruses [1, 2]. Type A influenza is the most serious type, specifically the highly pathogenic H5N1 [3–5], H1N1 [5–7], and the newly emerged lethal H7N9 [8, 9].

Hemagglutinin (HA) of influenza virus mediates both receptor binding and membrane fusion [10]. HA cleavage is important for viral infectivity; HA proteins are synthesized as HA0 precursor proteins during transport through the Golgi apparatus. HA0 is cleaved by host cell protease into HA1 and HA2 subunits [11, 12]. Cleaved HA proteins bind to cell receptor and then are endocytosed into the endosome where they undergo conformational changes and exposure of fusion peptide on HA2 subunit under low pH. Then, the fusion peptide

is inserted into the cell membrane and mediates the formation of fusion pore [13, 14]. Fusion relies on precise HA0 cleavage for a fusion-capable HA2 subunit. HA proteins of H5 highly pathogenic avian influenza viruses (HPAIV) have multibasic cleavage sites (R-X-R/K-R) which can be cleaved by ubiquitously expressed furin or PC5/6 protease to cause fatal systemic infections [15–17]. HA of most of the other mammalian and avian influenza viruses contains a single arginine (or lysine) at the cleavage site, so cleavage of these HAs is restricted to the respiratory tract in mammals and to the respiratory and intestinal tracts in avians and assumed to be processed extracellularly by trypsin-like proteases. Of these proteases, some type II transmembrane serine proteases (TTSPs) family members such as human airway trypsin-like (HAT) protease, transmembrane protease, serine 2 (TMPRSS2), transmembrane protease, serine 4 (TMPRSS4),

and mosaic serine protease large form (MSPL) play important roles in influenza viral infection. TTSPs are expressed in the airways and can cleave multiple strains of influenza HA protein. Böttcher and colleagues reported a cell-associated cleavage of influenza viruses HA with a monobasic cleavage site by HAT and TMPRSS2 [18]. Wang and colleagues reported that TMPRSS2 and HAT could cleave the HA of the H1, H3, and H5 subtypes [19]. Zmora and colleagues demonstrated that mosaic serine protease large form (MSPL) could activate HA protein of H1N1 and H3N2 influenza virus [20], while Okumura and colleagues confirmed that MSPL can cleave the HA protein of H5 HPAIV and support their multicycle replication [21].

Here, we established two MDCK cell lines that stably express TMPRSS2 and MSPL. Western blot and RT-PCR confirmed the presence of the target gene; FACS assay confirmed target gene expression in serially passaged cells. Cell fusion assay indicated that TMPRSS2 and MSPL cell lines could cleave the HA protein of H5 and H9 subtypes. Both cell lines can support multicycle growth of Re-5, Re-6, and Re-9 in absence of exogenous trypsin. Vaccine titers of these cell lines were comparable to those in MDCK cells plus TPCK-trypsin.

2. Materials and Methods

2.1. Viruses and Cells. Low-passage Madin-Darby canine kidney (MDCK) cells were maintained in DMEM containing 10% fetal bovine serum (FBS). Influenza viruses Re-5 [22], Re-6 [23], and Re-9 were provided by the National Animal Influenza Reference Laboratory. Viruses were generated with a "6 + 2" strategy: all three viruses contained 6 internal genes from A/Puerto Rico/8/1934 (H1N1). HA and NA genes of Re-5 were from A/Duck/Anhui/1/2005 (H5N1); HA and NA genes of Re-6 were from A/Duck/Guangdong/s1322/2010 (H5N1); and HA and NA genes of Re-9 were from A/Chicken/Hunan/S933/2008 (H9N2). To enhance safety, the multibasic amino acid cleavage site of the HA protein of Re-5 (RRRRKR) and Re-6 (RERRRKR) was changed to monobasic amino acids (RETR).

2.2. Generation of MDCK-TMPRSS2 and MDCK-MSPL Stable Cell Lines. Human TMPRSS2 (GenBank number U75329.1) and human MSPL (GenBank number AB048796.1) genes were synthesized by Genaray Biotech (Shanghai, China), and both genes were fused to a Flag tag (DYKDDDDK) at the 3' end of the ORF. Eukaryotic expression vector P4 was used to obtain the stable cell lines. P4 was derived from the pCAGGS vector and this was modified by inserting an enhanced GFP (eGFP) gene and G418-resistant gene (*neo*) to render it suitable for establishing a stable cell line. Either TMPRSS2-FLAG or MSPL-FLAG was inserted into the P4 vector multiple cloning site which is located upstream of the IRES and eGFP genes, allowing the inserted genes to be detected with FLAG and green fluorescence. The resultant plasmids, P4-TMPRSS2-FLAG and P4-MSPL-FLAG, were prepared and purified by Qiagen Maxi Prep plasmid kit (Qiagen, Valencia, CA) before transfection.

Newly recovered, low-passage MDCK cells were serially passaged 5 times. Cells were then electrotransfected following

the protocol of Nucleofector Kit for MDCK cells (Lonza, Cologne, Germany). Briefly, 5×10^5 MDCK cells were resuspended in 0.1 mL Solution I (provided by the kit). 5 μ g plasmid (P4-TMPRSS2-FLAG or P4-MSPL-FLAG) was added to Solution I within 15 min. The cell-plasmid mix was transferred to a specialized cuvette which was then loaded to an Amaxa Nucleofector II machine (Lonza, Cologne, Germany) for electrotransfection using the A24 program. After electrotransfection, cells were transferred to 6-well plates containing DMEM with 10% FBS. 48 h later, attached monolayer cells were trypsinized, diluted 1:100, and seeded into 24-well plates containing DMEM with 5% FBS and 800 μ g/mL G418. Medium was replaced every 3 days and cultured for 2 weeks after which G418-resistant cell colonies were collected and transferred to new plates. Cells were then trypsinized and loaded to a BD FACSAria cell sorter; single green fluorescence positive cells were sorted to 96-well plates and cultured under G418 pressure. Two weeks later, surviving cell colonies were transferred to 48-well plates and continued to be cultured in presence of G418. Cells were then scaled up and cultured for at least 10 passages to generate MDCK-TMPRSS2 and MDCK-MSPL cells.

2.3. Verification of TMPRSS2 and MSPL Expression in Cell Lines. For RT-PCR, total RNA of 5×10^6 MDCK-TMPRSS2 or MDCK-MSPL cells was extracted with a Qiagen RNeasy Mini Kit (Qiagen, Valencia, CA). RNA was reverse transcribed with a Promega Improm II RT kit (Promega, Madison, WI). The following primers were used to amplify TMPRSS2 or MSPL with PrimeStar HS polymerase (Takara, Dalian, China): TMPRSS2-F 5'-ATGGCTTTGAACTCAG-3' and TMPRSS2-R 5'-TTACTTATCGTCGTCATC-3'; MSPL-F: 5'-ATGGAGAGGGACAGCCAC-3' and MSPL-R 5'-TTACTTATCGTCGTCATC-3'.

For Western blot, MDCK-TMPRSS2, MDCK-MSPL, or MDCK cells were washed with PBS, lysed with RIPA buffer (containing 25 mM Tris-HCl pH 7.6, 150 mM NaCl, 1% NP-40, 1% desoxycholic acid sodium salt, and 0.1% SDS) (Thermo Fisher, Waltham, MA), and mixed with reducing protein loading buffer. Samples were separated with SDS-PAGE and subsequently transferred to a nitrocellulose (NC) membrane (GE Healthcare, Pittsburgh, PA). NC membranes were incubated with 1:100 diluted mouse anti-FLAG antibody (Abcam, Cambridge, MA) as primary antibody and 1:4000 diluted HRP labeled goat anti-mouse IgG (Sigma, St. Louis, MO) as secondary antibody. NC membranes were then incubated with ECL plus reagent (GE Healthcare, Pittsburgh, PA) and protein bands were visualized on a Kodak XAR films (Kodak).

For FACS assay, 2×10^6 of the 10th serially passaged MDCK-TMPRSS2 and MDCK-MSPL cells were collected and washed with PBS and then filtered through a 40- μ m cell strainer (BD Biosciences, San Jose, CA). Cells were then loaded to a BD FACSAria machine to measure green fluorescence. Data were analyzed with FlowJo (Treestar Inc., Ashland, OR) software and expressed as histogram.

2.4. Cell Fusion Assay. MDCK cells were seeded in 6-well plates, and 1 μ g of pCAGGS-TMPRSS2-FLAG, pCAGGS-MSPL-FLAG, and pCAGGS was cotransfected with 1 μ g of pCAGGS-Re-5HA, pCAGGS-Re-6HA, and pCAGGS-Re-9HA, respectively, using Lipofectamine 3000 reagent (Invitrogen, Carlsbad, CA). 48 h after transfection, cells were treated with acidic PBS (pH 5.0) at room temperature for 5 min and replenished with DMEM containing 5% FBS and cultured at 37°C for 1 h. When fusion was obvious, cells were fixed with 3% paraformaldehyde and stained with mouse anti-FLAG antibody and chicken anti-Re-5, anti-Re-6, or anti-Re-9 as primary antibody with FITC-labeled goat anti-mouse antibody (Abcam, Cambridge, CA) and TRITC-labeled rabbit anti-chicken antibody (Abcam, Cambridge, CA) for 30 min. Cells were washed and stained with DAPI and images were acquired with a Leica TCS SP5 confocal laser microscope (Leica Microsystems, Wetzlar, Germany).

2.5. Immunofluorescence Assay. MDCK-TMPRSS2, MDCK-MSPL, and MDCK cells were seeded in 12-well plates and grown in DMEM containing 5% FBS for 24 h. Cells were infected with Re-5, Re-6, and Re-9 at MOI = 0.01 for 1 h at 37°C. Inoculum was removed and cells were washed 3 times and placed in DMEM with 5% FBS. 36 h after incubation, cells were washed with PBS and fixed with 3% paraformaldehyde at room temperature for 15 min. Infected cells were detected with chicken anti-Re-5, anti-Re-6, and anti-Re-9 as primary antibodies and TRITC-labeled goat anti-chicken IgG (Sigma, St. Louis, MO) as secondary antibody. Stained cells were observed under a Zeiss Axioscop (Zeiss, Thornwood, NY) fluorescent microscope.

2.6. Viral Titration. MDCK-TMPRSS2, MDCK-MSPL, and MDCK cells were seeded in 6-well plates and grown in DMEM containing 5% FBS for 24 h. Cells were infected with Re-5, Re-6, and Re-9 at MOI = 0.01 in DMEM for 1 h at 37°C. Inoculum was removed and cells were washed 3 times after which DMEM with 5% FBS was added. TPCK-treated trypsin (1 μ g/mL) was added to media of infected MDCK cells. 100 μ L supernatant from each infected cell culture was sampled at 12, 24, 48, 72, and 96 h after infection. Sample TCID₅₀ values were determined, and growth kinetics for each virus from each cell line were obtained. Statistical analysis (*t*-test) was carried out by Excel software (Microsoft, Redmond, WA) for each virus between MDCK-TMPRSS2, MDCK-MSPL, MDCK + TPCK, and MDCK infection groups at the same time point.

3. Results

3.1. Stable Expression of TMPRSS2 and MSPL in MDCK Cells. MDCK-TMPRSS2 and MDCK-MSPL cells were established by electrotransfection of plasmids encoding either TMPRSS2 or MSPL. After transfection, cells were serially passaged in presence of 800 μ g/mL G418. Surviving cells were cultured and scaled up for experiments. To detect TMPRSS2 and MSPL expression, a FLAG tag was in-frame fused to the C-terminus of the TMPRSS2 or MSPL. An eGFP gene was placed upstream of the FLAG-tagged TMPRSS2 or MSPL,

which were linked by an IRES sequence. Thus TMPRSS2 or MSPL expression could be detected with FLAG antibody and green fluorescence. As shown in Figure 1, the morphology of TMPRSS2-MDCK and MSPL-MDCK cells was slightly different from MDCK cells. Under a fluorescent microscope, almost all TMPRSS2-MDCK and MSPL-MDCK cells were positive for green fluorescence except for MDCK cells.

RT-PCR was performed to test for the presence of *TMPRSS2* and *MSPL* genes in cell lines. Figure 2 depicts a specific band amplified from total RNA of MDCK-TMPRSS2 or MDCK-MSPL cells and this confirmed the presence of *TMPRSS2* and *MSPL* genes. To measure expression of TMPRSS2 and MSPL, Western blot and flow cytometry were performed. As shown in Figure 2(b), a 70 kDa band indicated MSPL, and a 55 kDa band indicated TMPRSS2 and no band was detected in MDCK cells. As shown in Figure 2(c), flow cytometry revealed over 99% of TMPRSS2-MDCK and MSPL-MDCK cells to be positive for green fluorescence. Normal MDCK cells were negative. These data confirmed the stable expression of TMPRSS2 and MSPL by the cell lines.

3.2. Proteolytic Cleavage of HA Protein of Re-5, Re-6, and Re-9 by TMPRSS2 and MSPL. To determine whether TMPRSS2 and MSPL could proteolytically activate the HA proteins of Re-5, Re-6, and Re-9, cotransfection and cell fusion assay were carried out. As shown in Figure 3, cell fusion was observed under a confocal microscope 48 h after transfection of TMPRSS2 + Re-5HA, TMPRSS2 + Re-6HA, and TMPRSS2 + Re-9HA, as well as MSPL + Re-5HA, MSPL + Re-6HA, and MSPL + Re-9HA. Cell fusion was also observed in the pCAGGS vector + HAs group in presence of TPCK-trypsin, whereas cell fusion failed in the pCAGGS vector + HAs group in absence of TPCK-trypsin. Fusion of HA relied on precise cleavage of HA and endosome acidification. These data showed that both TMPRSS2 and MSPL could proteolytically activate the HA protein of Re-5, Re-6, and Re-9.

3.3. Re-5, Re-6, and Re-9 Could Infect and Spread within MDCK-TMPRSS2 and MDCK-MSPL Cells in Absence of Exogenous Trypsin. Re-5, Re-6, and Re-9 were used to infect MDCK-TMPRSS2 or MDCK-MSPL cells (MOI = 0.01). 60 h after infection, cells were washed and fixed, and 1:100 diluted chicken anti-Re-5, anti-Re-6, or anti-Re-9 serum was used as primary antibody. 1:200 diluted TRITC-labeled rabbit anti-chicken IgG was used as secondary antibody to detect the virus in infected cells. As shown in Figure 4, Re-5, Re-6, and Re-9 could infect and spread within MDCK-TMPRSS2 and MDCK-MSPL cells in absence of TPCK-trypsin, which was similar to MDCK cells with TPCK-trypsin. These results confirmed that MDCK-TMPRSS2 and MDCK-MSPL cells could proteolytically cleave HA protein of Re-5, Re-6, and Re-9 and support multicycle replication of these viruses in the cell lines.

3.4. MDCK-TMPRSS2 and MDCK-MSPL Cells Support High-Titer Growth of Re-5, Re-6, and Re-9 in Absence of Exogenous Trypsin. Growth kinetics of Re-5, Re-6, and Re-9 on MDCK-TMPRSS2, MDCK-MSPL, and MDCK cells were measured. Growth curves of each virus in different cells were plotted.




FIGURE 1: Morphology of MDCK-TMPRSS2 and MDCK-MSPL cells. MDCK-TMPRSS2, MDCK-MSPL, and MDCK cells were observed under a fluorescent microscope. (a) Cells under a light field; (b) cells under a fluorescent field.

Figure 5 depicts the growth kinetic data. Re-5, Re-6, and Re-9 in MDCK-TMPRSS2 and MDCK-MSPL cells were comparable to those with MDCK plus TPCK-trypsin ($P > 0.05$), while the viral titers in MDCK-TMPRSS2, MDCK-MSPL, and MDCK with TPCK-trypsin were significantly higher than those of MDCK cells without TPCK-trypsin ($P < 0.01$). Thus, MDCK-TMPRSS2 and MDCK-MSPL cells can well support high-titer growth of influenza viruses in absence of exogenous trypsin, suggesting a future application for industrial vaccine production.

4. Discussion

Cleavage of HA by host protease is the prerequisite for influenza viral infection. Most influenza viral HA proteins possess monobasic amino acid cleavage sites. Thus, HA must be proteolytically cleaved into HA1 and HA2 subunits to acquire cell fusion ability and infectivity. Many trypsin-like proteases which can cleave HA protein have been reported (plasmin, trypsin, chymotrypsin, swine mast cell trypsin, chicken embryo trypsin-like protease, and coagulation factor Xa) [24, 25]. These proteases were isolated from tissues and biochemically characterized, but their molecular identity is unknown. Recently, human type II transmembrane serine proteases (TTSPs) have come under study and TTSPs such as HAT,

TMPRSS2, TMPRSS4, and MSPL were found to be capable of cleaving different subtypes of influenza viruses. TMPRSS2 and HAT could cleave the HA protein of H1, H2, H3, and H5 viruses and the 1918 "Spanish flu" virus [18, 19, 26]. It is noteworthy that MSPL can activate different subtypes of influenza virus HA protein possessing either monobasic or multibasic amino acid cleavage site. Zmora and colleagues demonstrated that MSPL can activate the HA protein of H1 and H3 influenza virus as well as the 1918 "Spanish flu" virus [20]. While Kido and colleagues reported that MSPL can cleave an H5 influenza HA-derived multibasic peptide, which was not efficiently cleaved by furin [21, 27], the study of Okumura and colleagues confirmed that MSPL can activate the HA protein of H5 HPAIV and support their multicycle replication [21]. The versatile cleaving ability of MSPL is the major reason we use it to construct the stable cell line.

We have established two stable MDCK cell lines, MDCK-TMPRSS2 and MDCK-MSPL, which could proteolytically activate influenza virus HA and support multicycle growth of the influenza viruses. Re-5, Re-6, and Re-9 were used to produce inactivated commercial vaccines for China's bird flu control and infection. Using the "6 + 2" strategy, the 6 internal genes of PR8 and HA and the NA genes of other influenza viruses were used to generate recombinant viruses. Because PR8-based viruses can propagate to high




FIGURE 2: Stable expression of TMPRSS2 and MSPL in cell lines. (a) RT-PCR of MDCK-TMPRSS2 and MDCK-MSPL cells. Lanes 1 and 3: beta-actin; Lane 2: TMPRSS2; Lane 4: MSPL. (b) Western blot of MDCK-TMPRSS2 and MDCK-MSPL cells. Cell lysates were separated on SDS-PAGE and transferred to nitrocellulose membranes; TMPRSS2-FLAG and MSPL-FLAG fusion proteins were detected with FLAG antibody. (c) Green fluorescence of MDCK-TMPRSS2 and MDCK-MSPL cells was detected by flow cytometry. Both MDCK-TMPRSS2 and MDCK-MSPL cells were positive for green fluorescence.

titer easily in chicken eggs, they are suitable for making inactivated vaccines. To increase the safety of Re-5 and Re-6, the multibasic cleavage sequence was changed to monobasic sequence.

Cell-based vaccines are gaining popularity over egg-based vaccines because cell lines are well characterized and comply with regulatory guidelines. Also, cell culture media are chemically defined and yield consistent results. At present, two cell lines, MDCK cells and Vero cells, are approved and used for influenza vaccine production [28–31]. Cell-based influenza vaccines can provide equivalent protection in animals and humans compared to vaccines produced in chicken eggs [31–33] and MDCK cells have been recognized to be the most suitable cells for producing influenza vaccines [33–39]. Usually, exogenous trypsin is needed for cell-based vaccine production. However, we report that the virus propagated in MDCK-TMPRSS2 and MDCK-MSPL cells could reach an equivalent titer compared with the MDCK cells plus

TPCK-trypsin. Thus, these cell lines can be used for cell-based vaccine production.

Böttcher and coworkers reported inducible MDCK cell lines which express TMPRSS2 and HAT under the control of tetracycline transcription system [40]. Their results showed that the cell lines can support multicycle replication of human H2 and H3 influenza viruses without trypsin. Different from their inducible cell lines, the cell lines we established consistently expressed TMPRSS2 or MSPL. Unexpectedly, permanent expression of these proteases was not toxic to the cells and both cell lines could be serially passaged for at least 20 passages. Even though these cells grew slightly slower than MDCK cells, when they grew to the suitable confluence for infection, the viral titer was comparable to the MDCK cells plus exogenous trypsin. Since Re-5, Re-6, and Re-9 are commercial inactivated AIV vaccines in China, we speculate that the cell lines could be applied in the large scale vaccine production in the future. They may also simplify the




FIGURE 3: Cell fusion induced by cotransfection of protease and influenza HA under acidic conditions. MDCK cells were cotransfected with pCAGGS-TMPRSS2 plus pCAGGS-(Re-5, Re-6, and Re-9) HA and pCAGGS-MSPL plus pCAGGS-(Re-5, Re-6, and Re-9) HA, respectively (a). pCAGGS plus pCAGGS-(Re-5, Re-6, and Re-9) HA supplemented with TPCK-trypsin and pCAGGS plus pCAGGS-(Re-5, Re-6, and Re-9) HA without TPCK-trypsin as control groups (b). See methods for cell treatments. Stained cells were observed under a Leica confocal laser microscope.




FIGURE 4: Re-5, Re-6, and Re-9 infected and spread within MDCK-TMPRSS2 and MDCK-MSPL cells. MDCK-TMPRSS2, MDCK-MSPL, MDCK + TPCK, and MDCK cells were infected with Re-5, Re-6, and Re-9, respectively (MOI = 0.01); cells were fixed 60 h after infection. Cells were incubated with Re-5, Re-6, and Re-9 antibodies and TRITC-labeled secondary antibody. Cells were observed under a Zeiss fluorescent microscope.

related researches or industrial applications to meet diverse demands.

Also, both MDCK-TMPRSS2 and MDCK-MSPL can cleave the HA protein of Re-5, Re-6, and Re-9 and

support multicycle replication of these viruses. MDCK-TMPRSS2 and MDCK-MSPL cells could therefore be useful for studying cleavage activation of influenza virus HA by TMPRSS2 and MSPL. Finally, the cell lines may




FIGURE 5: Growth kinetics of Re-5, Re-6, and Re-9 on MDCK-TMPRSS2 and MDCK-MSPL cells. MDCK-TMPRSS2, MDCK-MSPL, MDCK + TPCK, and MDCK cells were infected with Re-5, Re-6, and Re-9 as described and samples were taken at 12, 24, 48, 72, and 96 h after infection. Sample TCID₅₀ were determined and calculated using the method of Reed and Muench.

allow development of TMPRSS2 and MSPL-specific protease inhibitors.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Authors' Contribution

Zhiyuan Wen, Chao Wu, and Weiye Chen contribute equally to this work.

Acknowledgment

This study was supported by The National Key Technology R&D Program (2013BAD12B05).

References

- [1] M. Ozawa and Y. Kawaoka, "Taming influenza viruses," *Virus Research*, vol. 162, no. 1-2, pp. 8-11, 2011.
- [2] R. G. Webster, W. J. Bean, O. T. Gorman, T. M. Chambers, and Y. Kawaoka, "Evolution and ecology of influenza a viruses," *Microbiological Reviews*, vol. 56, no. 1, pp. 152-179, 1992.
- [3] H. Chen and Z. Bu, "Development and application of avian influenza vaccines in China," *Current Topics in Microbiology and Immunology*, vol. 333, no. 1, pp. 153-162, 2009.
- [4] G. Neumann, H. Chen, G. F. Gao, Y. Shu, and Y. Kawaoka, "H5N1 influenza viruses: outbreaks and biological properties," *Cell Research*, vol. 20, no. 1, pp. 51-61, 2010.
- [5] Y. Zhang, Q. Zhang, H. Kong et al., "H5N1 hybrid viruses bearing 2009/H1N1 virus genes transmit in guinea pigs by respiratory droplet," *Science*, vol. 340, no. 6139, pp. 1459-1463, 2013.
- [6] Y. Itoh, K. Shinya, M. Kiso et al., "In vitro and in vivo characterization of new swine-origin H1N1 influenza viruses," *Nature*, vol. 460, no. 7258, pp. 1021-1025, 2009.
- [7] T. Watanabe, M. Imai, S. Watanabe et al., "Characterization in vitro and in vivo of pandemic (H1N1) 2009 influenza viruses isolated from patients," *Journal of Virology*, vol. 86, no. 17, pp. 9361-9368, 2012.
- [8] T. Watanabe, M. Kiso, S. Fukuyama et al., "Characterization of H7N9 influenza A viruses isolated from humans," *Nature*, vol. 501, no. 7468, pp. 551-555, 2013.

- [9] Q. Zhang, J. Shi, G. Deng et al., "H7N9 influenza viruses are transmissible in ferrets by respiratory droplet," *Science*, vol. 341, no. 6144, pp. 410–414, 2013.
- [10] J. S. Rossman and R. A. Lamb, "Influenza virus assembly and budding," *Virology*, vol. 411, no. 2, pp. 229–236, 2011.
- [11] H. D. Klenk, R. Rott, M. Orlich, and J. Bloedorn, "Activation of influenza A viruses by trypsin treatment," *Virology*, vol. 68, no. 2, pp. 426–439, 1975.
- [12] D. A. Steinhauer, "Role of hemagglutinin cleavage for the pathogenicity of influenza virus," *Virology*, vol. 258, no. 1, pp. 1–20, 1999.
- [13] B. S. Hamilton, G. R. Whittaker, and S. Daniel, "Influenza virus-mediated membrane fusion: determinants of hemagglutinin fusogenic activity and experimental approaches for assessing virus fusion," *Viruses*, vol. 4, no. 7, pp. 1144–1168, 2012.
- [14] X. Sun and G. R. Whittaker, "Entry of influenza virus," *Advances in Experimental Medicine and Biology*, vol. 790, pp. 72–82, 2013.
- [15] T. Horimoto, K. Nakayama, S. P. Smeeckens, and Y. Kawaoka, "Proprotein-processing endoproteases PC6 and furin both activate hemagglutinin of virulent avian influenza viruses," *Journal of Virology*, vol. 68, no. 9, pp. 6074–6078, 1994.
- [16] H.-D. Klenk and W. Garten, "Host cell proteases controlling virus pathogenicity," *Trends in Microbiology*, vol. 2, no. 2, pp. 39–43, 1994.
- [17] A. Stieneke-Gröber, M. Vey, H. Angliker et al., "Influenza virus hemagglutinin with multibasic cleavage site is activated by furin, a subtilisin-like endoprotease," *The EMBO Journal*, vol. 11, no. 7, pp. 2407–2414, 1992.
- [18] E. Böttcher, T. Matrosovich, M. Beyerle, H.-D. Klenk, W. Garten, and M. Matrosovich, "Proteolytic activation of influenza viruses by serine proteases TMPRSS2 and HAT from human airway epithelium," *Journal of Virology*, vol. 80, no. 19, pp. 9896–9898, 2006.
- [19] W. Wang, E. N. Butler, V. Veguilla et al., "Establishment of retroviral pseudotypes with influenza hemagglutinins from H1, H3, and H5 subtypes for sensitive and specific detection of neutralizing antibodies," *Journal of Virological Methods*, vol. 153, no. 2, pp. 111–119, 2008.
- [20] P. Zmora, P. Blazejewski, A. Moldenhauer et al., "DESC1 and MSPL activate influenza A viruses and emerging coronaviruses for host cell entry," *Journal of Virology*, vol. 88, no. 20, pp. 12087–12097, 2014.
- [21] Y. Okumura, E. Takahashi, M. Yano et al., "Novel type II transmembrane serine proteases, MSPL and TMPRSS13, proteolytically activate membrane fusion activity of the hemagglutinin of highly pathogenic avian influenza viruses and induce their multicycle replication," *Journal of Virology*, vol. 84, no. 10, pp. 5089–5096, 2010.
- [22] G. Wang, T. Zhang, X. Li et al., "Serological evidence of H7, H5 and H9 avian influenza virus co-infection among herons in a city park in Jiangxi, China," *Scientific Reports*, vol. 4, article 6345, 2014.
- [23] V. G. Nguyen, H. K. Kim, H. J. Moon et al., "Evolutionary dynamics of a highly pathogenic type 2 porcine reproductive and respiratory syndrome virus: analyses of envelope protein-coding genes," *Transboundary and Emerging Diseases*, 2013.
- [24] B. Gotoh, F. Yamauchi, T. Ogasawara, and Y. Nagai, "Isolation of factor Xa from chick embryo as the amniotic endoprotease responsible for paramyxovirus activation," *FEBS Letters*, vol. 296, no. 3, pp. 274–278, 1992.
- [25] H. Kido, Y. Okumura, H. Yamada, T. Q. Le, and M. Yano, "Proteases essential for human influenza virus entry into cells and their inhibitors as potential therapeutic agents," *Current Pharmaceutical Design*, vol. 13, no. 4, pp. 405–414, 2007.
- [26] C. Chaipan, D. Kobasa, S. Bertram et al., "Proteolytic activation of the 1918 influenza virus hemagglutinin," *Journal of Virology*, vol. 83, no. 7, pp. 3200–3211, 2009.
- [27] H. Kido, Y. Okumura, E. Takahashi et al., "Host envelope glycoprotein processing proteases are indispensable for entry into human cells by seasonal and highly pathogenic avian influenza viruses," *Journal of Molecular and Genetic Medicine*, vol. 3, pp. 167–175, 2008.
- [28] J. M. Audsley and G. A. Tannock, "Cell-based influenza vaccines: progress to date," *Drugs*, vol. 68, no. 11, pp. 1483–1491, 2008.
- [29] Y. Genzel, I. Behrendt, S. König, H. Sann, and U. Reichl, "Metabolism of MDCK cells during cell growth and influenza virus production in large-scale microcarrier culture," *Vaccine*, vol. 22, no. 17–18, pp. 2202–2208, 2004.
- [30] O. Kistner, P. N. Barrett, W. Mundt, M. Reiter, S. Schober-Bendixen, and F. Dorner, "Development of a mammalian cell (Vero) derived candidate influenza virus vaccine," *Vaccine*, vol. 16, no. 9–10, pp. 960–968, 1998.
- [31] O. Kistner, M. K. Howard, M. Spruth et al., "Cell culture (Vero) derived whole virus (H5N1) vaccine based on wild-type virus strain induces cross-protective immune responses," *Vaccine*, vol. 25, no. 32, pp. 6028–6036, 2007.
- [32] K. Nerome, H. Kumihashi, R. Nerome et al., "Evaluation of immune responses to inactivated influenza vaccines prepared in embryonated chicken eggs and MDCK cells in a mouse model," *Developments in Biological Standardization*, vol. 98, pp. 53–74, 1999.
- [33] J. A. Tree, C. Richardson, A. R. Fooks, J. C. Clegg, and D. Looby, "Comparison of large-scale mammalian cell culture systems with egg culture for the production of influenza virus A vaccine strains," *Vaccine*, vol. 19, no. 25–26, pp. 3444–3450, 2001.
- [34] Y. Genzel, M. Fischer, and U. Reichl, "Serum-free influenza virus production avoiding washing steps and medium exchange in large-scale microcarrier culture," *Vaccine*, vol. 24, no. 16, pp. 3261–3272, 2006.
- [35] Y. Genzel, R. M. Olmer, B. Schäfer, and U. Reichl, "Wave microcarrier cultivation of MDCK cells for influenza virus production in serum containing and serum-free media," *Vaccine*, vol. 24, no. 35–36, pp. 6074–6087, 2006.
- [36] Y. Genzel and U. Reichl, "Continuous cell lines as a production system for influenza vaccines," *Expert Review of Vaccines*, vol. 8, no. 12, pp. 1681–1692, 2009.
- [37] A. Y.-C. Hu, T.-C. Weng, Y.-F. Tseng et al., "Microcarrier-based MDCK cell culture system for the production of influenza H5N1 vaccines," *Vaccine*, vol. 26, no. 45, pp. 5736–5740, 2008.
- [38] J. Liu, X. Shi, R. Schwartz, and G. Kemble, "Use of MDCK cells for production of live attenuated influenza vaccine," *Vaccine*, vol. 27, no. 46, pp. 6460–6463, 2009.
- [39] P. D. Minor, O. G. Engelhardt, J. M. Wood et al., "Current challenges in implementing cell-derived influenza vaccines: implications for production and regulation, July 2007, NIBSC, Potters Bar, UK," *Vaccine*, vol. 27, no. 22, pp. 2907–2913, 2009.
- [40] E. Böttcher, C. Freuer, T. Steinmetzer, H.-D. Klenk, and W. Garten, "MDCK cells that express proteases TMPRSS2 and HAT provide a cell system to propagate influenza viruses in the absence of trypsin and to study cleavage of HA and its inhibition," *Vaccine*, vol. 27, no. 45, pp. 6324–6329, 2009.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

