

## Review Article

# Medicinal Herbals with Antiplatelet Properties Benefit in Coronary Atherothrombotic Diseases

Nurul Huda Mohd Nor,<sup>1</sup> Fauziah Othman,<sup>1</sup>  
Eusni Rahayu Mohd Tohit,<sup>2</sup> and Sabariah Md Noor<sup>2</sup>

<sup>1</sup>Department of Human Anatomy, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, 43000 Serdang, Selangor, Malaysia

<sup>2</sup>Department of Pathology, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

Correspondence should be addressed to Sabariah Md Noor; [md\\_sabariah@upm.edu.my](mailto:md_sabariah@upm.edu.my)

Received 7 December 2015; Revised 10 February 2016; Accepted 17 February 2016

Academic Editor: Giovanni de Gaetano

Copyright © 2016 Nurul Huda Mohd Nor et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Coronary atherothrombotic diseases such as coronary artery disease, peripheral vascular disease, cerebrovascular disease, and heart failure are the serious concerns of the thrombus formed in blood vessels. Anticoagulant and antiplatelet drugs are the cornerstones of the management of these diseases. To prevent the recurrence of these diseases, double antiplatelet drugs such as aspirin and clopidogrel has been the standard management in most hospitals. However, aspirin resistance and clopidogrel inefficient effects due to noncompliance with double drugs regimen can cause a sinister effect on patients. Medicinal plants serve as a greater resource for new medication and their potential currently became a topic of interest to the researchers all over the world. Traditionally, certain herbs have been used as a treatment for heart diseases but have been investigated for their antiplatelet properties. This current review explained few traditional antithrombotic herbals and their antiplatelet properties *in vitro* and *in vivo* and this is to be deeply discussed in further research.

## 1. Introduction

Coronary atherothrombotic diseases (CADs) are the leading causes of death and disabilities in the world. Globally, it is estimated that about 17.3 million people had died from CADs in 2008 and, by 2030, approximately 23.6 million of the world population might die from CADs [1]. CADs consist of coronary artery disease, peripheral vascular disease, cerebrovascular disease, and heart failure. The risk factor of CADs is mainly due to atherosclerosis that leads to arterial thrombosis. The pathogenesis process encompasses chronic progressive atherosclerosis punctuated by acute processes, that is, (1) plaque rupture and (2) platelet thrombus formation in the areas of progressive stenosis. The final pathway to arterial thrombosis is atherosclerosis plaque rupture; thus, the effective way to treat or to prevent this occurrence is by limiting or eradicating platelet-dependent thrombus formation.

Oral antiplatelet drugs are the cornerstone of the therapy of CADs that consist of primary and secondary prevention strategies to manage these diseases. Patients with efficient antiplatelet therapy can reduce stroke and death rates by about 25% [2]. Currently, the standard practice in a clinical setting to prevent adverse events of CADs is a combination of aspirin and clopidogrel which inhibits cyclooxygenase and antagonizes P2Y<sub>12</sub>, respectively. Despite their proven effectiveness, there are cases of recurrent cardiovascular diseases among those who take double antiplatelet drugs. This might be due to incompliance and the adverse effect faced by the patients. Gastrointestinal bleeding and gastric ulcers are the most common adverse effects reported [3]. There are three points of argument of aspirin resistance that include the following: no evidence of antithrombotic effect being aspirin dosage related, ability to generate thromboxane A<sub>2</sub> despite the usual aspirin dosage, and low dose of aspirin causing blocking of more than 95% of platelet cyclooxygenase-1

activity. Thus, inability to benefit from regular aspirin dosage leads to the failure of aspirin as an antiplatelet therapy [4].

Medicinal plants serve as a greater resource for new medication and their potential currently became a topic of interest to the researchers all over the world. The World Health Organisation (WHO) has recommended medicinal plants to be used more effectively in healthcare system [1]. Recently, there is a trend of studies on the development of natural compound present in traditional medicinal herbs demonstrating antiplatelet properties. The currently known mechanisms of action that can be used to interfere with the primary haemostatic system that regulates receptors such as catecholamine, collagen, thrombin, prostacyclin, ADP, PGD<sub>2</sub>, PGE<sub>2</sub>, and serotonin, TXA<sub>2</sub>, cAMP, cGMP, and calcium have been identified to be the effective target of antiplatelet therapy [5]. In order to have a wider variety of treatments, screening of medicinal herbs that is postulated to have high potential of antiplatelet properties was performed.

## 2. Methods

The information about medicinal plants that are traditionally used in the management of coronary atherothrombotic disease was obtained from published papers and texts on ethnobotanical studies as well as those investigating the effects of plant(s) used with anticoagulation effect. A literature search on electronic databases such as Google Scholar, PubMed, and Scopus up to 2015 was carried out using “antiplatelet” and “herbal” as the keywords for primary search and then “plant name – accepted or synonyms”, “constituents”, and/or “mechanism” for the secondary search.

In order to highlight the medicinal plants traditionally used in CADs management with the potential of integration in the healthcare system, not all identified plants were included in this paper. Therefore, this review was not exhaustive of all plants used traditionally for treating CADs.

## 3. Results and Discussions

CAD can be prevented or treated by interfering with its mechanism of thrombosis. The primary goal of antithrombotic therapy is to inhibit platelet activation (primary haemostasis) and fibrin formation via cascade (secondary haemostasis) from happening. There are four steps in platelet activation pathways which are adhesion, aggregation, secretion, and procoagulation which lead to secondary haemostasis. However, this review focuses on primary haemostasis and the main method used in these various studies was light impedance aggregometry, either *in vivo* or *in vitro*. There are a lot of plants that exhibit antiplatelet properties. Ten plants from various families and regions are discussed as they possibly contain the compound responsible for those properties as listed in Table 1.

**3.1. *Curcuma longa* (Turmeric).** Turmeric is derived from the rhizomes of *Curcuma longa* and has been used in Southeast Asia for a long time as an Indian spice and treatment for several conditions such as stomachache, wounds, and

joint problems in Ayurvedic medicine [6]. Turmeric itself has been extensively used for its antioxidant, antimicrobial, anti-inflammatory, and antiproliferative properties. The most lipophilic component of *Curcuma longa* extracted called Curcuma oil (C. oil) has been utilized as antimicrobial, antifungal, antiviral, and anti-inflammatory agent and also used for wound healing activity.

A fraction of C. oil, namely, ar-turmerone, has been shown to inhibit collagen- (10 µg/ml), adenosine-5'-diphosphate- (ADP- (10 µM)), and thrombin- (0.64 U/ml) induced platelet aggregation. The effect of C. oil on platelet aggregation (*in vivo*) in rats significantly reduced collagen- (28%), ADP- (31%), and thrombin- (34%) induced platelet aggregation in comparison to aspirin (50 to 75%). While these three agonists had significantly inhibitory effect, calcium ionophore, PAF, and arachidonic acid were not affected at all [6]. This study is supported by another study done on ar-turmerone that showed that ar-turmerone inhibited platelet activation induced by collagen (2 µg/ml) gave 100% inhibition at concentration of 100 µg/ml and 80% and 60% inhibition at concentration of 10 µg/ml and 5 µg/ml, respectively [7]. In a study that used curcumin (500 µM), a major ingredient of *Curcuma longa* showed effective inhibition against AA-induced platelet aggregation (1.0 mM). A further study has noted that an increase in lipoxygenase- (LOX-) derived products was observed in curcumin-treated platelets [8]. These could be due to the redirection of AA to LOX pathway or/and potentiation of the enzyme. AA is a membrane-derived fatty acid metabolized by cyclooxygenase to prostaglandin (PG) endoperoxide intermediates such as PGH<sub>2</sub>. In platelets, endoperoxides are further metabolized to thromboxane A<sub>2</sub> and PGs [9]. This suggests that the antiplatelet effect of C. oil possibly affects the cyclooxygenase (COX) pathway in platelet at membrane/receptor level.

**3.2. *Berberis vulgaris*.** Scientifically named as *Berberis vulgaris*, the herb known as “barberry” is a native herb in Europe and Asia and is widely grown in Canada to Pennsylvania. In traditional Chinese medicine, barberry was mentioned more than 3000 years ago to have diverse medicinal properties including antimicrobial, antiemetic, antipyretic, and antipruritic properties. Scientifically, barberry may offer healthcare benefits as an antihypertensive and vasodilator agent [10] and play a prominent role in apoptosis in the treatment of hepatocarcinogenic rats [11].

A study has been done to investigate the antiplatelet effects of *Berberis vulgaris* on alkaloid component named berberine (BR) towards rabbits as *in vivo* model. In the study, platelet aggregation induced by adenosine diphosphate (ADP), collagen, AA, and calcium ionophore A23187 was differentially inhibited in which collagen-induced platelet aggregation was most potently inhibited and calcium ionophore A23187-induced platelet aggregation was least inhibited. The inhibition of platelet aggregation is dose dependent from 10 to 30 mg/kg, giving the maximum inhibition of 52.5% [12]. There is also another report of inhibitory effects on AA metabolism and calcium influx, partial agonist activity on platelet α<sub>2</sub>-adrenoceptors using *Berberis vulgaris* extract [13].

TABLE 1: Summary of inhibition of platelet activation by medicinal herbs.

Reference	Plants	Compound	Dose of plants' extract	Agonist and dosage	Postulated antiplatelet properties	In vivo or in vitro
[6–8]	Turmeric ( <i>Curcuma longa</i> )	Ar-turmerone, curcumin	100 µg/ml, 500 µM	Collagen (2 µg/ml), ADP (10 µM), and thrombin (0.64 U/ml)	Adhesion	In vivo
[12, 13]	Barberry ( <i>Berberis vulgaris</i> )	Berberine	30 mg/kg	Collagen	Adhesion	In vivo
[14]	<i>Osmanthus fragrans</i>		2.8 mg/ml	Collagen (1 mg/ml) and ADP (5 mM)	Adhesion, secretion	In vitro
[15]	Magnolia ( <i>Magnolia officinalis</i> )	Magnolol, honokiol		Collagen (10 µg/ml)	Adhesion	In vitro
[16]	<i>Cudrania tricuspidata</i> Bureau	Cudraticusxanthone A	10 µM	Thrombin (3 U/ml)	Aggregation	In vitro
[17, 18]	Dong quai ( <i>Angelica sinensis</i> )	Z-Ligustilide	40 mg/kg	ADP (5 µM)	Secretion	In vivo
[19–22]	Ginger ( <i>Zingiber officinale</i> )	Gingerol, paradol	25 µM	AA (20 mM) and AA (2 µM)	Secretion	In vitro
[23, 24]	Garlic ( <i>Allium sativum</i> )	Methanol extract	48–95 g/kg	ADP (10 µM), epinephrine (10 µg/ml), collagen (2 µg/ml), thrombin (0.1 U/ml), AA (1.1 mM), and PAF (10 µM)	Adhesion, secretion, and aggregation	In vitro, in vivo
[25–27]	<i>Scutellaria baicalensis</i> Georgi	Baicalin, wogonoside	45.6 µM, 30 µM	Thrombin (3 U/ml), collagen (1 µg/ml), and ADP (5 µM)	Adhesion, secretion	In vitro
[28–30]	<i>Fructus gardenia</i>	Iridoid glycosides, genipin	100 mg/k	Collagen (1 µg/ml)	Adhesion, vasoconstriction	In vivo

**3.3. *Osmanthus fragrans*.** This plant has been used as an additive in beverage and also in perfumes as it has intense fragrance [31]. Besides, it had been used traditionally as treatment's constituent for stomachache, rheumatism, and halitosis [14]. *Osmanthus fragrans* (OF) has high amount of phenolic acid which has been used as a natural source of antioxidant for controlling probing free radicals related to various diseases including chronic inflammation, cataracts, and atherosclerosis and other cardiovascular diseases [32].

One novel study on phytochemical investigation on OF seed, the purple-brown ripe fruit of OF, extracted using ethyl acetate and n-butyl alcohol showed the greatest effect on inhibition of collagen-induced platelet aggregation at concentration of 1.4 mg/ml and 2.8 mg/ml. Further study using the same extracts but using ADP showed equal effect compared to a positive drug at 2.8 mg/ml [14]. These findings suggest that OF seeds may have the potential to be developed as a novel inhibitor of platelet aggregation.

**3.4. *Magnolia officinalis*.** This traditional Chinese and Japanese medicinal herb comes from Magnoliaceae family and is being used for the treatment of thrombotic stroke, headache, and typhus fever [15]. It is believed to have multifunctional health benefits including general antioxidative effects through scavenging hydroxyl radicals that are generated by ultraviolet (UV) irradiation [33] and anticancer properties induced by apoptosis of HER2 overexpressed in breast cancer cells [34]. They are also capable of controlling pain [35]. Besides that, magnolol has an antiapoptotic effect through the regulation of bcl-XL gene and suppression of bcl-xS gene [36].

It contains two bioactive isomers called magnolol and honokiol. As for the cardiovascular protection effects, the herbs through magnolol suppress the platelet activation in aggregation step via inhibition of intracellular calcium mobilization and thromboxane formation [15, 37]. Study on inhibition potential of honokiol and magnolol against collagen (10  $\mu$ g/ml) showed that honokiol is 5 to 10 times more potent than magnolol at even very low concentration and it is concentration dependent. Honokiol and magnolol possess 80% and 55% inhibition on collagen-induced platelet aggregation at the highest concentration. Further test on treated platelets with magnolol and honokiol showed that both had reversible action from collagen-induced platelet aggregation. After conducting the centrifugation of platelet suspension, it is found that there is no presence of lactate dehydrogenase which confirms that both compounds did not cause breakdown of platelets [15].

**3.5. *Cudrania tricuspidata* Bureau.** This plant is also known as *Maclura tricuspidata* Bureau. It is a plant from the Moraceae family. It is a tree native to East Asia and the root has been used as a traditional Chinese medicine for treating dysmenorrhea, rheumatism, and jaundice. The wealthy source of flavonoids and xanthenes enhances this plant to be explored. It has been reported to possess antioxidant [38], anti-inflammatory, anticancer [39], antihepatoprotective [40], and anticoagulant properties.

The bioactive compound used, which exhibits the antiplatelet properties obtained from the roots of *Cudrania*

*tricuspidata*, is called cudraticusxanthone A (CTXA). Studies done using mouse platelet showed that CTXA significantly prevented the platelet aggregation induced by thrombin (3 U/ml) in a concentration-dependent manner. At concentration of 10  $\mu$ M, platelet aggregation reduced to 35% compared to control (80%). CTXA inhibited thrombin-catalyzed fibrin polymerization resulting in a reduction of plasminogen activator inhibitor type 1 (PAI-1) and platelet aggregation [16].

**3.6. *Angelica sinensis*.** *Angelica sinensis* is a type of herb from the family Apiaceae that grows in mountains of China, Korea, and Japan. *Angelica sinensis* is a Chinese traditional herb used for treating dysmenorrhea and it is also used as a food supplement in Korea. Radix *Angelica sinensis* is also officially listed in the Chinese pharmacopoeia. The health benefits are well documented [41]. Conventionally, this plant had been used in the treatment and prevention of thrombosis.

An arterial thrombus is mainly composed of platelet aggregation; hence, the platelets play a vital role in both initiation and growth of thrombi. An initial study has examined the effect of herbs extract named Z-ligustilide, characterized by 3-n-alkylphthalide constituents on atrial-type thrombus formation on a rat arteriovenous shunt model. After administration of the extract (40 mg/kg) for 3 consecutive days orally, thrombus formation was significantly reduced. Further tests on the treated rats showed that the maximum platelet aggregation by ADP (5  $\mu$ M) had decreased and the result was 3 times greater than aspirin at a dose of 40 mg/kg [17]. Another study done on rabbit showed the same inhibition of ADP-induced platelet aggregation at 10  $\mu$ M and it was comparable to aspirin at the same dosage. The possible mechanism of action is due to the inhibition of thromboxane A<sub>2</sub> formation or Ca<sup>2+</sup> activation in platelet [18].

**3.7. *Zingiber officinale* (Ginger).** *Zingiber officinale* from Zingiberaceae family is a medicinal herb that is used in our daily food especially in Asian countries. The benefit of this herb is well known for its antiulcer [42] and antibacterial properties [43]. The common name for this plant is ginger, a traditional medicine with a carminative effect.

This herb has been investigated for its antiplatelet properties. An initial study done on aqueous extract of ginger had shown that it inhibits platelet aggregation induced by AA, adenosine diphosphate, and collagen [19]. A research done on a compound of [8]-gingerol showed that selectively inhibited AA-induced platelet aggregation in washed platelet preparation started at 1 mM up to 20 mM AA concentration. It gave partial inhibition against collagen-induced platelet aggregation at 10  $\mu$ g/ml [20, 21]. Further studies on [6]- and [8]-gingerol have shown inhibition activity towards COX-1 in human platelet at 0.75 mM AA concentration and using rat basophilic leukemia (RBL-2H3) cells, gingerols at IC<sub>max</sub> approximately 20 to 25  $\mu$ M inhibit COX activity by measuring the product of AA metabolism by COX, prostaglandin D<sub>2</sub> [3]. These suggest the inhibition mechanism of gingerols possibly due to effect on AA-induced platelet activation. Besides that, another phenolic compound, [8]-paradol, a natural constituent of ginger, also has been found to be the



most potent inhibitor against AA-induced platelet aggregation at very low concentration ( $2\ \mu\text{M}$ ). Further test on inhibitory activity of this compound against COX-1 showed the strongest inhibitory activity with  $\text{IC}_{50}$  values of  $4\ \mu\text{M}$  [22]. Overall studies exhibited that ginger mechanism of platelet inhibition through cyclooxygenase-1/thromboxane synthase activity compared to aspirin activity.

**3.8. *Allium sativum*.** *Allium sativum* is also known as garlic. This medicinal herb had been narratively used by veterinarians to treat peripheral venous thrombosis in horses in the French literature a long time ago.

Study on cooked blanched garlic leaves juices on rabbit platelets aggregation induced by ADP and collagen showed significant inhibition with maximum ratio of 87.37% and 86.24%, respectively, *in vitro* and *in vivo*. The inhibition of aggregation pathway mainly blocks the combination of fibrinogen with  $\mu\text{Fib-R}$ , which finally results in the inhibition of platelet aggregation proved by the amount of platelet fibrinogen binding detected by flow cytometry [23]. Besides, study on the effect of methanol extract of garlic bulbs (EOG) and three pure components isolated from it (F1, F2, and F3) on human platelet aggregation induced by ADP ( $10\ \mu\text{M}$ ), epinephrine ( $10\ \mu\text{g/ml}$ ), collagen ( $2\ \mu\text{g/ml}$ ), thrombin ( $0.1\ \text{U/ml}$ ), arachidonate ( $1.1\ \text{mM}$ ), and PAF ( $10\ \mu\text{M}$ ) was conducted. The incubation of platelet rich plasma with EOG inhibits 100% platelet aggregation induced by all of the abovementioned agonists at  $68 \pm 5\ \text{g/ml}$ ,  $48 \pm 2\ \text{g/ml}$ ,  $52 \pm 7\ \text{g/ml}$ ,  $95 \pm 7\ \text{g/ml}$ ,  $55 \pm 3\ \text{g/ml}$ , and  $60 \pm 3\ \text{g/ml}$  concentration, respectively. However, for 3 other fractions isolated from it, F3 was shown to have about four times more potency than others using listed agonists at  $120 \pm 9\ \text{g/ml}$ ,  $60 \pm 8\ \text{g/ml}$ ,  $75 \pm 11\ \text{g/ml}$ ,  $100 \pm 12\ \text{g/ml}$ ,  $30 \pm 1\ \text{g/ml}$ , and  $65 \pm 7\ \text{g/ml}$  and is also able to induce ADP-rapid deaggregation when added to aggregated platelet. This can indicate the possible mechanism of antiplatelet maybe on the plasma membrane properties. Thrombin-induced release of ATP from platelet also was inhibited by 75–80% after F3 treatment [24].

**3.9. *Scutellaria baicalensis* Georgi.** It is one of the important herbs used in traditional Chinese medicine for treatment of febrile, cough, and liver problem. Its roots have been used for their anti-inflammatory and anticancer effects, for treating bacterial and viral infections, and for reducing the total cholesterol level and hypertension [44]. Several bioactive compounds have been isolated from the root of the plant, namely, baicalein, baicalin, wogonin, and oroxylin A. Baicalin possesses anti-HIV, antitumor, antioxidant, and free radical scavenging effects. Wogonin has antirespiratory syncytial virus, antihepatitis B virus, antitumor, antioxidant, and free radical scavenging effects while oroxylin A has antirespiratory syncytial virus activity [45]. Baicalin had been used in treatment of bronchitis, asthma, and skin allergy [25]. Studies have shown that baicalin inhibited thrombin-catalyzed fibrin polymerization and platelet functions, prolonged aPTT and PT significantly, and inhibited the activities and production of thrombin and FXa.

Further study done on baicalin showed that it suggestively inhibited thrombin- ( $3\ \text{U/ml}$ ) and collagen- ( $1\ \mu\text{g/ml}$ )

induced mouse platelet aggregation in a concentration-dependent manner with  $\text{IC}_{50}$  of baicalin in PRP being 45.6 and  $41.1\ \mu\text{M}$ , respectively [25]. A study also demonstrated that baicalin enhanced the phenylephrine-induced contraction through inhibiting the production or/and release of endothelial nitric oxide in the isolated rat mesenteric artery rings [46]. A treatment with active metabolite wogonin named flavonoid wogonoside (WGN) has been shown to inhibit thrombin-catalyzed fibrin polymerization and thrombin-induced platelet aggregation. The inhibition is concentration dependent and at  $30\ \mu\text{M}$  of WGN, it is about 50% inhibition with final concentration of thrombin at  $3\ \text{U/ml}$ . WGN was also shown to elicit anticoagulant effects in mice [26]. Further studies on the root extract of *Scutellaria baicalensis* Georgi have shown a significant inhibition activity of platelet aggregation ratio of rabbits induced by ADP ( $5\ \mu\text{M}$ ) with average inhibition rate of 45.52% at  $4.023\ \text{mg/ml}$  of extract and that of aspirin group was 54.96% [27].

**3.10. *Gardenia jasminoides* Ellis.** This plant had been known as Zhi Zi in Chinese herb directory. It has been traditionally used as an analgesic for several medical conditions such as pain due to liver problem, dysentery, and trauma for external application [47].

*Gardenia jasminoides* Ellis with its bioactive compounds, iridoid glycosides (IG), has been studied. IGs have been shown to have antithrombotic properties through the inhibition of platelet aggregation with a little effect on the coagulation time of peripheral blood. A study showed that IGs may significantly and dose-dependently reduce arterial thrombus load in a model of carotid artery thrombosis and inhibit collagen-induced platelet aggregation in rats at  $1\ \mu\text{g/ml}$ . Pretreatment with IGs for 3 days with dose of  $100\ \text{mg/kg}$  gave 13% platelet aggregation ratio comparable to aspirin (18%) at  $40\ \text{mg/kg}$ . [28]. Another compound of *Gardenia jasminoides* Ellis called genipin (GE) with different concentrations has shown significant inhibition on the collagen-induced platelet aggregation *in vitro*, but only a little effect on that induced by ADP and AA [29]. In another *in vivo* thrombosis model, it has also been confirmed that *Gardenia jasminoides* Ellis extract can reduce the weight or length of thrombus. Its mechanism of antiplatelet aggregation and inhibition on thromboembolism was probably related to the activation via inhibiting endothelin-1 (ET-1) expression as a potent vasoconstrictive peptide. GE was found to be very effective in reducing the nuclear factor kappa B (NF- $\kappa\text{B}$ ) by disturbing the I $\kappa\text{B}$  degradation. NF- $\kappa\text{B}$  is an inducer of cyclooxygenase-2 (COX-2) expression. By reducing NF- $\kappa\text{B}$  activity, the production of COX-2 and inducible nitric oxide synthase (iNOS) is downregulated [30, 48].

## 4. Conclusion

This review focused on the availability of medicinal herbs that possess antiplatelet properties. Most studies are still at the preliminary stage as many steps are required so that the herbs can be used as a substitution or can be combined with current antiplatelet agents such as aspirin and clopidogrel.

## Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

## Acknowledgments

This work is supported by Exploratory Research Grant Scheme, Ministry of Education Malaysia (5527163). The authors also acknowledge the Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, for providing library resource facilities and support.

## References

- [1] World Health Organization, *WHO Traditional Medicine Strategy 2002–2005*, WHO, 2002.
- [2] Antithrombotic Trialists' Collaboration, "Collaborative meta-analysis of randomised trials of antiplatelet therapy for prevention of death, myocardial infarction, and stroke in high risk patients," *British Medical Journal*, vol. 324, no. 7329, pp. 71–86, 2002.
- [3] K. L. K. Koo, A. J. Ammit, V. H. Tran, C. C. Duke, and B. D. Roufogalis, "Gingerols and related analogues inhibit arachidonic acid-induced human platelet serotonin release and aggregation," *Thrombosis Research*, vol. 103, no. 5, pp. 387–397, 2001.
- [4] J. W. Eikelboom, J. Hirsh, J. I. Weitz, M. Johnston, Q. Yi, and S. Yusuf, "Aspirin-resistant thromboxane biosynthesis and the risk of myocardial infarction, stroke, or cardiovascular death in patients at high risk for cardiovascular events," *Circulation*, vol. 105, no. 14, pp. 1650–1655, 2002.
- [5] Y. Liu, H.-J. Yin, D.-Z. Shi, and K.-J. Chen, "Chinese herb and formulas for promoting blood circulation and removing blood stasis and antiplatelet therapies," *Evidence-Based Complementary and Alternative Medicine*, vol. 2012, Article ID 184503, 8 pages, 2012.
- [6] P. Prakash, A. Misra, W. R. Surin et al., "Anti-platelet effects of Curcuma oil in experimental models of myocardial ischemia-reperfusion and thrombosis," *Thrombosis Research*, vol. 127, no. 2, pp. 111–118, 2011.
- [7] H. S. Lee, "Antiplatelet property of *Curcuma longa* L. rhizome-derived ar-turmerone," *Bioresource Technology*, vol. 97, no. 12, pp. 1372–1376, 2006.
- [8] K. C. Srivastava, A. Bordia, and S. K. Verma, "Curcumin, a major component of food spice turmeric (*Curcuma longa*) inhibits aggregation and alters eicosanoid metabolism in human blood platelets," *Prostaglandins, Leukotrienes and Essential Fatty Acids*, vol. 52, no. 4, pp. 223–227, 1995.
- [9] B. Samuelsson, G. Folco, E. Granström, H. Kindahl, and C. Malmsten, "Prostaglandins and thromboxanes: biochemical and physiological considerations," *Advances in Prostaglandin and Thromboxane Research*, vol. 4, pp. 1–25, 1978.
- [10] Z. Fatehi-Hassanabad, M. Jafarzadeh, A. Tarhini, and M. Fatehi, "The antihypertensive and vasodilator effects of aqueous extract from *Berberis vulgaris* fruit on hypertensive rats," *Phytotherapy Research*, vol. 19, no. 3, pp. 222–225, 2005.
- [11] G. Motaleb, P. Hanachi, O. Fauziah, and R. Asmah, "Effect of *Berberis vulgaris* fruit extract on alpha-fetoprotein gene expression and chemical carcinogen metabolizing enzymes activities in hepatocarcinogenesis rats," *Iranian Journal of Cancer Prevention*, vol. 1, no. 1, pp. 33–42, 2012.
- [12] C. G. Huang, Z. L. Chu, S. J. Wei, H. Jiang, and B. H. Jiao, "Effect of berberine on arachidonic acid metabolism in rabbit platelets and endothelial cells," *Thrombosis Research*, vol. 106, no. 4–5, pp. 223–227, 2002.
- [13] M. El Sayed, D. Ghareeb, E. Sarhan, and A. Khalil, "Therapeutic bio-screening of the bioactive ingredients of *Berberis vulgaris*," *Functional Plant Science and Biotechnology*, vol. 5, no. 1, pp. 63–68, 2011.
- [14] W. Tang, J. Cao, X. Zhang, and Y. Zhao, "*Osmanthus fragrans* seeds, a source of secoiridoid glucosides and its antioxidizing and novel platelet-aggregation inhibiting function," *Journal of Functional Foods*, vol. 14, no. 3, pp. 337–344, 2015.
- [15] C.-M. Teng, C.-C. Chen, F.-N. Ko et al., "Two antiplatelet agents from *Magnolia officinalis*," *Thrombosis Research*, vol. 50, no. 6, pp. 757–765, 1988.
- [16] H. Yoo, S.-K. Ku, W. Lee et al., "Antiplatelet, anticoagulant, and profibrinolytic activities of cudraticusxanthone A," *Archives of Pharmacol Research*, vol. 37, no. 8, pp. 1069–1078, 2014.
- [17] L. Zhang, J.-R. Du, J. Wang et al., "Z-ligustilide extracted from *Radix angelica sinensis* decreased platelet aggregation induced by ADP ex vivo and arterio-venous shunt thrombosis in vivo in rats," *Yakugaku Zasshi*, vol. 129, no. 7, pp. 855–859, 2009.
- [18] N.-Y. Yang, G.-S. Zhou, Y.-P. Tang et al., "Two new  $\alpha$ -pinene derivatives from *Angelica sinensis* and their anticoagulative activities," *Fitoterapia*, vol. 82, no. 4, pp. 692–695, 2011.
- [19] K. C. Srivastava, "Effects of aqueous extracts of onion, garlic and ginger on platelet aggregation and metabolism of arachidonic acid in the blood vascular system: in vitro study," *Prostaglandins, Leukotrienes and Medicine*, vol. 13, no. 2, pp. 227–235, 1984.
- [20] J.-H. Guh, F.-N. Ko, T.-T. Jong, and C.-M. Teng, "Antiplatelet effect of gingerol isolated from *Zingiber officinale*," *Journal of Pharmacy and Pharmacology*, vol. 47, no. 4, pp. 329–332, 1995.
- [21] K. C. Srivastava, "Isolation and effects of some ginger components on platelet aggregation and eicosanoid biosynthesis," *Prostaglandins, Leukotrienes and Medicine*, vol. 25, no. 2–3, pp. 187–198, 1986.
- [22] E. Nurtjahja-Tjendraputra, A. J. Ammit, B. D. Roufogalis, V. H. Tran, and C. C. Duke, "Effective anti-platelet and COX-1 enzyme inhibitors from pungent constituents of ginger," *Thrombosis Research*, vol. 111, no. 4–5, pp. 259–265, 2003.
- [23] X. H. Wang and Y. Di, "Mechanism of cooked blanched garlic leaves against platelet aggregation," *Journal of Experimental Hematology*, vol. 22, no. 3, pp. 753–757, 2014.
- [24] R. Apitz-Castro, S. Cabrera, M. R. Cruz, E. Ledezma, and M. K. Jain, "Effects of garlic extract and of three pure components isolated from it on human platelet aggregation, arachidonate metabolism, release reaction and platelet ultrastructure," *Thrombosis Research*, vol. 32, no. 2, pp. 155–169, 1983.
- [25] W. Lee, S.-K. Ku, and J.-S. Bae, "Antiplatelet, anticoagulant, and profibrinolytic activities of baicalin," *Archives of Pharmacol Research*, vol. 38, no. 5, pp. 893–903, 2015.
- [26] S.-K. Ku and J.-S. Bae, "Antithrombotic activities of wogonin and wogonoside via inhibiting platelet aggregation," *Fitoterapia*, vol. 98, pp. 27–35, 2014.
- [27] Y. Zhang, X. Wang, X. Wang et al., "Protective effect of flavonoids from *Scutellaria baicalensis* Georgi on cerebral ischemia injury," *Journal of Ethnopharmacology*, vol. 108, no. 3, pp. 355–360, 2006.

- [28] P. Wang, Q. Wang, C. Luo, C. Tan, and X. Yuan, "Iridoid glycosides extracted from Zhizi (Fructus Gardeniae) decrease collagen-induced platelet aggregation and reduce carotid artery thrombosis in an in vivo rat model," *Journal of Traditional Chinese Medicine*, vol. 33, no. 4, pp. 531–534, 2013.
- [29] Y. Suzuki, K. Kondo, Y. Ikeda, and K. Umemura, "Antithrombotic effect of geniposide and genipin in the mouse thrombosis model," *Planta Medica*, vol. 67, no. 9, pp. 807–810, 2001.
- [30] H. Liu, Y.-F. Chen, F. Li, and H.-Y. Zhang, "Fructus Gardenia (*Gardenia jasminoides* J. Ellis) phytochemistry, pharmacology of cardiovascular, and safety with the perspective of new drugs development," *Journal of Asian Natural Products Research*, vol. 15, no. 1, pp. 94–110, 2013.
- [31] C.-Y. Hung, Y.-C. Tsai, and K.-Y. Li, "Phenolic antioxidants isolated from the flowers of *Osmanthus fragrans*," *Molecules*, vol. 17, no. 9, pp. 10724–10737, 2012.
- [32] A.-N. Li, S. Li, H.-B. Li, D.-P. Xu, X.-R. Xu, and F. Chen, "Total phenolic contents and antioxidant capacities of 51 edible and wild flowers," *Journal of Functional Foods*, vol. 6, no. 1, pp. 319–330, 2014.
- [33] S. Fujita and J. Taira, "Biphenyl compounds are hydroxyl radical scavengers: their effective inhibition for UV-induced mutation in *Salmonella typhimurium* TA102," *Free Radical Biology and Medicine*, vol. 17, no. 3, pp. 273–277, 1994.
- [34] H. Liu, C. Zang, A. Emde et al., "Anti-tumor effect of honokiol alone and in combination with other anti-cancer agents in breast cancer," *European Journal of Pharmacology*, vol. 591, no. 1–3, pp. 43–51, 2008.
- [35] Y.-J. Lee, Y. M. Lee, C.-K. Lee, J. K. Jung, S. B. Han, and J. T. Hong, "Therapeutic applications of compounds in the *Magnolia* family," *Pharmacology & Therapeutics*, vol. 130, no. 2, pp. 157–176, 2011.
- [36] B. Jawan, S. Goto, T.-L. Pan et al., "The protective mechanism of magnolol, a Chinese herb drug, against warm ischemia-reperfusion injury of rat liver," *Journal of Surgical Research*, vol. 110, no. 2, pp. 378–382, 2003.
- [37] C.-M. Teng, F.-N. Ko, J.-P. Wang et al., "Antithrombotic and antithrombotic effect of some antiplatelet agents isolated from Chinese herbs," *Journal of Pharmacy and Pharmacology*, vol. 43, no. 9, pp. 667–669, 1991.
- [38] B. W. Lee, J. H. Lee, S.-T. Lee et al., "Antioxidant and cytotoxic activities of xanthones from *Cudrania tricuspidata*," *Bioorganic & Medicinal Chemistry Letters*, vol. 15, no. 24, pp. 5548–5552, 2005.
- [39] G.-S. Jeong, D.-S. Lee, and Y.-C. Kim, "Cudraticusxanthone A from *Cudrania tricuspidata* suppresses pro-inflammatory mediators through expression of anti-inflammatory heme oxygenase-1 in RAW264.7 macrophages," *International Immunopharmacology*, vol. 9, no. 2, pp. 241–246, 2009.
- [40] R.-B. An, D.-H. Sohn, and Y.-C. Kim, "Hepatoprotective compounds of the roots of *Cudrania tricuspidata* on tacrine-induced cytotoxicity in Hep G2 cells," *Biological and Pharmaceutical Bulletin*, vol. 29, no. 4, pp. 838–840, 2006.
- [41] M. Yang, G. C. F. Chan, R. Deng et al., "An herbal decoction of *Radix astragali* and *Radix angelicae* sinensis promotes hematopoiesis and thrombopoiesis," *Journal of Ethnopharmacology*, vol. 124, no. 1, pp. 87–97, 2009.
- [42] H. Wu, D. Ye, Y. Bai, and Y. Zhao, "Effect of dry ginger and roasted ginger on experimental gastric ulcers in rats," *China Journal of Chinese Materia Medica*, vol. 15, no. 5, pp. 278–318, 1990.
- [43] S. Malu, G. Obochi, E. Tawo, and B. Nyong, "Antibacterial activity and medicinal properties of ginger (*Zingiber officinale*)," *Global Journal of Pure and Applied Sciences*, vol. 15, no. 3–4, pp. 365–368, 2009.
- [44] S.-T. Huang, C.-Y. Wang, R.-C. Yang, C.-J. Chu, H.-T. Wu, and J.-H. S. Pang, "Wogonin, an active compound in *Scutellaria baicalensis*, induces apoptosis and reduces telomerase activity in the HL-60 leukemia cells," *Phytomedicine*, vol. 17, no. 1, pp. 47–54, 2010.
- [45] H.-B. Li and F. Chen, "Isolation and purification of baicalein, wogonin and oroxylin A from the medicinal plant *Scutellaria baicalensis* by high-speed counter-current chromatography," *Journal of Chromatography A*, vol. 1074, no. 1–2, pp. 107–110, 2005.
- [46] J. Sun, B. Kh Tan, S.-H. Huang, M. Whiteman, and Y.-Z. Zhu, "Effects of natural products on ischemic heart diseases and cardiovascular system," *Acta Pharmacologica Sinica*, vol. 23, no. 12, pp. 1142–1151, 2002.
- [47] S.-C. Wang, T.-Y. Tseng, C.-M. Huang, and T.-H. Tsai, "Gardenia herbal active constituents: applicable separation procedures," *Journal of Chromatography B*, vol. 812, no. 1–2, pp. 193–202, 2004.
- [48] Y.-J. Surh, K.-S. Chun, H.-H. Cha et al., "Molecular mechanisms underlying chemopreventive activities of anti-inflammatory phytochemicals: down-regulation of COX-2 and iNOS through suppression of NF- $\kappa$ B activation," *Mutation Research—Fundamental and Molecular Mechanisms of Mutagenesis*, vol. 480–481, pp. 243–268, 2001.



