

## Retraction

# Retracted: The Curative Effect of Shuangshen Decoction Combined with Immunological Preparations in the Treatment of Pediatric Nephrotic Syndrome and Its Influence on the Rate of Complicated Infection and Recurrence

### Computational and Mathematical Methods in Medicine

Received 31 October 2023; Accepted 31 October 2023; Published 1 November 2023

Copyright © 2023 Computational and Mathematical Methods in Medicine. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This article has been retracted by Hindawi following an investigation undertaken by the publisher [1]. This investigation has uncovered evidence of one or more of the following indicators of systematic manipulation of the publication process:

- (1) Discrepancies in scope
- (2) Discrepancies in the description of the research reported
- (3) Discrepancies between the availability of data and the research described
- (4) Inappropriate citations
- (5) Incoherent, meaningless and/or irrelevant content included in the article
- (6) Peer-review manipulation

The presence of these indicators undermines our confidence in the integrity of the article's content and we cannot, therefore, vouch for its reliability. Please note that this notice is intended solely to alert readers that the content of this article is unreliable. We have not investigated whether authors were aware of or involved in the systematic manipulation of the publication process.

Wiley and Hindawi regrets that the usual quality checks did not identify these issues before publication and have since put additional measures in place to safeguard research integrity.

We wish to credit our own Research Integrity and Research Publishing teams and anonymous and named external researchers and research integrity experts for contributing to this investigation.

The corresponding author, as the representative of all authors, has been given the opportunity to register their agreement or disagreement to this retraction. We have kept a record of any response received.

### References

- [1] N. Han, J. Xie, Q. Li, H. Li, X. Wang, and Y. Sheng, "The Curative Effect of Shuangshen Decoction Combined with Immunological Preparations in the Treatment of Pediatric Nephrotic Syndrome and Its Influence on the Rate of Complicated Infection and Recurrence," *Computational and Mathematical Methods in Medicine*, vol. 2022, Article ID 3355462, 7 pages, 2022.

## Research Article

# The Curative Effect of Shuangshen Decoction Combined with Immunological Preparations in the Treatment of Pediatric Nephrotic Syndrome and Its Influence on the Rate of Complicated Infection and Recurrence

Naiqin Han <sup>1</sup>, Jialing Xie <sup>2</sup>, Qiuying Li <sup>3</sup>, Haixia Li <sup>4</sup>, Xiuzhen Wang <sup>5</sup>,  
and Ying Sheng <sup>6</sup>

<sup>1</sup>Department of Pediatric, Zhangqiu District People's Hospital, Jinan 250200, Shandong Province, China

<sup>2</sup>Department of Pharmacy, Zhangqiu District People's Hospital, Jinan 250200, Shandong Province, China

<sup>3</sup>Department of Imaging, Zhangqiu District People's Hospital, Jinan 250200, Shandong Province, China

<sup>4</sup>Operation Room, Zhangqiu District People's Hospital, Jinan 250200, Shandong Province, China

<sup>5</sup>Department of Nutriology, Zhangqiu District People's Hospital, Jinan 250200, Shandong Province, China

<sup>6</sup>Department of Pediatric, The Third People's Hospital of Qingdao, Qingdao 266041, Shandong Province, China

Correspondence should be addressed to Ying Sheng; shengying@qd3hospital.cn

Received 23 November 2021; Revised 14 December 2021; Accepted 29 December 2021; Published 26 January 2022

Academic Editor: Osamah Ibrahim Khalaf

Copyright © 2022 Naiqin Han et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Background.** To explore the curative effect of Shuangshen Decoction combined with immunological preparations in the treatment of pediatric nephrotic syndrome and its influence on concurrent infection and recurrence rate. **Methods.** Ninety children with nephrotic syndrome were divided into the routine group and the combined group. The routine group received conventional treatment and immune agents, and the combined group was treated with Shuangshen Decoction on the basis of the routine group. The clinical indexes of the two groups were analyzed and followed up. The infection rate and recurrence rate were calculated. **Results.** The TCM syndrome scores in the combined group were significantly lower than those in the routine group. The total effective rate of the combined group was significantly higher than that of the routine group. The recurrence rate and infection rate of the combined group were significantly lower than those of the routine group. The incidence of adverse reactions in the combined group was significantly lower than that in the routine group. **Conclusion.** Shuangshen Decoction combined with immune preparations is effective in treating pediatric nephrotic syndrome and can reduce the incidence of adverse reactions, infection rate, and recurrence rate.

## 1. Introduction

Pediatric nephrotic syndrome is a common pediatric kidney disease in the clinic, which mainly manifests as massive proteinuria, hypoalbuminemia, severe edema, and hypercholesterolemia [1]. Prednisone should be used as soon as possible after the diagnosis of newly diagnosed children. Relapse patients need to adjust the dose and course of treatment of glucocorticoids [2]. However, children who frequently

relapse and receive long-term high-dose glucocorticoid therapy are prone to slow growth and delayed development [3]. Therefore, clinical treatment of primary nephropathy in children is based on strengthening the spleen and kidney and controlling the side effects of Western medicine as the main principles [4]. In the treatment of children with nephrotic syndrome, especially those who have not used hormones, corn silk, plantain, Lysimachia, and Pianch are commonly used [5]. Chinese medicine adjuvant therapy

TABLE 1: TCM syndrome score sheet (prepared based on relevant information and the clinical experience of the tutor).

| Main symptoms        | 0    | 2 points                  | 4 points                      | 6 points                     |
|----------------------|------|---------------------------|-------------------------------|------------------------------|
| Sore waist and knees | None | Occasionally              | Frequently                    | Unbearable                   |
| Chills               | None | Occasionally              | Frequently                    | Afraid of cold               |
| Poor appetite        | None | Reduce food intake by 1/4 | Reduce food intake by 1/4-1/2 | Little or no food            |
| Secondary symptoms   | 0    | 1 point                   | 2 points                      | 3 points                     |
| Tired                | None | Occasionally              | Frequently                    | Bedridden                    |
| Bloating             | None | Slightly full             | Fullness                      | Abdominal distension all day |
| Loose stools         | None | 1 time/d                  | 2-3 times/d                   | >4 times/d                   |
| Nocturia             | None | 2 times/night             | 3 times/night                 | >4 times/night               |
| Shortness of breath  | None | Lack of energy            | Lazy in words                 | Powerless speech             |
| Pale complexion      | None | Dull                      | Pale                          | Pale and dull                |

TABLE 2: Comparison of general information between the two groups before treatment.

| Index                                       | Routine group ( $n = 45$ ) | Combination group ( $n = 45$ ) | $X^2/t$ | $P$   |
|---|----------------------------|--------------------------------|---------|-------|
| Age (year)                                  | $6.42 \pm 1.16$            | $6.55 \pm 1.23$                | 0.516   | 0.607 |
| Course (d)                                  | $11.59 \pm 2.25$           | $11.68 \pm 2.30$               | 0.188   | 0.852 |
| Gender                                      |                            |                                | 0.182   | 0.670 |
| Male  | 25 (55.56)                 | 27 (60)                        |         |       |
| Female                                      | 20 (44.44)                 | 18 (40)                        |         |       |
| Disease progression                         |                            |                                | 0.830   | 0.362 |
| Active nephrotic syndrome                   | 33 (73.33)                 | 29 (64.44)                     |         |       |
| Nephrotic syndrome in remission             | 12 (26.67)                 | 16 (35.56)                     |         |       |
| BMI ( $\text{kg}/\text{m}^2$ )              | $17.52 \pm 2.44$           | $17.69 \pm 2.60$               | 0.320   | 0.750 |
| White blood cell ( $\times 10^9/\text{L}$ ) | $2.76 \pm 0.54$            | $2.85 \pm 0.62$                | 0.734   | 0.465 |
| Cholesterol (mmol/L)                        | $6.88 \pm 1.23$            | $6.59 \pm 1.17$                | 1.146   | 0.255 |
| Triglycerides (mmol/L)                      | $3.17 \pm 0.84$            | $3.20 \pm 0.78$                | 0.176   | 0.861 |

can alleviate the feedback inhibition effect of exogenous corticosteroids on adrenal cortex function [6]. Therefore, a comprehensive treatment plan combining traditional Chinese and Western medicine should be investigated for pediatric nephrotic syndrome.

According to the clinical manifestations of nephrotic syndrome, the traditional Chinese medicine (TCM) theory classifies it as the "edema category," mostly yin water. Weakness of the kidney, spleen, and lung is the main cause of nephrotic syndrome, which is mainly manifested as spleen and kidney deficiency [7]. Shuangshen Decoction contains a variety of herbs. Among them, Morus mulberry can dispel rheumatism, invigorate the liver and kidney, and strengthen muscles and bones [8]. Corn silk can clear the liver, promote choleric, diuresis, and reduce swelling. The keel can condense sweat, consolidate essence, and calm and soothe the nerves. Magnets can calm the nerves, calm the liver, and suppress the yang [9]. Polygonum multiflorum can moisten the intestines, help defecate and nourish the body. Ligusticum chuanxiong can promote blood circulation, relieve pain, and expel wind and dampness [10]. Epimedium can replenish life, nourish vital energy, strengthen muscles and bones, and nourish the kidney. Eucommia can nourish the

liver and kidney and strengthen bones and muscles [11]. However, there are few relevant research reports about the use of Shuangshen Decoction in the treatment of pediatric nephrotic syndrome.

Immunosuppressants are mainly used for patients with frequent relapses of nephrotic syndrome, glucocorticoid dependence, drug resistance, or severe side effects [12]. In small doses of glucocorticoids used every other day, immunosuppressive agents can be used at the same time. Currently, commonly used clinical immunosuppressants are cyclosporine A, tacrolimus (FK506), mycophenolate mofetil, and leflunomide [13]. The dual inhibitory effect of tacrolimus makes it not only able to prevent the occurrence of immunity but also have a good therapeutic effect on the immune response and autoimmune diseases that have appeared [14]. It has been reported that the efficacy of Chinese medicine combined with immune preparations in the treatment of pediatric nephrotic syndrome is significantly higher than that of Western medicine alone [15]. The combination of Chinese and Western medicine not only expands the treatment options but also has better curative effects. Based on this, this study will explore the efficacy of Shuangshen Decoction combined with immune preparations in the

treatment of pediatric nephrotic syndrome and its influence on the rate of complicated infection and recurrence in children.

## 2. Materials and methods

**2.1. Case Grouping.** 90 children with nephrotic syndrome (2018.06-2021.06) admitted to the Zhangqiu District People's Hospital were selected for a retrospective study. Forty-five children who received conventional treatment and immune preparations were set as the routine group. On the basis of the conventional group, 45 children treated with Shuangshen Decoction were included in the combined group. The study was approved by the Ethics Committee of the Zhangqiu District People's Hospital (2018-086).

### 2.2. Inclusion Criteria

- (1) It meets the diagnostic criteria of the "Guidelines for the Diagnosis and Treatment of Common Kidney Diseases in Children (1)" [16]. The child has obvious symptoms of edema with the 24-hour urine protein quantitative  $\geq 50 \text{ mg} \cdot (\text{kg} \cdot 24 \text{ h})^{-1}$ , serum albumin (ALB)  $\leq 25 \text{ g} \cdot \text{L}^{-1}$ , and hypercholesterolemia ( $> 5.7 \text{ mmol} \cdot \text{L}^{-1}$ ).
- (2) According to the "Practical Pediatrics of Traditional Chinese Medicine" [17], the child presented with syndrome of deficiency of both the lung and spleen and syndrome of spleen deficiency and dampness
- (3) It meets the indications for immune preparations. Immunosuppressive agents are mainly used in patients with frequent relapses of nephrotic syndrome, glucocorticoid dependence, drug resistance, or severe side effects
- (4) The family members of the patient agreed to this study

### 2.3. Exclusion Criteria

- (1) Combined with Henoch-Schonlein purpura nephritis, lupus nephritis, hereditary nephropathy, and hepatitis B-related nephritis
- (2) Urinary red blood cells  $> 10/\text{HP}$
- (3) Taking Chinese medicine for the past month
- (4) Combined with renal insufficiency

**2.4. Treatment. Routine group:** (1) Basic treatment measures were used to avoid infection and fatigue in children. Children's daily diet should be careful to avoid spicy, greasy, and irritating food. Salt should be limited during the edema period. (2) Symptomatic treatment measures. Appropriate diuretic drugs should be given, such as HMG-CoA reductase inhibitors or sodium alginate diester. Heparin anticoagulation therapy is given. Infected children need to choose antibiotics for anti-infective treatment [18–20]. (3) Tacrolimus capsule (specification: 1 mg; manufacturer: Sinopharm




FIGURE 1: The ALB level was compared between the routine group and the combined group. \* $P < 0.05$ , \*\* $P < 0.01$ .




FIGURE 2: The 24 h urine protein quantification was compared between the routine group and the combined group. \* $P < 0.05$ , \*\* $P < 0.01$ .

Chuankang Pharmaceutical Co., Ltd.; Sinopharm Zhunzi H20083943) immune preparation treatment. The first medication is 0.05-0.1 mg/kg on an empty stomach (12 h/time). The blood concentration of tacrolimus was rechecked one week later. According to the blood concentration and the

TABLE 3: Comparison of TCM syndrome scores between the two groups.

| Group                     | Routine ( $n = 45$ ) | Combination ( $n = 45$ ) | $t$    | $P$    |
|---------------------------|----------------------|--------------------------|--------|--------|
| Before treatment          | 15.15 $\pm$ 1.20     | 15.20 $\pm$ 1.22         |        |        |
| 3 months after treatment  | 12.47 $\pm$ 1.04     | 9.84 $\pm$ 0.72          | 13.948 | <0.001 |
| 6 months after treatment  | 9.91 $\pm$ 0.53      | 7.22 $\pm$ 0.41          | 26.930 | <0.001 |
| 12 months after treatment | 8.84 $\pm$ 0.48      | 4.97 $\pm$ 0.51          | 37.068 | <0.001 |

condition of the child, the dosage was adjusted appropriately to maintain the blood concentration of 5-10  $\mu\text{g/L}$ . The treatment cycle is 1 year.

**Combined group:** Shuangshen Decoction was added on the basis of the routine group. Shuangshen Decoction prescription: 30 g Morus alba, 30 g corn silk, 30 g raw keel (first decoction), 30 g magnet (first decoction), 24 g Polygonum multiflorum, 9 g Chuanxiong, 9 g Epimedium, and 9 g Eucommia. **Decoction method:** One dose of the medicine was put into a sand jar, and cold water was added to the surface. After soaking for 30 minutes, the medicine was heated to boiling water. Then, the medicine was cooked for 25 minutes on medium-low heat. The residue was filtered to get the juice. About 100-150 mL is the first medicine. Boiled water was added to 1 finger higher than the noodles. After the water boils, the medicine was cooked for 30 minutes. Then, the residue was filtered to get the juice, about 200 mL. After mixing the first and second medicines, the medicine juice was taken in three warm doses, one dose per day. >12-year-old children need 100-120 mL/time, 3 times a day. Children aged 7-12 need 80-100 mL/time, 3 times a day. 3-6-year-old children need 50-80 mL/time, 3 times a day. The medicine juice should be taken half an hour before or after a meal.

**2.5. Observation Index.** The patient's age, course of disease, gender, body mass index (BMI), disease progression, white blood cell count, cholesterol, triglycerides, and other general information were recorded. The patient's albumin (ALB) level was detected by immunoassay. And the patient's 24 h urine protein quantitative level was recorded. According to the severity of the primary and secondary symptoms, the patient's TCM syndrome score was evaluated. The specific information about the TCM syndrome score was shown in Table 1.

The clinical efficacy of patients was evaluated according to the "Standards for Diagnosis and Efficacy of TCM Diseases and Syndromes" [21] and changes in clinical signs, 24 h urine protein quantification, and recovery of ALB indicators. Significantly effective means the obvious improvement of TCM symptoms and clinical signs and 24 h urine protein quantification  $\leq 1.0$  g,  $\text{ALB} \geq 35$  g/L. Effective means that TCM symptoms and clinical signs have improved, and  $1.0\text{g} < 24$  h urine protein  $\leq 3.0$  g,  $30$  g/L  $\leq \text{ALB} < 35$  g/L. Invalid means that TCM symptoms and clinical signs have not improved or worsened, and 24 h urine protein quantitative  $> 3.0$  g and  $\text{ALB} < 30$  g/L. The total effective rate of treatment = (significantly effective + effective)/total  $\times 100\%$ .

The occurrence of infections and adverse reactions during the treatment of patients was recorded. The types of infections

mainly include upper respiratory tract infections, lower respiratory tract infections, urinary tract infections, and gastrointestinal infections. The patients were followed up for no less than 6 months to review the patient's recurrence.

**2.6. Statistical Analysis.** SPSS22.0 Software was used to calculate the differences between groups. The figure was drawn using GraphPad Prism 7 (GraphPad Software, San Diego, USA). Research data includes counts and measurements, expressed in the form of  $[n (\%)]$  and mean  $\pm$  standard deviation (SD), respectively. Chi-squared test and Student  $t$  test were used for difference analysis.  $P < 0.05$  indicates that there is a statistical difference between the groups.

### 3. Results

**3.1. Comparison of General Information.** There was no statistically significant difference in general information between the routine group and the combined group ( $P > 0.05$ , Table 2). Therefore, a controlled study can be conducted.

**3.2. Comparison of Serum Albumin and 24 h Urine Protein.** The ALB level of the two groups was significantly increased after treatment. However, the ALB level of the combined group was increased significantly after 1 and 3 months of treatment compared with the routine group ( $P < 0.05$ , Figure 1). The 24 h urine protein quantification after treatment in the two groups was significantly decreased. However, the 24 h urine protein quantification after 1-month and 3-month treatment in the combined group was significantly lower than that in the routine group ( $P < 0.05$ , Figure 2).


**3.3. Comparison of TCM Syndrome Points.** The scores of TCM syndromes in the combined group were significantly lower than those in the routine group after treatment ( $P < 0.05$ , Table 3). The difference is statistically significant.

**3.4. Comparison of Clinical Efficacy.** The total effective rate of the combined group was significantly higher than that of the routine group ( $P < 0.05$ , Table 4). The difference between the groups was statistically significant.

**3.5. Comparison of Infection Rate and Recurrence Rate.** The recurrence rate and infection rate of patients in the combined group were significantly lower than those in the routine group ( $P < 0.05$ , Figure 3). The incidence of adverse reactions such as nausea and vomiting, acne, hypertension, and coagulopathy in the combined group was significantly lower than that in the routine group ( $P < 0.05$ , Table 5).

TABLE 4: Comparison of the clinical efficacy of the two groups [ $n$  (%)].

| Group                | Routine<br>( $n = 45$ ) | Combination<br>( $n = 45$ ) | $\chi^2$ | $P$   |
|----------------------|-------------------------|-----------------------------|----------|-------|
| Invalid              | 11 (24.44)              | 3 (6.67)                    |          |       |
| Effective            | 19 (42.22)              | 20 (44.44)                  |          |       |
| Markedly effective   | 15 (33.33)              | 22 (48.89)                  |          |       |
| Total effective rate | 34 (75.56)              | 42 (93.33)                  | 5.414    | 0.020 |

FIGURE 3: The infection rate and recurrence rate were compared between the routine group and the combined group. \* $P < 0.05$ , \*\* $P < 0.01$ .TABLE 5: Comparison of the incidence of adverse reactions between the two groups [ $n$  (%)].

| Group               | Routine<br>( $n = 45$ ) | Combination<br>( $n = 45$ ) | $\chi^2$ | $P$   |
|---------------------|-------------------------|-----------------------------|----------|-------|
| Nausea and vomiting | 8 (17.78)               | 2 (4.44)                    | 4.050    | 0.044 |
| Acne                | 5 (11.11)               | 0 (0)                       | 5.294    | 0.021 |
| Hypertension        | 6 (13.33)               | 1 (2.22)                    | 3.873    | 0.049 |
| Coagulopathy        | 4 (8.89)                | 0 (0)                       | 4.186    | 0.041 |

#### 4. Discussion

Pediatric nephritic syndrome is second only to acute nephritis in childhood nephropathy. Conventional treatment drugs are mostly glucocorticoids and cytotoxic drugs. The short-term curative effect is remarkable. However, the long-term application of the drug has serious adverse reactions, which can easily cause symptoms such as infection, growth inhibition, water and sodium retention, and central obesity. These adverse reactions can cause the recurrence of pediatric nephritic syndrome, and it is difficult to completely cure [22]. In this study, we found that the ALB levels of the two groups after treatment were significantly increased. However, the increase in the ALB level of the combined group after 1 month and 3 months of treatment was significantly higher than that of the routine group. The 24 h urine protein quantification of the two groups was decreased significantly

after treatment. However, the 24 h urine protein quantification after 1-month and 3-month treatment in the combined group was significantly lower than that in the routine group. Xu et al. also reported similar results [23]. These findings indicate that Shuangshen Decoction combined with immune preparations have a more obvious regulatory effect on the quantitative level of ALB and 24 h urine protein in children. Therefore, Shuangshen Decoction combined with immune preparations is conducive to disease control and guiding prognosis.

The scores of TCM syndromes after treatment in the combined group were significantly lower than those in the routine group. Traditional Chinese medicine believes that children's nephrotic syndrome is mostly caused by edema, proteinuria, and other symptoms. Lungs dominate qi, and poor qi movement can cause edema. The spleen is the source of biochemistry, and it is a syndrome of phlegm and dampness block caused by the failure of the spleen [24]. The kidney is a visceral organ with water, and the lack of kidney qi causes abnormal urination or edema. Therefore, the main treatment is to warm the spleen and kidney, promote blood circulation to remove stasis, remove dampness, and promote qi [25]. Shuangshen Decoction contains *Morus vulgaris*, corn silk, keel, magnet, *Polygonum multiflorum*, *Chuanxiong*, *Epimedium*, and *Eucommia*. All medicines work together to nourish the kidney and clear away heat [26]. All these results demonstrate that Shuangshen Decoction is beneficial to the treatment of pediatric nephritic syndrome.

In order to improve the curative effect of pediatric nephrotic syndrome and reduce infection and recurrence, this study implemented a comprehensive treatment plan of immune preparations combined with Shuangshen Decoction. Immunosuppressive agents are often used clinically to treat patients with refractory nephropathy or severe hormonal side effects [27]. The dual inhibitory effect of tacrolimus not only makes it able to prevent the occurrence of immunity but also has a good therapeutic effect on the immune response and autoimmune diseases that have appeared [28]. In present study, the overall effect of tacrolimus combined with Shuangshen Decoction was good. Moreover, the recurrence rate and infection rate of patients in the combined group were significantly lower than those in the routine group. The incidence of adverse reactions in the combined group was significantly lower than that in the routine group. These results have not been reported in previous studies. Taken together, Shuangshen Decoction combined with immune preparations are safe and effective in the treatment of pediatric nephrotic syndrome and can also greatly reduce the probability of infection and recurrence.

#### 5. Conclusion

In summary, Shuangshen Decoction combined with immune preparations has a significant effect on the treatment of children with nephrotic syndrome. It can increase serum ALB and 24-hour urine protein levels and effectively reduce the incidence of adverse reactions, infection rates, and recurrence rates in children. Due to the small sample

size in this study, the sample size will still need to be expanded for further confirmation in the future.

## Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

## Conflicts of Interest

The authors declare that they have no competing interests.

## References

- [1] A. Bierzynska and M. A. Saleem, "Deriving and understanding the risk of post-transplant recurrence of nephrotic syndrome in the light of current molecular and genetic advances," *Pediatric Nephrology*, vol. 33, no. 11, pp. 2027–2035, 2018.
- [2] B. A. Kerlin, N. B. Blatt, B. Fuh et al., "Epidemiology and risk factors for thromboembolic complications of childhood nephrotic syndrome: a Midwest Pediatric Nephrology Consortium (MWPNC) study," *The Journal of pediatrics*, vol. 155, no. 1, pp. 105–110.e1, 2009.
- [3] H. X. Chen, Q. Cheng, F. Li et al., "Efficacy and safety of tacrolimus and low-dose prednisone in Chinese children with steroid-resistant nephrotic syndrome," *World journal of pediatrics: WJP*, vol. 16, no. 2, pp. 159–167, 2020.
- [4] N. Larkins, S. Kim, J. Craig, and E. Hodson, "Steroid-sensitive nephrotic syndrome: an evidence-based update of immunosuppressive treatment in children," *Archives of Disease in Childhood*, vol. 101, no. 4, pp. 404–408, 2016.
- [5] A. Sinha, A. Saha, M. Kumar et al., "Extending initial prednisolone treatment in a randomized control trial from 3 to 6 months did not significantly influence the course of illness in children with steroid-sensitive nephrotic syndrome," *Kidney International*, vol. 87, no. 1, pp. 217–224, 2015.
- [6] K. V. Lieberman and A. Pavlova-Wolf, "Adrenocorticotrophic hormone therapy for the treatment of idiopathic nephrotic syndrome in children and young adults: a systematic review of early clinical studies with contemporary relevance," *Journal of Nephrology*, vol. 30, no. 1, pp. 35–44, 2017.
- [7] X. Wen, W. Wang, M. Zheng, and B. Song, "The potential mechanism of Astragali radix in the treatment of children with nephrotic syndrome," *Translational pediatrics*, vol. 10, no. 9, pp. 2298–2306, 2021.
- [8] W. X. J. Lim, C. S. Gammon, P. von Hurst, L. Chepulis, and R. A. Page, "A narrative review of human clinical trials on the impact of phenolic-rich plant extracts on prediabetes and its subgroups," *Nutrients*, vol. 13, no. 11, 2021.
- [9] J. H. Ong, J. A. Koh, H. Cao et al., "Purification, identification and characterization of antioxidant peptides from corn silk tryptic hydrolysate: an integrated in vitro-in silico approach," *Antioxidants*, vol. 10, no. 11, 2021.
- [10] C. Z. Liu, X. L. Jiang, Q. Z. Cai, L. Y. Zhou, and Q. Yang, "Screening of plant growth-promoting rhizobacteria and its effect on seed germination of *Polygonum multiflorum*," *Zhongguo Zhong yao za zhi = Zhongguo zhongyao zazhi = China journal of Chinese materia medica*, vol. 46, no. 20, pp. 5247–5252, 2021.
- [11] S. B. Zhao, S. Y. Huang, Z. Pan et al., "The compatibility of coix leaves and epimedium against fatigue and hypoxia tolerance," *Zhongguo ying yong sheng li xue za zhi = Zhongguo yingyong shenglixue zazhi = Chinese journal of applied physiology*, vol. 37, no. 5, pp. 510–513, 2021.
- [12] T. C. von Groote, G. Williams, E. H. Au et al., "Immunosuppressive treatment for primary membranous nephropathy in adults with nephrotic syndrome," *The Cochrane database of systematic reviews*, vol. 11, 2021.
- [13] H. Yasuda, Y. Fukusumi, V. Ivanov, Y. Zhang, and H. Kawachi, "Tacrolimus ameliorates podocyte injury by restoring FK506 binding protein 12 (FKBP12) at actin cytoskeleton," *FASEB Journal : Official Publication of the Federation of American Societies for Experimental Biology*, vol. 35, no. 11, p. e21983, 2021.
- [14] K. Miura, T. Ando, S. Kanda et al., "Response to steroid and immunosuppressive therapies may predict post-transplant recurrence of steroid-resistant nephrotic syndrome," *Pediatric Transplantation*, p. e14103, 2021.
- [15] S. Fujinaga, D. Hirano, A. Mizutani et al., "Predictors of relapse and long-term outcome in children with steroid-dependent nephrotic syndrome after rituximab treatment," *Clinical and Experimental Nephrology*, vol. 21, no. 4, pp. 671–676, 2017.
- [16] A. Abedini, "Is medium dose of cyclosporine A effective in treatment of children with steroid-dependent nephrotic syndrome with reduction in corticosteroids dose consumption?," *Clinical and Experimental Nephrology*, vol. 23, no. 2, pp. 287–288, 2019.
- [17] S. Fujinaga and T. Nishino, "Is cytokine-release syndrome the cause of rituximab treatment-related infusion reactions in children with nephrotic syndrome?," *Impact of anti-rituximab antibodies, Pediatric nephrology*, vol. 33, no. 6, pp. 1097–1098, 2018.
- [18] S. Fujinaga and Y. Urushihara, "Impact of interrupted cyclosporine treatment on the development of chronic nephrotoxicity in children with steroid-dependent nephrotic syndrome," *Pediatric Nephrology*, vol. 32, no. 8, pp. 1469–1470, 2017.
- [19] U. Querfeld and L. T. Weber, "Treatment strategies for children with steroid-dependent nephrotic syndrome: in need of controlled studies," *Pediatric Nephrology*, vol. 33, no. 12, p. 2391, 2018.
- [20] H. Lin, L. Chen, S. Wen et al., "Early diagnosis and successful treatment of cytomegalovirus peritonitis in children with primary nephrotic syndrome: case series and literature review," *Renal Failure*, vol. 42, no. 1, pp. 776–784, 2020.
- [21] F. Nakubulwa, R. C. Lusoby, A. Batte et al., "Prevalence and predictors of ocular complications among children undergoing nephrotic syndrome treatment in a resource-limited setting," *BMC Ophthalmology*, vol. 21, no. 1, p. 55, 2021.
- [22] A. Pasini, E. Benetti, G. Conti et al., "The Italian Society for Pediatric Nephrology (SINePe) consensus document on the management of nephrotic syndrome in children: part I-diagnosis and treatment of the first episode and the first relapse," *Italian Journal of Pediatrics*, vol. 43, no. 1, p. 41, 2017.
- [23] L. Xu, F. Yang, Y. Ge, G. Guo, and Y. Wang, "Crosslinking porcine aortic valve by radical polymerization for the preparation of BHVs with improved cytocompatibility, mild immune response, and reduced calcification," *Journal of Biomaterials Applications*, vol. 35, no. 9, pp. 1218–1232, 2021.
- [24] D. Olczak-Kowalczyk, M. Roszkowska-Blaim, M. Dabkowska et al., "Enzymatic activity of *Candida* spp. from oral cavity and urine in children with nephrotic syndrome," *Advances in Experimental Medicine and Biology*, vol. 1022, pp. 63–70, 2017.

- [25] J. X. Liu, X. Han, Y. G. Xu, and R. Ma, "Mechanism of serum containing SSTG on calcium overload in cardiomyocytes injured by hypoxia and reoxygenation," *Zhongguo Zhong Yao Za Zhi*, vol. 31, no. 12, pp. 995–998, 2006.
- [26] J. X. Liu, X. Han, and X. B. Ma, "Effect of Shuangshen Tongguan recipe on nuclear factor-kappa B signal pathway and myocardial junction-mediated intercellular communication in acute myocardial ischemia/reperfusion injured model rats," *Zhongguo Zhong Xi Yi Jie He Za Zhi*, vol. 25, no. 3, pp. 228–231, 2005.
- [27] S. Qiu, H. Zhang, S. Yu et al., "Clinical manifestations, prognosis, and treat-to-target assessment of pediatric lupus nephritis," *Pediatric Nephrology*, 2021.
- [28] M. A. Mazumder, S. Gulati, A. S. Narula, D. Shehwar, and I. M. Mir, "Tacrolimus-induced acute pancreatitis and diabetic ketoacidosis (DKA) in pediatric kidney transplant recipient," *Pediatric Transplantation*, p. e14194, 2021.