

Retraction

Retracted: *CASC1* Expression in Bladder Cancer Is Regulated by Exosomal miRNA-150: A Comprehensive Pan-Cancer and Bioinformatics Study

Computational and Mathematical Methods in Medicine

Received 5 December 2023; Accepted 5 December 2023; Published 6 December 2023

Copyright © 2023 Computational and Mathematical Methods in Medicine. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This article has been retracted by Hindawi, as publisher, following an investigation undertaken by the publisher [1]. This investigation has uncovered evidence of systematic manipulation of the publication and peer-review process. We cannot, therefore, vouch for the reliability or integrity of this article.

Please note that this notice is intended solely to alert readers that the peer-review process of this article has been compromised.

Wiley and Hindawi regret that the usual quality checks did not identify these issues before publication and have since put additional measures in place to safeguard research integrity.

We wish to credit our Research Integrity and Research Publishing teams and anonymous and named external researchers and research integrity experts for contributing to this investigation.

The corresponding author, as the representative of all authors, has been given the opportunity to register their agreement or disagreement to this retraction. We have kept a record of any response received.

References

- [1] H. Luo, C. Xu, B. Ge, and T. Wang, “*CASC1* Expression in Bladder Cancer Is Regulated by Exosomal miRNA-150: A Comprehensive Pan-Cancer and Bioinformatics Study,” *Computational and Mathematical Methods in Medicine*, vol. 2022, Article ID 8100325, 18 pages, 2022.

Research Article

CASC1 Expression in Bladder Cancer Is Regulated by Exosomal miRNA-150: A Comprehensive Pan-Cancer and Bioinformatics Study

Huarong Luo,¹ Chengdang Xu,¹ Bujun Ge^{1,2} ,² and Tianru Wang¹ 

¹Department of Urology, Tongji Hospital, Tongji University School of Medicine, Shanghai, China

²Department of General Surgery, Tongji Hospital, Tongji University School of Medicine, Shanghai, China

Correspondence should be addressed to Bujun Ge; gebujun@126.com and Tianru Wang; 1811183@tongji.edu.cn

Received 3 March 2022; Revised 30 May 2022; Accepted 1 June 2022; Published 5 July 2022

Academic Editor: Min Tang

Copyright © 2022 Huarong Luo et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This study explored the role of cancer susceptibility 1 (CASC1) in tumorigenesis and development as well as the key pathways affecting bladder cancer progression. CASC1 was examined in various normal tissues in humans using the HPA database to quantify its expression level and subcellular localization. CASC1 is abundantly expressed in tumor tissues, primarily in cytoplasmic vesicles and stroma. TIMER2 was used to analyze the correlation between CASC1 expression levels and the types of infiltrates associated with immune cells and immunosuppressive cells. MDSC, Treg, M2, and CAF were significantly correlated with CASC1 expression in various tumors. Comparing patients with and without CASC1 mutation, those with CASC1 mutation had worse overall survival, progression-free survival, and disease-free survival. The correlation between has-miR-150 and CASC1 (for the case of bladder cancer) was then analyzed, and the related ceRNA network was mapped. A negative relationship between CASC1 expression and has-miR-150 expression was found in cases of bladder cancer. And the presence of miR-150-targeted CASC1 may be associated with bladder cancer progression. CASC1 is expressed at elevated levels in various tumor tissues, and it is associated with tumorigenesis and development. Exosomes containing miR-150-targeted CASC1 may affect the progression of bladder cancer.

1. Introduction

A total of 430,000 new cases of bladder cancer are reported every year, with 165,000 deaths related to the disease [1]. Based on the 2015 China Cancer Data, the incidence and mortality related to bladder cancer are on the rise every year [2, 3]. The prognosis for patients with bladder cancer remains poor despite the development of new treatments due to high recurrence and metastasis rates [4]. Therefore, the development of new therapeutic targets requires an understanding of the molecular mechanisms associated with bladder cancer.

Cancer susceptibility 1 (CASC1) regulates microtubule dynamics and plays an important role in mitosis in tumor cells [5]. It has been demonstrated that DEGs upregulated by bladder cancer are frequently linked to mitotic spindle assembly checkpoints [5]. RNAi inhibition of CASC1

expression can reduce tumor growth in vivo and increase survival rates for patients [6–9]. To improve the treatment methods and prognosis of bladder cancer patients, we need to explore the regulatory pathway of CASC1, which affects the progression of bladder cancer.

The specific mechanism associated with the regulation pathway of CASC1 that affects the progression of bladder cancer is still unknown. A study found that CASC1 influenced the function of pulmonary adenoma susceptibility 1 locus (PAS1), which is the regulator of lung tumors [10]. CASC1 was found to be upregulated in malignant tumors [11]. Proteins that target CASC1 and the genes related to CASC1 can be potentially important molecules that influence cancer incidence and development.

There is increasing evidence that the ceRNA regulatory network plays a role in cancer development and occurrence [12–14]. It might be possible to uncover the mechanism by

which *CASC1* influences bladder cancer progression by studying the ceRNA network. The bioinformatics analysis technique can be used to investigate the role of ceRNA regulatory network in cancer development. In previous research, bioinformatics analysis was used to explore how ceRNAs contribute to the development of drug-resistant non-small cell lung cancers (NSCLCs) [15]. Therefore, bioinformatics analysis of ceRNA regulatory networks could provide more insight into the mechanisms whereby *CASC1* causes bladder cancer.

MicroRNAs (miRNAs) influence various physiological and pathological processes in the body through ceRNA regulatory networks [16–20]. Moreover, miRNAs may be released by cells and loaded into exosomes, acting as messengers for transmitting information between cells and participating in tumor development [21, 22]. A number of studies have shown that exosomal miRNAs play a crucial role in the development of bladder cancer [23]. Researchers have found that miRNA-150 affects bladder cancer prognosis, and we speculate that *CASC1* may be associated with miRNA-150 [24]. By analyzing pan-cancer data and using bioinformatics, we are attempting to uncover how *CASC1* affects bladder cancer progression. As part of our research, we hope to identify potential therapeutic targets, prevent bladder cancer progression, and improve patient outcomes.

2. Methods

2.1. Mutants, Location-Related Diseases, and Tissue Specificity of *CASC1*. The Human Protein Atlas (HPA) (<https://www.proteinatlas.org/>) consists of a dataset analyzed to map proteins in human tissue, organ, and cell. Based on the HPA database, we investigated the expression levels of *CASC1* gene in different normal human tissues and the localization of *CASC1* protein in JURKAT, SuSa, and U-2 OS cell lines. The OPENTARGET platform (<https://www.targetvalidation.org/>) integrated genetics and multiple omics methods were used to identify the role of the genes in the progression and occurrence of the disease. We used the OPENTARGET platform to mine disease networks associated with *CASC1*. GPS-Prot (<http://gpsprot.org/index.php>) is an online platform that helps visualize protein-protein interactions (PPIs). We used the GPS-PORT database to reveal the PPI network associated with *CASC1* protein. PROTTER (<https://wlab.ethz.ch/protter/help/>) is an interactive protein characterization visualization tool, the data of which can be found in the UniprotKB database. The PROTTER database was used to visualize the topological structure of the *CASC1* protein.

2.2. Differential Expression and Prognostic Value of *CASC1* in Tumors. RNA-seq data was obtained from the TCGA and Genotype-Tissue Expression (GTEx) databases, and clinical matching data was acquired from the TCGA database [25, 26]. The Wilcoxon rank-sum test was conducted to compare the expression levels of *CASC1* observed in tumor and normal groups. It was also conducted to compare the expression levels observed at different pathological stages. And cancer samples of each type were divided into

two high–low expression groups according to the expression level of *CASC1*.

The Kaplan-Meier (KM) plot was plotted using the *Survminer* (version 0.4.9) package, and the log-rank test was conducted to compare the survival differences between the two groups. The *survival* package (version 3.2-10) was used to conduct the univariate Cox proportional risk regression analysis. All analyses were based on R software (version 3.6.3).

2.3. The Relationship between *CASC1* Expression Levels and Tumor Immunity. TIMER2 (<http://timer.cistrome.org/>) was used to analyze the correlation between *CASC1* expression level and the six types of immune cells, such as B cells, CD8+ T cells, CD4+ T cells, macrophages, neutrophils, and dendritic cells (DC). Additionally, we studied the relationship between *CASC1* expression levels and four immunosuppressive cells (that promote T cell rejection), including marrow-derived suppressor cells (MDSCs), cancer-associated fibroblasts (CAFs), tumor-associated macrophages M2 subtypes (M2-tams), and regulatory T cells (Treg). A Spearman's rank correlation test was used to calculate the partial correlation (*cor*) and *p* values. Additionally, the TIDE database (<http://tide.dfci.harvard.edu/>) was used to compare the efficiency of *CASC1* and standard biomarkers in predicting the immune state during treatment response to immune checkpoint inhibitors (ICBs).

2.4. Mutation Characteristics of *CASC1*. CBioPortal (<http://www.cbioportal.org>) is a cancer genomics database, which was used to observe the mutation frequencies and mutation sites of *CASC1* associated with various cancer types. Furthermore, we investigated the relationship between *CASC1* mutations and patient survival rates (OS, DFS, and PFS). The expression levels of *CASC1* were also studied in relation to the different copy number variation types.

2.5. Analysis of Related Genes and Functions Associated with *CASC1* and Bladder Cancer. STRING (<https://string-db.org/>) database was used to obtain the names of the organism and protein. Detailed information about protein interactions was obtained from STRING database. A minimum interaction score of 0.4 was required (medium confidence).

GEPIA2 (<http://gepia2.cancer-pku.cn/#index>) is an online tool for understanding TCGA gene expression and survival analysis. For bladder cancer, we used the GEPIA2 Similar Gene Detection module to identify the top 100 genes associated with *CASC1*. Using the correlation analysis module, the correlation scatter diagram (Pearson correlation coefficient) representing the relationship between *CASC1* and the five most related genes was generated. We further analyzed the first 100 genes associated with *CASC1* using the clusterProfiler package. GO terms with *q*-values less than 0.05 were considered significantly enriched. The scatter plot representing the correlation between has-miR-150 and *CASC1* (for bladder cancer) was created using ggplot2, ggExtra, and ggpubr.

In order to predict lncRNAs targeted by has-miR-150, we used the StarBase database (version 3), and Cytoscape







FIGURE 1: Continued.



(c)



(d)

FIGURE 1: Continued.




FIGURE 1: Mutants, location-related diseases, and tissue specificity of *CASCI*. (a) Cancer susceptibility candidate gene 1 (*CASCI*) protein topology revealed that it consisted of two natural variants, Arg33Ser (from arginine to serine) and Ala633Glu (from alanine to glutamate); (b) *Cascl*-related disease networks; (c) immunofluorescence staining images of *CASCI* in JURKAT, SuSa, and U-2 OS cells; (d) expression of *CASCI* in healthy human tissues; (e) functional companion of *CASCI*.

(version 3.8.0) to plot the ceRNA network. For *CASCI* and miR-150 expression, we divided the samples by the median expression levels calculated for bladder cancer. The clusterProfiler was used to perform the KEGG gene set enrichment analysis (GSEA).


3. Results

3.1. Mutants, Location-Related Diseases, and Tissue Specificity of *CASCI*. The protein topology of *CASCI* presented two natural variants (Arg33Ser and Ala633Glu) (Figure 1(a)). *CASCI* was primarily associated with neurological diseases, respiratory diseases, tumors, and congenital diseases (Figure 1(b)). According to immunofluorescence images of *CASCI* obtained from the HPA database, *CASCI* is primarily distributed in the intracellular cytoplasmic matrix and vesicles (Figure 1(c)). The expression levels of *CASCI* in human tissues (mRNA) are shown in Figure 1(d). High levels of expression were detected in testes, fallopian tubes, and choroid plexuses. *CASCI* has also been demonstrated to interact with several proteins (MYO9A, HADHA, RUNDC1, LRIF1, BARD1, and PPP1CA) (Figure 1(e)).


3.2. Differential Expression and Prognostic Value of *CASCI* in Tumors. *CASCI* expression in tumor and normal samples from the TCGA and GTEx databases was compared (Figure 2(a)). The results revealed that *CASCI* was highly expressed in tumor tissues associated with UCEC, THYM, PAAD, OV, LGG, LAML, KIRP, GBM, DLBC, CHOL, and BRCA. In order to study the prognostic value of *CASCI*, 33 TCGA cancer types were analyzed using the univariate

Cox proportional risk regression method (Figure 2(b)). The Log-rank test was also used to compare the survival differences for the groups with high and low levels of *CASCI* expression (Figure 2(c)). ACC, CESC, GBMLGG, KIRC, MESO, SKCM, UCS, and UVM had significantly different survival rates between high and low *CASCI* expression groups. In GBMLGG and UCS, the survival prognosis was better in the group with low *CASCI* expression than in the group with high *CASCI* expression. For ACC, CESC, KIRC, MESO, SKCM, and UVM, the high *CASCI* expression group had a better survival prognosis than the low expression group. A difference in expression levels of *CASCI* was also assessed at various tumor pathological stages (Figure 2(d)). There were significant differences in the expression levels of BRCA, KIRC, THCA, and LUSC at different stages of disease. In brief, *CASCI* appeared to become overexpressed during the early stages of tumor development.

3.3. Relationship between *CASCI* Expression Levels and Tumor Immunity. *CASCI* expression was correlated with tumor immune cell infiltration (including CD4+ T cells, B cells, CD8+ T cells, neutrophils, macrophages, and DCs) in only 4 out of 39 TCGA tumor types (LGG, LIHC, PRAD, and LUSC) (Figure 3(a)). For HNSC, KIRC, and SKCM, the level of immune cell infiltration strongly correlated with *CASCI* expression. Furthermore, the extent of immune cell infiltration observed in the case of THYM correlated negatively with the expression levels of *CASCI*. Additionally, we examined the relationship between the level of expression of *CASCI* and the infiltration of four immunosuppressive cells (M2-TAMs, CAFs, MDSCs, and Treg cells) (Figure 3(b)). *CASCI* expression in KIRP, LIHC, OV,




(a)



(b)

FIGURE 2: Continued.



(c)

FIGURE 2: Continued.




FIGURE 2: Differential expression and prognostic value of *CASC1* in tumors. (a) Differential expression of *CASC1* in 30 tumor tissues compared to normal/paracancer tissues (\log_2 (TPM+1)). These data were obtained from the TCGA and GTEx databases. (b) Forest map presents the results obtained following the univariate cox regression method for the analysis of *CASC1* in 34 types of TCGA tumors. (c) OS KM-plot for high and low expression groups of *CASC1* (grouped according to median value). (d) The expression of *CASC1* differed with the tumor pathologic stage. ns, $p \geq 0.05$; * $p < 0.05$; ** $p < 0.01$; and *** $p < 0.001$.

SKCM-Primary, THCA, and THYM was significantly correlated with the extent of tumor invasion recorded for MDSC. *CASC1* expression in ESCA, HNSC, LUAD, LUSC, PAAD, SKCM, STAD, UCEC, UCS, and UVM is significantly inversely correlated with Treg level. There was a significant positive correlation between the expression levels of *CASC1* in BRCA, COAD, HNSC, LGG, LIHC, LUAD, PAAD, PRAD, and THYM and the extent of tumor invasion recorded for M2. *CASC1* expression was positively correlated with the extent of tumor invasion in ACC, BRCA, CESC, CHOL, COAD, ESCA, HNSC, LUAD, LUSC, and MESO. *CASC1* was also analyzed according to the efficacy predictive power of its ICB subgroups (Figure 3(c)). Seven subcohorts of the ICB showed AUC values greater than 0.5. This demonstrated *CASC1* as a predictive marker and its superiority over other biomarkers. Therefore, *CASC1* exhibited a better predictive power in the Uppaluri2020_PD1_HNSC_post.

3.4. Mutation Characteristics of *CASC1*. Genetic variation significantly influences the initiation of cancer and immune tolerance. Based on TCGA database (10,967 samples), we used cBioPortal to detect genetic variations in *CASC1* and to count the mutation frequency in *CASC1*. *CASC1* mutations were found to exceed 8% in esophagogastric adenocarcinoma, ovarian epithelial tumor, and non-seminomatous germ cell tumors (Figure 4(a)). In total, 140 mutated loci (110 missense, 23 truncating, 5 splicing, and 2 SV/Fusion) were identified (Figure 4(b)). In patients with *CASC1* mutations, overall survivals ($p < 0.001$) (Figure 4(c)), disease-free survivals ($p < 0.001$) (Figure 4(d)), and progression-free survivals ($p < 0.001$) (Figure 4(e)) were negatively correlated with those without *CASC1* mutations. Furthermore, there were significant differences in the expression of mRNA associated with different CNA subtypes of *CASC1* (Figure 4(f)). Compared to amplification, the level of expression was relatively higher in deep deletion.


3.5. Analysis of Related Genes and Functions of *CASC1* in Bladder Cancer. *CASC1* interaction proteins and genes asso-

ciated with *CASC1* expression may influence important molecular mechanisms involved in cancer genesis and progression. STRING identified a total of 10 *CASC1* interacting proteins (Figure 5(a)). We identified the first 100 genes co-expressed with *CASC1* associated with bladder cancer using GEPIA2. In the first five genes, we found EFCAB12 ($R = 0.59$), CFAP126 ($R = 0.56$), RIBC1 ($R = 0.54$), SPAG8 ($R = 0.54$), and DNAA1 ($R = 0.53$) (Figures 5(b)–(f)). By analyzing the first 100 genes using GO enrichment analysis, we found that 30 GO terms exhibited a significant correlation. We visualized the first 10 significant biological processes (BP) and cellular components (CC) (Figure 5(g)). The level of expression of has-miR-150 was reported to correlate negatively with *CASC1* in bladder cancer ($R = -0.11$) (Figure 5(h)).


The miRWALK database confirmed that has-miR-150 targets *CASC1*. All potential targets for has-miR-150 were predicted using the StarBase database. A network of 66 targeted long noncoding RNAs (lncRNAs), including has-miR-150 and *CASC1*, was constructed (Figure 6(a)). The samples were divided into high and low expression groups based on their median expression levels of *CASC1* and has-miR-150. KEGG analysis was also performed, and 37 significantly enriched pathways were associated with *CASC1* and 67 significantly enriched pathways with has-miR-150. It was found that 10 KEGG pathways were associated with *CASC1* and has-miR-150 (Figure 6(b)). In GSEA, pathways associated with *CASC1* and has-miR-150 were visualized (Figures 6(c)–(d)). They were complement and coagulation cascades, drug metabolism cytochrome p450, hematopoietic cell lineage, Leishmania infection, metabolism of xenobiotics by cytochrome p450, natural killer cell-mediated cytotoxicity, proteasome, ribosome, systemic lupus erythematosus, and viral myocarditis.

4. Discussion

CASC1 was detected through RNA-seq analysis to be critical for bladder cancer progression. And a negative correlation was observed between *CASC1* and has-miR-150 expression



(a)



(b)

FIGURE 3: Continued.










FIGURE 3: Relationship between *CASCI* expression levels and tumor immunity. (a) Correlations between *CASCI* and six types of infiltrating immune cells in 39 types of tumors. (b) Correlations between *CASCI* and four types of infiltrating immunosuppressive cells in 39 types of tumors. (c) Area under the subject operating characteristic curve (AUC) in the bar chart was used for predicting response status to immune checkpoint inhibitors (ICB).




(c)




(d)

FIGURE 4: Continued.



Overall
 ■ Altered group
 ■ Unaltered group



CNA
 ■ Amplification
 ■ Deep deletion
 ■ Diploid
 ■ Gain
 ■ Shallow deletion

(f)

FIGURE 4: Mutation characteristics of *CASC1*. (a) Characterization of the mutation of *CASC1* in multiple TCGA cancer types. (b) Mutation site of *CASC1*. (c) Relationship between the *CASC1* mutation status and overall survival (OS). (d) Association between the *CASC1* mutation status and disease-free survival (DFS); (e) Association between the *CASC1* mutation status and progression-free survival (PFS). (f) Differences in the expression levels recorded for different copy number types of *CASC1*.









FIGURE 5: Continued.




FIGURE 5: Analysis of CASCI related genes. (a) PPI network associated with CASCI obtained by analyzing the STRING database. (b)–(f) First 5 genes associated with CASCI for the case of bladder cancer. (g) Results obtained by conducting GO enrichment analysis for the top 100 genes related to CASCI associated with bladder cancer. (h) Scatter plot of correlation between CASCI and hsa-miR-150 for the case of bladder cancer.

in bladder cancer. CASCI targeting with has-mir-150 was associated with bladder cancer progression.

We analyzed the relationship between CASCI and the extent of immune invasion and found a strong correlation between the extent of immune cell invasion of various tumors and CASCI. The infiltration of immune cells (CD8 + T cells, B cells, neutrophils, CD4+ T cells, DC, and macrophages) associated with LGG, LIHC, PRAD, and LUSC cor-

related positively with the expression of CASCI. In addition, tumor infiltration corresponding to the four immunosuppressive cells that promote the process of T cell rejection (MDSCs, M2-TAMs, CAFs, and Treg cells) was also significantly positively correlated with the CASCI expression level. The overall survival, progression-free survival, and disease-free survival for patients with the CASCI mutation were poorer than those without CASCI mutation. The results





FIGURE 6: Continued.




FIGURE 6: Analysis of *CASC1* functions in bladder cancer. (a) ceRNA network of hsa-miR-150 associated with *CASC1*, where diamond nodes represent lncRNAs. (b) Venn diagram corresponding to the has-miR-150-related GSEA pathway and *CASC1*-related GSEA pathway. (c) Significant GSEA pathways in *CASC1* enrichment analysis results. (d) Significant pathways in hsa-miR-150 enrichment analysis results.

agreed well with previously reported results. It has been previously reported that *CASC1* is associated with the mitosis and microtubule polymerization of tumor cells. *CASC1* and *KRAS* co-amplify in lung tumors. This is crucial to realize microtubule polymerization and for meeting spindle assembly checkpoints [6]. *CASC1*-deficient pulmonary tumor mice are sensitive to chemotherapy [27]. The *Ras* gene is a common gene that is associated with bladder cancer [28]. Inhibition of the expression of *CASC1* is associated with increased survival of tumor patients with *Ras* mutation. The genetic link between *CASC1* and *KRAS* may mitigate mitotic damage in the context of *KRAS* by increasing the expression level of microtubule-regulatory proteins associated with tumor genes. *KRAS* influences the survival prognosis of cancer patients [29]. The extent of *KRAS* mutation can be controlled by the cis-acting elements present in the *Pas1* locus that is associated with *CASC1* [30]. The *KRAS* mutation-specific T cells can kill *KRAS* mutated tumor cells, thereby inhibiting tumor growth [31]. Thus, we can speculate that a high level of *CASC1* may result in the generation of an abnormal immune microenvironment.

CASC1 is expressed at higher levels in many types of tumors, including bladder cancer, especially at the early stages of the disease. *CASC1*, a new regulator of mitotic spindle assembly in tumor cells, is necessary for tumor growth in vivo [6]. In the case of bladder cancer, *CASC1*-related KEGG pathways include complement and coagula-

tion cascades, drug metabolism cytochrome p450, hematopoietic cell lineage, Leishmania infection, metabolism of xenobiotics by cytochrome p450, natural killer cell-mediated cytotoxicity, proteasome, ribosome, systemic lupus erythematosus, and viral myocarditis. Complement has a significant effect on the working mechanism of numerous anti-cancer antibodies [32]. As reported by Chen et al., CYP450 regulates proliferation, cell cycle, and apoptosis [33]. Previous studies have shown that CYP450 is highly expressed in cancer tissues [34, 35]. CYP24A1 dysfunction of the CYP450 members was found to be associated with prostate cancer progression by Khan et al. [36]. There is a positive correlation between the degree of NK cell infiltration in tumors and prognosis [37]. Therefore, we hypothesized that complement and coagulation cascades, drug metabolism cytochrome p450, and natural killer cell-mediated cytotoxicity were the potential pathways through which *CASC1* influenced the progression of bladder cancer.

lncRNAs and microRNAs have been reported to be predictive biomarkers for invasion, progression, and metastasis prognosis in advanced bladder cancer [8]. Wang et al. constructed a ceRNA regulatory network containing 5 muscle-invasive bladder cancer-specific lncRNAs, 9 miRNAs, and 32 mRNAs [38]. The new lncRNAs were identified as potential prognostic biomarkers for muscle-invasive bladder cancer [38]. They are more susceptible to miRNA regulation because they are mainly found in cytoplasmic vesicles and intracellular matrix [39, 40]. We are the first to report that

CASC1 can potentially be regulated by miR-150. It has been suggested that miR-150 may play a role in various types of tumor development. It is also associated with the regulation of ovarian cancer and the poor prognosis of NSCLC patients [41, 42]. In the case of breast cancer, miR-150 is regulated by lncRNA MAFG-AS1. This influences the proliferation and migration of breast cancer cells [43]. There was a significant negative correlation between CASC1 and the expression level of has-miR-150. The miRWALK database was analyzed to identify CASC1 that is has-miR-150 targeted. In our study, we hypothesized that the ceRNA regulatory network formed by miR-150 targeting CASC1 may lead to the progression of bladder cancer.

Using bioinformatics tools, we found that miR-150 substantially influenced the development and progression of bladder cancer by targeting CASC1. CASC1 expression was found to be high in bladder cancer patients, and has-miR-150 expression level was negatively correlated with the expression of CASC1. CASC1 is also associated with immune and immunosuppressive cell infiltration into tumors. Cancer patients with CASC1 mutations had a worse survival prognosis than cancer patients without CASC1 mutations. This can be potentially attributed to the involvement of CASC1 in the tumor-regulation mechanism. In order to determine the role of the CASC1-related ceRNA regulatory network in bladder cancer progression, we explored the pathogenesis of bladder cancer further. The results can help develop methods that can be used to prevent the progression of bladder cancer and improve the prognosis of patients.

Due to the fact that it is a retrospective study based on information available in different databases, its application is limited. Experimental confirmation may be lacking, and potential bias may occur. It is necessary to conduct in vitro and in vivo experiments, as well as analyze clinical data to confirm the inhibition and treatment of bladder cancer by has-miR-150 and CASC1 in the future.

5. Conclusion

There were high levels of CASC1 expression in various tumor tissues, which were associated with immune cell infiltration and immunosuppressive cells, which also affected patient prognosis. Exosomes containing miR-150-targeted CASC1 may affect the progression of bladder cancer. Based on these findings, a molecular pathway can be developed to explain bladder cancer progression.

Data Availability

The datasets generated and analyzed during the present study are available from the corresponding author on reasonable request.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgments

We thank Bullet Edits Limited for the linguistic editing and proofreading of the manuscript.

References

- [1] S. Antoni, J. Ferlay, I. Soerjomataram, A. Znaor, A. Jemal, and F. Bray, "Bladder cancer incidence and mortality: a global overview and recent trends," *European Urology*, vol. 71, no. 1, pp. 96–108, 2017.
- [2] W. Chen, R. Zheng, P. D. Baade et al., "Cancer Statistics in China, 2015: cancer statistics in China, 2015," *CA: a Cancer Journal for Clinicians*, vol. 66, no. 2, pp. 115–132, 2016.
- [3] R. Siegel, J. Ma, Z. Zou, and A. Jemal, "Cancer Statistics, 2014: cancer statistics, 2014," *CA: a Cancer Journal for Clinicians*, vol. 64, no. 1, pp. 9–29, 2014.
- [4] O. Sanli, J. Dobruch, M. A. Knowles et al., "Bladder cancer," *Nature reviews Disease primers*, vol. 3, no. 1, pp. 1–19, 2017.
- [5] Y. Liu, S. Xiong, S. Liu et al., "Analysis of gene expression in bladder cancer: possible involvement of mitosis and complement and coagulation cascades signaling pathway," *Journal of Computational Biology*, vol. 27, no. 6, pp. 987–998, 2020.
- [6] R. Sinnott, L. Winters, B. Larson et al., "Mechanisms promoting escape from mitotic stress-induced tumor cell death," *Cancer Research*, vol. 74, no. 14, pp. 3857–3869, 2014.
- [7] J. Luo, M. J. Emanuele, D. Li et al., "A genome-wide RNAi screen identifies multiple synthetic lethal interactions with the Ras oncogene," *Cell*, vol. 137, no. 5, pp. 835–848, 2009.
- [8] S. Ouerhani and A. B. A. Elgaaied, "The mutational spectrum of HRAS, KRAS, NRAS and FGFR3 genes in bladder cancer," *Cancer Biomarkers*, vol. 10, no. 6, pp. 259–266, 2012.
- [9] I. Ahmad, R. Patel, Y. Liu et al., "Ras mutation cooperates with β -catenin activation to drive bladder tumorigenesis," *Cell Death & Disease*, vol. 2, no. 3, pp. e124–e124, 2011.
- [10] A. Dassano, F. Colombo, G. Trincucci et al., "Mouse pulmonary adenoma susceptibility 1 locus is an expression QTL modulating Kras-4A," *PLoS Genetics*, vol. 10, no. 4, p. e1004307, 2014.
- [11] J. Melendez-Zajgla, G. E. Mercado-Celis, J. Gaytan-Cervantes et al., "Genomics of a pediatric ovarian fibrosarcoma. Association with the DICER1 syndrome," *Scientific Reports*, vol. 8, no. 1, p. 3252, 2018.
- [12] H. Xie, J. Zhao, J. Wan et al., "Long non-coding RNA AC245100.4 promotes prostate cancer tumorigenesis via the MicroRNA-145-5p/RBBP5 Axis," *Oncology Reports*, vol. 45, no. 2, pp. 619–629, 2020.
- [13] Q. Cao, Z. Dong, S. Liu, G. An, B. Yan, and L. Lei, "Construction of a metastasis-associated CeRNA network reveals a prognostic signature in lung cancer," *Cancer Cell International*, vol. 20, no. 1, p. 208, 2020.
- [14] J. Li, Y. Qian, C. Zhang et al., "LncRNA LINC00473 is involved in the progression of invasive pituitary adenoma by upregulating KMT5A via CeRNA-mediated MiR-502-3p evasion," *Cell Death & Disease*, vol. 12, no. 6, p. 580, 2021.
- [15] X. Kong, S. Hu, Y. Yuan et al., "Analysis of LncRNA, MiRNA and mRNA-associated CeRNA networks and identification of potential drug targets for drug-resistant non-small cell lung cancer," *Journal of Cancer*, vol. 11, no. 11, pp. 3357–3368, 2020.

- [16] X. Yang, T. Ye, H. Liu et al., "Expression profiles, biological functions and clinical significance of CircRNAs in bladder cancer," *Molecular Cancer*, vol. 20, no. 1, p. 4, 2021.
- [17] R. Dai, Y. Zhou, Z. Chen, Z. Zou, P. Liu, and X. Gao, "The analysis of a CeRNA network and the correlation between LncRNA, MiRNA, and mRNA in bladder cancer," *Translational Cancer Research*, vol. 9, no. 2, pp. 869–881, 2020.
- [18] L. Du, X. Wang, Y. Yin et al., "Identification of a potentially functional CircRNA-MiRNA-MRNA CeRNA regulatory network in bladder cancer by analysis of microarray data," *Transl. Androl. Urol.*, vol. 10, no. 1, pp. 24–36, 2021.
- [19] Z. Luo, Y. Sun, B. Qi et al., "Human bone marrow mesenchymal stem cell-derived extracellular vesicles inhibit shoulder stiffness via let-7a/Tgfb1 axis," *Bioact. Mater.*, vol. 17, pp. 344–359, 2022.
- [20] Y. Sun, X. Sun, S. Liu, L. Liu, and J. Chen, "The overlap between regeneration and fibrosis in injured skeletal muscle is regulated by phosphatidylinositol 3-kinase/Akt signaling pathway - a bioinformatic analysis based on LncRNA microarray," *Gene*, vol. 672, pp. 79–87, 2018.
- [21] J. S. You and P. A. Jones, "Cancer genetics and epigenetics: two sides of the same coin?," *Cancer Cell*, vol. 22, no. 1, pp. 9–20, 2012.
- [22] N. Cherradi, "MicroRNAs as potential biomarkers in adrenocortical cancer: progress and challenges," *Frontiers in Endocrinology*, vol. 6, p. doi:10.3389/fendo.2015.00195, 2016.
- [23] X. Zhou, P. Kurywchak, K. Wolf-Dennen et al., "Unique somatic variants in DNA from urine exosomes of individuals with bladder cancer," *Mol. Ther. - Methods Clin. Dev.*, vol. 22, pp. 360–376, 2021.
- [24] J. Li, A.-S. Wang, S. Wang et al., "LncSNHG14 promotes the development and progression of bladder cancer by targeting MiRNA-150-5p," *European Review for Medical and Pharmacological Sciences*, vol. 23, no. 3, pp. 1022–1029, 2019.
- [25] J. Vivian, A. A. Rao, F. A. Nothaft et al., "Toil enables reproducible, open source," *Nature biotechnology*, vol. 35, no. 4, pp. 314–316, 2017.
- [26] G. Fang and S. Liu, "Low expression of novel biomarker RCSD1 predicts poor prognosis of lung adenocarcinoma," *Life Research*, vol. 5, no. 1, p. 6, 2022.
- [27] P. Liu, Y. Wang, H. Vikis et al., "Candidate lung tumor susceptibility genes identified through whole-genome association analyses in inbred mice," *Nature Genetics*, vol. 38, no. 8, pp. 888–895, 2006.
- [28] T. Jiang, T. Liu, L. Li et al., "Knockout of phospholipase Ce attenuates N-butyl-N-(4-Hydroxybutyl) nitrosamine-induced bladder tumorigenesis," *Molecular Medicine Reports*, vol. 13, no. 3, pp. 2039–2045, 2016.
- [29] G. Manenti, F. Galbiati, A. Pettinicchio et al., "A V141L polymorphism of the human LRMP gene is associated with survival of lung cancer patients," *Carcinogenesis*, vol. 27, no. 7, pp. 1386–1390, 2006.
- [30] G. Manenti, G. Trincucci, A. Pettinicchio, E. Amendola, M. Scarfò, and T. A. Dragani, "Cis-acting genomic elements of the Pas1 locus control Kras mutability in lung tumors," *Oncogene*, vol. 27, no. 43, pp. 5753–5758, 2008.
- [31] Y. Wan, Y. Zhang, G. Wang, P. M. Mwangi, H. Cai, and R. Li, "Recombinant KRAS G12D protein vaccines elicit significant anti-tumor effects in mouse CT26 tumor models," *Frontiers in Oncology*, vol. 10, p. 1326, 2020.
- [32] M. Elvington, Y. Huang, B. P. Morgan et al., "A targeted complement-dependent strategy to improve the outcome of MAb therapy, and characterization in a murine model of metastatic cancer," *Blood*, vol. 119, no. 25, pp. 6043–6051, 2012.
- [33] T. C. Chen, T. Sakaki, K. Yamamoto, and A. Kittaka, "The roles of cytochrome P450 enzymes in prostate cancer development and treatment," *Anticancer Research*, vol. 32, no. 1, pp. 291–298, 2012.
- [34] L. Annovazzi, M. Mellai, V. Caldera, G. Valente, L. Tessitore, and D. Schiffer, "MTOR, S6 and AKT expression in relation to proliferation and apoptosis/autophagy in glioma," *Anticancer Research*, vol. 29, no. 8, pp. 3087–3094, 2009.
- [35] Q. Wang, X. Wang, and C.-B. Zhang, "Lentivirus mediated GOLPH3 ShRNA inhibits growth and metastasis of esophageal squamous cancer," *Asian Pacific Journal of Cancer Prevention*, vol. 14, no. 9, pp. 5391–5396, 2013.
- [36] N. A. Khan, K. H. Stopsack, E. H. Allott et al., "Intratymoral Sterol-27-hydroxylase (CYP27A1) expression in relation to cholesterol synthesis and vitamin D signaling and its association with lethal prostate cancer," *Cancer Epidemiology, Biomarkers & Prevention*, vol. 28, no. 6, pp. 1052–1058, 2019.
- [37] S. Rusakiewicz, M. Semeraro, M. Sarabi et al., "Immune infiltrates are prognostic factors in localized gastrointestinal stromal tumors," *Cancer Research*, vol. 73, no. 12, pp. 3499–3510, 2013.
- [38] H. Wang, L. Niu, S. Jiang et al., "Comprehensive analysis of aberrantly expressed profiles of LncRNAs and MiRNAs with associated CeRNA network in muscle-invasive bladder cancer," *Oncotarget*, vol. 7, no. 52, pp. 86174–86185, 2016.
- [39] A. Kotipalli, R. Gutti, and C. K. Mitra, "Dynamics of MiRNA biogenesis and nuclear transport," *Journal of Integrative Bioinformatics*, vol. 13, no. 5, pp. 22–34, 2016.
- [40] V. Sackmann, M. S. Sinha, C. Sackmann et al., "Inhibition of NSMase2 reduces the transfer of oligomeric α -synuclein irrespective of hypoxia," *Frontiers in Molecular Neuroscience*, vol. 12, p. 200, 2019.
- [41] Q. Zhang, X. Zhou, M. Wan et al., "FoxP3-MiR-150-5p/3p suppresses ovarian tumorigenesis via an IGF1R/IRS1 pathway feedback loop," *Cell Death & Disease*, vol. 12, no. 3, p. 275, 2021.
- [42] Q.-W. Yin, X.-F. Sun, G.-T. Yang, X.-B. Li, M.-S. Wu, and J. Zhao, "Increased expression of MicroRNA-150 is associated with poor prognosis in non-small cell lung cancer," *International Journal of Clinical and Experimental Pathology*, vol. 8, pp. 842–846, 2015.
- [43] H. Jia, D. Wu, Z. Zhang, and S. Li, "Regulatory effect of the MAFG-AS1/MiR-150-5p/MYB Axis on the proliferation and migration of breast cancer cells," *International Journal of Oncology*, vol. 58, no. 1, pp. 33–44, 2020.