

## Retraction

# Retracted: MiR-129-5p Inactivates NF- $\kappa$ B Pathway to Block Rheumatoid Arthritis Development via Targeting BRD4

### Journal of Healthcare Engineering

Received 11 July 2023; Accepted 11 July 2023; Published 12 July 2023

Copyright © 2023 Journal of Healthcare Engineering. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This article has been retracted by Hindawi following an investigation undertaken by the publisher [1]. This investigation has uncovered evidence of one or more of the following indicators of systematic manipulation of the publication process:

- (1) Discrepancies in scope
- (2) Discrepancies in the description of the research reported
- (3) Discrepancies between the availability of data and the research described
- (4) Inappropriate citations
- (5) Incoherent, meaningless and/or irrelevant content included in the article
- (6) Peer-review manipulation

The presence of these indicators undermines our confidence in the integrity of the article's content and we cannot, therefore, vouch for its reliability. Please note that this notice is intended solely to alert readers that the content of this article is unreliable. We have not investigated whether authors were aware of or involved in the systematic manipulation of the publication process.

In addition, our investigation has also shown that one or more of the following human-subject reporting requirements has not been met in this article: ethical approval by an Institutional Review Board (IRB) committee or equivalent, patient/participant consent to participate, and/or agreement to publish patient/participant details (where relevant).

Wiley and Hindawi regrets that the usual quality checks did not identify these issues before publication and have since put additional measures in place to safeguard research integrity.

We wish to credit our own Research Integrity and Research Publishing teams and anonymous and named external researchers and research integrity experts for contributing to this investigation.

The corresponding author, as the representative of all authors, has been given the opportunity to register their agreement or disagreement to this retraction. We have kept a record of any response received.

### References

- [1] Z. Wu and D. Chen, "MiR-129-5p Inactivates NF- $\kappa$ B Pathway to Block Rheumatoid Arthritis Development via Targeting BRD4," *Journal of Healthcare Engineering*, vol. 2022, Article ID 8330659, 8 pages, 2022.

## Research Article

# MiR-129-5p Inactivates NF- $\kappa$ B Pathway to Block Rheumatoid Arthritis Development via Targeting BRD4

Zhaoli Wu<sup>1</sup> and Disi Chen<sup>2</sup> 

<sup>1</sup>Department of Integrated TCM & Western Medicine, Hunan Provincial People's Hospital, Changsha 410005, Hunan, China

<sup>2</sup>Department of Respiratory Medicine, Hunan Provincial People's Hospital, Changsha 410005, Hunan, China

Correspondence should be addressed to Disi Chen; [chendisi@hnsrmyy.org.cn](mailto:chendisi@hnsrmyy.org.cn)

Received 18 March 2022; Revised 2 April 2022; Accepted 5 April 2022; Published 19 April 2022

Academic Editor: Kalidoss Rajakani

Copyright © 2022 Zhaoli Wu and Disi Chen. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Object.** Rheumatoid arthritis (RA) is an osteoarticular disease which seriously affects the motor abilities of the patients. MicroRNA disorder has been confirmed as a biomarker event in RA development, and several studies have identified that miR-129-5p is related with the progression of multiple bone diseases. The study attempted to investigate the regulation mechanism of miR-129-5p in RA development. **Methods.** The abundance of miR-129-5p was detected in the samples including normal tissues and RA tissues and cell lines including human fibroblast-like synoviocytes (hFLSs) and human rheumatoid arthritis fibroblast-like synoviocytes (RA-FLSs). The CCK-8 assay, flow cytometry, Transwell, and ELISA were used to observe the effects of miR-129-5p on the phenotype of RA-FLSs. Moreover, the potential targets of miR-129-5p were identified with TargetScan and dual-luciferase reporter gene assay. Besides, the abundances of the proteins were analyzed with western blot. **Results.** Decreased miR-129-5p was observed in RA tissues and cells. Increased miR-129-5p obviously blocked the proliferation, inflammatory stress, and migration and remarkably promoted cellular apoptosis. Moreover, BRD4 was confirmed as targets of miR-129-5p, and BRD4 upregulation could partly rescue the inhibition of miR-129-5p on aggressive behaviors of RA-FLSs. Besides, the finding of this study also proved that upregulated miR-129-5p could impede the NF- $\kappa$ B pathway via targeting BRD4. **Conclusion.** This study suggests that miR-129-5p suppresses the activation of NF- $\kappa$ B pathway to block the progression of RA via targeting BRD4.

## 1. Introduction


Rheumatoid arthritis (RA) is still an intractable disease with high incidence in middle-aged and senior people. The patients with RA always exhibit persistent inflammatory reactions in synovial tissue, neovascularization, and joint destruction. Moreover, persistent inflammation may also induce multiple severe complications such as lung and cardiovascular disease [1, 2]. The aggressive behaviors of RA fibroblast-like synoviocytes (RA-FLS) are direct reason causing RA development. RA-FLSs are characterized with the high aggressive behaviors including high proliferative and migratory abilities [3, 4]. Moreover, the matrix metalloproteinases (MMPs) produced by RA-FLSs can promote the joint injury which may cause irreversible joint dysfunction for the patients [5]. Therefore, the intervention of suppressing the aggressive phenotype of RA-FLSs has been

suggested as a promising strategy for RA treatment in clinical.

The disorder of microRNA profile acts as a biomarker event for RA progression [6]. Moreover, accumulating reports have indicated that miRNAs directly regulate RA development via influencing the proliferation, inflammatory level, and cellular molecular signal transduction of RA-FLSs [7, 8]. MiR-129-5p, located on p13, functions as a tumor obstructor to restrain malignant behaviors of cancer cells. MiR-129-5p plays a critical role in bone development, and miR-129-5p has been confirmed to promoter the differentiation of bone marrow mesenchymal stem cells [9]. Moreover, several reports have indicated that aberrant abundance of miR-129-5p is associated with multiple bone diseases. The research has revealed that miR-129-5p is associated with osteoarthritis development [10]. Moreover, the study has also found that miR-129-5p downregulation may

TABLE 1: Primer sequence of genes.

miR-129-5p-F	5'-: ACCCAGTGC GATTGT-3'
miR-129-5p-R	5'-: ACTGTACTGGAAGATGGA-3'
BRD4-F	5'-ACAACAAGCCTGGAGATGACA-3'
BRD4-R	5'-GTTTGGTACCGTGGAAACGC-3'
U6-F	5'-CTCGCTTCGGCAGCACA-3'
U6-R	5'-AACGCTTCACGAATTTGCGT-3'

FIGURE 1: The related abundance of miR-129-5p in RA tissues and cells. (a) Normal tissues vs. RA tissues. (b) hFLSs vs. RA-FLSs. \*\*  $P < 0.01$ .

promote the progression of RA [11]. However, the research studies on regulation mechanism of miR-129-5p in RA are still necessary.

This study was aimed to investigate the functions and molecular mechanism of miR-129-5p in RA development.

## 2. Materials and Methods

**2.1. Sample Collection.** The specimens including RA and normal tissues were stored at  $-70^{\circ}\text{C}$ . This experiment has been approved by the Hospital Ethics Committee.

**2.2. Cell Culture and Transfection.** The human rheumatoid arthritis fibroblast-like synoviocytes (RA-FLSs) were purchased from Beijing Donggeboye Biotechnology Co., Ltd (Beijing, China). The cell medium was DMEM supplying 10% FBS and 1% penicillin/streptomycin. The cells were cultured in the conditions including  $37^{\circ}\text{C}$  and 5%  $\text{CO}_2$ . Moreover, the cell subculture was performed when cell density was at 90%.

For cell transfection, 4  $\mu\text{g}$  of DNA (or 100 pmol RNA) and 10  $\mu\text{L}$  Lipofectamine 2000 were mixed with 250  $\mu\text{L}$  medium and incubated for 5 min, respectively. Subsequently, the medium containing DNA (or RNA) was mixed and incubated with the medium containing Lipofectamine 2000 for 20 min at  $25^{\circ}\text{C}$ . Finally, 500  $\mu\text{L}$  of mixture was added into each well, and then cells were incubated for 24 h.

**2.3. qRT-PCR.** RA-FLSs were treated with TRIzol reagent for RNA extraction, and the related concentration was measured with spectrophotometry. Subsequently, PrimeScript<sup>®</sup> RT reagent Kit (Thermo Fisher, Massachusetts, USA) was applied for cDNA reverse transcription.

Briefly, 10  $\mu\text{L}$  of the reaction system for qRT-PCR was performed with following conditions: for miRNA,  $95^{\circ}\text{C}$  for 5 min followed by 40 cycles of  $95^{\circ}\text{C}$  for 15 s and  $60^{\circ}\text{C}$  for 1 min; for mRNA,  $95^{\circ}\text{C}$  for 5 min, followed by 40 cycles of  $95^{\circ}\text{C}$  for 15 s,  $62^{\circ}\text{C}$  for 30 s, and  $72^{\circ}\text{C}$  for 30 s. The abundance of the related miRNA or mRNA was calculated with  $2^{-\Delta\Delta\text{Ct}}$ . The primers of the genes are listed in Table 1.

**2.4. Western Blotting.** The abundance of the proteins in RA-FLSs was observed by western blot. In brief, the cells were treated with the RIPA buffer for 30 min on the ice, and then total proteins were extracted from the cells with centrifugation (13,200 rpm for 15 min). The BCA kit was applied for concentration measurement of proteins. The proteins, mixed with loading buffer, were boiled for 5 min. The proteins were separated with SDS-PAGE and then translated on PVDF membranes. The membranes were incubated with the primary antibodies (anti-BED4, ab243862 abcam; anti-P65, ab32536, abcam) at  $4^{\circ}\text{C}$  overnight after immersing in 5% fat-free milk for 2 h. TBST was used to wash membranes for three times, and second antibodies were used for membrane incubation at  $25^{\circ}\text{C}$  for 2 h. Finally, the abundance of the proteins was observed by a chemiluminescence detection system after removing redundant second antibodies by TBST.

**2.5. Flow Cytometry.**  $2 \times 10^3$  cells RA-FLSs were suspended with 200  $\mu\text{L}$  ice Annexin V-FITC binding buffer. Subsequently, 10  $\mu\text{L}$  of propidium iodide (PI 20  $\mu\text{g}/\text{ml}$ ) and 5  $\mu\text{L}$  of Annexin V-FITC (10  $\mu\text{g}/\text{ml}$ ) were used for cell staining in the dark for 5 min. The apoptosis of the cells was detected with a flow cytometry equipment (BD Biosciences, State of New Jersey, USA).




FIGURE 2: MiR-129-5p effectively suppressed the aggressive behaviors of RA-FLSs. (a) Cell viability detected by CCK-8 assay. (b) Inflammatory factor levels detected by ELISA. (c) Cell migration observed by Transwell. (d) Cell apoptosis detected by flow cytometry assay. \*\*  $P < 0.01$ .

**2.6. CCK-8 Assay.**  $5 \times 10^3$  cells were seeded into 96-well plates. After transfection for 24 h, 48 h, 72 h, and 96 h, the cells were added with  $10 \mu\text{L}$  of CCK-8 solution and then incubated for 4 h under lightless condition. Finally, the OD450 value of the cells was detected with microplate reader.

**2.7. Dual-Luciferase Report Genes Assay.** The sequences of BRD4 wild type (BRD4-wt) and BRD4 mutant type (BRD4-mut) with mutant sites in the binding region with miR-129-5p were inserted into pmirGLO vector (Youbao Biotechnology Co., Ltd, Chongqing, China), respectively. After that, miR-




FIGURE 3: BRD4 was a downstream target of miR-129-5p. (a) Dual-luciferase report gene assay. (b) The mRNA abundance of BRD4. \*\*  $P < 0.01$ .

NC/miR-129-5p mimics and BRD4-wt/BRD4-mut were used for the transfection of RA-FLSs. Finally, after culturing for 48 h, the luciferase activity in the cells was measured.

**2.8. Transwell Assay.**  $2 \times 10^4$  cells were cultured in upper chamber of Transwell plates to normal growth stage. After transfection for 72 h, the cells were cultured with  $200 \mu\text{L}$  serum-free medium, and  $500 \mu\text{L}$  DMEM plus 10% FBS was supplied into lower chamber. The cells were cultured at  $37^\circ\text{C}$  for 24 h and then were fixed with formaldehyde for 30 min. Subsequently, the cells in upper chamber were removed with a cotton ball. Finally, the migratory cells were calculated in 3 randomly visual regions.

**2.9. ELISA.** The ELISA kits (Shanghai Fusheng Biotechnology Co., Ltd, Shanghai, China) were applied for the detection of inflammatory factors including IL-6 and IL- $1\beta$ . The detection process was performed following the instructions of the kits, strictly.

**2.10. Data Analysis.** The experiments were repeated for 3 times, independently. SPSS 20.0 was applied for data analysis, and the difference of the data was judged with Chi-squared test or ANOVA with Tukey's post hoc-test. Moreover, Graphpad Prism 8 was used for data visualization.

### 3. Results

**3.1. Decreased miR-129-5p Was Observed in RA-FLSs.** To investigate the connection of miR-129-5p and RA development, the abundance of miR-129-5p was detected in RA tissues and cells. The results revealed that miR-129-5p was remarkably upregulated in the RA samples (Figure 1(a);  $P < 0.01$ ). Besides, compared with the human fibroblast-like synoviocytes (hFLSs), decreased miR-129-5p was observed

in the RA-FLSs (Figure 1(b);  $P < 0.01$ ). Those evidences hinted that miR-129-5p upregulation was involved in the progression of RA.

**3.2. Increased miR-129-5p Suppressed the Aggressive Behaviors of RA-FLSs.** To illustrate the functions of miR-129-5p in RA-FLSs, miR-129-5p was upregulated in RA-FLSs to observe the changes in phenotype of RA-FLSs. The results proved that miR-129-5p upregulation effectively suppressed the proliferation and migration and induced the apoptosis of RA-FLSs (Figures 2(a), 2(c), and 2(d);  $P < 0.01$ ). Moreover, miR-129-5p downregulation also effectively improved the inflammatory levels in RA-FLSs (Figure 2(b);  $P < 0.01$ ). Those proofs suggested that miR-129-5p can inhibit the inflammatory apoptosis of RA-FLSs.

**3.3. BRD4 Was Directly Regulated by miR-129-5p.** To reveal the regulation mechanism of miR-129-5p in RA, miRDB and TargetScan database were used to screen the downstream factors of miR-129-5p. The prediction result showed that miR-129-5p could directly bind with BRD4. Moreover, the luciferase reporter gene assay also confirmed that miR-129-5p mimics reduced luciferase activity of RA-FLSs transfected with the wt-BRD4 (Figure 3(a);  $P < 0.01$ ). Moreover, reduced BRD4 was also found in the RA-FLSs compared with that in normal FLSs (Figure 3(b);  $P < 0.01$ ). Those observations supported that BRD4 was directly regulated by miR-129-5p.

**3.4. BRD4 Reversed the Effects of miR-129-5p on RA-FLSs.** To verify whether miR-129-5p can inhibit the progression of RA via regulating BRD4, the expressed vectors of miR-129-5p and BRD4 were used for RA-FLSs transfection to observe the changes in phenotype of RA-FLSs. The results showed that suppressed proliferation, invasion, and inflammatory








FIGURE 4: MiR-129-5p effectively suppressed the aggressive behaviors of RA-FLSs via targeting BRD4. (a) The viability of RA-FLSs observed by CCK-8 assay. (b) Inflammatory factor levels detected by ELISA. (c) Cell migration observed by Transwell. (d) Cell apoptosis detected by flow cytometry assay. \*\*  $P < 0.01$ .




FIGURE 5: MiR-129-5p inhibited NF- $\kappa$ B pathway via targeting BRD4. (a) The related abundance of miR-129-5p. (b–e) The abundances of BRD4, P65, and p-P65 measured with western blot. \*\*  $P < 0.01$ .

responses induced by increased miR-129-5p were reversed by BRD4 upregulation (Figures 4(a)–4(d);  $P < 0.01$ ). Those observations hinted that miR-129-5p affected the behaviors of RA-FLSs via inhibiting BRD4.

**3.5. MiR-129-5p Inactivated the NF- $\kappa$ B Pathway via Suppressing BRD4.** To delve the molecular mechanism of miR-129-5p in RA development, the activity of NF- $\kappa$ B pathway was analyzed in RA-FLSs after transfecting with the expressed vectors of miR-129-5p and BRD4. The results confirmed that increased miR-129-5p distinctly reduced the abundances of BRD4, P65, and p-P65 (Figure 5;  $P < 0.01$ ). In addition, BRD4 upregulation partly rescued the inactivation of miR-129-5p on NF- $\kappa$ B pathway (Figure 5;  $P < 0.01$ ).

#### 4. Discussion

RA is still a spiny question for modern medical therapeutic system and may induce irreversible motor dysfunctions of the patients [12]. Considerable research

studies have proved that the aggressive behaviors of RA-FLSs are direct reason causing the persistent pathological changes in joints [13, 14]. Disorder miRNA profile has been found as a biomarker event in RA-FLSs [15]. This study investigated the connection of miR-129-5p with RA, revealed the functions and regulation of miR-129-5p on the aggressive behaviors of RA-FLSs, and provided some references for the RA treatment.

Accumulating research studies have indicated that abnormal miRNA profile is a distinguishing feature of RA, and some miRNAs have been confirmed to involve the development of this disease [16]. In this study, decreased miR-129-5p was observed in RA-FLSs compared with normal cells. MiR-129-5p serves as an intruder role to impede the formation and development of multiple diseases. The research has confirmed that miR-129-5p is obviously reduced in the tissues of the patients with osteoarthritis, and miR-129-5p upregulation could remarkably improve the IL-1 $\beta$ -mediated osteoarthritis via suppressing HMGB1 [17]. Moreover, miR-129-5p has been confirmed to block the proliferation and induce the apoptosis of RA-FLSs via

directly regulating the expression of IGF-1 [11]. The finding of this study also proved that miR-129-5p functioned as a suppressor to inhibit the proliferation and invasion of RA-FLSs. MiR-129-5p has been verified to involve the release of inflammatory factors, and increased miR-129-5p can effectively alleviate the levels of inflammatory response in multiple diseases including acute kidney and intestinal injury [18, 19]. The observation in this study also confirmed that increased miR-129-5p could effectively relieve the IL-6 and IL-8 levels. Thus, it suggests that miR-129-5p plays an inhibitor role in progression of RA.

MiRNAs are characterized with restraining the translation process to influence the phenotype of cells via interacting with the related mRNAs. For RA, the miRNAs also serve as important regulators in the pathological changes of the cells. The study has indicated that miR-130a reduces the proliferation and inflammation of RA-FLSs via targeting NDRG2 [20]. Wang et al. have discovered that miR-129-5p functions as a inhibitor role in multiple myeloma, and increased miR-129-5p could restrain multiple malignant behaviors of tumor cells via targeting JAG1 [21]. The major observation in this research was that miR-129-5p was involved in RA development via directly targeting BRD4. BRD4 upregulations have been found in multiple diseases. For bone disease, increased BRD4 also mediates the inflammatory reactions and aberrant proliferation of cells [21]. BRD4 has been confirmed as a promising therapeutic target for osteoarthritis, and decreased BRD4 could effectively improve the inflammatory stress in osteoarthritis [22]. Moreover, this study also confirmed that BRD4 could effectively reverse the effects of miR-129-5p on RA-FLSs.

RA development is related with the dysfunctions of multiple cellular pathways including JAK/STAT, PI3K/AKT, NF- $\kappa$ B, and so on. Crispino et al. have indicated the abnormally activated JAK/STAT pathway in RA-FLSs, and the inhibition of JAK/STAT pathway can effectively alleviate the pain and symptoms of the patients with RA [23]. This study also observed the aberrant activation of NF- $\kappa$ B pathway in RA-FLSs. Several reports have indicated that activated NF- $\kappa$ B pathway is involved in promoting the multiple aggressive behaviors of RA-FLSs, such as excessive proliferation and violent invasion [24]. Inactivating NF- $\kappa$ B pathway has been considered as a promising strategy for restraining the RA development. This study confirmed that increased miR-129-5p could effectively reduce the activity of NF- $\kappa$ B pathway. MiR-129-5p has been confirmed to take part in attenuating the lipopolysaccharide-mediated acute kidney injury via regulating the activity of NF- $\kappa$ B pathway [18]. Moreover, it was also found that miR-129-5p inactivated NF- $\kappa$ B pathway in RA-FLSs was related with BRD4. For RA, several research studies have also proved that BRD4 upregulation is critical reason for NF- $\kappa$ B pathway which may promote the progression of this arthropathy [25]. For osteoarthritis, decreased BRD4 could reduce the activity of NF- $\kappa$ B and thus block the progression of the disease. The study has also indicated that BRD4 can inhibit the ubiquitination of RelA to increase the stability of NF- $\kappa$ B [26]. Therefore, it implies that miR-129-5p regulates the activity of NF- $\kappa$ B pathways via targeting BRD4.

## 5. Conclusion

In summary, this study suggests that miR-129-5p could regulate the activity of NF- $\kappa$ B pathways to block the aggressive behaviors of RA-FLSs by targeting BRD4.

## Data Availability

The data used to support the findings of this study are available on reasonable request from the corresponding author.

## Conflicts of Interest

The authors declare that they have no conflicts of interest.

## References

- [1] H. Radner, T. Lesperance, N. A. Accortt, and D. H. Solomon, "Incidence and prevalence of cardiovascular risk factors among patients with rheumatoid arthritis, psoriasis, or psoriatic arthritis," *Arthritis Care & Research*, vol. 69, no. 10, pp. 1510–1518, 2017.
- [2] J. A. Zamora-Legoff, M. L. Krause, C. S. Crowson, J. H. Ryu, and E. L. Matteson, "Patterns of interstitial lung disease and mortality in rheumatoid arthritis," *Rheumatology*, vol. 56, no. 3, pp. 344–350, 2017.
- [3] W. Yang, X. Wei, Y. Jiao et al., "STAT3/HIF-1 $\alpha$ /fascin-1 axis promotes RA FLSs migration and invasion ability under hypoxia," *Molecular Immunology*, vol. 142, pp. 83–94, 2022.
- [4] Y. Zou, C. Shen, T. Shen et al., "LncRNA *THRIL* is involved in the proliferation, migration, and invasion of rheumatoid fibroblast-like synoviocytes," *Annals of Translational Medicine*, vol. 9, no. 17, p. 1368, 2021 Sep.
- [5] M. Magnol, B. Eleonore, R. Claire et al., "Use of eHealth by patients with rheumatoid arthritis: observational, cross-sectional, multicenter study," *Journal of Medical Internet Research*, vol. 23, no. 1, Article ID e19998, 2021.
- [6] R. Erfan, O. G. Shaker, M. A. F. Khalil et al., "Long non-coding RNA PVT1 and its target miRNA-146a as potential prognostic biomarkers in rheumatoid arthritis patients," *Life*, vol. 11, no. 12, p. 1382, 2021.
- [7] J. Hao, Y. Chen, and Y. Yu, "Circular RNA circ\_0008360 inhibits the proliferation, migration, and inflammation and promotes apoptosis of fibroblast-like synoviocytes by regulating miR-135b-5p/HDAC4 Axis in rheumatoid arthritis," *Inflammation*, vol. 45, no. 1, pp. 196–211, 2022.
- [8] J. Tang, S. Yi, and Y. Liu, "Long non-coding RNA PVT1 can regulate the proliferation and inflammatory responses of rheumatoid arthritis fibroblast-like synoviocytes by targeting microRNA-145-5p," *Human Cell*, vol. 33, no. 4, pp. 1081–1090, 2020.
- [9] C. Zhao, Y. Gu, Y. Wang et al., "miR-129-5p promotes osteogenic differentiation of BMSCs and bone regeneration via repressing *Dkk3*," *Stem Cells International*, vol. 2021, Article ID 7435605, 18 pages, 2021.
- [10] Q. Cheng, X. Chen, H. Wu, and Y. Du, "Three hematologic/immune system-specific expressed genes are considered as the potential biomarkers for the diagnosis of early rheumatoid arthritis through bioinformatics analysis," *Journal of Translational Medicine*, vol. 19, no. 1, p. 18, 2021.
- [11] Y. Zhang, N. Yan, X. Wang, Y. Chang, and Y. Wang, "MiR-129-5p regulates cell proliferation and apoptosis via IGF-1R/



- Src/ERK/Egr-1 pathway in RA-fibroblast-like synoviocytes,” *Bioscience Reports*, vol. 39, no. 12, Article ID BSR20192009, 2019.
- [12] S. Yuan and S. Larsson, “Causal associations of iron status with gout and rheumatoid arthritis, but not with inflammatory bowel disease,” *Clinical Nutrition*, vol. 39, no. 10, pp. 3119–3124, 2020.
- [13] K. Yip and I. Navarro-Millán, “Racial, ethnic, and healthcare disparities in rheumatoid arthritis,” *Current Opinion in Rheumatology*, vol. 33, no. 2, pp. 117–121, 2021.
- [14] F. I. L. Clanchy, F. Borghese, J. Bystrom et al., “TLR expression profiles are a function of disease status in rheumatoid arthritis and experimental arthritis,” *Journal of Autoimmunity*, vol. 118, Article ID 102597, 2021.
- [15] F. Liu, X.-X. Feng, S.-L. Zhu et al., “Long non-coding RNA SNHG1 regulates rheumatoid synovial invasion and proliferation by interaction with PTBP1,” *International Immunopharmacology*, vol. 90, Article ID 102597, 2021.
- [16] Y. Gu, H. Zhou, H. Yu et al., “miR-99a regulates CD4+ T cell differentiation and attenuates experimental autoimmune encephalomyelitis by mTOR-mediated glycolysis,” *Molecular Therapy - Nucleic Acids*, vol. 26, pp. 1173–1185, 2021.
- [17] M. Qiu, D. Liu, and Q. Fu, “MiR-129-5p shuttled by human synovial mesenchymal stem cell-derived exosomes relieves IL-1 $\beta$  induced osteoarthritis via targeting HMGB1,” *Life Sciences*, vol. 269, Article ID 118987, 2021.
- [18] X. Huang, X. Hou, L. Chuan et al., “miR-129-5p alleviates LPS-induced acute kidney injury via targeting HMGB1/TLRs/NF-kappaB pathway,” *International Immunopharmacology*, vol. 89, no. Pt A, Article ID 107016, 2020 Dec.
- [19] X. Du, D. Tian, J. Wei et al., “MEG3 alleviated LPS-induced intestinal injury in sepsis by modulating miR-129-5p and surfactant protein D,” *Mediators of Inflammation*, vol. 2020, Article ID 8232734, 11 pages, 2020.
- [20] X. Su, H. Zhang, H. Wang, and P. Sun, “MiR-130a/Ndr2 Axis inhibits the proliferation of fibroblast-like synoviocytes in rheumatoid arthritis,” *Inflammation*, vol. 43, no. 6, pp. 2048–2060, 2020.
- [21] Y. Wang, Q. Lin, C. Song, R. Ma, and X. Li, “Depletion of circ\_0007841 inhibits multiple myeloma development and BTZ resistance via miR-129-5p/JAG1 axis,” *Cell Cycle*, vol. 19, no. 23, pp. 3289–3302, 2020.
- [22] Z. Huang, R. Yang, L. Zhang et al., “BRD4 inhibition alleviates mechanical stress-induced TMJ OA-like pathological changes and attenuates TREM1-mediated inflammatory response,” *Clinical Epigenetics*, vol. 13, no. 1, p. 10, 2021.
- [23] N. Crispino and F. Ciccia, “JAK/STAT pathway and nociceptive cytokine signalling in rheumatoid arthritis and psoriatic arthritis,” *Clinical & Experimental Rheumatology*, vol. 39, no. 3, pp. 668–675, 2021.
- [24] R. Wang, J. Liu, Z. Wang et al., “Mangiferin exert protective effects on joints of adjuvant-induced arthritis rats by regulating the MAPKs/NF- $\kappa$ B pathway of fibroblast-like synoviocytes,” *International Immunopharmacology*, vol. 101, no. Pt B, Article ID 108352, 2021.
- [25] L.-m. Zhang, J.-J. Zhou, and C.-l. Luo, “CYLD suppression enhances the pro-inflammatory effects and hyperproliferation of rheumatoid arthritis fibroblast-like synoviocytes by enhancing NF- $\kappa$ B activation,” *Arthritis Research and Therapy*, vol. 20, no. 1, p. 219, 2018.
- [26] Z. Zou, B. Huang, X. Wu et al., “Brd4 maintains constitutively active NF- $\kappa$ B in cancer cells by binding to acetylated RelA,” *Oncogene*, vol. 33, no. 18, pp. 2395–2404, 2014.