

Research Article

Weighted Composition Operators on Some Weighted Spaces in the Unit Ball

Xiaohong Fu and Xiangling Zhu

Department of Mathematics, Jiaying University Guangdong, Meizhou 514015, China

Correspondence should be addressed to Xiaohong Fu, jyufxh@163.com

Received 18 May 2008; Accepted 28 May 2008

Recommended by Stevo Stevic

Let \mathbb{B}_n be the unit ball of \mathbb{C}^n , $H(\mathbb{B}_n)$ the space of all holomorphic functions in \mathbb{B}_n . Let $u \in H(\mathbb{B}_n)$ and α be a holomorphic self-map of \mathbb{B}_n . For $f \in H(\mathbb{B}_n)$, the weighted composition operator uC_α is defined by $(uC_\alpha f)(z) = u(z)f(\alpha(z))$, $z \in \mathbb{B}_n$. The boundedness and compactness of the weighted composition operator on some weighted spaces on the unit ball are studied in this paper.

Copyright © 2008 X. Fu and X. Zhu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let \mathbb{B}_n be the unit ball of \mathbb{C}^n , $H(\mathbb{B}_n)$ the space of all holomorphic functions in \mathbb{B}_n , and $H^\infty = H^\infty(\mathbb{B}_n)$ the space of all bounded holomorphic functions in the unit ball. For $f \in H(\mathbb{B}_n)$, let

$$\mathfrak{R}f(z) = \sum_{j=1}^n z_j \frac{\partial f}{\partial z_j}(z) \quad (1.1)$$

be the radial derivative of f .

A positive continuous function μ on $[0, 1)$ is called normal if there exist positive numbers α and β , $0 < \alpha < \beta$, and $\delta \in [0, 1)$ such that (see, e.g., [1, 2])

$$\begin{aligned} \frac{\mu(r)}{(1-r)^\alpha} \text{ is decreasing on } [\delta, 1), & \quad \lim_{r \rightarrow 1} \frac{\mu(r)}{(1-r)^\alpha} = 0, \\ \frac{\mu(r)}{(1-r)^\beta} \text{ is increasing on } [\delta, 1), & \quad \lim_{r \rightarrow 1} \frac{\mu(r)}{(1-r)^\beta} = \infty. \end{aligned} \quad (1.2)$$

An $f \in H(\mathbb{B}_n)$ is said to belong to the weighted-type space, denoted by $H_\mu^\infty = H_\mu^\infty(\mathbb{B}_n)$, if

$$\|f\|_{H_\mu^\infty} = \sup_{z \in \mathbb{B}_n} \mu(|z|) |f(z)| < \infty, \quad (1.3)$$

where μ is normal on $[0, 1)$ (see [3]). H_μ^∞ is a Banach space with the norm $\|\cdot\|_{H_\mu^\infty}$.

The little weighted-type space, denoted by $H_{\mu,0}^{\infty}$, is the subspace of H_{μ}^{∞} consisting of those $f \in H_{\mu}^{\infty}$ such that

$$\lim_{|z| \rightarrow 1} \mu(|z|) |f(z)| = 0. \quad (1.4)$$

When $\mu(r) = (1-r^2)^{\alpha}$, the induced spaces H_{μ}^{∞} and $H_{\mu,0}^{\infty}$ become the (classical) weighted spaces H_{α}^{∞} and $H_{\alpha,0}^{\infty}$ respectively.

An $f \in H(\mathbb{B}_n)$ is said to belong to the logarithmic-type space H_{\log}^{∞} if

$$\|f\|_{H_{\log}^{\infty}} = \sup_{z \in \mathbb{B}_n} \frac{|f(z)|}{\log(e/(1-|z|^2))} < \infty. \quad (1.5)$$

It is easy to see that H_{\log}^{∞} becomes a Banach space under the norm $\|\cdot\|_{H_{\log}^{\infty}}$, and that the following inclusions hold:

$$H^{\infty} \subset \mathcal{B} \subset H_{\log}^{\infty} \subset H_{\alpha}^{\infty}, \quad \alpha > 0, \quad (1.6)$$

where \mathcal{B} is the Bloch space defined by

$$\mathcal{B} = \left\{ f \in H(\mathbb{B}_n) : \sup_{z \in \mathbb{B}_n} (1-|z|^2) |\Re f(z)| < \infty \right\}. \quad (1.7)$$

For some information on the Bloch and related spaces see, for example, [4–13] and the references therein. For some information on the space H_{\log}^{∞} in the unit disk see [14].

Let $u \in H(\mathbb{B}_n)$, and let φ be a holomorphic self-map of \mathbb{B}_n . For $f \in H(\mathbb{B}_n)$, the weighted composition operator uC_{φ} is defined by

$$(uC_{\varphi}f)(z) = u(z)f(\varphi(z)), \quad z \in \mathbb{B}_n. \quad (1.8)$$

The weighted composition operator can be regarded as a generalization of a multiplication operator and a composition operator, which is defined by $(C_{\varphi}f)(z) = f(\varphi(z))$. The work in [15] contains much information on this topic.

In the setting of the unit ball, Zhu studied the boundedness and compactness of the weighted composition operator between Bergman-type spaces and H^{∞} in [16]. More general results can be found in [17, 18]. Some necessary and sufficient conditions for the weighted composition operator to be bounded and compact between the Bloch space and H^{∞} are given in [19]. In the setting of the unit polydisk, some necessary and sufficient conditions for a weighted composition operator to be bounded and compact between the Bloch space and $H^{\infty}(\mathbb{B}_1^n)$ are given in [20, 21] (see also [22] for the case of composition operators). Other related results can be found, for example, in [3, 23–32].

In this paper, we study the weighted composition operator from H_{\log}^{∞} to the spaces H_{μ}^{∞} and H_{\log}^{∞} . Some necessary and sufficient conditions for the weighted composition operator uC_{φ} to be bounded and compact are given.

Throughout the paper, constants are denoted by C ; they are positive and may not be the same in every occurrence.

2. Main results and proofs

In this section, we give our main results and their proofs. Before stating these results, we need some auxiliary results, which are incorporated in the lemmas which follow.

Lemma 2.1. *Assume that $u \in H(\mathbb{B}_n)$, φ is a holomorphic self-map of \mathbb{B}_n , and μ is a normal function on $[0, 1)$. Then, $uC_\varphi : H_{\log}^\infty \rightarrow H_\mu^\infty$ is compact if and only if $uC_\varphi : H_{\log}^\infty \rightarrow H_\mu^\infty$ is bounded, and for any bounded sequence $(f_k)_{k \in \mathbb{N}}$ in H_{\log}^∞ which converges to zero uniformly on compact subsets of \mathbb{B}_n as $k \rightarrow \infty$, one has $\|uC_\varphi f_k\|_{H_\mu^\infty} \rightarrow 0$ as $k \rightarrow \infty$.*

The proof of Lemma 2.1 follows by standard arguments (see, e.g., [15, Proposition 3.11] as well as the proofs of the corresponding results in [7, 22, 33, 34]). Hence, we omit the details.

Lemma 2.2. *Assume that μ is normal. A closed set K in $H_{\mu,0}^\infty$ is compact if and only if it is bounded and satisfies*

$$\limsup_{|z| \rightarrow 1} \mu(|z|) |f(z)| = 0. \quad (2.1)$$

The proof of Lemma 2.2 is similar to the proof of Lemma 1 in [35]. We omit the details. Now, we are in a position to state and prove our main results.

Theorem 2.3. *Assume that $u \in H(\mathbb{B}_n)$, φ is a holomorphic self-map of \mathbb{B}_n , and μ is normal on $[0, 1)$. Then, $uC_\varphi : H_{\log}^\infty \rightarrow H_\mu^\infty$ is bounded if and only if*

$$M = \sup_{z \in \mathbb{B}_n} \mu(|z|) |u(z)| \log \frac{e}{1 - |\varphi(z)|^2} < \infty. \quad (2.2)$$

Proof. Assume that $uC_\varphi : H_{\log}^\infty \rightarrow H_\mu^\infty$ is bounded. For $a \in \mathbb{B}_n$, set

$$f_a(z) = \log \frac{e}{1 - \langle z, a \rangle}. \quad (2.3)$$

It is easy to see that $f_a \in H_{\log}^\infty$ and $\sup_{a \in \mathbb{B}_n} \|f_a\|_{H_{\log}^\infty} < \infty$.

For any $b \in \mathbb{B}_n$, we have

$$\begin{aligned} \infty &> \|uC_\varphi f_{\varphi(b)}\|_{H_\mu^\infty} \\ &= \sup_{z \in \mathbb{B}_n} \mu(|z|) |(uC_\varphi f_{\varphi(b)})(z)| \\ &= \sup_{z \in \mathbb{B}_n} \mu(|z|) |u(z)| |f_{\varphi(b)}(\varphi(z))| \\ &\geq \mu(|b|) |u(b)| \log \frac{e}{1 - |\varphi(b)|^2}, \end{aligned} \quad (2.4)$$

which implies (2.2).

Conversely, assume that (2.2) holds. Then, for any $f \in H_{\log}^\infty$, we have

$$\mu(|z|) |(uC_\varphi f)(z)| = \mu(|z|) |u(z)| |f(\varphi(z))| \leq \mu(|z|) |u(z)| \log \frac{e}{1 - |\varphi(z)|^2} \|f\|_{H_{\log}^\infty}. \quad (2.5)$$

Taking the supremum in (2.5) over \mathbb{B}_n and using condition (2.2), the boundedness of the operator $uC_\varphi : H_{\log}^\infty \rightarrow H_\mu^\infty$ follows, as desired. \square

Theorem 2.4. Assume that $u \in H(\mathbb{B}_n)$, φ is a holomorphic self-map of \mathbb{B}_n , and μ is a normal function on $[0, 1)$. Then, $uC_\varphi : H_{\log}^\infty \rightarrow H_\mu^\infty$ is compact if and only if $u \in H_\mu^\infty$ and

$$\lim_{|\varphi(z)| \rightarrow 1} \mu(|z|) |u(z)| \log \frac{e}{1 - |\varphi(z)|^2} = 0. \quad (2.6)$$

Proof. Assume that $uC_\varphi : H_{\log}^\infty \rightarrow H_\mu^\infty$ is compact. Then, it is obvious that $uC_\varphi : H_{\log}^\infty \rightarrow H_\mu^\infty$ is bounded. Taking the function $f(z) = 1 \in H_{\log}^\infty$, we see that $u \in H_\mu^\infty$. Let $(\varphi(z_k))_{k \in \mathbb{N}}$ be a sequence in \mathbb{B}_n such that $\lim_{k \rightarrow \infty} |\varphi(z_k)| = 1$. Set

$$f_k(z) = \left(\log \frac{e}{1 - \langle z, \varphi(z_k) \rangle} \right)^2 \left(\log \frac{e}{1 - |\varphi(z_k)|^2} \right)^{-1}, \quad k \in \mathbb{N}. \quad (2.7)$$

It is easy to see that $\sup_{k \in \mathbb{N}} \|f_k\|_{H_{\log}^\infty} < \infty$. Moreover, $f_k \rightarrow 0$ uniformly on compact subsets of \mathbb{B}_n as $k \rightarrow \infty$. By Lemma 2.1,

$$\lim_{k \rightarrow \infty} \|uC_\varphi f_k\|_{H_\mu^\infty} = 0. \quad (2.8)$$

We have

$$\|uC_\varphi f_k\|_{H_\mu^\infty} = \sup_{z \in \mathbb{B}_n} \mu(|z|) |u(z) f_k(\varphi(z))| \geq \mu(|z_k|) |u(z_k)| \log \frac{e}{1 - |\varphi(z_k)|^2}, \quad (2.9)$$

which together with (2.8) implies that

$$\lim_{k \rightarrow \infty} \mu(|z_k|) |u(z_k)| \log \frac{e}{1 - |\varphi(z_k)|^2} = 0. \quad (2.10)$$

This proves that (2.6) holds.

Conversely, assume that $u \in H_\mu^\infty$ and (2.6) holds. From this, it follows that (2.2) holds; hence $uC_\varphi : H_{\log}^\infty \rightarrow H_\mu^\infty$ is bounded. In order to prove that $uC_\varphi : H_{\log}^\infty \rightarrow H_\mu^\infty$ is compact, according to Lemma 2.1, it suffices to show that if $(f_k)_{k \in \mathbb{N}}$ is a bounded sequence in H_{\log}^∞ converging to 0 uniformly on compact subsets of \mathbb{B}_n , then

$$\lim_{k \rightarrow \infty} \|uC_\varphi f_k\|_{H_\mu^\infty} = 0. \quad (2.11)$$

Let $(f_k)_{k \in \mathbb{N}}$ be a bounded sequence in H_{\log}^∞ such that $f_k \rightarrow 0$ uniformly on compact subsets of \mathbb{B}_n as $k \rightarrow \infty$. By (2.6), we have that for any $\varepsilon > 0$, there is a constant $\delta \in (0, 1)$ such that

$$\mu(|z|) |u(z)| \log \frac{e}{1 - |\varphi(z)|^2} < \varepsilon \quad (2.12)$$

whenever $\delta < |\varphi(z)| < 1$. Let $K = \{w \in \mathbb{B}_n : |w| \leq \delta\}$. Equation (2.12) along with the fact that $u \in H_\mu^\infty$ implies

$$\begin{aligned}
\|uC_\varphi f_k\|_{H_\mu^\infty} &= \sup_{z \in \mathbb{B}_n} \mu(|z|) |(uC_\varphi f_k)(z)| \\
&= \sup_{z \in \mathbb{B}_n} \mu(|z|) |u(z) f_k(\varphi(z))| \\
&\leq \left(\sup_{\{z \in \mathbb{B}_n: |\varphi(z)| \leq \delta\}} + \sup_{\{z \in \mathbb{B}_n: \delta \leq |\varphi(z)| < 1\}} \right) \mu(|z|) |u(z)| |f_k(\varphi(z))| \\
&\leq \|u\|_{H_\mu^\infty} \sup_{w \in K} |f_k(w)| + \sup_{\{z \in \mathbb{B}_n: \delta \leq |\varphi(z)| < 1\}} \mu(|z|) |u(z)| \log \frac{e}{1 - |\varphi(z)|^2} \|f_k\|_{H_{\log}^\infty} \\
&\leq \|u\|_{H_\mu^\infty} \sup_{w \in K} |f_k(w)| + C\varepsilon.
\end{aligned} \tag{2.13}$$

Observe that K is a compact subset of \mathbb{B}_n so that

$$\lim_{k \rightarrow \infty} \sup_{w \in K} |f_k(w)| = 0. \tag{2.14}$$

With the aid of the above inequality, we can deduce that

$$\lim_{k \rightarrow \infty} \sup \|uC_\varphi f_k\|_{H_\mu^\infty} \leq C\varepsilon \tag{2.15}$$

by letting $k \rightarrow \infty$. Since ε is an arbitrary positive number, it follows that the last limit is equal to zero. Therefore, $uC_\varphi : H_{\log}^\infty \rightarrow H_{\mu,0}^\infty$ is compact. The proof is complete. \square

Theorem 2.5. *Assume that $u \in H(\mathbb{B}_n)$, φ is a holomorphic self-map of \mathbb{B}_n , and μ is a normal function on $[0, 1)$. Then, $uC_\varphi : H_{\log}^\infty \rightarrow H_{\mu,0}^\infty$ is compact if and only if $u \in H_{\mu,0}^\infty$ and*

$$\lim_{|\varphi(z)| \rightarrow 1} \mu(|z|) |u(z)| \log \frac{e}{1 - |\varphi(z)|^2} = 0. \tag{2.16}$$

Proof. Assume that $uC_\varphi : H_{\log}^\infty \rightarrow H_{\mu,0}^\infty$ is compact. Then, it is clear that $uC_\varphi : H_{\log}^\infty \rightarrow H_\mu^\infty$ is compact, and hence (2.16) holds. In addition, taking the function given by $f(z) = 1$, we get $u \in H_{\mu,0}^\infty$.

Conversely, suppose that $u \in H_{\mu,0}^\infty$ and (2.16) holds. In the proof of the implication we follow the lines, for example, of the proof of Lemma 4.2 in [24]. From (2.16), it follows that for every $\varepsilon > 0$, there exists a $\delta \in (0, 1)$ such that

$$\mu(|z|) |u(z)| \log \frac{e}{1 - |\varphi(z)|^2} < \varepsilon \tag{2.17}$$

when $\delta < |\varphi(z)| < 1$. From the assumption $u \in H_{\mu,0}^\infty$, we have that for the above ε , there exists an $r \in (0, 1)$ such that when $r < |z| < 1$,

$$\mu(|z|) |u(z)| < \frac{\varepsilon}{\log(e/(1 - \delta^2))}. \tag{2.18}$$

Therefore, if $r < |z| < 1$ and $\delta < |\varphi(z)| < 1$, we obtain

$$\mu(|z|)|u(z)| \log \frac{e}{1 - |\varphi(z)|^2} < \varepsilon. \quad (2.19)$$

If $|\varphi(z)| \leq \delta$ and $r < |z| < 1$, we have that

$$\mu(|z|)|u(z)| \log \frac{e}{1 - |\varphi(z)|^2} \leq \mu(|z|)|u(z)| \log \frac{e}{1 - \delta^2} < \varepsilon. \quad (2.20)$$

Combining (2.19) with (2.20), we get

$$\lim_{|z| \rightarrow 1} \mu(|z|)|u(z)| \log \frac{e}{1 - |\varphi(z)|^2} = 0. \quad (2.21)$$

On the other hand, from (1.5) we have that

$$\mu(|z|)|(uC_\varphi f)(z)| \leq \mu(|z|)|u(z)| \log \frac{e}{1 - |\varphi(z)|^2} \|f\|_{H_{\log}^\infty}.$$

Taking the supremum in the above inequality over all $f \in H_{\log}^\infty$ such that $\|f\|_{H_{\log}^\infty} \leq 1$, then letting $|z| \rightarrow 1$, by (2.21) it follows that

$$\lim_{|z| \rightarrow 1} \sup_{\|f\|_{H_{\log}^\infty} \leq 1} \mu(|z|)|(uC_\varphi(f))(z)| = 0. \quad (2.22)$$

From this and by employing Lemma 2.2, we see that $uC_\varphi : H_{\log}^\infty \rightarrow H_{\mu,0}^\infty$ is compact. The proof is complete. \square

Similar to the proofs of Theorems 2.3 and 2.4, we easily get the following two results. We also omit their proofs.

Theorem 2.6. *Assume that $u \in H(\mathbb{B}_n)$ and φ is a holomorphic self-map of \mathbb{B}_n . Then, the following statements hold.*

(a) $uC_\varphi : H_{\log}^\infty \rightarrow H_{\log}^\infty$ is bounded if and only if

$$\sup_{z \in \mathbb{B}_n} \frac{|u(z)| \log(e/(1 - |\varphi(z)|^2))}{\log(e/(1 - |z|^2))} < \infty. \quad (2.23)$$

(b) $uC_\varphi : H_{\log}^\infty \rightarrow H_{\log}^\infty$ is compact if and only if $u \in H_{\log}^\infty$ and

$$\lim_{|\varphi(z)| \rightarrow 1} \frac{|u(z)| \log(e/(1 - |\varphi(z)|^2))}{\log(e/(1 - |z|^2))} = 0. \quad (2.24)$$

Theorem 2.7. *Assume that $u \in H(\mathbb{B}_n)$ and φ is a holomorphic self-map of \mathbb{B}_n . Then, the following statements hold.*

(a) $uC_\varphi : H_{\log}^\infty \rightarrow H^\infty$ is bounded if and only if

$$\sup_{z \in \mathbb{B}_n} |u(z)| \log \frac{e}{1 - |\varphi(z)|^2} < \infty. \quad (2.25)$$

(b) $uC_\varphi : H_{\log}^\infty \rightarrow H^\infty$ is compact if and only if $u \in H^\infty$ and

$$\lim_{|\varphi(z)| \rightarrow 1} |u(z)| \log \frac{e}{1 - |\varphi(z)|^2} = 0. \quad (2.26)$$




Acknowledgment

X. Fu is supported in part by the Natural Science Foundation of Guangdong Province (no. 73006147).

References

- [1] Z. Hu, "Extended Cesàro operators on mixed norm spaces," *Proceedings of the American Mathematical Society*, vol. 131, no. 7, pp. 2171–2179, 2003.
- [2] A. L. Shields and D. L. Williams, "Bonded projections, duality, and multipliers in spaces of analytic functions," *Transactions of the American Mathematical Society*, vol. 162, pp. 287–302, 1971.
- [3] S. Stević, "Norm of weighted composition operators from Bloch space to H_μ^∞ on the unit ball," *Ars Combinatoria*, vol. 88, pp. 125–127, 2008.
- [4] D. D. Clahane and S. Stević, "Norm equivalence and composition operators between Bloch/Lipschitz spaces of the ball," *Journal of Inequalities and Applications*, vol. 2006, Article ID 61018, 11 pages, 2006.
- [5] S. Li, "Fractional derivatives of Bloch-type functions," *Siberian Mathematical Journal*, vol. 46, no. 2, pp. 308–314, 2005.
- [6] S. Li, "Derivative-free characterizations of Bloch spaces," *Journal of Computational Analysis and Applications*, vol. 10, no. 2, pp. 253–258, 2008.
- [7] S. Li and S. Stević, "Riemann-Stieltjes-type integral operators on the unit ball in \mathbb{C}^n ," *Complex Variables and Elliptic Equations*, vol. 52, no. 6, pp. 495–517, 2007.
- [8] S. Li and S. Stević, "Some characterizations of the Besov space and the α -Bloch space," *Journal of Mathematical Analysis and Applications*. In press.
- [9] S. Li and H. Wulan, "Characterizations of α -Bloch spaces on the unit ball," *Journal of Mathematical Analysis and Applications*, vol. 343, no. 1, pp. 58–63, 2008.
- [10] S. Stević, "On an integral operator on the unit ball in \mathbb{C}^n ," *Journal of Inequalities and Applications*, no. 1, pp. 81–88, 2005.
- [11] S. Stević, "On Bloch-type functions with Hadamard gaps," *Abstract and Applied Analysis*, vol. 2007, Article ID 39176, 8 pages, 2007.
- [12] R. M. Timoney, "Bloch functions in several complex variables—I," *The Bulletin of the London Mathematical Society*, vol. 12, no. 4, pp. 241–267, 1980.
- [13] K. Zhu, *Spaces of Holomorphic Functions in the Unit Ball*, vol. 226 of *Graduate Texts in Mathematics*, Springer, New York, NY, USA, 2005.
- [14] D. Girela, J. Peláez, F. Pérez-González, and J. Rättyä, "Carleson measures for the Bloch space," preprint, 2008.
- [15] C. C. Cowen and B. D. MacCluer, *Composition Operators on Spaces of Analytic Functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, Fla, USA, 1995.
- [16] X. Zhu, "Weighted composition operators between H^∞ and Bergman type spaces," *Korean Mathematical Society*, vol. 21, no. 4, pp. 719–727, 2006.
- [17] S. Stević, "Weighted composition operators between mixed norm spaces and H_α^∞ spaces in the unit ball," *Journal of Inequalities and Applications*, vol. 2007, Article ID 28629, 9 pages, 2007.
- [18] X. Tang, "Weighted composition operators between Bers-type spaces and Bergman spaces," *Applied Mathematics: A Journal of Chinese Universities*, vol. 22, no. 1, pp. 61–68, 2007.
- [19] S. Li and S. Stević, "Weighted composition operators between H^∞ and α -Bloch space in the unit ball," to appear in *Taiwanese Journal of Mathematics*.
- [20] S. Li and S. Stević, "Weighted composition operators from H^∞ to the Bloch space on the polydisc," *Abstract and Applied Analysis*, vol. 2007, Article ID 48478, 13 pages, 2007.
- [21] S. Li and S. Stević, "Weighted composition operators from α -Bloch space to H^∞ on the polydisc," *Numerical Functional Analysis and Optimization*, vol. 28, no. 7-8, pp. 911–925, 2007.
- [22] S. Stević, "Composition operators between H^∞ and α -Bloch spaces on the polydisc," *Zeitschrift für Analysis und ihre Anwendungen*, vol. 25, no. 4, pp. 457–466, 2006.
- [23] S. Li and S. Stević, "Composition followed by differentiation between Bloch type spaces," *Journal of Computational Analysis and Applications*, vol. 9, no. 2, pp. 195–205, 2007.
- [24] S. Li and S. Stević, "Weighted composition operators from Bergman-type spaces into Bloch spaces," *Proceedings of the Indian Academy of Sciences. Mathematical Sciences*, vol. 117, no. 3, pp. 371–385, 2007.

- [25] S. Li and S. Stević, "Generalized composition operators on Zygmund spaces and Bloch type spaces," *Journal of Mathematical Analysis and Applications*, vol. 338, no. 2, pp. 1282–1295, 2008.
- [26] S. Li and S. Stević, "Products of composition and integral type operators from H^∞ to the Bloch space," *Complex Variables and Elliptic Equations*, vol. 53, no. 5, pp. 463–474, 2008.
- [27] S. Li and S. Stević, "Products of Volterra type operator and composition operator from H^∞ and Bloch spaces to the Zygmund space," *Journal of Mathematical Analysis and Applications*, vol. 345, no. 1, pp. 40–52, 2008.
- [28] S. Ohno, K. Stroethoff, and R. Zhao, "Weighted composition operators between Bloch-type spaces," *The Rocky Mountain Journal of Mathematics*, vol. 33, no. 1, pp. 191–215, 2003.
- [29] A. Montes-Rodríguez, "Weighted composition operators on weighted Banach spaces of analytic functions," *Journal of the London Mathematical Society*, vol. 61, no. 3, pp. 872–884, 2000.
- [30] S.-I. Ueki and L. Luo, "Compact weighted composition operators and multiplication operators between Hardy spaces," *Abstract and Applied Analysis*, vol. 2008, Article ID 196498, 12 pages, 2008.
- [31] S. Ye, "Weighted composition operator between the little α -Bloch spaces and the logarithmic Bloch," *Journal of Computational Analysis and Applications*, vol. 10, no. 2, pp. 243–252, 2008.
- [32] X. Zhu, "Generalized weighted composition operators from Bloch type spaces to weighted Bergman spaces," *Indian Journal of Mathematics*, vol. 49, no. 2, pp. 139–150, 2007.
- [33] S. Stević, "Boundedness and compactness of an integral operator on a weighted space on the polydisc," *Indian Journal of Pure and Applied Mathematics*, vol. 37, no. 6, pp. 343–355, 2006.
- [34] S. Stević, "Boundedness and compactness of an integral operator in a mixed norm space on the polydisk," *Sibirskii Matematicheskii Zhurnal*, vol. 48, no. 3, pp. 694–706, 2007.
- [35] K. Madigan and A. Matheson, "Compact composition operators on the Bloch space," *Transactions of the American Mathematical Society*, vol. 347, no. 7, pp. 2679–2687, 1995.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

