

Research Article

Approximate Riesz Algebra-Valued Derivations

Faruk Polat

Department of Mathematics, Faculty of Science, Çankırı Karatekin University, 18000 Çankırı, Turkey

Correspondence should be addressed to Faruk Polat, faruk.polat@gmail.com

Received 7 May 2012; Accepted 26 August 2012

Academic Editor: Janusz Brzdęk

Copyright © 2012 Faruk Polat. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let F be a Riesz algebra with an extended norm $\|\cdot\|_u$ such that $(F, \|\cdot\|_u)$ is complete. Also, let $\|\cdot\|_v$ be another extended norm in F weaker than $\|\cdot\|_u$ such that whenever (a) $x_n \rightarrow x$ and $x_n \cdot y \rightarrow z$ in $\|\cdot\|_v$, then $z = x \cdot y$; (b) $y_n \rightarrow y$ and $x \cdot y_n \rightarrow z$ in $\|\cdot\|_v$, then $z = x \cdot y$. Let ε and $\delta > 0$ be two nonnegative real numbers. Assume that a map $f : F \rightarrow F$ satisfies $\|f(x+y) - f(x) - f(y)\|_u \leq \varepsilon$ and $\|f(x \cdot y) - x \cdot f(y) - f(x) \cdot y\|_v \leq \delta$ for all $x, y \in F$. In this paper, we prove that there exists a unique derivation $d : F \rightarrow F$ such that $\|f(x) - d(x)\|_u \leq \varepsilon$, $(x \in F)$. Moreover, $x \cdot (f(y) - d(y)) = 0$ for all $x, y \in F$.

1. Introduction

Let E and E' be Banach spaces and let $\delta > 0$. A function $f : E \rightarrow E'$ is called δ -additive if $\|f(x+y) - f(x) - f(y)\| < \delta$ for all $x, y \in E$. The well-known problem of stability of functional equation $f(x+y) = f(x) + f(y)$ started with the following question of Ulam [1]. Does there exist for each $\varepsilon > 0$, a $\delta > 0$ such that, to each δ -additive function f of E into E' there corresponds an additive function l of E into E' satisfying the inequality $\|f(x) - l(x)\| \leq \varepsilon$ for each $x \in E$? In 1941, Hyers [2] answered this question in the affirmative way and showed that δ may be taken equal to ε . The answer of Hyers is presented in a great number of articles and books. For the theory of the stability of functional equations see Hyers et al [3].

Let F be an algebra. A mapping $d : F \rightarrow F$ is called a derivation if and only if it satisfies the following functional equations:

$$d(a+b) = d(a) + d(b), \quad (1.1)$$

$$d(ab) = ad(b) + d(a)b, \quad (1.2)$$

for all $a, b \in F$.

The stability of derivations was first studied by Jun and Park [4]. Further, approximate derivations were investigated by a number of mathematicians (see, e.g., [5–7]).

The aim of the present paper is to examine the stability problem of derivations for Riesz algebras with extended norms.

2. Preliminaries

A vector space F with a partial order \leq satisfying the following two conditions:

- (1) $x \leq y \Rightarrow \alpha x + z \leq \alpha y + z$ for all $z \in F$ and $0 \leq \alpha \in \mathbb{R}$,
- (2) for all $x, y \in F$, the supremum $x \vee y$ and infimum $x \wedge y$ exist in F (hence, the modulus $|x| := x \vee (-x)$ exists for each $x \in F$),

is called a Riesz space or vector lattice. Typical examples of Riesz spaces are provided by the function spaces. $C(K)$ the spaces of real valued continuous functions on a topological space K , l_p real valued absolutely summable sequences, c the spaces of real valued convergent sequences, and c_0 the spaces of real valued sequences converging to zero are natural examples of Riesz spaces under the pointwise ordering. A Riesz space F is called Archimedean if $0 \leq u, v \in F$ and $nu \leq v$ for each $n \in \mathbb{N}$ imply $u = 0$. A subset S in a Riesz space F is said to be solid if it follows from $|u| \leq |v|$ in F and $v \in S$ that $u \in S$. A solid linear subspace of a Riesz space F is called an ideal. Every subset D of a Riesz space F is included in a smallest ideal F_D , called ideal generated by D . A principal ideal of a Riesz space F is any ideal generated by a singleton $\{u\}$. This ideal will be denoted by I_u . It is easy to see that

$$I_u = \{v \in F : \lambda \geq 0 \text{ such that } |v| \leq \lambda|u|\}. \quad (2.1)$$

Let F be a Riesz space and $0 \leq u \in F$. Firstly, we give the following definition.

Definition 2.1. (1) The sequence (x_n) in F is said to be u -uniformly convergent to the element $x \in F$ whenever, for every $\varepsilon > 0$, there exists n_0 such that $|x_{n_0+k} - x| \leq \varepsilon u$ holds for each k .

(2) The sequence (x_n) in F is said to be relatively uniformly convergent to x whenever x_n converges u -uniformly to $x \in F$ for some $0 \leq u \in F$.

When dealing with relative uniform convergence in an Archimedean Riesz space F , it is natural to associate with every positive element $u \in F$ an extended norm $\|\cdot\|_u$ in F by the following formula:

$$\|x\|_u = \inf\{\lambda \geq 0 : |x| \leq \lambda u\} \quad (x \in F). \quad (2.2)$$

Note that $\|x\|_u < \infty$ if and only if $x \in I_u$. Also $|x| \leq \delta u$ if and only if $\|x\|_u \leq \delta$.

A Banach lattice is a vector lattice F that is simultaneously a Banach space whose norm is monotone in the following sense.

For all $x, y \in F$, $|x| \leq |y|$ implies $\|x\| \leq \|y\|$. Hence, $\|x\| = \||x|\|$ for all $x \in F$.

The sequence (x_n) in $(F, \|\cdot\|_u)$ is called an extended u -normed Cauchy sequence, if for every $\varepsilon > 0$ there exists k such that $\|x_{n+k} - x_{m+k}\|_u < \varepsilon$ for all m, n . If every extended u -normed Cauchy sequence is convergent in F , then F is called an extended u -normed Banach lattice.

A Riesz space F is called a Riesz algebra or a lattice ordered algebra if there exists an associative multiplication in F with the usual algebra properties such that $0 \leq u \cdot v$ for all $0 \leq u, v \in F$.

For more detailed information about Riesz spaces, the reader can consult the book *Riesz Spaces* by Luxemburg and Zaanen [8]. In the sequel, all the Riesz spaces are assumed to be Archimedean.

3. Main Result

Recently, Polat [9] generalized the Hyers' result [2] to Riesz spaces with extended norms and proved the following.

Theorem 3.1. *Let E be a linear space and F a Riesz space equipped with an extended norm $\|\cdot\|_u$ such that the space $(F, \|\cdot\|_u)$ is complete. If, for some $\delta > 0$, a map $f : E \rightarrow (F, \|\cdot\|_u)$ is δ -additive, then limit $l(x) = \lim_{n \rightarrow \infty} f(2^n x)/2^n$ exists for each $x \in E$. $l(x)$ is the unique additive function satisfying the inequality $\|f(x) - l(x)\|_u \leq \delta$ for all $x \in E$.*

By using Theorem 3.1, we give the main result of the paper as follows.

Theorem 3.2. *Let F be a Riesz algebra with an extended norm $\|\cdot\|_u$ such that $(F, \|\cdot\|_u)$ is complete. Also, let $\|\cdot\|_v$ be another extended norm in F weaker than $\|\cdot\|_u$ such that whenever*

- (a) $x_n \rightarrow x$ and $x_n \cdot y \rightarrow z$ in $\|\cdot\|_v$, then $z = x \cdot y$;
- (b) $y_n \rightarrow y$ and $x \cdot y_n \rightarrow z$ in $\|\cdot\|_v$, then $z = x \cdot y$.

Let ε and δ be two nonnegative real numbers. Assume that a map $f : F \rightarrow F$ satisfies

$$\|f(x + y) - f(x) - f(y)\|_u \leq \varepsilon, \quad (3.1)$$

$$\|f(x \cdot y) - x \cdot f(y) - f(x) \cdot y\|_v \leq \delta, \quad (3.2)$$

for all $x, y \in F$. Then, there exists a unique derivation $d : F \rightarrow F$ such that $\|f(x) - d(x)\|_u \leq \varepsilon$, ($x \in F$). Moreover, $x \cdot (f(y) - d(y)) = 0$ for all $x, y \in F$.

Proof. By Condition (3.1), Theorem 3.1 shows that there exists a unique additive function $d : F \rightarrow F$ such that

$$\|f(x) - d(x)\|_u \leq \varepsilon, \quad (3.3)$$

for each $x \in F$. It is enough to show that d satisfies Condition (1.2). The inequality (3.3) implies that

$$\|f(nx) - d(nx)\|_u \leq \varepsilon \quad (x \in F, n \in \mathbb{N}). \quad (3.4)$$

By the additivity of d , we then have

$$\left\| \frac{1}{n} f(nx) - d(x) \right\|_u \leq \frac{1}{n} \varepsilon \quad (x \in F, n \in \mathbb{N}), \quad (3.5)$$

which means that

$$d(x) = \lim_{n \rightarrow \infty} \frac{1}{n} f(nx), \quad (x \in F), \quad (3.6)$$

with respect to $\|\cdot\|_u$ norm and so is with respect to $\|\cdot\|_v$ norm. Condition (3.2) implies that the function $r : F \times F \rightarrow F$ defined by $r(x, y) = f(x \cdot y) - x \cdot f(y) - f(x) \cdot y$ is bounded. Hence

$$\lim_{n \rightarrow \infty} \frac{1}{n} r(nx, y) = 0, \quad (x, y \in F), \quad (3.7)$$

with respect to $\|\cdot\|_v$ norm. Applying (3.6) and (3.7), we have

$$d(x \cdot y) = x \cdot f(y) + d(x) \cdot y, \quad (x, y \in F). \quad (3.8)$$

Indeed, we have the following with respect to $\|\cdot\|_v$ norm,

$$\begin{aligned} d(x \cdot y) &= \lim_{n \rightarrow \infty} \frac{1}{n} f(n(x \cdot y)) = \lim_{n \rightarrow \infty} \frac{1}{n} f((nx) \cdot y) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} (nx \cdot f(y) + f(nx) \cdot y + r(nx, y)) \\ &= \lim_{n \rightarrow \infty} \left(x \cdot f(y) + \frac{f(nx)}{n} \cdot y + \frac{r(nx, y)}{n} \right) \\ &= x \cdot f(y) + d(x) \cdot y, \quad (x, y \in F). \end{aligned} \quad (3.9)$$

Let $x, y \in F$ and $n \in \mathbb{N}$ be fixed. Then using (3.8) and additivity of d , we have

$$\begin{aligned} x \cdot f(ny) + nd(x) \cdot y &= x \cdot f(ny) + d(x) \cdot ny = d(x \cdot ny) \\ &= d(nx \cdot y) = nx \cdot f(y) + d(nx) \cdot y \\ &= nx \cdot f(y) + nd(x) \cdot y. \end{aligned} \quad (3.10)$$

Therefore,

$$x \cdot f(y) = x \cdot \frac{f(ny)}{n}, \quad (x, y \in F, n \in \mathbb{N}). \quad (3.11)$$

Sending n to infinity, by (3.6), we see that

$$x \cdot f(y) = x \cdot d(y), \quad (x, y \in F). \quad (3.12)$$

Combining this formula with (3.8), we have that d satisfies (1.2) which is the desired result. Moreover, the last formula yields $x \cdot (f(y) - d(y)) = 0$ for all $x, y \in F$. \square

References

- [1] S. M. Ulam, *A Collection of Mathematical Problems*, Interscience Publishers, London, UK, 1960.
- [2] D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, pp. 222–224, 1941.
- [3] D. H. Hyers, G. Isac, and Th. M. Rassias, *Stability of Functional Equations in Several Variables*, Birkhauser Boston, Boston, Mass, USA, 1998.
- [4] K.-W. Jun and D.-W. Park, "Almost derivations on the Banach algebra $C^n[0, 1]$," *Bulletin of the Korean Mathematical Society*, vol. 33, no. 3, pp. 359–366, 1996.
- [5] M. S. Moslehian, "Ternary derivations, stability and physical aspects," *Acta Applicandae Mathematicae*, vol. 100, no. 2, pp. 187–199, 2008.
- [6] M. E. Gordji and M. S. Moslehian, "A trick for investigation of approximate derivations," *Mathematical Communications*, vol. 15, no. 1, pp. 99–105, 2010.
- [7] A. Fošner, "On the generalized Hyers-Ulam stability of module left (m, n) derivations," *Aequationes Mathematicae*, vol. 84, no. 1-2, pp. 91–98, 2012.
- [8] W. A. J. Luxemburg and A. C. Zaanen, *Riesz Spaces*, vol. 1, North-Holland Publishing Company, Amsterdam, The Netherlands, 1971.
- [9] F. Polat, "Some generalizations of Ulam-Hyers stability functional equations to Riesz algebras," *Abstract and Applied Analysis*, vol. 2012, Article ID 653508, 9 pages, 2012.

