

Research Article

Coupled Coincidence Point Theorem in Partially Ordered Metric Spaces via Implicit Relation

Nguyen Manh Hung,¹ Erdal Karapınar,² and Nguyen Van Luong¹

¹ Department of Natural Sciences, Hong Duc University, Thanh Hoa 41000, Vietnam

² Department of Mathematics, Atilim University, 06586 Incek, Ankara, Turkey

Correspondence should be addressed to Erdal Karapınar, erdalkarapinar@yahoo.com

Received 28 February 2012; Revised 19 April 2012; Accepted 19 April 2012

Academic Editor: Stefan Siegmund

Copyright © 2012 Nguyen Manh Hung et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We prove a coupled coincidence point theorem for mappings $F : X \times X \rightarrow X$ and $g : X \rightarrow X$, where F has the mixed g -monotone property, in partially ordered metric spaces via implicit relations. Our result extends and improves several results in the literature. Examples are also given to illustrate our work.

1. Introduction and Preliminaries

The notion of coupled fixed point was introduced by Guo and Lakshmikantham [1] in 1987. Later, Bhaskar and Lakshmikantham [2] defined the notions of mixed monotone mapping and proved some coupled fixed point theorems for the mixed monotone mappings. In this pioneer paper [2], they also discussed the existence and uniqueness of solution for a periodic boundary value problem. We start with recalling these basic concepts.

Definition 1.1 (see [2]). Let (X, \leq) be a partially ordered set and $F : X \times X \rightarrow X$. The mapping F is said to have the mixed monotone property if $F(x, y)$ is monotone nondecreasing in x and is monotone nonincreasing in y , that is, for any $x, y \in X$,

$$\begin{aligned} x_1, x_2 \in X, \quad x_1 \leq x_2 &\implies F(x_1, y) \leq F(x_2, y), \\ y_1, y_2 \in X, \quad y_1 \leq y_2 &\implies F(x, y_1) \geq F(x, y_2). \end{aligned} \tag{1.1}$$

Definition 1.2 (see [2]). An element $(x, y) \in X \times X$ is called a coupled fixed point of the mapping $F : X \times X \rightarrow X$ if

$$x = F(x, y), \quad y = F(y, x). \quad (1.2)$$

The main results of Bhaskar and Lakshmikantham in [2] are the following theorems.

Theorem 1.3 (see [2]). Let (X, \leq) be a partially ordered set and suppose there exists a metric d on X such that (X, d) is a complete metric space. Let $F : X \times X \rightarrow X$ be a continuous mapping having the mixed monotone property on X . Assume that there exists a $k \in [0, 1)$ with

$$d(F(x, y), F(u, v)) \leq \frac{k}{2} [d(x, u) + d(y, v)], \quad (1.3)$$

for all $x \geq u$ and $y \leq v$. If there exist two elements $x_0, y_0 \in X$ with

$$x_0 \leq F(x_0, y_0), \quad y_0 \geq F(y_0, x_0), \quad (1.4)$$

then there exist $x, y \in X$ such that

$$x = F(x, y), \quad y = F(y, x). \quad (1.5)$$

Theorem 1.4 (see [2]). Let (X, \leq) be a partially ordered set and suppose there exists a metric d on X such that (X, d) is a complete metric space. Assume that X has the following property:

- (i) if a nondecreasing sequence $\{x_n\} \rightarrow x$, then $x_n \leq x$ for all n ,
- (ii) if a nonincreasing sequence $\{y_n\} \rightarrow y$, then $y \leq y_n$ for all n .

Let $F : X \times X \rightarrow X$ be a mapping having the mixed monotone property on X . Assume that there exists a $k \in [0, 1)$ with

$$d(F(x, y), F(u, v)) \leq \frac{k}{2} [d(x, u) + d(y, v)], \quad (1.6)$$

for all $x \geq u$ and $y \leq v$. If there exist two elements $x_0, y_0 \in X$ with

$$x_0 \leq F(x_0, y_0), \quad y_0 \geq F(y_0, x_0), \quad (1.7)$$

then there exist $x, y \in X$ such that

$$x = F(x, y), \quad y = F(y, x). \quad (1.8)$$

Afterwards, a number of coupled coincidence/fixed point theorems and their application to integral equations, matrix equations, and periodic boundary value problem have been established (e.g., see [3–28] and references therein). In particular, Lakshmikantham and Ćirić

[7] established coupled coincidence and coupled fixed point theorems for two mappings $F : X \times X \rightarrow X$ and $g : X \rightarrow X$, where F has the mixed g -monotone property and the functions F and g commute, as an extension of the fixed point results in [2]. Choudhury and Kundu in [15] introduced the concept of compatibility and proved the result established in [7] under a different set of conditions. Precisely, they established their result by assuming that F and g are compatible mappings. For the sake of completeness, we remind these characterizations.

Definition 1.5 (see [7]). Let (X, \leq) be a partially ordered set and let $F : X \times X \rightarrow X$ and $g : X \rightarrow X$ are two mappings. We say F has the mixed g -monotone property if $F(x, y)$ is g -nondecreasing in its first argument and is g -nonincreasing in its second argument, that is, for any $x, y \in X$,

$$\begin{aligned} x_1, x_2 \in X, \quad gx_1 \leq gx_2 &\implies F(x_1, y) \leq F(x_2, y), \\ y_1, y_2 \in X, \quad gy_1 \leq gy_2 &\implies F(x, y_1) \geq F(x, y_2). \end{aligned} \tag{1.9}$$

Definition 1.6 (see [7]). An element $(x, y) \in X \times X$ is called a coupled coincident point of the mappings $F : X \times X \rightarrow X$ and $g : X \rightarrow X$ if

$$gx = F(x, y), \quad gy = F(y, x). \tag{1.10}$$

Definition 1.7 (see [15]). The mappings F and g where $F : X \times X \rightarrow X, g : X \rightarrow X$ are said to be compatible if

$$\begin{aligned} \lim_{n \rightarrow \infty} d(gF(x_n, y_n), F(gx_n, gy_n)) &= 0, \\ \lim_{n \rightarrow \infty} d(gF(y_n, x_n), F(gy_n, gx_n)) &= 0, \end{aligned} \tag{1.11}$$

where $\{x_n\}$ and $\{y_n\}$ are sequences in X such that $\lim_{n \rightarrow \infty} F(x_n, y_n) = \lim_{n \rightarrow \infty} gx_n = x$ and $\lim_{n \rightarrow \infty} F(y_n, x_n) = \lim_{n \rightarrow \infty} gy_n = y$ for all $x, y \in X$ are satisfied.

Luong and Thuan [11] slightly extended the concept of compatible mappings into the context of partially ordered metric spaces, namely, O -compatible mappings and proved some coupled coincidence point theorems for such mappings in partially ordered generalized metric spaces.

The concept of O -compatible mappings is stated as follows.

Definition 1.8 (cf. [11]). Let (X, \leq, d) be a partially ordered metric space. The mappings $F : X \times X \rightarrow X$ and $g : X \rightarrow X$ are said to be O -compatible if

$$\begin{aligned} \lim_{n \rightarrow \infty} d(gF(x_n, y_n), F(gx_n, gy_n)) &= 0, \\ \lim_{n \rightarrow \infty} d(gF(y_n, x_n), F(gy_n, gx_n)) &= 0, \end{aligned} \tag{1.12}$$

where $\{x_n\}$ and $\{y_n\}$ are sequences in X such that $\{gx_n\}, \{gy_n\}$ are monotone and

$$\begin{aligned}\lim_{n \rightarrow \infty} F(x_n, y_n) &= \lim_{n \rightarrow \infty} gx_n = x, \\ \lim_{n \rightarrow \infty} F(y_n, x_n) &= \lim_{n \rightarrow \infty} gy_n = y,\end{aligned}\tag{1.13}$$

for all $x, y \in X$ are satisfied.

Let (X, \leq, d) be a partially metric space. If $F : X \times X \rightarrow X$ and $g : X \rightarrow X$ are compatible then they are O -compatible. However, the converse is not true. The following example shows that there exist mappings that are O -compatible but not compatible.

Example 1.9 (see [11]). Let $X = \{0\} \cup [1/2, 2]$ with the usual metric $d(x, y) = |x - y|$, for all $x, y \in X$. We consider the following order relation on X :

$$x, y \in X \quad x \leq y \iff x = y \quad \text{or} \quad (x, y) \in \{(0, 0), (0, 1), (1, 1)\}.\tag{1.14}$$

Let $F : X \times X \rightarrow X$ be given by

$$F(x, y) = \begin{cases} 0, & \text{if } x, y \in \{0\} \cup \left[\frac{1}{2}, 1\right], \\ 1, & \text{otherwise,} \end{cases}\tag{1.15}$$

and $g : X \rightarrow X$ be defined by

$$gx = \begin{cases} 0, & \text{if } x = 0, \\ 1, & \text{if } \frac{1}{2} \leq x \leq 1, \\ 2 - x, & \text{if } 1 < x \leq \frac{3}{2}, \\ \frac{1}{2}, & \text{if } \frac{3}{2} < x \leq 2. \end{cases}\tag{1.16}$$

Then F and g are O -compatible but not compatible.

Indeed, let $\{x_n\}, \{y_n\}$ in X such that $\{gx_n\}, \{gy_n\}$ are monotone and

$$\begin{aligned}\lim_{n \rightarrow \infty} F(x_n, y_n) &= \lim_{n \rightarrow \infty} gx_n = x, \\ \lim_{n \rightarrow \infty} F(y_n, x_n) &= \lim_{n \rightarrow \infty} gy_n = y,\end{aligned}\tag{1.17}$$

for some $x, y \in X$. Since $F(x_n, y_n) = F(y_n, x_n) \in \{0, 1\}$ for all n , $x = y \in \{0, 1\}$. The case $x = y = 1$ is impossible. In fact, if $x = y = 1$. Then since $\{gx_n\}, \{gy_n\}$ are monotone, $gx_n = gy_n = 1$ for all $n \geq n_1$, for some n_1 . That is $x_n, y_n \in [1/2, 1]$ for all $n \geq n_1$. This implies $F(x_n, y_n) = F(y_n, x_n) = 0$, for all $n \geq n_1$, which is a contradiction. Thus $x = y = 0$. That implies $gx_n = gy_n = 0$ for all $n \geq n_2$, for some n_2 . That is $x_n = y_n = 0$ for all $n \geq n_2$. Thus, for all $n \geq n_2$,

$$d(gF(x_n, y_n), F(gx_n, gy_n)) = d(gF(y_n, x_n), F(gy_n, gx_n)) = 0.\tag{1.18}$$

Hence

$$\begin{aligned} \lim_{n \rightarrow \infty} d(gF(x_n, y_n), F(gx_n, gy_n)) &= 0, \\ \lim_{n \rightarrow \infty} d(gF(y_n, x_n), F(gy_n, gx_n)) &= 0 \end{aligned} \tag{1.19}$$

hold. Therefore F and g are O -compatible.

Now let $\{x_n\}, \{y_n\}$ in X be defined by

$$x_n = y_n = 1 + \frac{1}{n+1}, \quad n = 1, 2, 3, \dots \tag{1.20}$$

We have

$$\begin{aligned} F(x_n, y_n) = F(y_n, x_n) &= F\left(1 + \frac{1}{n+1}, 1 + \frac{1}{n+1}\right) = 1, \\ gx_n = gy_n = g\left(1 + \frac{1}{n+1}\right) &= 1 - \frac{1}{n+1} \rightarrow 1 \quad \text{as } n \rightarrow \infty, \end{aligned} \tag{1.21}$$

but

$$\begin{aligned} d(gF(x_n, y_n), F(gx_n, gy_n)) &= d\left(F\left(1 - \frac{1}{n+1}, 1 - \frac{1}{n+1}\right), g1\right) \\ &= d(0, 1) = 1 \not\rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned} \tag{1.22}$$

Thus, F and g are not compatible.

Implicit relation on metric spaces has been used in many articles (see, e.g., [29–31] and references therein). In this paper, we use the following implicit relation to prove a coupled coincidence point theorem for mappings $F : X \times X \rightarrow X$ and $g : X \rightarrow X$, where F has the mixed g -monotone property and F, g are O -compatible.

Let Φ denote all functions $\varphi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ which satisfy

- (i) φ is continuous,
- (ii) $\varphi(t) < t$ for each $t > 0$.

Obviously, if $\varphi \in \Phi$ then $\varphi(0) = 0$.

Let \mathbb{H} denote all continuous functions $H : (\mathbb{R}^+)^5 \rightarrow \mathbb{R}$ which satisfy

- (H1) $H(t_1, t_2, t_3, t_4, t_5)$ is nonincreasing in t_2 and t_5 ,
- (H2) there exists a function $\varphi \in \Phi$ such that

$$H(u, u + v, v, w, u + v) \leq 0 \quad \text{implies } u \leq \varphi(\max\{v, w\}). \tag{1.23}$$

It is easy to check that the following functions are in \mathbb{H} :

- (i) $H_1(t_1, t_2, t_3, t_4, t_5) = t_1 - \alpha t_2 - \beta t_3 - \gamma t_4 - \theta t_5$, where $\alpha, \beta, \gamma, \theta$ are nonnegative real numbers satisfying $2\alpha + \beta + \gamma + 2\theta < 1$;

- (ii) $H_2(t_1, t_2, t_3, t_4, t_5) = t_1 - \alpha \max\{t_2/2, t_3, t_4, t_5/2\}$, where $\alpha \in (0, 1)$;
 (iii) $H_3(t_1, t_2, t_3, t_4, t_5) = t_1 - \varphi(\max\{t_3, t_4\})$, where $\varphi \in \Phi$.

In this paper, we prove a coupled coincidence point theorem for mappings satisfying such implicit relations.

2. Coupled Coincidence Point Theorem

Now we are going to prove our main result.

Theorem 2.1. *Let (X, d, \preceq) be a partially ordered complete metric space. Suppose $F : X \times X \rightarrow X$ and $g : X \rightarrow X$ are mappings such that F has the mixed g -monotone property. Assume that there exists $H \in \mathbb{H}$ such that*

$$H \left(\begin{array}{l} d(F(x, y), F(u, v)), d(F(x, y), gx) + d(F(u, v), gu), \\ d(gx, gu), d(gy, gv), d(F(x, y), gu) + d(F(u, v), gx) \end{array} \right) \leq 0, \quad (2.1)$$

for all $x, y, u, v \in X$ with $gx \geq gu$ and $gy \leq gv$. Suppose $F(X \times X) \subseteq g(X)$, g is continuous and g is O -compatible with F . Suppose either

- (a) F is continuous or;
 (b) X has the following property:
- (i) if a nondecreasing sequence $\{x_n\} \rightarrow x$, then $gx_n \leq gx$ for all n ,
 (ii) if a nonincreasing sequence $\{y_n\} \rightarrow y$, then $gy \leq gy_n$ for all n .

If there exist two elements $x_0, y_0 \in X$ with

$$gx_0 \leq F(x_0, y_0), \quad gy_0 \geq F(y_0, x_0), \quad (2.2)$$

then F and g have a coupled coincidence point in X .

Proof. Let $x_0, y_0 \in X$ be such that $gx_0 \leq F(x_0, y_0)$ and $gy_0 \geq F(y_0, x_0)$. Since $F(X \times X) \subseteq g(X)$, we construct the sequences $\{x_n\}$ and $\{y_n\}$ in X as follows:

$$gx_{n+1} = F(x_n, y_n), \quad gy_{n+1} = F(y_n, x_n) \quad \forall n \geq 0. \quad (2.3)$$

By using the mathematical induction and the mixed g -monotone property of F , we can show that

$$gx_n \leq gx_{n+1}, \quad gy_n \geq gy_{n+1}, \quad \forall n \geq 0. \quad (2.4)$$

If there is some $n_0 \in \mathbb{N}^*$ such that $gx_{n_0} = gx_{n_0+1}$ and $gy_{n_0} = gy_{n_0+1}$ then

$$gx_{n_0} = gx_{n_0+1} = F(x_{n_0}, y_{n_0}), \quad gy_{n_0} = gy_{n_0+1} = F(y_{n_0}, x_{n_0}), \quad (2.5)$$

that means (x_{n_0}, y_{n_0}) is a coupled coincidence point of F and g . Thus we may assume that $\max\{d(gx_{n+1}, gx_n), d(gy_{n+1}, gy_n)\} > 0$ for all n .

Since $gx_{n+1} \geq gx_n$ and $gy_{n+1} \leq gy_n$, from (2.1), we have

$$H\left(d(F(x_{n+1}, y_{n+1}), F(x_n, y_n)), d(F(x_{n+1}, y_{n+1}), gx_{n+1}) + d(F(x_n, y_n), gx_n), d(gx_{n+1}, gx_n), d(gy_{n+1}, gy_n), d(F(x_{n+1}, y_{n+1}), gx_n) + d(F(x_n, y_n), gx_{n+1}))\right) \leq 0 \quad (2.6)$$

or

$$H\left(d(gx_{n+2}, gx_{n+1}), d(gx_{n+2}, gx_{n+1}) + d(gx_{n+1}, gx_n), d(gx_{n+1}, gx_n), d(gy_{n+1}, gy_n), d(gx_{n+2}, gx_n)\right) \leq 0. \quad (2.7)$$

By the properties of H , we have

$$H\left(d(gx_{n+2}, gx_{n+1}), d(gx_{n+2}, gx_{n+1}) + d(gx_{n+1}, gx_n), d(gx_{n+1}, gx_n), d(gy_{n+1}, gy_n), d(gx_{n+2}, gx_n) + d(gx_{n+1}, gx_n)\right) \leq 0, \quad (2.8)$$

which implies that

$$d(gx_{n+2}, gx_{n+1}) \leq \varphi(\max\{d(gx_{n+1}, gx_n), d(gy_{n+1}, gy_n)\}). \quad (2.9)$$

Similarly, one can show that

$$d(gy_{n+2}, gy_{n+1}) \leq \varphi(\max\{d(gx_{n+1}, gx_n), d(gy_{n+1}, gy_n)\}). \quad (2.10)$$

From (2.9) and (2.10), we have

$$\max\{d(gx_{n+2}, gx_{n+1}), d(gy_{n+2}, gy_{n+1})\} \leq \varphi(\max\{d(gx_{n+1}, gx_n), d(gy_{n+1}, gy_n)\}), \quad (2.11)$$

which implies

$$\max\{d(gx_{n+2}, gx_{n+1}), d(gy_{n+2}, gy_{n+1})\} < \max\{d(gx_{n+1}, gx_n), d(gy_{n+1}, gy_n)\}. \quad (2.12)$$

This means that $\{d_n := \max\{d(gx_{n+1}, gx_n), d(gy_{n+1}, gy_n)\}\}$ is a decreasing sequence of positive real numbers. So there is a $d \geq 0$ such that

$$\lim_{n \rightarrow \infty} d_n = \lim_{n \rightarrow \infty} \max\{d(gx_{n+1}, gx_n), d(gy_{n+1}, gy_n)\} = d. \quad (2.13)$$

We will show that $d = 0$. Assume, to the contrary, that $d > 0$. Taking $n \rightarrow \infty$ in (2.11), we have

$$d \leq \lim_{n \rightarrow \infty} \varphi(d_n) = \varphi(d) < d, \quad (2.14)$$

which is a contradiction. Thus $d = 0$.

In what follows, we will show that $\{gx_n\}$ and $\{gy_n\}$ are Cauchy sequences. Suppose, to the contrary that at least one of $\{gx_n\}$ or $\{gy_n\}$ is not a Cauchy sequence. This means that there exists an $\varepsilon > 0$ for which we can find subsequences $\{gx_{n(k)}\}, \{gx_{m(k)}\}$ of $\{gx_n\}$ and $\{gy_{n(k)}\}, \{gy_{m(k)}\}$ of $\{gy_n\}$ with $n(k) > m(k) \geq k$ such that

$$\max\{d(gx_{n(k)}, gx_{m(k)}), d(gy_{n(k)}, gy_{m(k)})\} \geq \varepsilon. \quad (2.15)$$

Further, corresponding to $m(k)$, we can choose $n(k)$ in such a way that it is the smallest integer with $n(k) > m(k) \geq k$ and satisfies (2.15). Then

$$\max\{d(gx_{n(k)-1}, gx_{m(k)}), d(gy_{n(k)-1}, gy_{m(k)})\} < \varepsilon. \quad (2.16)$$

Using the triangle inequality and (2.16), we have

$$\begin{aligned} d(gx_{n(k)}, gx_{m(k)}) &\leq d(gx_{n(k)}, gx_{n(k)-1}) + d(gx_{n(k)-1}, gx_{m(k)}) \\ &< d(gx_{n(k)}, gx_{n(k)-1}) + \varepsilon, \\ d(gy_{n(k)}, gy_{m(k)}) &\leq d(gy_{n(k)}, gy_{n(k)-1}) + d(gy_{n(k)-1}, gy_{m(k)}) \\ &< d(gy_{n(k)}, gy_{n(k)-1}) + \varepsilon. \end{aligned} \quad (2.17)$$

From (2.15) and (2.17), we have

$$\begin{aligned} \varepsilon &\leq \max\{d(gx_{n(k)}, gx_{m(k)}), d(gy_{n(k)}, gy_{m(k)})\} \\ &< \max\{d(gx_{n(k)}, gx_{n(k)-1}), d(gy_{n(k)}, gy_{n(k)-1})\} + \varepsilon. \end{aligned} \quad (2.18)$$

Letting $k \rightarrow \infty$ in the inequalities above and using (2.13) we get

$$\lim_{k \rightarrow \infty} \max\{d(gx_{n(k)}, gx_{m(k)}), d(gy_{n(k)}, gy_{m(k)})\} = \varepsilon. \quad (2.19)$$

By the triangle inequality

$$\begin{aligned} d(gx_{n(k)}, gx_{m(k)}) &\leq d(gx_{n(k)}, gx_{n(k)-1}) + d(gx_{n(k)-1}, gx_{m(k)-1}) + d(gx_{m(k)-1}, gx_{m(k)}), \\ d(gy_{n(k)}, gy_{m(k)}) &\leq d(gy_{n(k)}, gy_{n(k)-1}) + d(gy_{n(k)-1}, gy_{m(k)-1}) + d(gy_{m(k)-1}, gy_{m(k)}). \end{aligned} \quad (2.20)$$

From the last two inequalities and (2.15), we have

$$\begin{aligned} \varepsilon &\leq \max\{d(gx_{n(k)}, gx_{m(k)}), d(gy_{n(k)}, gy_{m(k)})\} \\ &\leq \max\{d(gx_{n(k)}, gx_{n(k)-1}), d(gy_{n(k)}, gy_{n(k)-1})\} \\ &\quad + \max\{d(gx_{m(k)-1}, gx_{m(k)}), d(gy_{m(k)-1}, gy_{m(k)})\} \\ &\quad + \max\{d(gx_{n(k)-1}, gx_{m(k)-1}), d(gy_{n(k)-1}, gy_{m(k)-1})\}. \end{aligned} \tag{2.21}$$

Again, by the triangle inequality,

$$\begin{aligned} d(gx_{n(k)-1}, gx_{m(k)-1}) &\leq d(gx_{n(k)-1}, gx_{m(k)}) + d(gx_{m(k)}, gx_{m(k)-1}) \\ &< d(gx_{m(k)}, gx_{m(k)-1}) + \varepsilon, \\ d(gy_{n(k)-1}, gy_{m(k)-1}) &\leq d(gy_{n(k)-1}, gy_{m(k)}) + d(gy_{m(k)}, gy_{m(k)-1}) \\ &< d(gy_{m(k)}, gy_{m(k)-1}) + \varepsilon. \end{aligned} \tag{2.22}$$

Therefore,

$$\begin{aligned} &\max\{d(gx_{n(k)-1}, gx_{m(k)-1}), d(gy_{n(k)-1}, gy_{m(k)-1})\} \\ &< \max\{d(gx_{m(k)}, gx_{m(k)-1}), d(gy_{m(k)}, gy_{m(k)-1})\} + \varepsilon. \end{aligned} \tag{2.23}$$

From (2.21) and (2.23), we have

$$\begin{aligned} \varepsilon - \max\{d(gx_{n(k)}, gx_{n(k)-1}), d(gy_{n(k)}, gy_{n(k)-1})\} \\ - \max\{d(gx_{m(k)-1}, gx_{m(k)}), d(gy_{m(k)-1}, gy_{m(k)})\} \\ \leq \max\{d(gx_{n(k)-1}, gx_{m(k)-1}), d(gy_{n(k)-1}, gy_{m(k)-1})\} \\ < \max\{d(gx_{m(k)}, gx_{m(k)-1}), d(gy_{m(k)}, gy_{m(k)-1})\} + \varepsilon. \end{aligned} \tag{2.24}$$

Taking $k \rightarrow \infty$ in the inequalities above and using (2.13), we get

$$\lim_{k \rightarrow \infty} \max\{d(gx_{n(k)-1}, gx_{m(k)-1}), d(gy_{n(k)-1}, gy_{m(k)-1})\} = \varepsilon. \tag{2.25}$$

From (2.19) and (2.25), the sequences $\{d(gx_{n(k)}, gx_{m(k)})\}$, $\{d(gy_{n(k)}, gy_{m(k)})\}$, $\{d(gx_{n(k)-1}, gx_{m(k)-1})\}$, and $\{d(gy_{n(k)-1}, gy_{m(k)-1})\}$ have subsequences converging to $\varepsilon_1, \varepsilon_2, \varepsilon_3$ and ε_4 , respectively, and $\max\{\varepsilon_1, \varepsilon_2\} = \max\{\varepsilon_3, \varepsilon_4\} = \varepsilon > 0$. We may assume that

$$\begin{aligned} \lim_{k \rightarrow \infty} d(gx_{n(k)}, gx_{m(k)}) &= \varepsilon_1, & \lim_{k \rightarrow \infty} d(gy_{n(k)}, gy_{m(k)}) &= \varepsilon_2, \\ \lim_{k \rightarrow \infty} d(gx_{n(k)-1}, gx_{m(k)-1}) &= \varepsilon_3, & \lim_{k \rightarrow \infty} d(gy_{n(k)-1}, gy_{m(k)-1}) &= \varepsilon_4. \end{aligned} \tag{2.26}$$

We first assume that $\varepsilon_1 = \max\{\varepsilon_1, \varepsilon_2\} = \varepsilon$. Since $n(k) > m(k)$, $gx_{n(k)-1} \geq gx_{m(k)-1}$ and $gy_{n(k)-1} \leq gy_{m(k)-1}$. From (2.1), we have

$$H \left(\begin{array}{c} d(F(x_{n(k)-1}, y_{n(k)-1}), F(x_{m(k)-1}, y_{m(k)-1})), d(F(x_{n(k)-1}, y_{n(k)-1}), gx_{n(k)-1}) \\ + d(F(x_{m(k)-1}, y_{m(k)-1}), gx_{m(k)-1}), d(gx_{n(k)-1}, gx_{m(k)-1}), d(gy_{n(k)-1}, gy_{m(k)-1}), \\ d(F(x_{n(k)-1}, y_{n(k)-1}), gx_{m(k)-1}) + d(F(x_{m(k)-1}, y_{m(k)-1}), gx_{n(k)-1}) \end{array} \right) \leq 0 \quad (2.27)$$

or

$$H \left(\begin{array}{c} d(gx_{n(k)}, gx_{m(k)}), d(gx_{n(k)}, gx_{n(k)-1}) + d(gx_{m(k)}, gx_{m(k)-1}), \\ d(gx_{n(k)-1}, gx_{m(k)-1}), d(gy_{n(k)-1}, gy_{m(k)-1}), d(gx_{n(k)}, gx_{m(k)-1}) + d(gx_{m(k)}, gx_{n(k)-1}) \end{array} \right) \leq 0 \quad (2.28)$$

or

$$H \left(\begin{array}{c} d(gx_{n(k)}, gx_{m(k)}), d(gx_{n(k)}, gx_{n(k)-1}) + d(gx_{m(k)}, gx_{m(k)-1}), \\ d(gx_{n(k)-1}, gx_{m(k)-1}), d(gy_{n(k)-1}, gy_{m(k)-1}), d(gx_{n(k)}, gx_{m(k)}) \\ + d(gx_{m(k)}, gx_{m(k)-1}) + d(gx_{m(k)}, gx_{m(k)-1}) + d(gx_{m(k)-1}, gx_{n(k)-1}) \end{array} \right) \leq 0. \quad (2.29)$$

Letting $k \rightarrow \infty$, we have

$$H(\varepsilon_1, 0, \varepsilon_3, \varepsilon_4, \varepsilon_1 + \varepsilon_3) \leq 0. \quad (2.30)$$

Thus,

$$H(\varepsilon_1, \varepsilon_1 + \varepsilon_3, \varepsilon_3, \varepsilon_4, \varepsilon_1 + \varepsilon_3) \leq 0, \quad (2.31)$$

which implies $\varepsilon = \varepsilon_1 \leq \varphi(\max\{\varepsilon_3, \varepsilon_4\}) = \varphi(\varepsilon) < \varepsilon$. That is a contradiction.

Using the same argument as above for the case $\varepsilon_2 = \max\{\varepsilon_1, \varepsilon_2\} = \varepsilon$, we also get a contradiction. Thus $\{gx_n\}$ and $\{gy_n\}$ are Cauchy sequences. Since X is complete, there exist $x, y \in X$ such that

$$\lim_{n \rightarrow \infty} gx_n = x, \quad \lim_{n \rightarrow \infty} gy_n = y. \quad (2.32)$$

Thus

$$\lim_{n \rightarrow \infty} F(x_n, y_n) = \lim_{n \rightarrow \infty} gx_n = x, \quad \lim_{n \rightarrow \infty} F(y_n, x_n) = \lim_{n \rightarrow \infty} gy_n = y. \quad (2.33)$$

Since F and g are O -compatible, from (2.33), we have

$$\lim_{n \rightarrow \infty} d(gF(x_n, y_n), F(gx_n, gy_n)) = 0, \quad (2.34)$$

$$\lim_{n \rightarrow \infty} d(gF(y_n, x_n), F(gy_n, gx_n)) = 0. \quad (2.35)$$

Now, suppose that assumption (a) holds. We have

$$d(gx, F(g(x_n), g(y_n))) \leq d(gx, gF(x_n, y_n)) + d(gF(x_n, y_n), F(gx_n, gy_n)). \quad (2.36)$$

Taking the limit as $n \rightarrow \infty$ in (2.36) and by (2.32), (2.34) and the continuity of F and g we get $d(gx, F(x, y)) = 0$.

Similarly, we can show that $d(gy, F(y, x)) = 0$. Therefore, $gx = F(x, y)$ and $gy = F(y, x)$.

Finally, suppose that assumption (b) holds. Since $\{gx_n\}$ is nondecreasing sequence and $gx_n \rightarrow x$ and $\{gy_n\}$ is nonincreasing sequence and $gy_n \rightarrow y$, by the assumption, we have $ggx_n \leq gx$ and $ggy_n \geq gy$ for all n .

Since g is continuous, from (2.32), (2.34), and (2.35) we have

$$\begin{aligned} \lim_{n \rightarrow \infty} ggx_n &= gx = \lim_{n \rightarrow \infty} gF(x_n, y_n) = \lim_{n \rightarrow \infty} F(gx_n, gy_n), \\ \lim_{n \rightarrow \infty} ggy_n &= gy = \lim_{n \rightarrow \infty} gF(y_n, x_n) = \lim_{n \rightarrow \infty} F(gy_n, gx_n). \end{aligned} \quad (2.37)$$

We have

$$H\left(d(F(gx_n, gy_n), F(x, y)), d(F(gx_n, gy_n), ggx_n) + d(F(x, y), gx), d(ggx_n, gx), d(ggy_n, gy), d(F(gx_n, gy_n), gx) + d(F(x, y), ggy_n)\right) \leq 0. \quad (2.38)$$

Letting $n \rightarrow \infty$ and using (2.37), we have

$$H(d(gx, F(x, y)), d(gx, F(x, y)), 0, 0, d(gx, F(x, y))) \leq 0, \quad (2.39)$$

which implies that $d(gx, F(x, y)) \leq \varphi(\max\{0, 0\}) = 0$. Hence $gx = F(x, y)$. Similarly, one can show that $gy = F(y, x)$.

Thus proved that F and g have a coupled coincidence point in X . \square

Example 2.2 (see, e.g., [11]). Let $(X, d, \leq), F$ and g be defined as in Example 1.9. Then

(i) X is complete and X has the property

- (a) if a nondecreasing sequence $\{x_n\} \rightarrow x$, then $gx_n \leq gx$ for all n ,
- (b) if a nonincreasing sequence $\{y_n\} \rightarrow y$, then $gy \leq gy_n$ for all n ;

(ii) $F(X \times X) = \{0, 1\} \subset \{0\} \cup [1/2, 1] = g(X)$;

(iii) g is continuous and g and F are O -compatible;

(iv) there exist $x_0 = 0, y_0 = 1$ such that $gx_0 \leq F(x_0, y_0)$ and $gy_0 \geq F(y_0, x_0)$;

(v) F has the mixed g -monotone property. Indeed, for every $y \in X$, let $x_1, x_2 \in X$ such that $gx_1 \leq gx_2$

- (a) if $gx_1 = gx_2$ then $x_1, x_2 = 0$ or $x_1, x_2 \in [1/2, 1]$ or $x_1, x_2 \in (1, 3/2]$ or $x_1, x_2 \in (3/2, 2]$. Thus,

$$F(x_1, y) = 0 = F(x_2, y) \quad \text{if } y \in \{0\} \cup \left[\frac{1}{2}, 1\right], \quad x_1, x_2 = 0 \text{ or } x_1, x_2 \in \left[\frac{1}{2}, 1\right], \quad (2.40)$$

otherwise $F(x_1, y) = 1 = F(x_2, y)$,

- (b) if $gx_1 < gx_2$, then $gx_1 = 0$ and $gx_2 = 1$, that is, $x_1 = 0$ and $x_2 \in [1/2, 1]$. Thus

$$F(x_1, y) = 0 = F(x_2, y) \quad \text{if } y \in \{0\} \cup \left[\frac{1}{2}, 1\right], \quad F(x_1, y) = 1 = F(x_2, y) \quad \text{if } y \in (1, 2], \quad (2.41)$$

therefore, F is the g -nondecreasing in its first argument. Similarly, F is the g -nonincreasing in its second argument;

- (vi) for $x, y, u, v \in X$, if $gx \geq gu$ and $gy \leq gv$ then $d(F(x, y), F(u, v)) = 0$. Indeed,

- (a) if $gx > gu$ and $gy < gv$ then $y = u = 0$ and $x, v \in [1/2, 1]$. Thus $d(F(x, y), F(u, v)) = d(0, 0) = 0$,
 (b) if $gx = gu$ and $gy < gv$ then $y = 0$ and $v \in [1/2, 1]$. Thus if $x = u = 0$ or $x, u \in [1/2, 1]$ then $d(F(x, y), F(u, v)) = d(0, 0) = 0$, otherwise $d(F(x, y), F(u, v)) = d(1, 1) = 0$. Similarly, if $gx > gu$ and $gy = gv$ then $d(F(x, y), F(u, v)) = 0$,
 (c) if $gx = gu$ and $gy = gv$ then both x, u are in one of the sets $\{0\}$, $[1/2, 1]$, $(1, 3/2]$ or $(3/2, 2]$ and both y, v are also in one of the sets $\{0\}$, $[1/2, 1]$, $(1, 3/2]$ or $(3/2, 2]$. Thus $d(F(x, y), F(u, v)) = d(0, 0) = 0$ if $x = u = 0$ or $x, u \in [1/2, 1]$ and $y = v = 0$ or $y, v \in [1/2, 1]$, otherwise, $d(F(x, y), F(u, v)) = d(1, 1) = 0$.

Therefore, all the conditions of Theorem 2.1 are satisfied with $H(t_1, t_2, t_3, t_4, t_5) = t_1 - \max\{t_3, t_4\}/2$. Applying Theorem 2.1, we conclude that F and g have a coupled coincidence point.

Note that, we cannot apply the result of Choudhury and Kundu [15], the result of Choudhury et al. [32] as well as the result of Lakshmikantham and Ćirić [7] to this example.

References

- [1] D. Guo and V. Lakshmikantham, "Coupled fixed points of nonlinear operators with applications," *Nonlinear Analysis*, vol. 11, no. 5, pp. 623–632, 1987.
- [2] T. G. Bhaskar and V. Lakshmikantham, "Fixed point theorems in partially ordered metric spaces and applications," *Nonlinear Analysis, Theory, Methods and Applications*, vol. 65, no. 7, pp. 1379–1393, 2006.
- [3] J. Harjani, B. Lopez, and K. Sadarangani, "Fixed point theorems for mixed monotone operators and applications to integral equations," *Nonlinear Analysis, Theory, Methods and Applications*, vol. 74, no. 5, pp. 1749–1760, 2011.
- [4] E. Karapinar, N. V. Luong, and N. X. Thuan, "Coupled coincidence points for mixed monotone operators in partially ordered metric spaces," *Arabian Journal of Mathematics*. In press.
- [5] E. Karapinar, "Couple fixed point theorems for nonlinear contractions in cone metric spaces," *Computers and Mathematics with Applications*, vol. 59, no. 12, pp. 3656–3668, 2010.
- [6] E. Karapinar, "Couple fixed point on cone metric spaces," *Gazi University Journal of Science*, vol. 24, no. 1, pp. 51–58, 2011.

- [7] V. Lakshmikantham and L. Ćirić, "Coupled fixed point theorems for nonlinear contractions in partially ordered metric spaces," *Nonlinear Analysis, Theory, Methods and Applications*, vol. 70, no. 12, pp. 4341–4349, 2009.
- [8] N. V. Luong and N. X. Thuan, "Coupled fixed points in partially ordered metric spaces and application," *Nonlinear Analysis, Theory, Methods and Applications*, vol. 74, no. 3, pp. 983–992, 2011.
- [9] N. V. Luong and N. X. Thuan, "Coupled fixed point theorems in partially ordered metric spaces," *Bulletin of Mathematical Analysis and Applications*, vol. 2, no. 4, pp. 16–24, 2010.
- [10] N. V. Luong and N. X. Thuan, "Coupled fixed point theorems for mixed monotone mappings and an application to integral equations," *Computers and Mathematics with Applications*, vol. 62, no. 11, pp. 4238–4248, 2011.
- [11] N. V. Luong and N. X. Thuan, "Coupled points in ordered generalized metric spaces and application to integro-differential equations," Submitted to *Analele Stiintifice ale Universitatii "Ovidius" Constanta, Seria Matematica*.
- [12] H. Aydi, E. Karapinar, and W. Shatanawi, "Coupled fixed point results for (ψ, ϕ) -weakly contractive condition in ordered partial metric spaces," *Computers and Mathematics with Applications*, vol. 62, no. 12, pp. 4449–4460, 2011.
- [13] V. Berinde and M. Borcut, "Tripled fixed point theorems for contractive type mappings in partially ordered metric spaces," *Nonlinear Analysis, Theory, Methods and Applications*, vol. 74, no. 15, pp. 7347–7355, 2011.
- [14] V. Berinde, "Coupled fixed point theorems for ϕ -contractive mixed monotone mappings in partially ordered metric spaces," *Nonlinear Analysis: Theory, Methods and Applications*, vol. 75, pp. 3218–3228, 2012.
- [15] B. S. Choudhury and A. Kundu, "A coupled coincidence point result in partially ordered metric spaces for compatible mappings," *Nonlinear Analysis, Theory, Methods and Applications*, vol. 73, no. 8, pp. 2524–2531, 2010.
- [16] B. Samet, "Coupled fixed point theorems for a generalized Meir-Keeler contraction in partially ordered metric spaces," *Nonlinear Analysis, Theory, Methods and Applications*, vol. 72, no. 12, pp. 4508–4517, 2010.
- [17] L. Gholizadeh, R. Saadati, W. Shatanawi, and S. M. Vaezpour, "Contractive mapping in generalized, ordered metric spaces with application in integral equations," *Mathematical Problems in Engineering*, vol. 2011, Article ID 380784, 14 pages, 2011.
- [18] W. Shatanawi, Z. Mustafa, and N. Tahat, "Some coincidence point theorems for nonlinear contraction in ordered metric spaces," *Fixed Point Theory and Applications*, vol. 2011, article 68, 2011.
- [19] W. Shatanawi and B. Samet, "On (ψ, ϕ) -weakly contractive condition in partially ordered metricspaces," *Computers and Mathematics with Applications*, vol. 62, pp. 3204–3214, 2011.
- [20] W. Shatanawi, "Some fixed point theorems in ordered G-metric spaces and applications," *Abstract and Applied Analysis*, Article ID 126205, 11 pages, 2011.
- [21] Y. J. Cho, Z. Kadelburg, R. Saadati, and W. Shatanawi, "Coupled fixed point theorems under weak contractions," *Discrete Dynamics in Nature and Society*, vol. 2012, Article ID 184534, 9 pages, 2012.
- [22] H. Aydi, B. Damjanović, B. Samet, and W. Shatanawi, "Coupled fixed point theorems for nonlinear contractions in partially ordered G-metric spaces," *Mathematical and Computer Modelling*, vol. 54, pp. 2443–2450, 2011.
- [23] Y. J. Cho, B. E. Rhoades, R. Saadati, B. Samet, and W. Shatanawi, "Nonlinear coupled fixed point theorems in ordered generalized metric spaces with integral type," *Fixed Point Theory and Applications*, vol. 2012, article 8, 2012.
- [24] W. Shatanawi, "On w -compatible mappings and common coupled coincidence point in cone metric spaces," *Applied Mathematics Letters*, vol. 25, pp. 925–931, 2012.
- [25] W. Shatanawi, B. Samet, and M. Abbas, "Coupled fixed point theorems for mixed monotone mappings in ordered partial metric spaces," *Mathematical and Computer Modelling*, vol. 55, pp. 680–687, 2012.
- [26] H. K. Nashine and W. Shatanawi, "Coupled common fixed point theorems for a pair of commuting mappings in partially ordered complete metric spaces," *Computers and Mathematics with Applications*, vol. 62, pp. 1984–1993, 2011.
- [27] W. Shatanawi and Z. Mustafa, "On coupled random fixed point results in partially ordered metric spaces," *Matematicki Vesnik*, vol. 64, pp. 139–146, 2012.

- [28] V. Berinde, "Approximating fixed points of implicit almost contractions," *Hacettepe Journal of Mathematics and Statistics*, vol. 40, no. 1, pp. 93–102, 2012.
- [29] I. Altun and D. Turkoglu, "Some fixed point theorems for weakly compatible multivalued mappings satisfying an implicit relation," *Filomat*, vol. 22, pp. 13–23, 2008.
- [30] A. Djoudi and A. Aliouche, "A general common fixed point theorem for reciprocally continuous mappings satisfying an implicit relation," *The Australian Journal of Mathematical Analysis and Applications*, vol. 3, no. 1, pp. 1–7, 2006.
- [31] V. Popa, "A general coincidence theorem for compatible multivalued mappings satisfying an implicit relation," *Demonstratio Mathematica*, vol. 33, no. 1, pp. 159–164, 2000.
- [32] B. S. Choudhury, N. Metiya, and A. Kundu, "Coupled coincidence point theorems in ordered metric spaces," *Annali dell'Universita di Ferrara*, vol. 57, no. 1, pp. 1–16, 2011.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

