Hindawi Publishing Corporation
Abstract and Applied Analysis
Volume 2012, Article ID 817436, 9 pages
doi:10.1155/2012/817436

Research Article

Strong Convergence of a Modified Extragradient Method to the Minimum-Norm Solution of Variational Inequalities

Yonghong Yao,[1] Muhammad Aslam Noor,[2,3] and Yeong-Cheng Liou[4]

[1] *Department of Mathematics, Tianjin Polytechnic University, Tianjin 300387, China*
[2] *Mathematics Department, COMSATS Institute of Information Technology, Islamabad 44000, Pakistan*
[3] *Mathematics Department, College of Science, King Saud University, Riyadh 11451, Saudi Arabia*
[4] *Department of Information Management, Cheng Shiu University, Kaohsiung 833, Taiwan*

Correspondence should be addressed to Yonghong Yao, yaoyonghong@yahoo.cn

Received 18 August 2011; Accepted 14 October 2011

Academic Editor: Khalida Inayat Noor

We suggest and analyze a modified extragradient method for solving variational inequalities, which is convergent strongly to the minimum-norm solution of some variational inequality in an infinite-dimensional Hilbert space.

1. Introduction

Let C be a closed convex subset of a real Hilbert space H. A mapping $A : C \to H$ is called α-inverse-strongly monotone if there exists a positive real number α such that

$$\langle Au - Av, u - v \rangle \geq \alpha \|Au - Av\|^2, \quad \forall u, v \in C. \tag{1.1}$$

The variational inequality problem is to find $u \in C$ such that

$$\langle Au, v - u \rangle \geq 0, \quad \forall v \in C. \tag{1.2}$$

The set of solutions of the variational inequality problem is denoted by $\mathrm{VI}(C, A)$. It is well known that variational inequality theory has emerged as an important tool in studying a wide

class of obstacle, unilateral, and equilibrium problems, which arise in several branches of pure and applied sciences in a unified and general framework. Several numerical methods have been developed for solving variational inequalities and related optimization problems; see [1–36] and the references therein.

It is well known that variational inequalities are equivalent to the fixed point problem. This alternative formulation has been used to study the existence of a solution of the variational inequality as well as to develop several numerical methods. Using this equivalence, one can suggest the following iterative method.

Algorithm 1.1. For a given $u_0 \in C$, calculate the approximate solution u_{n+1} by the iterative scheme

$$u_{n+1} = P_C[u_n - \lambda A u_n], \quad n = 0, 1, 2, \ldots. \tag{1.3}$$

It is well known that the convergence of Algorithm 1.1 requires that the operator A must be both strongly monotone and Lipschitz continuous. These restrict conditions rules out its applications in several important problems. To overcome these drawbacks, Korpelevič suggested in [8] an algorithm of the form

$$\begin{aligned} y_n &= P_C[x_n - \lambda A x_n], \\ x_{n+1} &= P_C[x_n - \lambda A y_n], \quad n \geq 0. \end{aligned} \tag{1.4}$$

Noor [2] further suggested and analyzed the following new iterative methods for solving the variational inequality (1.2).

Algorithm 1.2. For a given $u_0 \in C$, calculate the approximate solution u_{n+1} by the iterative scheme

$$\begin{aligned} u_{n+1} &= P_C[w_n - \lambda A w_n], \\ w_n &= P_C[u_n - \lambda A u_n], \quad n = 0, 1, 2, \ldots, \end{aligned} \tag{1.5}$$

which is known as the modified extragradient method. For the convergence analysis of Algorithm 1.2, see Noor [1, 2] and the references therein. We would like to point out that Algorithm 1.2 is quite different from the method of Korpelevič [8]. However, Algorithm 1.2 fails, in general, to converge strongly in the setting of infinite-dimensional Hilbert spaces.

In this paper, we suggest and consider a very simple modified extragradient method which is convergent strongly to the minimum-norm solution of variational inequality (1.2) in an infinite-dimensional Hilbert space. This new method includes the method of Noor [2] as a special case.

2. Preliminaries

Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$, and let C be a closed convex subset of H. It is well known that, for any $u \in H$, there exists a unique $u_0 \in C$ such that

$$\|u - u_0\| = \inf\{\|u - x\| : x \in C\}. \tag{2.1}$$

We denote u_0 by $P_C u$, where P_C is called the *metric projection* of H onto C. The metric projection P_C of H onto C has the following basic properties:

(i) $\|P_C x - P_C y\| \leq \|x - y\|$ for all $x, y \in H$;

(ii) $\langle x - y, P_C x - P_C y \rangle \geq \|P_C x - P_C y\|^2$ for every $x, y \in H$;

(iii) $\langle x - P_C x, y - P_C x \rangle \leq 0$ for all $x \in H$, $y \in C$.

We need the following lemma for proving our main results.

Lemma 2.1 (see [15]). *Assume that $\{a_n\}$ is a sequence of nonnegative real numbers such that*

$$a_{n+1} \leq (1 - \gamma_n) a_n + \delta_n, \tag{2.2}$$

where $\{\gamma_n\}$ is a sequence in $(0, 1)$ and $\{\delta_n\}$ is a sequence such that

(1) $\sum_{n=1}^{\infty} \gamma_n = \infty$;

(2) $\limsup_{n \to \infty} \delta_n / \gamma_n \leq 0$ *or* $\sum_{n=1}^{\infty} |\delta_n| < \infty$.
Then $\lim_{n \to \infty} a_n = 0$.

3. Main Result

In this section we will state and prove our main result.

Theorem 3.1. *Let C be a closed convex subset of a real Hilbert space H. Let $A : C \to H$ be an α-inverse-strongly monotone mapping. Suppose that $\mathrm{VI}(C, A) \neq \emptyset$. For given $x_0 \in C$ arbitrarily, define a sequence $\{x_n\}$ iteratively by*

$$\begin{aligned} y_n &= P_C[(1 - \alpha_n)(x_n - \lambda A x_n)], \\ x_{n+1} &= P_C(y_n - \lambda A y_n), \quad n \geq 0, \end{aligned} \tag{3.1}$$

where $\{\alpha_n\}$ is a sequence in $(0, 1)$ and $\lambda \in [a, b] \subset (0, 2\alpha)$ is a constant. Assume the following conditions are satisfied:

$(C1)$: $\lim_{n \to \infty} \alpha_n = 0$;

$(C2)$: $\sum_{n=1}^{\infty} \alpha_n = \infty$;

$(C3)$: $\lim_{n \to \infty} (\alpha_{n+1} / \alpha_n) = 1$.

Then the sequence $\{x_n\}$ generated by (3.1) converges strongly to $P_{\mathrm{VI}(C,A)}(0)$ which is the minimum-norm element in $\mathrm{VI}(C, A)$.

We will divide our detailed proofs into several conclusions.

Proof. Take $x^* \in \mathrm{VI}(C, A)$. First we need to use the following facts:

(1) $x^* = P_C(x^* - \lambda A x^*)$ for all $\lambda > 0$; in particular,

$$x^* = P_C[x^* - \lambda(1-\alpha_n)Ax^*] = P_C[\alpha_n x^* + (1-\alpha_n)(x^* - \lambda A x^*)], \quad \forall n \geq 0; \tag{3.2}$$

(2) $I - \lambda A$ is nonexpansive and for all $x, y \in C$

$$\|(I-\lambda A)x - (I-\lambda A)y\|^2 \leq \|x-y\|^2 + \lambda(\lambda - 2\alpha)\|Ax - Ay\|^2. \tag{3.3}$$

From (3.1), we have

$$\begin{aligned}
\|y_n - x^*\| &= \|P_C[(1-\alpha_n)(x_n - \lambda A x_n)] - P_C[\alpha_n x^* + (1-\alpha_n)(x^* - \lambda A x^*)]\| \\
&\leq \|\alpha_n(-x^*) + (1-\alpha_n)[(x_n - \lambda A x_n) - (x^* - \lambda A x^*)]\| \\
&\leq \alpha_n\|x^*\| + (1-\alpha_n)\|(I-\lambda A)x_n - (I-\lambda A)x^*\| \\
&\leq \alpha_n\|x^*\| + (1-\alpha_n)\|x_n - x^*\|.
\end{aligned} \tag{3.4}$$

Thus,

$$\begin{aligned}
\|x_{n+1} - x^*\| &= \|P_C(y_n - \lambda A y_n) - P_C(x^* - \lambda A x^*)\| \\
&\leq \|(y_n - \lambda A y_n) - (x^* - \lambda A x^*)\| \\
&\leq \|y_n - x^*\| \\
&\leq \alpha_n\|x^*\| + (1-\alpha_n)\|x_n - x^*\| \\
&\leq \max\{\|x^*\|, \|x_0 - x^*\|\}.
\end{aligned} \tag{3.5}$$

Therefore, $\{x_n\}$ is bounded and so are $\{y_n\}$, $\{Ax_n\}$, and $\{Ay_n\}$.

From (3.1), we have

$$\begin{aligned}
\|x_{n+1} - x_n\| &= \|P_C(y_n - \lambda A y_n) - P_C(y_{n-1} - \lambda A y_{n-1})\| \\
&\leq \|(y_n - \lambda A y_n) - (y_{n-1} - \lambda A y_{n-1})\| \\
&\leq \|y_n - y_{n-1}\| \\
&= \|P_C[(1-\alpha_n)(x_n - \lambda A x_n)] - P_C[(1-\alpha_{n-1})(x_{n-1} - \lambda A x_{n-1})]\| \\
&\leq \|(1-\alpha_n)[(I-\lambda A)x_n - (I-\lambda A)x_{n-1}] - (\alpha_n - \alpha_{n-1})(I-\lambda A)x_{n-1}\| \\
&\leq (1-\alpha_n)\|(I-\lambda A)x_n - (I-\lambda A)x_{n-1}\| + |\alpha_n - \alpha_{n-1}|\|(I-\lambda A)x_{n-1}\| \\
&\leq (1-\alpha_n)\|x_n - x_{n-1}\| + |\alpha_n - \alpha_{n-1}|M,
\end{aligned} \tag{3.6}$$

where $M > 0$ is a constant such that $\sup_n\{\|(I-\lambda A)x_n\|, \|(I-\lambda A)x_n\|(\|(I-\lambda A)x_n\| + 2\|x_n - x^*\|)\} \le M$. Hence, by Lemma 2.1, we obtain

$$\lim_{n\to\infty}\|x_{n+1}-x_n\| = 0. \tag{3.7}$$

From (3.4), (3.5) and the convexity of the norm, we deduce

$$\begin{aligned}
\|x_{n+1}-x^*\|^2 &\le \|\alpha_n(-x^*) + (1-\alpha_n)[(x_n-\lambda Ax_n)-(x^*-\lambda Ax^*)]\|^2 \\
&\le \alpha_n\|x^*\|^2 + (1-\alpha_n)\|(I-\lambda A)x_n-(I-\lambda A)x^*\|^2 \\
&\le \alpha_n\|x^*\|^2 + (1-\alpha_n)\left[\|x_n-x^*\|^2 + \lambda(\lambda-2\alpha)\|Ax_n-Ax^*\|^2\right] \\
&\le \alpha_n\|x^*\|^2 + \|x_n-x^*\|^2 + (1-\alpha_n)a(b-2\alpha)\|Ax_n-Ax^*\|^2.
\end{aligned} \tag{3.8}$$

Therefore, we have

$$\begin{aligned}
(1-\alpha_n)a(2\alpha-b)\|Ax_n-Ax^*\|^2 &\le \alpha_n\|x^*\|^2 + \|x_n-x^*\|^2 - \|x_{n+1}-x^*\|^2 \\
&\le \alpha_n\|x^*\|^2 + (\|x_n-x^*\| + \|x_{n+1}-x^*\|)\times\|x_n-x_{n+1}\|.
\end{aligned} \tag{3.9}$$

Since $\alpha_n \to 0$ and $\|x_n - x_{n+1}\| \to 0$ as $n \to \infty$, we obtain $\|Ax_n - Ax^*\| \to 0$ as $n \to \infty$.

By the property (ii) of the metric projection P_C, we have

$$\begin{aligned}
\|y_n-x^*\|^2 &= \|P_C[(1-\alpha)(x_n-\lambda Ax_n)] - P_C(x^*-\lambda Ax^*)\|^2 \\
&\le \langle (1-\alpha_n)(x_n-\lambda Ax_n)-(x^*-\lambda Ax^*), y_n-x^*\rangle \\
&= \frac{1}{2}\Big\{\|(x_n-\lambda Ax_n)-(x^*-\lambda Ax^*)-\alpha_n(I-\lambda A)x_n\|^2 + \|y_n-x^*\|^2 \\
&\qquad - \|(x_n-\lambda Ax_n)-(x^*-\lambda Ax^*)-(y_n-x^*)-\alpha_n(I-\lambda A)x_n\|^2\Big\} \\
&\le \frac{1}{2}\Big\{\|(x_n-\lambda Ax_n)-(x^*-\lambda Ax^*)\|^2 + \alpha_n M + \|y_n-x^*\|^2 \\
&\qquad -\|(x_n-y_n)-\lambda(Ax_n-Ax^*)-\alpha_n(I-\lambda A)x_n\|^2\Big\} \\
&\le \frac{1}{2}\Big\{\|x_n-x^*\|^2 + \alpha_n M + \|y_n-x^*\|^2 - \|x_n-y_n\|^2 \\
&\qquad + 2\lambda\langle x_n-y_n, Ax_n-Ax^*\rangle + 2\alpha_n\langle (I-\lambda A)x_n, x_n-y_n\rangle \\
&\qquad -\|\lambda(Ax_n-Ax^*)+\alpha_n(I-\lambda A)x_n\|^2\Big\} \\
&\le \frac{1}{2}\Big\{\|x_n-x^*\|^2 + \alpha_n M + \|y_n-x^*\|^2 - \|x_n-y_n\|^2 \\
&\qquad +2\lambda\|x_n-y_n\|\|Ax_n-Ax^*\| + 2\alpha_n\|(I-\lambda A)x_n\|\|x_n-y_n\|\Big\}.
\end{aligned} \tag{3.10}$$

It follows that

$$\begin{aligned}\|y_n - x^*\|^2 &\le \|x_n - x^*\|^2 + \alpha_n M - \|x_n - y_n\|^2 \\ &\quad + 2\lambda\|x_n - y_n\|\|Ax_n - Ax^*\| + 2\alpha_n\|(I - \lambda A)x_n\|\|x_n - y_n\|,\end{aligned} \tag{3.11}$$

and hence

$$\begin{aligned}\|x_{n+1} - x^*\|^2 &\le \|y_n - x^*\|^2 \\ &\le \|x_n - x^*\|^2 + \alpha_n M - \|x_n - y_n\|^2 + 2\lambda\|x_n - y_n\|\|Ax_n - Ax^*\| \\ &\quad + 2\alpha_n\|(I - \lambda A)x_n\|\|x_n - y_n\|\end{aligned} \tag{3.12}$$

which implies that

$$\begin{aligned}\|x_n - y_n\|^2 &\le (\|x_n - x^*\| + \|x_{n+1} - x^*\|)\|x_{n+1} - x_n\| + \alpha_n M + 2\lambda\|x_n - y_n\|\|Ax_n - Ax^*\| \\ &\quad + 2\alpha_n\|(I - \lambda A)x_n\|\|x_n - y_n\|.\end{aligned} \tag{3.13}$$

Since $\alpha_n \to 0$, $\|x_n - x_{n+1}\| \to 0$, and $\|Ax_n - Ax^*\| \to 0$, we derive $\|x_n - y_n\| \to 0$.

Next we show that

$$\limsup_{n\to\infty}\langle z_0, z_0 - y_n\rangle \le 0, \tag{3.14}$$

where $z_0 = P_{\mathrm{VI}(C,A)}(0)$. To show it, we choose a subsequence $\{y_{n_i}\}$ of $\{y_n\}$ such that

$$\limsup_{n\to\infty}\langle z_0, z_0 - y_n\rangle = \lim_{i\to\infty}\langle z_0, z_0 - y_{n_i}\rangle. \tag{3.15}$$

As $\{y_{n_i}\}$ is bounded, we have that a subsequence $\{y_{n_{ij}}\}$ of $\{y_{n_i}\}$ converges weakly to z.

Next we show that $z \in \mathrm{VI}(C, A)$. We define a mapping T by

$$Tv = \begin{cases} Av + N_C v, & v \in C, \\ \emptyset, & v \notin C. \end{cases} \tag{3.16}$$

Then T is maximal monotone (see [16]). Let $(v, w) \in G(T)$. Since $w - Av \in N_C v$ and $y_n \in C$, we have $\langle v - y_n, w - Av\rangle \ge 0$. On the other hand, from $y_n = P_C[(1-\alpha_n)(x_n - \lambda Ax_n)]$, we have

$$\langle v - y_n, y_n - (1 - \alpha_n)(x_n - \lambda Ax_n)\rangle \ge 0, \tag{3.17}$$

that is,

$$\left\langle v - y_n, \frac{y_n - x_n}{\lambda} + Ax_n + \frac{\alpha_n}{\lambda}(I - \lambda A)x_n\right\rangle \ge 0. \tag{3.18}$$

Therefore, we have

$$\begin{aligned}
\langle v - y_{n_i}, w\rangle &\geq \langle v - y_{n_i}, Av\rangle \\
&\geq \langle v - y_{n_i}, Av\rangle - \left\langle v - y_{n_i}, \frac{y_{n_i} - x_{n_i}}{\lambda} + Ax_{n_i} + \frac{\alpha_{n_i}}{\lambda}(I - \lambda A)x_{n_i}\right\rangle \\
&= \left\langle v - y_{n_i}, Av - Ax_{n_i} - \frac{y_{n_i} - x_{n_i}}{\lambda} - \frac{\alpha_{n_i}}{\lambda}(I - \lambda A)x_{n_i}\right\rangle \\
&= \langle v - y_{n_i}, Av - Ay_{n_i}\rangle + \langle v - y_{n_i}, Ay_{n_i} - Ax_{n_i}\rangle \\
&\quad - \left\langle v - y_{n_i}, \frac{y_{n_i} - x_{n_i}}{\lambda} + \frac{\alpha_{n_i}}{\lambda}(I - \lambda A)x_{n_i}\right\rangle \\
&\geq \langle v - y_{n_i}, Ay_{n_i} - Ax_{n_i}\rangle - \left\langle v - y_{n_i}, \frac{y_{n_i} - x_{n_i}}{\lambda} + \frac{\alpha_{n_i}}{\lambda}(I - \lambda A)x_{n_i}\right\rangle.
\end{aligned} \tag{3.19}$$

Noting that $\alpha_{n_i} \to 0$, $\|y_{n_i} - x_{n_i}\| \to 0$, and A is Lipschitz continuous, we obtain $\langle v - z, w\rangle \geq 0$. Since T is maximal monotone, we have $z \in T^{-1}(0)$, and hence $z \in \mathrm{VI}(C, A)$. Therefore,

$$\limsup_{n\to\infty}\langle z_0, z_0 - y_n\rangle = \lim_{i\to\infty}\langle z_0, z_0 - y_{n_i}\rangle = \langle z_0, z_0 - z\rangle \leq 0. \tag{3.20}$$

Finally, we prove $x_n \to z_0$. By the property (ii) of metric projection P_C, we have

$$\begin{aligned}
\|y_n - z_0\|^2 &= \|P_C[(1-\alpha_n)(x_n - \lambda Ax_n)] - P_C[\alpha_n z_0 + (1-\alpha_n)(z_0 - \lambda Az_0)]\|^2 \\
&\leq \langle \alpha_n(-z_0) + (1-\alpha_n)[(x_n - \lambda Ax_n) - (z_0 - \lambda Az_0)], y_n - z_0\rangle \\
&\leq \alpha_n\langle z_0, z_0 - y_n\rangle + (1-\alpha_n)\|(x_n - \lambda Ax_n) - (z_0 - \lambda Az_0)\|\|y_n - z_0\| \\
&\leq \alpha_n\langle z_0, z_0 - y_n\rangle + (1-\alpha_n)\|x_n - z_0\|\|y_n - z_0\| \\
&\leq \alpha_n\langle z_0, z_0 - y_n\rangle + \frac{1-\alpha_n}{2}\left(\|x_n - z_0\|^2 + \|y_n - z_0\|^2\right).
\end{aligned} \tag{3.21}$$

Hence,

$$\|y_n - z_0\|^2 \leq (1-\alpha_n)\|x_n - z_0\|^2 + 2\alpha_n\langle z_0, z_0 - y_n\rangle. \tag{3.22}$$

Therefore,

$$\|x_{n+1} - z_0\|^2 \leq \|y_n - z_0\|^2 \leq (1-\alpha_n)\|x_n - z_0\|^2 + 2\alpha_n\langle z_0, z_0 - y_n\rangle. \tag{3.23}$$

We apply Lemma 2.1 to the last inequality to deduce that $x_0 \to z_0$. This completes the proof. □

Remark 3.2. Our Algorithm (3.1) is similar to Noor's modified extragradient method; see [2]. However, our algorithm has strong convergence in the setting of infinite-dimensional Hilbert spaces.

Acknowledgments

Y. Yao was supported in part by Colleges and Universities Science and Technology Development Foundation (20091003) of Tianjin, NSFC 11071279 and NSFC 71161001-G0105. Y.-C. Liou was partially supported by the Program TH-1-3, Optimization Lean Cycle, of Sub-Projects TH-1 of Spindle Plan Four in Excellence Teaching and Learning Plan of Cheng Shiu University and was supported in part by NSC 100-2221-E-230-012.

References

[1] M. Aslam Noor, "Some developments in general variational inequalities," *Applied Mathematics and Computation*, vol. 152, no. 1, pp. 199–277, 2004.
[2] M. A. Noor, "A class of new iterative methods for general mixed variational inequalities," *Mathematical and Computer Modelling*, vol. 31, no. 13, pp. 11–19, 2000.
[3] R. E. Bruck,, "On the weak convergence of an ergodic iteration for the solution of variational inequalities for monotone operators in Hilbert space," *Journal of Mathematical Analysis and Applications*, vol. 61, no. 1, pp. 159–164, 1977.
[4] J.-L. Lions and G. Stampacchia, "Variational inequalities," *Communications on Pure and Applied Mathematics*, vol. 20, pp. 493–519, 1967.
[5] W. Takahashi, "Nonlinear complementarity problem and systems of convex inequalities," *Journal of Optimization Theory and Applications*, vol. 24, no. 3, pp. 499–506, 1978.
[6] J. C. Yao, "Variational inequalities with generalized monotone operators," *Mathematics of Operations Research*, vol. 19, no. 3, pp. 691–705, 1994.
[7] H. K. Xu and T. H. Kim, "Convergence of hybrid steepest-descent methods for variational inequalities," *Journal of Optimization Theory and Applications*, vol. 119, no. 1, pp. 185–201, 2003.
[8] G. M. Korpelevič, "An extragradient method for finding saddle points and for other problems," *Èkonomika i Matematicheskie Metody*, vol. 12, no. 4, pp. 747–756, 1976.
[9] W. Takahashi and M. Toyoda, "Weak convergence theorems for nonexpansive mappings and monotone mappings," *Journal of Optimization Theory and Applications*, vol. 118, no. 2, pp. 417–428, 2003.
[10] L.-C. Ceng and J.-C. Yao, "An extragradient-like approximation method for variational inequality problems and fixed point problems," *Applied Mathematics and Computation*, vol. 190, no. 1, pp. 205–215, 2007.
[11] Y. Yao and J.-C. Yao, "On modified iterative method for nonexpansive mappings and monotone mappings," *Applied Mathematics and Computation*, vol. 186, no. 2, pp. 1551–1558, 2007.
[12] A. Bnouhachem, M. Aslam Noor, and Z. Hao, "Some new extragradient iterative methods for variational inequalities," *Nonlinear Analysis*, vol. 70, no. 3, pp. 1321–1329, 2009.
[13] M. A. Noor, "New extragradient-type methods for general variational inequalities," *Journal of Mathematical Analysis and Applications*, vol. 277, no. 2, pp. 379–394, 2003.
[14] B. S. He, Z. H. Yang, and X. M. Yuan, "An approximate proximal-extragradient type method for monotone variational inequalities," *Journal of Mathematical Analysis and Applications*, vol. 300, no. 2, pp. 362–374, 2004.
[15] H.-K. Xu, "Iterative algorithms for nonlinear operators," *Journal of the London Mathematical Society*, vol. 66, no. 1, pp. 240–256, 2002.
[16] R. T. Rockafellar, "Monotone operators and the proximal point algorithm," *SIAM Journal on Control and Optimization*, vol. 14, no. 5, pp. 877–898, 1976.
[17] Y. Censor, A. Motova, and A. Segal, "Perturbed projections and subgradient projections for the multiple-sets split feasibility problem," *Journal of Mathematical Analysis and Applications*, vol. 327, no. 2, pp. 1244–1256, 2007.
[18] Y. Censor, A. Gibali, and S. Reich, "The subgradient extragradient method for solving variational inequalities in Hilbert space," *Journal of Optimization Theory and Applications*, vol. 148, no. 2, pp. 318–335, 2011.
[19] Y. Yao and N. Shahzad, "Strong convergence of a proximal point algorithm with general errors," *Optimization Letters*. In press.
[20] Y. Yao and N. Shahzad, "New methods with perturbations for non-expansive mappings in hilbert spaces," *Fixed Point Theory and Applications*, vol. 2011, article 79, 2011.

[21] Y. Yao, R. Chen, and H.-K. Xu, "Schemes for finding minimum-norm solutions of variational inequalities," *Nonlinear Analysis*, vol. 72, no. 7-8, pp. 3447–3456, 2010.
[22] M. Aslam Noor, "Some developments in general variational inequalities," *Applied Mathematics and Computation*, vol. 152, no. 1, pp. 199–277, 2004.
[23] M. A. Noor, "Projection-proximal methods for general variational inequalities," *Journal of Mathematical Analysis and Applications*, vol. 318, no. 1, pp. 53–62, 2006.
[24] M. A. Noor, "Differentiable non-convex functions and general variational inequalities," *Applied Mathematics and Computation*, vol. 199, no. 2, pp. 623–630, 2008.
[25] M. Aslam Noor and Z. Huang, "Wiener-Hopf equation technique for variational inequalities and nonexpansive mappings," *Applied Mathematics and Computation*, vol. 191, no. 2, pp. 504–510, 2007.
[26] Y. Yao and M. A. Noor, "Convergence of three-step iterations for asymptotically nonexpansive mappings," *Applied Mathematics and Computation*, vol. 187, no. 2, pp. 883–892, 2007.
[27] Y. Yao and M. A. Noor, "On viscosity iterative methods for variational inequalities," *Journal of Mathematical Analysis and Applications*, vol. 325, no. 2, pp. 776–787, 2007.
[28] Y. Yao and M. A. Noor, "On modified hybrid steepest-descent methods for general variational inequalities," *Journal of Mathematical Analysis and Applications*, vol. 334, no. 2, pp. 1276–1289, 2007.
[29] Y. Yao and M. A. Noor, "On convergence criteria of generalized proximal point algorithms," *Journal of Computational and Applied Mathematics*, vol. 217, no. 1, pp. 46–55, 2008.
[30] M. A. Noor and Y. Yao, "Three-step iterations for variational inequalities and nonexpansive mappings," *Applied Mathematics and Computation*, vol. 190, no. 2, pp. 1312–1321, 2007.
[31] Y. Yao and M. A. Noor, "On modified hybrid steepest-descent method for variational inequalities," *Carpathian Journal of Mathematics*, vol. 24, no. 1, pp. 139–148, 2008.
[32] Y. Yao, M. A. Noor, R. Chen, and Y.-C. Liou, "Strong convergence of three-step relaxed hybrid steepest-descent methods for variational inequalities," *Applied Mathematics and Computation*, vol. 201, no. 1-2, pp. 175–183, 2008.
[33] Y. Yao, M. A. Noor, and Y.-C. Liou, "A new hybrid iterative algorithm for variational inequalities," *Applied Mathematics and Computation*, vol. 216, no. 3, pp. 822–829, 2010.
[34] Y. Yao, M. Aslam Noor, K. Inayat Noor, Y.-C. Liou, and H. Yaqoob, "Modified extragradient methods for a system of variational inequalities in Banach spaces," *Acta Applicandae Mathematicae*, vol. 110, no. 3, pp. 1211–1224, 2010.
[35] Y. Yao, M. A. Noor, K. I. Noor, and Y. -C. Liou, "On an iterative algorithm for variational inequalities in banach spaces," *Mathematical Communications*, vol. 16, no. 1, pp. 95–104, 2011.
[36] M. A. Noor, E. Al-Said, K. I. Noor, and Y. Yao, "Extragradient methods for solving nonconvex variational inequalities," *Journal of Computational and Applied Mathematics*, vol. 235, no. 9, pp. 3104–3108, 2011.