

## Research Article

# Oscillation Theorems for Second-Order Quasilinear Neutral Functional Differential Equations

Shurong Sun,<sup>1,2</sup> Tongxing Li,<sup>1,3</sup> Zhenlai Han,<sup>1</sup> and Hua Li<sup>1</sup>

<sup>1</sup> School of Mathematical Sciences, University of Jinan, Shandong Jinan 250022, China

<sup>2</sup> Department of Mathematics and Statistics, Missouri University of Science and Technology,  
 Rolla, MO 65409-0020, USA

<sup>3</sup> School of Control Science and Engineering, Shandong University, Shandong Jinan 250061, China

Correspondence should be addressed to Shurong Sun, sshrong@163.com

Received 15 March 2012; Revised 14 May 2012; Accepted 14 May 2012

Academic Editor: Agacik Zafer

Copyright © 2012 Shurong Sun et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

New oscillation criteria are established for the second-order nonlinear neutral functional differential equations of the form  $(r(t)|z'(t)|^{\alpha-1}z'(t))' + f(t, x[\sigma(t)]) = 0$ ,  $t \geq t_0$ , where  $z(t) = x(t) + p(t)x(\tau(t))$ ,  $p \in C^1([t_0, \infty), [0, \infty))$ , and  $\alpha \geq 1$ . Our results improve and extend some known results in the literature. Some examples are also provided to show the importance of these results.

## 1. Introduction

This paper is concerned with the oscillation problem of the second-order nonlinear functional differential equation of the following form:

$$(r(t)|z'(t)|^{\alpha-1}z'(t))' + f(t, x[\sigma(t)]) = 0, \quad t \geq t_0, \quad (1.1)$$

where  $\alpha \geq 1$  is a constant,  $z(t) = x(t) + p(t)x[\tau(t)]$ .

Throughout this paper, we will assume the following hypotheses:

- (A<sub>1</sub>)  $r \in C^1([t_0, \infty), \mathbb{R})$ ,  $r(t) > 0$  for  $t \geq t_0$ ,
- (A<sub>2</sub>)  $p \in C^1([t_0, \infty), [0, \infty))$ ,
- (A<sub>3</sub>)  $\tau \in C^2([t_0, \infty), \mathbb{R})$ ,  $\tau'(t) > 0$ ,  $\lim_{t \rightarrow \infty} \tau(t) = \infty$ ,
- (A<sub>4</sub>)  $\sigma \in C([t_0, \infty), \mathbb{R})$ ,  $\lim_{t \rightarrow \infty} \sigma(t) = \infty$ ,  $\tau \circ \sigma = \sigma \circ \tau$ ;

(A<sub>5</sub>)  $f(t, u) \in C([t_0, \infty) \times \mathbb{R}, \mathbb{R})$ , and there exists a function  $q \in C([t_0, \infty), [0, \infty))$  such that

$$f(t, u) \operatorname{sgn} u \geq q(t)|u|^\alpha, \quad \text{for } u \neq 0, t \geq t_0. \quad (1.2)$$

By a solution of (1.1), we mean a function  $x \in C([T_x, \infty), \mathbb{R})$  for some  $T_x \geq t_0$  which has the property that  $r(t)|z'(t)|^{\alpha-1}z'(t) \in C^1([T_x, \infty), \mathbb{R})$  and satisfies (1.1) on  $[T_x, \infty)$ . As is customary, a solution of (1.1) is called oscillatory if it has arbitrarily large zeros on  $[t_0, \infty)$ ; otherwise, it is called nonoscillatory. Equation (1.1) is said to be oscillatory if all of its nonconstant solutions are oscillatory.

We note that neutral delay differential equations find numerous applications in electric networks. For instance, they are frequently used for the study of distributed networks containing lossless transmission lines which rise in high-speed computers where the lossless transmission lines are used to interconnect switching circuits; see [1]. Therefore, there is constant interest in obtaining new sufficient conditions for the oscillation or nonoscillation of the solutions of variational types of the second-order equations, see, e.g., papers [2–17].

Known oscillation criteria require various restrictions on the coefficients of the studied neutral differential equations.

Agarwal et al. [2], Chern et al. [3], Džurina and Stavroulakis [4], Kusano et al. [5, 6], Mirzov [7], and Sun and Meng [8] observed some similar properties between

$$\left( r(t)|x'(t)|^{\alpha-1}x'(t) \right)' + q(t)|x[\sigma(t)]|^{\alpha-1}x[\sigma(t)] = 0 \quad (1.3)$$

and the corresponding linear equation

$$(r(t)x'(t))' + q(t)x(t) = 0. \quad (1.4)$$

Liu and Bai [10], Xu and Meng [11, 12], and Dong [13] established some oscillation criteria for (1.3) with neutral term under the assumption that

$$\int_{t_0}^{\infty} \frac{1}{r^{1/\alpha}(t)} dt = \infty. \quad (1.5)$$

Han et al. [14] examined the oscillation of second-order linear neutral differential equation

$$\left( r(t)[x(t) + p(t)x(\tau(t))] \right)' + q(t)x[\sigma(t)] = 0, \quad t \geq t_0, \quad (1.6)$$

where  $\tau'(t) = \tau_0 > 0$ ,  $0 \leq p(t) \leq p_0 < \infty$ , and obtained some oscillation criteria for (1.6) when

$$\int_{t_0}^{\infty} \frac{1}{r(t)} dt = \infty. \quad (1.7)$$

Han et al. [15] studied the oscillation of (1.6) under the case  $0 \leq p(t) \leq 1$  and

$$\int_{t_0}^{\infty} \frac{1}{r(t)} dt < \infty. \quad (1.8)$$

Tripathy [16] considered the nonlinear dynamic equation of the form

$$\left( r(t) \left[ (x(t) + p(t)x(t-\tau))^{\Delta} \right]^{\gamma} \right)^{\Delta} + q(t)|x(t-\delta)|^{\gamma} \operatorname{sgn} x(t-\delta) = 0, \quad (1.9)$$

where  $0 \leq p(t) \leq p_0 < \infty$ ,  $\gamma$  is a the ratios of two positive odd integers, and obtained some oscillation criteria under the following conditions:

$$\int_{t_0}^{\infty} \left( \frac{1}{r(t)} \right)^{\gamma} dt = \infty. \quad (1.10)$$

Džurina [17] was concerned with the oscillation behavior of the solutions of the second-order neutral differential equations as follows

$$\left( a(t) \left[ (x(t) + p(t)x(\tau(t)))^{\gamma} \right]^{\gamma} \right)' + q(t)x^{\beta}(\sigma(t)) = 0, \quad (1.11)$$

where  $0 \leq p(t) \leq p_0 < \infty$ ,  $\gamma$  is a the ratios of two positive odd integers, and obtained some new results under the following conditions

$$\int_{t_0}^{\infty} \left( \frac{1}{a(t)} \right)^{\gamma} dt = \infty. \quad (1.12)$$

Our purpose of this paper is to establish some new oscillation criteria for (1.1), and we will also consider the cases (1.5) and

$$\int_{t_0}^{\infty} \frac{1}{r^{1/\alpha}(t)} dt < \infty. \quad (1.13)$$

To the best of my knowledge, there is no result for the oscillation of (1.1) under the conditions both  $0 \leq p(t) \leq p_0 < \infty$  and (1.13).

In this paper, we will use a new inequality to establish some oscillation criteria for (1.1) for the first time. Some examples will be given to show the importance of these results. In Sections 3 and 4, for the sake of convenience, we denote that

$$\begin{aligned} Q(t) &:= \min\{q(t), q[\tau(t)]\}, & d_+(t) &:= \max\{0, d(t)\}, & \xi(t) &:= \frac{\alpha p'[\sigma(t)]\sigma'(t)}{p[\sigma(t)]} - \frac{\tau''(t)}{\tau'(t)}, \\ \zeta(t) &:= \frac{(\rho'(t))_+}{\rho(t)} + \xi(t), & \varphi(t) &:= \left( \frac{\rho'_+(t)}{\rho(t)} \right)^{\alpha+1} + \frac{p^{\alpha}[\sigma(t)](\zeta_+(t))^{\alpha+1}}{\tau'(t)}, & \delta(t) &:= \int_{\eta(t)}^{\infty} \frac{ds}{r^{1/\alpha}(s)}. \end{aligned} \quad (1.14)$$

## 2. Lemma

In this section, we give the following lemma, which we will use in the proofs of our main results.

**Lemma 2.1.** *Assume that  $\alpha \geq 1$ ,  $a, b \in \mathbb{R}$ . If  $a \geq 0$ ,  $b \geq 0$ , then one has*

$$a^\alpha + b^\alpha \geq \frac{1}{2^{\alpha-1}}(a+b)^\alpha. \quad (2.1)$$

*Proof.* (i) Suppose that  $a = 0$  or  $b = 0$ . Then we have (2.1). (ii) Suppose that  $a > 0$ ,  $b > 0$ . Define the function  $g$  by  $g(x) = x^\alpha$ ,  $x \in (0, \infty)$ . Then  $g''(x) = \alpha(\alpha-1)x^{\alpha-2} \geq 0$  for  $x > 0$ . Thus,  $g$  is a convex function. By the definition of convex function, for  $\lambda = 1/2$ ,  $a, b \in (0, \infty)$ , we have

$$g\left(\frac{a+b}{2}\right) \leq \frac{g(a) + g(b)}{2}, \quad (2.2)$$

that is,

$$a^\alpha + b^\alpha \geq \frac{1}{2^{\alpha-1}}(a+b)^\alpha. \quad (2.3)$$

This completes the proof.  $\square$

## 3. Oscillation Criteria for the Case (1.5)

In this section, we will establish some oscillation criteria for (1.1) under the case (1.5).

**Theorem 3.1.** *Suppose that (1.5) holds,  $\sigma \in C([t_0, \infty), \mathbb{R})$ ,  $\sigma'(t) > 0$ ,  $\sigma(t) \leq t$ , and  $\sigma(t) \leq \tau(t)$  for  $t \geq t_0$ . Furthermore, assume that there exists a function  $\rho \in C([t_0, \infty), (0, \infty))$  such that*

$$\limsup_{t \rightarrow \infty} \int_{t_0}^t \rho(s) \left\{ \frac{Q(s)}{2^{\alpha-1}} - \frac{r[\sigma(s)]\varphi(s)}{(\alpha+1)^{\alpha+1}(\sigma'(s))^\alpha} \right\} ds = \infty. \quad (3.1)$$

*Then (1.1) is oscillatory.*

*Proof.* Let  $x$  be a nonoscillatory solution of (1.1). Without loss of generality, we assume that there exists  $t_1 \geq t_0$  such that  $x(t) > 0$ ,  $x[\tau(t)] > 0$  and  $x[\sigma(t)] > 0$  for all  $t \geq t_1$ . By applying (1.1), for all sufficiently large  $t$ , we obtain that

$$\begin{aligned} & \left( r(t) |z'(t)|^{\alpha-1} z'(t) \right)' + q(t) x^\alpha[\sigma(t)] + q[\tau(t)] p^\alpha[\sigma(t)] x^\alpha(\sigma[\tau(t)]) \\ & + \frac{p^\alpha[\sigma(t)]}{\tau'(t)} \left( r[\tau(t)] |z'[\tau(t)]|^{\alpha-1} z'[\tau(t)] \right)' \leq 0. \end{aligned} \quad (3.2)$$

Using (2.1) and the definition of  $z$ , we conclude that

$$\left( r(t) |z'(t)|^{\alpha-1} z'(t) \right)' + \frac{1}{2^{\alpha-1}} Q(t) z^\alpha[\sigma(t)] + \frac{p^\alpha[\sigma(t)]}{\tau'(t)} \left( r[\tau(t)] |z'[\tau(t)]|^{\alpha-1} z'[\tau(t)] \right)' \leq 0. \quad (3.3)$$

In view of (1.1), we obtain that

$$\left( r(t) |z'(t)|^{\alpha-1} z'(t) \right)' \leq -q(t) x^\alpha[\sigma(t)] \leq 0, \quad t \geq t_1. \quad (3.4)$$

Thus,  $r(t) |z'(t)|^{\alpha-1} z'(t)$  is decreasing function. Now we have two possible cases for  $z'(t)$  :  
 (i)  $z'(t) < 0$  eventually and (ii)  $z'(t) > 0$  eventually.

(i) Suppose that  $z'(t) < 0$  for  $t \geq t_2 \geq t_1$ . Then, from (3.4), we get

$$r(t) |z'(t)|^{\alpha-1} z'(t) \leq r(t_2) |z'(t_2)|^{\alpha-1} z'(t_2), \quad t \geq t_2, \quad (3.5)$$

which implies that

$$z(t) \leq z(t_2) - r^{1/\alpha}(t_2) |z'(t_2)| \int_{t_2}^t r^{-1/\alpha}(s) ds. \quad (3.6)$$

Letting  $t \rightarrow \infty$ , by (1.5), we find  $z(t) \rightarrow -\infty$ , which is a contradiction.

(ii) Suppose that  $z'(t) > 0$  for  $t \geq t_2 \geq t_1$ . We define a Riccati substitution

$$\omega(t) = \rho(t) \frac{r(t)(z'(t))^\alpha}{(z[\sigma(t)])^\alpha}, \quad t \geq t_2. \quad (3.7)$$

Then  $\omega(t) > 0$ . From (3.4), we have

$$z'[\sigma(t)] \geq \left( \frac{r(t)}{r[\sigma(t)]} \right)^{1/\alpha} z'(t). \quad (3.8)$$

Differentiating (3.7), we find that

$$\begin{aligned} \omega'(t) &= \rho'(t) \frac{r(t)(z'(t))^\alpha}{(z[\sigma(t)])^\alpha} + \rho(t) \frac{(r(t)(z'(t))^\alpha)'}{(z[\sigma(t)])^\alpha} \\ &\quad - \alpha \rho(t) \frac{r(t)(z'(t))^\alpha z^{\alpha-1}[\sigma(t)] z'[\sigma(t)] \sigma'(t)}{(z[\sigma(t)])^{2\alpha}}. \end{aligned} \quad (3.9)$$

Therefore, by (3.7), (3.8), and (3.9), we see that

$$\omega'(t) \leq \frac{\rho'(t)}{\rho(t)} \omega(t) + \rho(t) \frac{(r(t)(z'(t))^\alpha)'}{(z[\sigma(t)])^\alpha} - \frac{\alpha \sigma'(t)}{\rho^{1/\alpha}(t) r^{1/\alpha}[\sigma(t)]} \omega^{(\alpha+1)/\alpha}(t). \quad (3.10)$$

Similarly, we introduce a Riccati substitution

$$v(t) = \rho(t) \frac{r[\tau(t)](z'[\tau(t)])^\alpha}{(z[\sigma(t)])^\alpha}, \quad t \geq t_2. \quad (3.11)$$

Then  $v(t) > 0$ . From (3.4), we have

$$z'[\sigma(t)] \geq \left( \frac{r[\tau(t)]}{r[\sigma(t)]} \right)^{1/\alpha} z'[\tau(t)]. \quad (3.12)$$

Differentiating (3.11), we find that

$$\begin{aligned} v'(t) &= \rho'(t) \frac{r[\tau(t)](z'[\tau(t)])^\alpha}{(z[\sigma(t)])^\alpha} + \rho(t) \frac{(r[\tau(t)](z'[\tau(t)])^\alpha)'}{(z[\sigma(t)])^\alpha} \\ &\quad - \alpha \rho(t) \frac{r[\tau(t)](z'[\tau(t)])^\alpha z^{\alpha-1}[\sigma(t)] z'[\sigma(t)] \sigma'(t)}{(z[\sigma(t)])^{2\alpha}}. \end{aligned} \quad (3.13)$$

Therefore, by (3.11), (3.12), and (3.13), we see that

$$v'(t) \leq \frac{\rho'(t)}{\rho(t)} v(t) + \rho(t) \frac{(r[\tau(t)](z'[\tau(t)])^\alpha)'}{(z[\sigma(t)])^\alpha} - \frac{\alpha \sigma'(t)}{\rho^{1/\alpha}(t) r^{1/\alpha}[\sigma(t)]} v^{(\alpha+1)/\alpha}(t). \quad (3.14)$$

Thus, from (3.10) and (3.14), we have

$$\begin{aligned} \omega'(t) + \frac{p^\alpha[\sigma(t)]}{\tau'(t)} v'(t) &\leq \rho(t) \left\{ \frac{(r(t)(z'(t))^\alpha)'}{(z[\sigma(t)])^\alpha} + \frac{p^\alpha[\sigma(t)]}{\tau'(t)} \frac{(r[\tau(t)](z'[\tau(t)])^\alpha)'}{(z[\sigma(t)])^\alpha} \right\} + \frac{\rho'(t)}{\rho(t)} \omega(t) \\ &\quad - \frac{\alpha \sigma'(t)}{\rho^{1/\alpha}(t) r^{1/\alpha}[\sigma(t)]} \omega^{(\alpha+1)/\alpha}(t) + \frac{p^\alpha[\sigma(t)]}{\tau'(t)} \frac{\rho'(t)}{\rho(t)} v(t) \\ &\quad - \frac{p^\alpha[\sigma(t)]}{\tau'(t)} \frac{\alpha \sigma'(t)}{\rho^{1/\alpha}(t) r^{1/\alpha}[\sigma(t)]} v^{(\alpha+1)/\alpha}(t). \end{aligned} \quad (3.15)$$

It is follows from (3.3) that

$$\begin{aligned} \omega'(t) + \frac{p^\alpha[\sigma(t)]}{\tau'(t)} v'(t) &\leq -\frac{1}{2^{\alpha-1}} \rho(t) Q(t) + \frac{\rho'_+(t)}{\rho(t)} \omega(t) \\ &\quad - \frac{\alpha \sigma'(t)}{\rho^{1/\alpha}(t) r^{1/\alpha}[\sigma(t)]} \omega^{(\alpha+1)/\alpha}(t) + \frac{p^\alpha[\sigma(t)]}{\tau'(t)} \frac{\rho'_+(t)}{\rho(t)} v(t) \\ &\quad - \frac{p^\alpha[\sigma(t)]}{\tau'(t)} \frac{\alpha \sigma'(t)}{\rho^{1/\alpha}(t) r^{1/\alpha}[\sigma(t)]} v^{(\alpha+1)/\alpha}(t). \end{aligned} \quad (3.16)$$

Integrating the above inequality from  $t_2$  to  $t$ , we obtain that

$$\begin{aligned}
& \omega(t) - \omega(t_2) + \frac{p^\alpha[\sigma(t)]}{\tau'(t)}v(t) - \frac{p^\alpha[\sigma(t_2)]}{\tau'(t_2)}v(t_2) \\
& \leq - \int_{t_2}^t \frac{1}{2^{\alpha-1}} \rho(s)Q(s)ds \\
& \quad + \int_{t_2}^t \left[ \frac{\rho'_+(s)}{\rho(s)}\omega(s) - \frac{\alpha\sigma'(s)}{\rho^{1/\alpha}(s)r^{1/\alpha}[\sigma(s)]}\omega^{(\alpha+1)/\alpha}(s) \right] ds \\
& \quad + \int_{t_2}^t \frac{p^\alpha[\sigma(s)]}{\tau'(s)} \left\{ \left[ \frac{\rho'_+(s)}{\rho(s)} + \xi(s) \right]_+ v(s) - \frac{\alpha\sigma'(s)}{\rho^{1/\alpha}(s)r^{1/\alpha}[\sigma(s)]}v^{(\alpha+1)/\alpha}(s) \right\} ds.
\end{aligned} \tag{3.17}$$

Define

$$A := \left[ \frac{\alpha\sigma'(t)}{\rho^{1/\alpha}(t)r^{1/\alpha}[\sigma(t)]} \right]^{\alpha/(\alpha+1)} \omega(t), \quad B := \left[ \frac{\rho'_+(t)}{\rho(t)} \frac{\alpha}{\alpha+1} \left[ \frac{\alpha\sigma'(t)}{\rho^{1/\alpha}(t)r^{1/\alpha}[\sigma(t)]} \right]^{-\alpha/(\alpha+1)} \right]^\alpha. \tag{3.18}$$

Using the following inequality:

$$\frac{\alpha+1}{\alpha}AB^{1/\alpha} - A^{(\alpha+1)/\alpha} \leq \frac{1}{\alpha}B^{(\alpha+1)/\alpha}, \quad \text{for } A \geq 0, B \geq 0 \text{ are constants,} \tag{3.19}$$

we have

$$\frac{\rho'_+(t)}{\rho(t)}\omega(t) - \frac{\alpha\sigma'(t)}{\rho^{1/\alpha}(t)r^{1/\alpha}[\sigma(t)]}\omega^{(\alpha+1)/\alpha}(t) \leq \frac{1}{(\alpha+1)^{\alpha+1}} \frac{r[\sigma(t)](\rho'_+(t))^{\alpha+1}}{(\rho(t)\sigma'(t))^\alpha}. \tag{3.20}$$

On the other hand, define

$$A := \left[ \frac{\alpha\sigma'(t)}{\rho^{1/\alpha}(t)r^{1/\alpha}[\sigma(t)]} \right]^{\alpha/(\alpha+1)} v(t), \quad B := \left[ \zeta_+(t) \frac{\alpha}{\alpha+1} \left[ \frac{\alpha\sigma'(t)}{\rho^{1/\alpha}(t)r^{1/\alpha}[\sigma(t)]} \right]^{-\alpha/(\alpha+1)} \right]^\alpha. \tag{3.21}$$

So we have

$$\zeta_+(t)v(t) - \frac{\alpha\sigma'(t)}{\rho^{1/\alpha}(t)r^{1/\alpha}[\sigma(t)]}v^{(\alpha+1)/\alpha}(t) \leq \frac{1}{(\alpha+1)^{\alpha+1}} \frac{r[\sigma(t)](\zeta_+(t))^{\alpha+1}\rho(t)}{(\sigma'(t))^\alpha}. \tag{3.22}$$

Thus, from (3.17), we get

$$\begin{aligned} & \omega(t) - \omega(t_2) + \frac{p^\alpha[\sigma(t)]}{\tau'(t)}v(t) - \frac{p^\alpha[\sigma(t_2)]}{\tau'(t_2)}v(t_2) \\ & \leq - \int_{t_2}^t \rho(s) \left\{ \frac{Q(s)}{2^{\alpha-1}} - \frac{r[\sigma(s)]}{(\alpha+1)^{\alpha+1}(\sigma'(s))^\alpha} \left[ \left( \frac{\rho'_+(s)}{\rho(s)} \right)^{\alpha+1} + \frac{p^\alpha[\sigma(s)](\zeta_+(s))^{\alpha+1}}{\tau'(s)} \right] \right\} ds, \end{aligned} \quad (3.23)$$

which contradicts (3.1). This completes the proof.  $\square$

When  $p(t) \leq p_0 < \infty$ ,  $\tau'(t) \geq \tau_0 > 0$ , where  $p_0, \tau_0$  are constants, we obtain the following result.

**Theorem 3.2.** Suppose that (1.5) holds,  $p(t) \leq p_0 < \infty$ ,  $\tau'(t) \geq \tau_0 > 0$ ,  $\sigma \in C([t_0, \infty), \mathbb{R})$ ,  $\sigma'(t) > 0$ ,  $\sigma(t) \leq t$ ,  $\sigma(t) \leq \tau(t)$  for  $t \geq t_0$ . Further, assume that there exists a function  $\rho \in C([t_0, \infty), (0, \infty))$  such that

$$\limsup_{t \rightarrow \infty} \int_{t_0}^t \left[ \frac{\rho(s)Q(s)}{2^{\alpha-1}} - \frac{1}{(\alpha+1)^{\alpha+1}} \left( 1 + \frac{p_0^\alpha}{\tau_0} \right) \frac{r[\sigma(s)](\rho'_+(s))^{\alpha+1}}{(\rho(s)\sigma'(s))^\alpha} \right] ds = \infty. \quad (3.24)$$

Then (1.1) is oscillatory.

*Proof.* Let  $x$  be a nonoscillatory solution of (1.1). Without loss of generality, we assume that there exists  $t_1 \geq t_0$  such that  $x(t) > 0$ ,  $x[\tau(t)] > 0$  and  $x[\sigma(t)] > 0$  for all  $t \geq t_1$ . Using (1.1), for all sufficiently large  $t$ , we obtain that

$$\begin{aligned} & \left( r(t)|z'(t)|^{\alpha-1}z'(t) \right)' + q(t)x^\alpha[\sigma(t)] + p_0^\alpha q[\tau(t)]x^\alpha(\sigma[\tau(t)]) \\ & + \frac{p_0^\alpha}{\tau_0} \left( r[\tau(t)]|z'[\tau(t)]|^{\alpha-1}z'[\tau(t)] \right)' \leq 0. \end{aligned} \quad (3.25)$$

By applying (2.1) and the definition of  $z$ , we conclude that

$$\begin{aligned} & \left( r(t)|z'(t)|^{\alpha-1}z'(t) \right)' + \frac{1}{2^{\alpha-1}}Q(t)z^\alpha[\sigma(t)] \\ & + \frac{p_0^\alpha}{\tau_0} \left( r[\tau(t)]|z'[\tau(t)]|^{\alpha-1}z'[\tau(t)] \right)' \leq 0. \end{aligned} \quad (3.26)$$

The remainder of the proof is similar to that of Theorem 3.1 and hence is omitted.  $\square$

**Theorem 3.3.** Suppose that (1.5) holds,  $\tau(t) \leq t$ ,  $\sigma(t) \geq \tau(t)$  for  $t \geq t_0$ . Furthermore, assume that there exists a function  $\rho \in C([t_0, \infty), (0, \infty))$  such that

$$\limsup_{t \rightarrow \infty} \int_{t_0}^t \rho(s) \left\{ \frac{Q(s)}{2^{\alpha-1}} - \frac{r[\tau(s)]\varphi(s)}{(\alpha+1)^{\alpha+1}(\tau'(s))^\alpha} \right\} ds = \infty. \quad (3.27)$$

Then (1.1) is oscillatory.



*Proof.* Let  $x$  be a nonoscillatory solution of (1.1). Without loss of generality, we assume that there exists  $t_1 \geq t_0$  such that  $x(t) > 0$ ,  $x[\tau(t)] > 0$  and  $x[\sigma(t)] > 0$  for all  $t \geq t_1$ . Proceeding as in the proof of Theorem 3.1, we get (3.3) and (3.4). In view of (3.4),  $r(t)|z'(t)|^{\alpha-1}z'(t)$  is decreasing function. Now we have two possible cases for  $z'(t)$ : (i)  $z'(t) < 0$  eventually and (ii)  $z'(t) > 0$  eventually.

(i) Suppose that  $z'(t) < 0$  for  $t \geq t_2 \geq t_1$ . Then, similar to the proof of case (i) of Theorem 3.1, we obtain a contradiction.

(ii) Suppose that  $z'(t) > 0$  for  $t \geq t_2 \geq t_1$ . We define a Riccati substitution

$$\omega(t) = \rho(t) \frac{r(t)(z'(t))^\alpha}{(z[\tau(t)])^\alpha}, \quad t \geq t_2. \quad (3.28)$$

Then  $\omega(t) > 0$ . From (3.4), we have

$$z'[\tau(t)] \geq \left( \frac{r(t)}{r[\tau(t)]} \right)^{1/\alpha} z'(t). \quad (3.29)$$

Differentiating (3.28), we find that

$$\omega'(t) = \rho'(t) \frac{r(t)(z'(t))^\alpha}{(z[\tau(t)])^\alpha} + \rho(t) \frac{(r(t)(z'(t))^\alpha)'}{(z[\tau(t)])^\alpha} - \alpha \rho(t) \frac{r(t)(z'(t))^\alpha z^{\alpha-1}[\tau(t)] z'[\tau(t)] \tau'(t)}{(z[\tau(t)])^{2\alpha}}. \quad (3.30)$$

Therefore, by (3.28), (3.29), and (3.30), we see that

$$\omega'(t) \leq \frac{\rho'(t)}{\rho(t)} \omega(t) + \rho(t) \frac{(r(t)(z'(t))^\alpha)'}{(z[\tau(t)])^\alpha} - \frac{\alpha \tau'(t)}{\rho^{1/\alpha}(t) r^{1/\alpha}[\tau(t)]} \omega^{(\alpha+1)/\alpha}(t). \quad (3.31)$$

Similarly, we introduce a Riccati substitution

$$v(t) = \rho(t) \frac{r[\tau(t)](z'[\tau(t)])^\alpha}{(z[\tau(t)])^\alpha}, \quad t \geq t_2. \quad (3.32)$$

Then  $v(t) > 0$ . Differentiating (3.32), we find that

$$\begin{aligned} v'(t) &= \rho'(t) \frac{r[\tau(t)](z'[\tau(t)])^\alpha}{(z[\tau(t)])^\alpha} + \rho(t) \frac{(r[\tau(t)](z'[\tau(t)])^\alpha)'}{(z[\tau(t)])^\alpha} \\ &\quad - \alpha \rho(t) \frac{r[\tau(t)](z'[\tau(t)])^\alpha z^{\alpha-1}[\tau(t)] z'[\tau(t)] \tau'(t)}{(z[\tau(t)])^{2\alpha}}. \end{aligned} \quad (3.33)$$

Therefore, by (3.32) and (3.33), we see that

$$v'(t) = \frac{\rho'(t)}{\rho(t)} v(t) + \rho(t) \frac{(r[\tau(t)](z'[\tau(t)])^\alpha)'}{(z[\tau(t)])^\alpha} - \frac{\alpha \tau'(t)}{\rho^{1/\alpha}(t) r^{1/\alpha}[\tau(t)]} v^{(\alpha+1)/\alpha}(t). \quad (3.34)$$

Thus, from (3.31) and (3.33), we have

$$\begin{aligned} \omega'(t) + \frac{p^\alpha[\sigma(t)]}{\tau'(t)} v'(t) &\leq \rho(t) \left\{ \frac{(r(t)(z'(t))^\alpha)' }{(z[\tau(t)])^\alpha} + \frac{p^\alpha[\sigma(t)]}{\tau'(t)} \frac{(r[\tau(t)](z'[\tau(t)])^\alpha)' }{(z[\tau(t)])^\alpha} \right\} + \frac{\rho'(t)}{\rho(t)} \omega(t) \\ &\quad - \frac{\alpha\tau'(t)}{\rho^{1/\alpha}(t)r^{1/\alpha}[\tau(t)]} \omega^{(\alpha+1)/\alpha}(t) + \frac{p^\alpha[\sigma(t)]}{\tau'(t)} \frac{\rho'(t)}{\rho(t)} v(t) \\ &\quad - \frac{p^\alpha[\sigma(t)]}{\tau'(t)} \frac{\alpha\tau'(t)}{\rho^{1/\alpha}(t)r^{1/\alpha}[\tau(t)]} v^{(\alpha+1)/\alpha}(t). \end{aligned} \quad (3.35)$$

It follows from (3.3) that

$$\begin{aligned} \omega'(t) + \frac{p^\alpha[\sigma(t)]}{\tau'(t)} v'(t) &\leq -\frac{1}{2^{\alpha-1}} \rho(t) Q(t) + \frac{\rho'_+(t)}{\rho(t)} \omega(t) \\ &\quad - \frac{\alpha\tau'(t)}{\rho^{1/\alpha}(t)r^{1/\alpha}[\tau(t)]} \omega^{(\alpha+1)/\alpha}(t) + \frac{p^\alpha[\sigma(t)]}{\tau'(t)} \frac{\rho'_+(t)}{\rho(t)} v(t) \\ &\quad - \frac{p^\alpha[\sigma(t)]}{\tau'(t)} \frac{\alpha\tau'(t)}{\rho^{1/\alpha}(t)r^{1/\alpha}[\tau(t)]} v^{(\alpha+1)/\alpha}(t). \end{aligned} \quad (3.36)$$

Integrating the above inequality from  $t_2$  to  $t$ , we obtain that

$$\begin{aligned} \omega(t) - \omega(t_2) + \frac{p^\alpha[\sigma(t)]}{\tau'(t)} v(t) - \frac{p^\alpha[\sigma(t_2)]}{\tau'(t_2)} v(t_2) \\ \leq - \int_{t_2}^t \frac{1}{2^{\alpha-1}} \rho(s) Q(s) ds \\ + \int_{t_2}^t \left[ \frac{\rho'_+(s)}{\rho(s)} \omega(s) - \frac{\alpha\tau'(s)}{\rho^{1/\alpha}(s)r^{1/\alpha}[\tau(s)]} \omega^{(\alpha+1)/\alpha}(s) \right] ds \\ + \int_{t_2}^t \frac{p^\alpha[\sigma(s)]}{\tau'(s)} \left\{ \left[ \frac{\rho'_+(s)}{\rho(s)} + \xi(s) \right]_+ v(s) - \frac{\alpha\tau'(s)}{\rho^{1/\alpha}(s)r^{1/\alpha}[\tau(s)]} v^{(\alpha+1)/\alpha}(s) \right\} ds. \end{aligned} \quad (3.37)$$

Define

$$A := \left[ \frac{\alpha\tau'(t)}{\rho^{1/\alpha}(t)r^{1/\alpha}[\tau(t)]} \right]^{\alpha/(\alpha+1)} \omega(t), \quad B := \left[ \frac{\rho'_+(t)}{\rho(t)} \frac{\alpha}{\alpha+1} \left[ \frac{\alpha\tau'(t)}{\rho^{1/\alpha}(t)r^{1/\alpha}[\tau(t)]} \right]^{-\alpha/(\alpha+1)} \right]^\alpha. \quad (3.38)$$

Using (3.19), we have

$$\frac{\rho'_+(t)}{\rho(t)}\omega(t) - \frac{\alpha\tau'(t)}{\rho^{1/\alpha}(t)r^{1/\alpha}[\tau(t)]}\omega^{(\alpha+1)/\alpha}(t) \leq \frac{1}{(\alpha+1)^{\alpha+1}} \frac{r[\tau(t)](\rho'_+(t))^{\alpha+1}}{(\rho(t)\tau'(t))^\alpha}. \quad (3.39)$$

On the other hand, define

$$A := \left[ \frac{\alpha\tau'(t)}{\rho^{1/\alpha}(t)r^{1/\alpha}[\tau(t)]} \right]^{\alpha/(\alpha+1)} v(t), \quad B := \left[ \zeta_+(t) \frac{\alpha}{\alpha+1} \left[ \frac{\alpha\tau'(t)}{\rho^{1/\alpha}(t)r^{1/\alpha}[\tau(t)]} \right]^{-\alpha/(\alpha+1)} \right]^\alpha. \quad (3.40)$$

So we have

$$\zeta_+(t)v(t) - \frac{\alpha\tau'(t)}{\rho^{1/\alpha}(t)r^{1/\alpha}[\tau(t)]}v^{(\alpha+1)/\alpha}(t) \leq \frac{1}{(\alpha+1)^{\alpha+1}} \frac{r[\tau(t)](\zeta_+(t))^{\alpha+1}\rho(t)}{(\tau'(t))^\alpha}. \quad (3.41)$$

Thus, from (3.37), we get

$$\begin{aligned} & \omega(t) - \omega(t_2) + \frac{p^\alpha[\sigma(t)]}{\tau'(t)}v(t) - \frac{p^\alpha[\sigma(t_2)]}{\tau'(t_2)}v(t_2) \\ & \leq - \int_{t_2}^t \rho(s) \left\{ \frac{Q(s)}{2^{\alpha-1}} - \frac{r[\tau(s)]}{(\alpha+1)^{\alpha+1}(\tau'(s))^\alpha} \left[ \left( \frac{\rho'_+(s)}{\rho(s)} \right)^{\alpha+1} + \frac{p^\alpha[\sigma(s)](\zeta_+(s))^{\alpha+1}}{\tau'(s)} \right] \right\} ds, \end{aligned} \quad (3.42)$$

which contradicts (3.27). This completes the proof.  $\square$

When  $p(t) \leq p_0 < \infty$ ,  $\tau'(t) \geq \tau_0 > 0$ , where  $p_0, \tau_0$  are constants, we obtain the following result.

**Theorem 3.4.** Suppose that (1.5) holds,  $p(t) \leq p_0 < \infty$ ,  $\tau'(t) \geq \tau_0 > 0$ ,  $\tau(t) \leq t$ ,  $\sigma(t) \geq \tau(t)$  for  $t \geq t_0$ . Furthermore, assume that there exists a function  $\rho \in C([t_0, \infty), (0, \infty))$  such that

$$\limsup_{t \rightarrow \infty} \int_{t_0}^t \left[ \frac{\rho(s)Q(s)}{2^{\alpha-1}} - \frac{1}{(\alpha+1)^{\alpha+1}} \left( 1 + \frac{p_0^\alpha}{\tau_0} \right) \frac{r[\tau(s)](\rho'_+(s))^{\alpha+1}}{(\tau_0\rho(s))^\alpha} \right] ds = \infty. \quad (3.43)$$

Then (1.1) is oscillatory.

*Proof.* Let  $x$  be a nonoscillatory solution of (1.1). Without loss of generality, we assume that there exists  $t_1 \geq t_0$  such that  $x(t) > 0$ ,  $x[\tau(t)] > 0$  and  $x[\sigma(t)] > 0$  for all  $t \geq t_1$ . Using (1.1) and the definition of  $z$ , we obtain (3.26) for all sufficiently large  $t$ . The remainder of the proof is similar to that of Theorem 3.3 and hence is omitted.  $\square$

#### 4. Oscillation Criteria for the Case (1.13)

In this section, we will establish some oscillation criteria for (1.1) under the case (1.13).

In the following, we assume that  $p_0, \tau_0$  are constants.

**Theorem 4.1.** Suppose that (1.13) holds,  $p(t) \leq p_0 < \infty$ ,  $\tau'(t) \geq \tau_0 > 0$ ,  $\sigma(t) \leq t$ ,  $\sigma \in C([t_0, \infty), \mathbb{R})$ ,  $\sigma'(t) > 0$ ,  $\sigma(t) \leq \tau(t)$  for  $t \geq t_0$ . Further, assume that there exists a function  $\rho \in C([t_0, \infty), (0, \infty))$  such that (3.24) holds. If there exists a function  $\eta \in C^1([t_0, \infty), \mathbb{R})$ ,  $\eta(t) \geq t$ ,  $\eta'(t) > 0$  for  $t \geq t_0$  such that

$$\limsup_{t' \rightarrow \infty} \int_{t_0}^{t'} \left[ \frac{Q(s)}{2^{\alpha-1}} \delta^\alpha(s) - \left( 1 + \frac{p_0^\alpha}{\tau_0} \right) \left( \frac{\alpha}{\alpha+1} \right)^{\alpha+1} \frac{\eta'(s)}{\delta(s) r^{1/\alpha} [\eta(s)]} \right] ds = \infty, \quad (4.1)$$

then (1.1) is oscillatory.

*Proof.* Let  $x$  be a nonoscillatory solution of (1.1). Without loss of generality, we assume that there exists  $t_1 \geq t_0$  such that  $x(t) > 0$ ,  $x[\tau(t)] > 0$  and  $x[\sigma(t)] > 0$  for all  $t \geq t_1$ . Proceeding as in the proof of Theorem 3.2, we get (3.26). In view of (1.1), we have (3.4). Thus,  $r(t)|z'(t)|^{\alpha-1}z'(t)$  is decreasing function. Now we have two possible cases for  $z'(t)$ : (i)  $z'(t) < 0$  eventually and (ii)  $z'(t) > 0$  eventually.

(i) Suppose that  $z'(t) > 0$  for  $t \geq t_2 \geq t_1$ . Then, by Theorem 3.2, we obtain a contradiction with (3.24).

(ii) Suppose that  $z'(t) < 0$  for  $t \geq t_2 \geq t_1$ . We define the function  $u$  by

$$u(t) = -\frac{r(t)(-z'(t))^\alpha}{z^\alpha[\eta(t)]}, \quad t \geq t_2. \quad (4.2)$$

Then  $u(t) < 0$ . Noting that  $r(t)(-z'(t))^\alpha$  is increasing, we get

$$r^{1/\alpha}(s)z'(s) \leq r^{1/\alpha}(t)z'(t), \quad s \geq t \geq t_2. \quad (4.3)$$

Dividing the above inequality by  $r^{1/\alpha}(s)$ , and integrating it from  $\eta(t)$  to  $t'$ , we obtain that

$$z(t') \leq z[\eta(t)] + r^{1/\alpha}(t)z'(t) \int_{\eta(t)}^{t'} \frac{ds}{r^{1/\alpha}(s)}. \quad (4.4)$$

Letting  $t' \rightarrow \infty$ , we have

$$0 \leq z[\eta(t)] + r^{1/\alpha}(t)z'(t)\delta(t), \quad (4.5)$$

that is,

$$-\delta(t) \frac{r^{1/\alpha}(t)z'(t)}{z[\eta(t)]} \leq 1. \quad (4.6)$$

Hence, by (4.2), we get

$$-\delta^\alpha(t)u(t) \leq 1. \quad (4.7)$$

Similarly, we define the function  $v$  by

$$v(t) = -\frac{r[\tau(t)](-z'[\tau(t)])^\alpha}{z^\alpha[\eta(t)]}, \quad t \geq t_2. \quad (4.8)$$

Then  $v(t) < 0$ . Noting that  $r(t)(-z'(t))^\alpha$  is increasing, we get the following:

$$r(t)(-z'(t))^\alpha \geq r[\tau(t)](-z'[\tau(t)])^\alpha. \quad (4.9)$$

Thus  $0 < -v(t) \leq -u(t)$ . So by (4.7), we see that

$$-\delta^\alpha(t)v(t) \leq 1. \quad (4.10)$$

Differentiating (4.2), we obtain that

$$u'(t) = \frac{(-r(t)(-z'(t))^\alpha)' z^\alpha[\eta(t)] + \alpha r(t)(-z'(t))^\alpha z^{\alpha-1}[\eta(t)] z'[\eta(t)] \eta'(t)}{z^{2\alpha}[\eta(t)]}, \quad (4.11)$$

by (3.4), and we have  $z'[\eta(t)] \leq (r(t)/r[\eta(t)])^{1/\alpha} z'(t)$ , so

$$u'(t) \leq \frac{(-r(t)(-z'(t))^\alpha)'}{z^\alpha[\eta(t)]} - \alpha \frac{\eta'(t)}{r^{1/\alpha}[\eta(t)]} (-u(t))^{(\alpha+1)/\alpha}. \quad (4.12)$$

Similarly, we see that

$$v'(t) \leq \frac{(-r[\tau(t)](-z'[\tau(t)])^\alpha)'}{z^\alpha[\eta(t)]} - \alpha \frac{\eta'(t)}{r^{1/\alpha}[\eta(t)]} (-v(t))^{(\alpha+1)/\alpha}. \quad (4.13)$$

Therefore, by (4.12) and (4.13), we get the following:

$$\begin{aligned} u'(t) + \frac{p_0^\alpha}{\tau_0} v'(t) &\leq \frac{(-r(t)(-z'(t))^\alpha)'}{z^\alpha[\eta(t)]} + \frac{p_0^\alpha}{\tau_0} \frac{(-r[\tau(t)](-z'[\tau(t)])^\alpha)'}{z^\alpha[\eta(t)]} \\ &\quad - \alpha \frac{\eta'(t)}{r^{1/\alpha}[\eta(t)]} (-u(t))^{(\alpha+1)/\alpha} - \frac{\alpha p_0^\alpha}{\tau_0} \frac{\eta'(t)}{r^{1/\alpha}[\eta(t)]} (-v(t))^{(\alpha+1)/\alpha}. \end{aligned} \quad (4.14)$$

Using (3.26) and (4.14), we obtain that

$$u'(t) + \frac{p_0^\alpha}{\tau_0} v'(t) \leq -\frac{Q(t)}{2^{\alpha-1}} - \alpha \frac{\eta'(t)}{r^{1/\alpha}[\eta(t)]} (-u(t))^{(\alpha+1)/\alpha} - \frac{\alpha p_0^\alpha}{\tau_0} \frac{\eta'(t)}{r^{1/\alpha}[\eta(t)]} (-v(t))^{(\alpha+1)/\alpha}. \quad (4.15)$$

Multiplying (4.15) by  $\delta^\alpha(t)$ , and integrating it from  $t_2$  to  $t'$ , we have

$$\begin{aligned} & u(t')\delta^\alpha(t') - u(t_2)\delta^\alpha(t_2) + \alpha \int_{t_2}^{t'} \frac{\delta^{\alpha-1}(s)\eta'(s)u(s)}{r^{1/\alpha}[\eta(s)]} ds + \alpha \int_{t_2}^{t'} \frac{\eta'(s)\delta^\alpha(s)}{r^{1/\alpha}[\eta(s)]} (-u(s))^{(\alpha+1)/\alpha} ds \\ & + \frac{p_0^\alpha}{\tau_0} v(t')\delta^\alpha(t') - \frac{p_0^\alpha}{\tau_0} v(t_2)\delta^\alpha(t_2) + \frac{\alpha p_0^\alpha}{\tau_0} \int_{t_2}^{t'} \frac{\delta^{\alpha-1}(s)\eta'(s)v(s)}{r^{1/\alpha}[\eta(s)]} ds \\ & + \frac{\alpha p_0^\alpha}{\tau_0} \int_{t_2}^{t'} \frac{\eta'(s)\delta^\alpha(s)}{r^{1/\alpha}[\eta(s)]} (-v(s))^{(\alpha+1)/\alpha} ds + \int_{t_2}^{t'} \frac{Q(s)}{2^{\alpha-1}} \delta^\alpha(s) ds \leq 0. \end{aligned} \quad (4.16)$$

Using (3.19), (4.7), and (4.10), we find that

$$\begin{aligned} \int_{t_2}^{t'} \left[ \frac{Q(s)}{2^{\alpha-1}} \delta^\alpha(s) - \left( 1 + \frac{p_0^\alpha}{\tau_0} \right) \left( \frac{\alpha}{\alpha+1} \right)^{\alpha+1} \frac{\eta'(s)}{\delta(s)r^{1/\alpha}[\eta(s)]} \right] ds & \leq u(t_2)\delta^\alpha(t_2) + \frac{p_0^\alpha}{\tau_0} v(t_2)\delta^\alpha(t_2) \\ & + 1 + \frac{p_0^\alpha}{\tau_0}. \end{aligned} \quad (4.17)$$

Letting  $t' \rightarrow \infty$ , we obtain a contradiction with (4.1). This completes the proof.  $\square$

From Theorems 3.4 and 4.1, we have the following result.

**Theorem 4.2.** Suppose that (1.13) holds,  $p(t) \leq p_0 < \infty$ ,  $\tau'(t) \geq \tau_0 > 0$ ,  $\tau(t) \leq t$ ,  $\sigma(t) \geq \tau(t)$  for  $t \geq t_0$ . Further, assume that there exists a function  $\rho \in C([t_0, \infty), (0, \infty))$  such that (3.43) holds. If there exists a function  $\eta \in C^1([t_0, \infty), \mathbb{R})$ ,  $\eta(t) \geq t$ ,  $\eta'(t) > 0$ ,  $\sigma(t) \leq \eta(t)$  for  $t \geq t_0$  such that (4.1) holds, then (1.1) is oscillatory.

## 5. Examples

In this section, we will give some examples to illustrate the main results.

*Example 5.1.* Study the second-order neutral differential equation

$$\left[ \left| (x(t) + tx(t - \lambda_1))' \right|^{\alpha-1} (x(t) + tx(t - \lambda_1))' \right]' + \beta |x(t - \lambda_2)|^{\alpha-1} x(t - \lambda_2) = 0, \quad t \geq t_0, \quad (5.1)$$

where  $\alpha \geq 1$ ,  $0 < \lambda_1 \leq \lambda_2 < 1$ ,  $\beta > 0$  are constants.

Let  $r(t) = 1$ ,  $p(t) = t$ ,  $\rho(t) = 1$ . It is easy to see that all the conditions of Theorem 3.1 hold. Hence, (5.1) is oscillatory.

*Example 5.2.* Consider the second-order quasilinear neutral differential equation

$$\left[ \left| (x(t) + p(t)x(\lambda_1 t))' \right|^{\alpha-1} (x(t) + p(t)x(\lambda_1 t))' \right]' + \frac{\beta}{t^{\alpha+1}} |x(\lambda_2 t)|^{\alpha-1} x(\lambda_2 t) = 0, \quad t \geq t_0, \quad (5.2)$$

where  $\alpha \geq 1$ ,  $\beta > 0$  are constants,  $\lambda_1, \lambda_2 \in (0, 1)$ ,  $\lambda_2 \leq \lambda_1$ .

Let  $r(t) = 1$ ,  $0 \leq p(t) \leq p_0 < \infty$ ,  $q(t) = \beta/t^{\alpha+1}$ ,  $\tau(t) = \lambda_1 t$ ,  $\sigma(t) = \lambda_2 t$ , and  $\rho(t) = t^\alpha$ . Then, we have

$$\begin{aligned} \limsup_{t \rightarrow \infty} \int_{t_0}^t \left[ \frac{\rho(s)Q(s)}{2^{\alpha-1}} - \frac{1}{(\alpha+1)^{\alpha+1}} \left( 1 + \frac{p_0^\alpha}{\tau_0} \right) \frac{r[\sigma(s)]((\rho'(s)_+)^{\alpha+1})}{(\rho(s)\sigma'(s))^\alpha} \right] ds \\ = \left[ \frac{\beta}{2^{\alpha-1}} - \frac{\alpha^{\alpha+1}}{(\alpha+1)^{\alpha+1}\lambda_2^\alpha} \left( 1 + \frac{p_0^\alpha}{\lambda_1} \right) \right] \limsup_{t \rightarrow \infty} \int_{t_0}^t \frac{ds}{s} = \infty, \end{aligned} \quad (5.3)$$

if  $\beta > 2^{\alpha-1}\alpha^{\alpha+1}(1 + p_0^\alpha/\lambda_1)/[(\alpha+1)^{\alpha+1}\lambda_2^\alpha]$ . Hence, by Theorem 3.2, (5.2) is oscillatory if

$$\beta > \frac{2^{\alpha-1}\alpha^{\alpha+1}}{(\alpha+1)^{\alpha+1}\lambda_2^\alpha} \left( 1 + \frac{p_0^\alpha}{\lambda_1} \right). \quad (5.4)$$

*Example 5.3.* Investigate the second-order neutral differential equation

$$\left[ x(t) + \frac{(t-1)(t-4\pi)}{t} x(t-4\pi) \right]'' + (t-2\pi)x(t-2\pi) = 0, \quad t \geq t_0. \quad (5.5)$$

Let  $r(t) = 1$ ,  $p(t) = (t-1)(t-4\pi)/t$ ,  $q(t) = t-2\pi$ , and  $\rho(t) = 1$ . It is easy to see that all the conditions of Theorem 3.3 hold. Hence, (5.5) is oscillatory, for example,  $x(t) = \sin t/t$  is a solution of (5.5).

*Example 5.4.* Discuss the second-order quasilinear neutral differential equation

$$\left[ \left| (x(t) + p(t)x(\lambda_1 t))' \right|^{\alpha-1} (x(t) + p(t)x(\lambda_1 t))' \right]' + \frac{\beta}{t^{\alpha+1}} |x(\lambda_2 t)|^{\alpha-1} x(\lambda_2 t) = 0, \quad t \geq t_0, \quad (5.6)$$

where  $\alpha \geq 1$ ,  $\beta > 0$  are constants,  $\lambda_1 \in (0, 1)$ ,  $\lambda_2 \in [\lambda_1, \infty)$ .

Let  $r(t) = 1$ ,  $0 \leq p(t) \leq p_0 < \infty$ ,  $q(t) = \beta/t^{\alpha+1}$ ,  $\tau(t) = \lambda_1 t$ ,  $\sigma(t) = \lambda_2 t$ ,  $\rho(t) = t^\alpha$ . Then, we have

$$\begin{aligned} \limsup_{t \rightarrow \infty} \int_{t_0}^t \left[ \frac{\rho(s)Q(s)}{2^{\alpha-1}} - \frac{1}{(\alpha+1)^{\alpha+1}} \left( 1 + \frac{p_0^\alpha}{\tau_0} \right) \frac{r[\tau(s)](\rho'(s)_+)^{\alpha+1}}{\tau_0(\rho(s))^\alpha} \right] ds \\ = \left[ \frac{\beta}{2^{\alpha-1}} - \frac{\alpha^{\alpha+1}}{(\alpha+1)^{\alpha+1}\lambda_1^\alpha} \left( 1 + \frac{p_0^\alpha}{\lambda_1} \right) \right] \limsup_{t \rightarrow \infty} \int_{t_0}^t \frac{ds}{s} = \infty \end{aligned} \quad (5.7)$$

if  $\beta > 2^{\alpha-1}\alpha^{\alpha+1}(1 + p_0^\alpha/\lambda_1)/[(\alpha+1)^{\alpha+1}\lambda_1^\alpha]$ . Hence, by Theorem 3.4, (5.6) is oscillatory if

$$\beta > \frac{2^{\alpha-1}\alpha^{\alpha+1}}{(\alpha+1)^{\alpha+1}\lambda_1^\alpha} \left( 1 + \frac{p_0^\alpha}{\lambda_1} \right). \quad (5.8)$$

*Example 5.5.* Examine the second-order quasilinear neutral differential equation

$$\left[ t^{2\alpha} \left| (x(t) + p(t)x(\lambda_1 t))' \right|^{\alpha-1} (x(t) + p(t)x(\lambda_1 t))' \right]' + \beta t^{\alpha-1} |x(\lambda_2 t)|^{\alpha-1} x(\lambda_2 t) = 0, \quad t \geq t_0, \quad (5.9)$$

where  $\alpha \geq 1$ ,  $\beta > 0$  are constants,  $\lambda_1, \lambda_2 \in (0, 1)$ ,  $\lambda_2 \leq \lambda_1$ .

Let  $r(t) = t^{2\alpha}$ ,  $0 \leq p(t) \leq p_0 < \infty$ ,  $q(t) = \beta t^{\alpha-1}$ ,  $\tau(t) = \lambda_1 t$ ,  $\sigma(t) = \lambda_2 t$ , and  $\rho(t) = 1$ . Then,  $Q(t) = \beta \lambda_1^{\alpha-1} t^{\alpha-1}$ . It is easy to see that (3.24) holds. On the other hand, taking  $\eta(t) = t$ , then  $\delta(t) = 1/t$ . Therefore, one has

$$\begin{aligned} & \limsup_{t' \rightarrow \infty} \int_{t_0}^{t'} \left[ \frac{Q(s)}{2^{\alpha-1}} \delta^\alpha(s) - \left( 1 + \frac{p_0^\alpha}{\tau_0} \right) \left( \frac{\alpha}{\alpha+1} \right)^{\alpha+1} \frac{\eta'(s)}{\delta(s) r^{1/\alpha} [\eta(s)]} \right] ds \\ &= \left[ \frac{\beta}{2^{\alpha-1}} \lambda_1^{\alpha-1} - \frac{1}{2} \left( 1 + \frac{p_0^\alpha}{\lambda_1} \right) \left( \frac{\alpha}{\alpha+1} \right)^{\alpha+1} \right] \limsup_{t' \rightarrow \infty} \int_{t_0}^{t'} \frac{ds}{s} = \infty \end{aligned} \quad (5.10)$$

if  $\beta > 2^{\alpha-2} (1 + p_0^\alpha / \lambda_1) (\alpha / \alpha + 1)^{\alpha+1} / \lambda_1^{\alpha-1}$ . Thus, by Theorem 4.1, (5.9) oscillates if

$$\beta > \frac{2^{\alpha-2}}{\lambda_1^{\alpha-1}} \left( 1 + \frac{p_0^\alpha}{\lambda_1} \right) \left( \frac{\alpha}{\alpha+1} \right)^{\alpha+1}. \quad (5.11)$$

## 6. Conclusions

Inequality technique plays an important role in studying the oscillatory behavior of differential equations; in this paper, we establish a new inequality (2.1); by using (2.1) and Riccati substitution, we establish some new oscillation criteria for (1.1). Theorem 3.1 can be applied to the case  $\tau(t) \geq t$ . Specially, taking  $\alpha = 1$ , our results include and improve the results in [15]; for example, and Theorem 4.1 includes [15, Theorem 3.1], Theorem 4.2 includes [15, Theorem 3.2]. The method can be applied on the second-order Emden-Fowler neutral differential equations

$$\left[ r(t) (x(t) + p(t)x(\tau(t)))' \right]' + q(t) |x(\delta(t))|^{\alpha-1} x(\delta(t)) = 0, \quad t \geq t_0, \quad (6.1)$$

where  $\alpha \geq 1$ . It would be interesting to find another method to investigate (1.1) when  $\tau \circ \sigma \neq \sigma \circ \tau$ .

## Acknowledgment

The authors sincerely thank the reviewers for their valuable suggestions and useful comments that have led to the present improved version of the original manuscript. This paper is supported by the Natural Science Foundation of China (11071143, 60904024, 61174217), Natural Science Outstanding Youth Foundation of Shandong Province (JQ201119), Shandong Provincial Natural Science Foundation (ZR2010AL002, ZR2009AL003), and by Natural Science Foundation of Educational Department of Shandong Province (J11LA01).



## References

- [1] J. Hale, *Theory of Functional Differential Equations*, Springer, New York, NY, USA, 1977.
- [2] R. P. Agarwal, S.-L. Shieh, and C.-C. Yeh, "Oscillation criteria for second-order retarded differential equations," *Mathematical and Computer Modelling*, vol. 26, no. 4, pp. 1–11, 1997.
- [3] J.-L. Chern, W.-C. Lian, and C.-C. Yeh, "Oscillation criteria for second order half-linear differential equations with functional arguments," *Publicationes Mathematicae Debrecen*, vol. 48, no. 3-4, pp. 209–216, 1996.
- [4] J. Džurina and I. P. Stavroulakis, "Oscillation criteria for second-order delay differential equations," *Applied Mathematics and Computation*, vol. 140, no. 2-3, pp. 445–453, 2003.
- [5] T. Kusano and N. Yoshida, "Nonoscillation theorems for a class of quasilinear differential equations of second order," *Journal of Mathematical Analysis and Applications*, vol. 189, no. 1, pp. 115–127, 1995.
- [6] T. Kusano and Y. Naito, "Oscillation and nonoscillation criteria for second order quasilinear differential equations," *Acta Mathematica Hungarica*, vol. 76, no. 1-2, pp. 81–99, 1997.
- [7] D. D. Mirzov, "The oscillation of the solutions of a certain system of differential equations," *Matematicheskie Zametki*, vol. 23, no. 3, pp. 401–404, 1978.
- [8] Y. G. Sun and F. W. Meng, "Note on the paper of Džurina and Stavroulakis," *Applied Mathematics and Computation*, vol. 174, no. 2, pp. 1634–1641, 2006.
- [9] M. T. Senel and T. Candan, "Oscillation of second order nonlinear neutral differential equation," *Journal of Computational Analysis and Applications*, vol. 14, no. 6, pp. 1112–1117, 2012.
- [10] L. Liu and Y. Bai, "New oscillation criteria for second-order nonlinear neutral delay differential equations," *Journal of Computational and Applied Mathematics*, vol. 231, no. 2, pp. 657–663, 2009.
- [11] R. Xu and F. Meng, "Some new oscillation criteria for second order quasi-linear neutral delay differential equations," *Applied Mathematics and Computation*, vol. 182, no. 1, pp. 797–803, 2006.
- [12] R. Xu and F. Meng, "Oscillation criteria for second order quasi-linear neutral delay differential equations," *Applied Mathematics and Computation*, vol. 192, no. 1, pp. 216–222, 2007.
- [13] J.-G. Dong, "Oscillation behavior of second order nonlinear neutral differential equations with deviating arguments," *Computers and Mathematics with Applications*, vol. 59, no. 12, pp. 3710–3717, 2010.
- [14] Z. Han, T. Li, S. Sun, and W. Chen, "On the oscillation of second-order neutral delay differential equations," *Advances in Difference Equations*, Article ID 289340, 8 pages, 2010.
- [15] Z. Han, T. Li, S. Sun, and Y. Sun, "Remarks on the paper," *Applied Mathematics and Computation*, vol. 215, no. 11, pp. 3998–4007, 2010.
- [16] A. K. Tripathy, "Some oscillation results for second order nonlinear dynamic equations of neutral type," *Nonlinear Analysis. Theory, Methods and Applications A*, vol. 71, no. 12, pp. e1727–e1735, 2009.
- [17] J. Džurina, "Oscillation theorems for second-order nonlinear neutral differential equations," *Computers and Mathematics with Applications*, vol. 62, no. 12, pp. 4472–4478, 2011.

