

Research Article

Some Relations of the Twisted q -Genocchi Numbers and Polynomials with Weight α and Weak Weight β

J. Y. Kang, H. Y. Lee, and N. S. Jung

*Department of Mathematics, Hannam University,
Daejeon 306-791, Republic of Korea*

Correspondence should be addressed to J. Y. Kang, rkdwjddnr2002@yahoo.co.kr

Received 12 February 2012; Accepted 9 March 2012

Academic Editor: Natig Atakishiyev

Copyright © 2012 J. Y. Kang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Recently many mathematicians are working on Genocchi polynomials and Genocchi numbers. We define a new type of twisted q -Genocchi numbers and polynomials with weight α and weak weight β and give some interesting relations of the twisted q -Genocchi numbers and polynomials with weight α and weak weight β . Finally, we find relations between twisted q -Genocchi zeta function and twisted Hurwitz q -Genocchi zeta function.

1. Introduction

The Genocchi numbers and polynomials possess many interesting properties and arising in many areas of mathematics and physics. Recently, many mathematicians have studied in the area of the q -Genocchi numbers and polynomials (see [1–16]).

Throughout this paper we use the following notations. By \mathbb{Z}_p we denote the ring of p -adic rational integers, \mathbb{Q}_p denotes the field of p -adic rational numbers, \mathbb{C}_p denotes the completion of algebraic closure of \mathbb{Q}_p , \mathbb{N} denotes the set of natural numbers, \mathbb{Z} denotes the ring of rational integers, \mathbb{Q} denotes the field of rational numbers, \mathbb{C} denotes the set of complex numbers, and $\mathbb{Z}^+ = \mathbb{N} \cup \{0\}$. Let v_p be the normalized exponential valuation of \mathbb{C}_p with $|p|_p = p^{-v_p(p)} = p^{-1}$. When one talks of q -extension, q is considered in many ways such as an indeterminate, a complex number $q \in \mathbb{C}$, or p -adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, one normally assume that $|q| < 1$. If $q \in \mathbb{C}_p$, we normally assume that $|q - 1|_p < p^{-1/(p-1)}$ so that $q^x = \exp(x \log q)$ for $|x|_p \leq 1$. Throughout this paper we use the following notation:

$$[x]_q = \frac{1 - q^x}{1 - q}, \quad [x]_{-q} = \frac{1 - (-q)^x}{1 + q} \quad (1.1)$$

(cf. [1–13]).

Hence, $\lim_{q \rightarrow 1} [x] = x$ for any x with $|x|_p \leq 1$ in the present p -adic case.

For

$$f \in UD(\mathbb{Z}_p) = \{f \mid f : \mathbb{Z}_p \longrightarrow \mathbb{C}_p \text{ is uniformly differentiable function}\}, \quad (1.2)$$

the fermionic p -adic q -integral on \mathbb{Z}_p is defined by Kim as follows:

$$I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_{-q}} \sum_{x=0}^{p^N-1} f(x) (-q)^x \quad (1.3)$$

(cf. [11–14]).

If we take $f_1(x) = f(x+1)$ in (1.1), then we easily see that

$$qI_{-q}(f_1) + I_{-q}(f) = [2]_q f(0). \quad (1.4)$$

From (1.4), we obtain

$$q^n I_{-q}(f_n) + (-1)^{n-1} I_{-q}(f) = [2]_q \sum_{l=0}^{n-1} (-1)^{n-1-l} q^l f(l), \quad (1.5)$$

where $f_n(x) = f(x+n)$ (cf. [5–9]).

Let $C_{p^n} = \{w \mid w^{p^n} = 1\}$ be the cyclic group of order p^n and let

$$T_p = \lim_{n \rightarrow \infty} C_{p^n} = C_{p^\infty} = \cup_{n \geq 0} C_{p^n} \quad (1.6)$$

be the locally constant space. For $w \in T_p$, we denote by $\phi_w : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ the locally constant function $x \mapsto w^x$.

As well-known definition, the Genocchi polynomials are defined by

$$F(t) = \frac{2t}{e^t + 1} = e^{Gt} = \sum_{n=0}^{\infty} G_n \frac{t^n}{n!}, \quad (1.7)$$

$$F(t, x) = \frac{2t}{e^t + 1} e^{xt} = e^{G(x)t} = \sum_{n=0}^{\infty} G_n(x) \frac{t^n}{n!},$$

with the usual convention of replacing $G^n(x)$ by $G_n(x)$. $G_n(0) = G_n$ are called the n th Genocchi numbers (cf. [2–5, 14]).

These numbers and polynomials are interpolated by the Genocchi zeta function and Hurwitz-type Genocchi zeta function, respectively:

$$\begin{aligned}\zeta_G(s) &= 2 \sum_{n=1}^{\infty} \frac{(-1)^n}{n^s}, \\ \zeta_G(s, x) &= 2 \sum_{n=0}^{\infty} \frac{(-1)^n}{(n+x)^s}.\end{aligned}\tag{1.8}$$

Our aim in this paper is to define twisted q -Genocchi numbers $G_{n,q,w}^{(\alpha,\beta)}$ and polynomials $G_{n,q,w}^{(\alpha,\beta)}(x)$ with weight α and weak weight β . We investigate some properties which are related to $G_{n,q,w}^{(\alpha,\beta)}$ and $G_{n,q,w}^{(\alpha,\beta)}(x)$. We also derive the existence of a specific interpolation function which interpolate $G_{n,q,w}^{(\alpha,\beta)}$ and $G_{n,q,w}^{(\alpha,\beta)}(x)$ at negative integers.

2. Generating Functions of Twisted q -Genocchi Numbers and Polynomials with Weight α and Weak Weight β

Our primary goal of this section is to define twisted q -Genocchi numbers $G_{n,q,w}^{(\alpha,\beta)}$ and polynomials $G_{n,q,w}^{(\alpha,\beta)}(x)$ with weight α and weak weight β . We also find generating functions of $G_{n,q,w}^{(\alpha,\beta)}$ and $G_{n,q,w}^{(\alpha,\beta)}(x)$.

Definition 2.1. For $\alpha, \beta \in \mathbb{Q}$ and $q \in \mathbb{C}_p$ with $|1 - q|_p \leq 1$,

$$G_{n,q,w}^{(\alpha,\beta)} = n \int_{\mathbb{Z}_p} \phi_w(x) [x]_{q^\alpha}^{n-1} d\mu_{-q^\beta}(x).\tag{2.1}$$

We call $G_{n,q,w}^{(\alpha,\beta)}$ twisted q -Genocchi numbers with weight α and weak weight β .

By using p -adic q -integral on \mathbb{Z}_p , we obtain

$$\begin{aligned}n \int_{\mathbb{Z}_p} \phi_w(x) [x]_{q^\alpha}^{n-1} d\mu_{-q^\beta}(x) &= n \lim_{N \rightarrow \infty} \frac{1}{[p^N]_{-q^\beta}} \sum_{x=0}^{p^N-1} w^x [x]_{q^\alpha}^{n-1} (-q^\beta)^x \\ &= n [2]_{q^\beta} \sum_{m=0}^{\infty} (-1)^m q^{\beta m} w^m [m]_{q^\alpha}^{n-1}.\end{aligned}\tag{2.2}$$

From (2.1) and (2.2), we have

$$G_{n,q,w}^{(\alpha,\beta)} = n [2]_{q^\beta} \left(\frac{1}{1 - q^\alpha} \right)^{n-1} \sum_{l=0}^{n-1} \binom{n-1}{l} (-1)^l \frac{1}{1 + w q^{\beta + \alpha l}}.\tag{2.3}$$

We set

$$F_{q,w}^{(\alpha,\beta)}(t) = \sum_{n=0}^{\infty} G_{n,q,w}^{(\alpha,\beta)} \frac{t^n}{n!}.\tag{2.4}$$

By using the previous equation and (2.3), we have

$$\begin{aligned} F_{q,w}^{(\alpha,\beta)}(t) &= \sum_{n=0}^{\infty} \left(n[2]_{q^\beta} \left(\frac{1}{1-q^\alpha} \right)^{n-1} \sum_{l=0}^{n-1} \binom{n-1}{l} (-1)^l \frac{1}{1+wq^{\beta+al}} \right) \frac{t^n}{n!} \\ &= [2]_{q^\beta} t \sum_{m=0}^{\infty} (-1)^m q^{\beta m} w^m e^{[m]_{q^\alpha} t}. \end{aligned} \quad (2.5)$$

Thus twisted q -Genocchi numbers $G_{n,q,w}^{(\alpha,\beta)}$ with weight α and weak weight β are defined by means of the generating function:

$$F_{q,w}^{(\alpha,\beta)}(t) = [2]_{q^\beta} t \sum_{n=0}^{\infty} (-1)^n q^{\beta n} w^n e^{[n]_{q^\alpha} t}. \quad (2.6)$$

By using (2.2), we have

$$\sum_{n=0}^{\infty} G_{n,q,w}^{(\alpha,\beta)} \frac{t^n}{n!} = t \int_{\mathbb{Z}_p} \phi_w(x) e^{[x]_{q^\alpha} t} d\mu_{-q^\beta}(x). \quad (2.7)$$

From (2.5) and (2.7), we have

$$\sum_{n=0}^{\infty} G_{n,q,w}^{(\alpha,\beta)} \frac{t^n}{n!} = [2]_{q^\beta} t \sum_{m=0}^{\infty} (-1)^m q^{\beta m} w^m e^{[m]_{q^\alpha} t}. \quad (2.8)$$

Next, we introduce twisted q -Genocchi polynomials $G_{n,q,w}^{(\alpha,\beta)}(x)$ with weight α and weak weight β .

Definition 2.2. For $\alpha, \beta \in \mathbb{Q}$ and $q \in \mathbb{C}_p$ with $|1-q|_p \leq 1$,

$$G_{n,q,w}^{(\alpha,\beta)}(x) = n \int_{\mathbb{Z}_p} \phi_w(y) [x+y]_{q^\alpha}^{n-1} d\mu_{-q^\beta}(y). \quad (2.9)$$

We call $G_{n,q,w}^{(\alpha,\beta)}(x)$ twisted q -Genocchi polynomials with weight α and weak weight β .

By using p -adic q -integral, we have

$$\begin{aligned} n \int_{\mathbb{Z}_p} \phi_w(y) [x+y]_{q^\alpha}^{n-1} d\mu_{-q^\beta}(y) &= n \lim_{N \rightarrow \infty} \frac{1}{[p^N]_{-q^\beta}} \sum_{y=0}^{p^N-1} w^y [x+y]_{q^\alpha}^{n-1} (-q^\beta)^y \\ &= n[2]_{q^\beta} \sum_{m=0}^{\infty} (-1)^m q^{\beta m} w^m [x+m]_{q^\alpha}^{n-1}. \end{aligned} \quad (2.10)$$

By using (2.9) and (2.10), we obtain

$$G_{n,q,w}^{(\alpha,\beta)}(x) = n[2]_{q^\beta} \left(\frac{1}{1-q^\alpha} \right)^{n-1} \sum_{l=0}^{n-1} \binom{n-1}{l} (-1)^l q^{\alpha x l} \frac{1}{1+wq^{\beta+al}}. \quad (2.11)$$

We set

$$F_{q,w}^{(\alpha,\beta)}(t, x) = \sum_{n=0}^{\infty} G_{n,q,w}^{(\alpha,\beta)}(x) \frac{t^n}{n!}. \quad (2.12)$$

By using the previous equation and (2.11), we have

$$\begin{aligned} F_{q,w}^{(\alpha,\beta)}(t, x) &= \sum_{n=0}^{\infty} \left(n[2]_{q^\beta} \left(\frac{1}{1-q^\alpha} \right)^{n-1} \sum_{l=0}^{n-1} \binom{n-1}{l} (-1)^l q^{\alpha x l} \frac{1}{1+wq^{\beta+al}} \right) \frac{t^n}{n!} \\ &= [2]_{q^\beta} t \sum_{m=0}^{\infty} (-1)^m q^{\beta m} w^m e^{[x+m]_{q^\alpha} t}. \end{aligned} \quad (2.13)$$

Thus twisted q -Genocchi polynomials $G_{n,q,w}^{(\alpha,\beta)}(x)$ with weight α and weak weight β are defined by means of the generating function:

$$F_{q,w}^{(\alpha,\beta)}(t, x) = [2]_{q^\beta} t \sum_{m=0}^{\infty} (-1)^m q^{\beta m} w^m e^{[x+m]_{q^\alpha} t}. \quad (2.14)$$

By using (2.9), we have

$$\sum_{n=0}^{\infty} G_{n,q,w}^{(\alpha,\beta)}(x) \frac{t^n}{n!} = t \int_{\mathbb{Z}_p} \phi_w(y) e^{[x+y]_{q^\alpha} t} d\mu_{-q^\beta}(y). \quad (2.15)$$

By (2.13) and (2.15) we have

$$\sum_{n=0}^{\infty} G_{n,q,w}^{(\alpha,\beta)}(x) \frac{t^n}{n!} = [2]_{q^\beta} t \sum_{m=0}^{\infty} (-1)^m q^{\beta m} w^m e^{[x+m]_{q^\alpha} t}. \quad (2.16)$$

Remark 2.3. In (2.14), we simply identify that

$$\begin{aligned} \lim_{q \rightarrow 1} F_{q,w}^{(\alpha,\beta)}(t, x) &= 2t \sum_{n=0}^{\infty} (-1)^n w^n e^{(x+n)t} \\ &= F_w(t, x). \end{aligned} \quad (2.17)$$

Observe that if $q \rightarrow 1$, then $F_{q,w}^{(\alpha,\beta)}(t) \rightarrow F_w(t)$ and $F_{q,w}^{(\alpha,\beta)}(t, x) \rightarrow F_w(t, x)$. Note that if $q \rightarrow 1$ and $w = 1$, then $G_{n,q,w}^{(\alpha,\beta)} \rightarrow G_n$ and $G_{n,q,w}^{(\alpha,\beta)}(x) \rightarrow G_n(x)$.

3. Some Relations between Twisted q -Genocchi Numbers and Polynomials with Weight α and Weak Weight β

By (2.11), we have the following complement relation.

Theorem 3.1. *One has the property of complement*

$$G_{n,q^{-1},w^{-1}}^{(\alpha,\beta)}(1-x) = (-1)^n w q^{\alpha(n-1)} G_{n,q,w}^{(\alpha,\beta)}(x). \quad (3.1)$$

Also, by (2.11), we have the following distribution relation.

Theorem 3.2. *For any positive integer m (=odd), one has*

$$G_{n,q,w}^{(\alpha,\beta)}(x) = \frac{[2]_{q^\beta}}{[2]_{q^{\beta m}}} [m]_{q^\alpha}^{n-1} \sum_{i=0}^{m-1} (-1)^i w^i q^{\beta i} G_{n,q,w^m}^{(\alpha,\beta)}\left(\frac{i+x}{m}\right), \quad n \in \mathbb{Z}^+. \quad (3.2)$$

Let $f(x) = tw^x e^{[x]_{q^\alpha} t}$. Then by (1.5), left-hand side is in the following form:

$$q^{\beta n} I_{-q^\beta}(f_n) + (-1)^{n-1} I_{-q^\beta}(f) = \sum_{m=0}^{\infty} \left(q^{\beta n} w^n G_{m,q,w}^{(\alpha,\beta)}(n) + (-1)^{n-1} G_{m,q,w}^{(\alpha,\beta)} \right) \frac{t^m}{m!}. \quad (3.3)$$

And right-hand side in (1.5) is in the following form:

$$[2]_{q^\beta} \sum_{l=0}^{n-1} (-1)^{n-1-l} q^{\beta l} f(l) = \sum_{m=0}^{\infty} [2]_{q^\beta} m \sum_{l=0}^{n-1} (-1)^{n-1-l} q^{\beta l} w^l [l]_{q^\alpha}^{m-1} \frac{t^m}{m!}. \quad (3.4)$$

By (3.3) and (3.4), one easily sees that

$$q^{\beta n} w^n G_{m,q,w}^{(\alpha,\beta)}(n) + (-1)^{n-1} G_{m,q,w}^{(\alpha,\beta)} = [2]_{q^\beta} m \sum_{l=0}^{n-1} (-1)^{n-1-l} q^{\beta l} w^l [l]_{q^\alpha}^{m-1}. \quad (3.5)$$

Hence, we have the following theorem.

Theorem 3.3 (Let $m \in \mathbb{Z}^+$). *If $n \equiv 0 \pmod{2}$, then*

$$q^{\beta n} w^n G_{m,q,w}^{(\alpha,\beta)}(n) - G_{m,q,w}^{(\alpha,\beta)} = [2]_{q^\beta} m \sum_{l=0}^{n-1} (-1)^{l+1} q^{\beta l} w^l [l]_{q^\alpha}^{m-1}. \quad (3.6)$$

If $n \equiv 1 \pmod{2}$, then

$$q^{\beta n} w^n G_{m,q,w}^{(\alpha,\beta)}(n) + G_{m,q,w}^{(\alpha,\beta)} = [2]_{q^\beta} m \sum_{l=0}^{n-1} (-1)^l q^{\beta l} w^l [l]_{q^\alpha}^{m-1}. \quad (3.7)$$

Since $[x + y]_{q^\alpha} = [x]_{q^\alpha} + q^{\alpha x}[y]_{q^\alpha}$, one easily obtains that

$$\begin{aligned} G_{n+1,q,w}^{(\alpha,\beta)}(x) &= (n+1) \int_{\mathbb{Z}_p} \phi_w(y) [x+y]_{q^\alpha}^n d\mu_{-q^\beta}(y) \\ &= (n+1)[2]_{q^\beta} \sum_{m=0}^{\infty} (-1)^m q^{\beta m} w^m [x+m]_{q^\alpha}^n. \end{aligned} \quad (3.8)$$

From (1.4), one notes that

$$\begin{aligned} [2]_{q^\beta} t &= q^\beta \int_{\mathbb{Z}_p} t w^{(x+1)} e^{[x+1]_{q^\alpha} t} d\mu_{-q^\beta}(x) + \int_{\mathbb{Z}_p} t w^x e^{[x]_{q^\alpha} t} d\mu_{-q^\beta}(x) \\ &= \sum_{n=0}^{\infty} \left(q^\beta w G_{n,q,w}^{(\alpha,\beta)}(1) + G_{n,q,w}^{(\alpha,\beta)} \right) \frac{t^n}{n!}. \end{aligned} \quad (3.9)$$

By using comparing coefficients of $t^n/n!$ in the previous equation, we easily obtain the following theorem.

Theorem 3.4. For $n \in \mathbb{Z}^+$, one has

$$q^\beta w G_{n,q,w}^{(\alpha,\beta)}(1) + G_{n,q,w}^{(\alpha,\beta)} = \begin{cases} [2]_{q^\beta}, & \text{if } n = 1, \\ 0, & \text{if } n \neq 1. \end{cases} \quad (3.10)$$

By (3.8) and (3.10), we have the following corollary.

Corollary 3.5. For $n \in \mathbb{Z}^+$, one has

$$q^{\beta-\alpha} w \left(q^\alpha G_{q,w}^{(\alpha,\beta)} + 1 \right)^n + G_{n,q,w}^{(\alpha,\beta)} = \begin{cases} [2]_{q^\beta}, & \text{if } n = 1, \\ 0, & \text{if } n \neq 1, \end{cases} \quad (3.11)$$

with the usual convention of replacing $(G_{q,w}^{(\alpha,\beta)})^n$ by $G_{n,q,w}^{(\alpha,\beta)}$.

4. The Analogue of the Genocchi Zeta Function

By using q -Genocchi numbers and polynomials with weight α and weak weight β , q -Genocchi zeta function and Hurwitz q -Genocchi zeta functions are defined. These functions interpolate the q -Genocchi numbers and q -Genocchi polynomials with weight α and weak weight β , respectively. In this section we assume that $q \in \mathbb{C}$ with $|q| < 1$.

From (2.5), we note that

$$\left. \frac{d^{k+1}}{dt^{k+1}} F_{q,w}^{(\alpha,\beta)}(t) \right|_{t=0} = (k+1)[2]_{q^\beta} \sum_{m=0}^{\infty} (-1)^m q^{\beta m} w^m [m]_{q^\alpha}^k = G_{k+1,q,w}^{(\alpha,\beta)} \quad (k \in \mathbb{N}). \quad (4.1)$$

By using the previous equation, we are now ready to define q -Genocchi zeta functions.

Definition 4.1. Let $s \in \mathbb{C}$. One has

$$\zeta_{q,w}^{(\alpha,\beta)}(s) = [2]_{q^\beta} \sum_{n=1}^{\infty} \frac{(-1)^n q^{\beta n} w^n}{[n]_{q^\alpha}^s}. \quad (4.2)$$

Note that $\zeta_{q,w}^{(\alpha,\beta)}(s)$ is a meromorphic function on \mathbb{C} . Observe that if $q \rightarrow 1$, then $\lim_{q \rightarrow 1} \zeta_{q,w}^{(\alpha,\beta)}(s) = \zeta_w(s)$.

Theorem 4.2. Relation between $\zeta_{q,w}^{(\alpha,\beta)}(s)$ and $G_{k,q,w}^{(\alpha,\beta)}$ is given by

$$\zeta_{q,w}^{(\alpha,\beta)}(-k) = \frac{G_{k+1,q,w}^{(\alpha,\beta)}}{k+1}. \quad (4.3)$$

Observe that $\zeta_{q,w}^{(\alpha,\beta)}(s)$ interpolates $G_{k,q,w}^{(\alpha,\beta)}$ at nonnegative integers.

By using (2.14), one notes that

$$\left. \frac{d^{k+1}}{dt^{k+1}} F_{q,w}^{(\alpha,\beta)}(t, x) \right|_{t=0} = (k+1) [2]_{q^\beta} \sum_{m=0}^{\infty} (-1)^m q^{\beta m} w^m [x+m]_{q^\alpha}^k = G_{k+1,q,w}^{(\alpha,\beta)}(x), \quad (4.4)$$

$$\left(\frac{d}{dt} \right)^{k+1} \left(\sum_{n=0}^{\infty} G_{n,q,w}^{(\alpha,\beta)}(x) \frac{t^n}{n!} \right) \Big|_{t=0} = G_{k+1,q,w}^{(\alpha,\beta)}(x), \quad \text{for } k \in \mathbb{N}. \quad (4.5)$$

By (4.5), we are now ready to define the twisted Hurwitz q -Genocchi zeta functions.

Definition 4.3. Let $s \in \mathbb{C}$. One has

$$\zeta_{q,w}^{(\alpha,\beta)}(s, x) = [2]_{q^\beta} \sum_{n=0}^{\infty} \frac{(-1)^n q^{\beta n} w^n}{[x+n]_{q^\alpha}^s}. \quad (4.6)$$

Note that $\zeta_{q,w}^{(\alpha,\beta)}(s, x)$ is a meromorphic function on \mathbb{C} . Observe that if $q \rightarrow 1$, then $\lim_{q \rightarrow 1} \zeta_{q,w}^{(\alpha,\beta)}(s, x) = \zeta_w(s, x)$.

Theorem 4.4. Relation between $\zeta_{q,w}^{(\alpha,\beta)}(s, x)$ and $G_{k,q,w}^{(\alpha,\beta)}(x)$ is given by

$$\zeta_{q,w}^{(\alpha,\beta)}(-k, x) = \frac{G_{k+1,q,w}^{(\alpha,\beta)}(x)}{k+1}. \quad (4.7)$$

Observe that $\zeta_{q,w}^{(\alpha,\beta)}(-k, x)$ interpolates $G_{k,q,w}^{(\alpha,\beta)}(x)$ at nonnegative integers.

References

- [1] M. Cenkci, M. Can, and V. Kurt, " q -adic interpolation functions and Kummer-type congruences for q -twisted and q -generalized twisted Euler numbers," *Advanced Studies in Contemporary Mathematics*, vol. 9, no. 2, pp. 203–216, 2004.
- [2] M. Cenkci, M. Can, and V. Kurt, " q -extensions of Genocchi numbers," *Journal of the Korean Mathematical Society*, vol. 43, no. 1, pp. 183–198, 2006.
- [3] L.-C. Jang, "A study on the distribution of twisted q -Genocchi polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 18, no. 2, pp. 181–189, 2009.
- [4] L.-C. Jang, K.-W. Hwang, and Y.-H. Kim, "A note on (h, q) -Genocchi polynomials and numbers of higher order," *Advances in Difference Equations*, Article ID 309480, 6 pages, 2010.
- [5] T. Kim, "On the multiple q -Genocchi and Euler numbers," *Russian Journal of Mathematical Physics*, vol. 15, no. 4, pp. 481–486, 2008.
- [6] T. Kim, "On the q -extension of Euler and Genocchi numbers," *Journal of Mathematical Analysis and Applications*, vol. 326, no. 2, pp. 1458–1465, 2007.
- [7] T. Kim, " q -Volkenborn integration," *Russian Journal of Mathematical Physics*, vol. 9, no. 3, pp. 288–299, 2002.
- [8] T. Kim, " q -Euler numbers and polynomials associated with p -adic q -integrals," *Journal of Nonlinear Mathematical Physics*, vol. 14, no. 1, pp. 15–27, 2007.
- [9] T. Kim, J. Choi, Y.-H. Kim, and C. S. Ryoo, "Note on the weighted p -adic q -Euler measure on \mathbb{Z}_p ," *Advanced Studies in Contemporary Mathematics*, vol. 21, pp. 35–40, 2011.
- [10] B. A. Kupershmidt, "Reflection symmetries of q -Bernoulli polynomials," *Journal of Nonlinear Mathematical Physics*, vol. 12, supplement 1, pp. 412–422, 2005.
- [11] C. S. Ryoo, T. Kim, and L.-C. Jang, "Some relationships between the analogs of Euler numbers and polynomials," *Journal of Inequalities and Applications*, Article ID 86052, 22 pages, 2007.
- [12] C. S. Ryoo, "On the generalized Barnes type multiple q -Euler polynomials twisted by ramified roots of unity," *Proceedings of the Jangjeon Mathematical Society*, vol. 13, no. 2, pp. 255–263, 2010.
- [13] C. S. Ryoo, "A note on the weighted q -Euler numbers and polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 21, no. 1, pp. 47–54, 2011.
- [14] C. S. Ryoo, "Calculating zeros of the twisted Genocchi polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 17, no. 2, pp. 147–159, 2008.
- [15] Y. Simsek, V. Kurt, and D. Kim, "New approach to the complete sum of products of the twisted (h, q) -Bernoulli numbers and polynomials," *Journal of Nonlinear Mathematical Physics*, vol. 14, no. 1, pp. 44–56, 2007.
- [16] Y. Simsek, "Theorems on twisted L -function and twisted Bernoulli numbers," *Advanced Studies in Contemporary Mathematics*, vol. 12, pp. 237–246, 2006.

