

## Research Article

# LMI-Based Stability Criterion of Impulsive T-S Fuzzy Dynamic Equations via Fixed Point Theory

Ruofeng Rao<sup>1</sup> and Zhilin Pu<sup>1,2</sup>

<sup>1</sup> *Institution of Mathematics, Yibin University, Yibin, Sichuan 644007, China*

<sup>2</sup> *College of Mathematics and Software Science, Sichuan Normal University, Chengdu 610066, China*

Correspondence should be addressed to Ruofeng Rao; ruofengrao@163.com

Received 27 June 2013; Accepted 6 August 2013

Academic Editor: Chuanzhi Bai

Copyright © 2013 R. Rao and Z. Pu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

By formulating a contraction mapping and the matrix exponential function, the authors apply linear matrix inequality (LMI) technique to investigate and obtain the LMI-based stability criterion of a class of time-delay Takagi-Sugeno (T-S) fuzzy differential equations. To the best of our knowledge, it is the first time to obtain the LMI-based stability criterion derived by a fixed point theory. It is worth mentioning that LMI methods have high efficiency and other advantages in largescale engineering calculations. And the feasibility of LMI-based stability criterion can efficiently be computed and confirmed by computer Matlab LMI toolbox. At the end of this paper, a numerical example is presented to illustrate the effectiveness of the proposed methods.

## 1. Introduction

In this paper, we consider a class of delayed impulsive differential equations, which admits some biomathematics, physics, and engineering backgrounds, including the famous cellular neural networks proposed by Chua and Yang in 1988 [1, 2]. In practice, both time delays and impulse are unavoidable and may cause undesirable dynamic network behaviors such as oscillation and instability. So the stability analysis for delayed impulsive neural networks has become a topic of great theoretic and practical importance in recent years [3–25]. However, the research skill of the above literature is mainly based on Lyapunov theory. And there are many difficulties in applications of corresponding theory to the specific problems [26–32]. Recently, Burton and other authors have applied fixed point theory to investigate the stability of deterministic systems and obtained some more applicable results [11, 26–42]. For example, in [11], the authors used Leray-Schauders fixed point theorem to obtain the stability criteria of neural networks. Besides, the contraction-mapping theory is also an important fixed point theory in studying the stability of dynamics equations (see, e.g., [11, 33, 39–42]). On the other hand, fuzzy logic theory has shown to be an appealing and efficient approach to

dealing with the analysis and synthesis problems for complex nonlinear system [20–25]. In practice, the fuzzy model is far more important than stochastic model. Among various kinds of fuzzy methods, Takagi-Sugeno (T-S) fuzzy models provide a successful method to describe certain complex nonlinear systems using some local linear subsystems. To the best of our knowledge, few authors have used the fixed point theorem to study the stability of Takagi-Sugeno fuzzy differential equations with impulses. In addition, the LMI-based stability criterion of neural networks has never been investigated or obtained via any of the fixed point theories. Such a situation motivates our present study. Motivated by the above related literature [3–9, 11, 26–42], we will not only apply the fixed point theory to study the impulsive Takagi-Sugeno fuzzy dynamics equations but also try to obtain the LMI-based stability criterion by applying the contraction-mapping theory. To the best of our knowledge, it is the first time to obtain the LMI-based stability criterion derived by a fixed point theory. It is worth mentioning that LMI methods have high efficiency and other advantages in large-scale engineering calculations. And the feasibility of LMI-based stability criterion can efficiently be computed and confirmed by computer Matlab LMI toolbox. In the end of this paper, a numerical example is presented to illustrate the

effectiveness of the proposed methods. Finally, a conclusion is given in the final chapter.

## 2. Preliminaries

Let us consider the following delayed differential equations:

$$\begin{aligned} \frac{dx(t)}{dt} = & -Bx(t) + Cf(x(t)) \\ & + Dg(x(t - \tau(t))), \quad t \geq 0, t \neq t_k, \end{aligned} \quad (1)$$

equipped with the impulsive condition

$$x(t_k^+) - x(t_k) = \rho(x(t_k)), \quad k = 1, 2, \dots, \quad (2)$$

and the initial condition

$$x(\theta) = \phi(\theta), \quad \theta \in [-\tau, 0], \quad (3)$$

where  $x(t) = (x_1(t), x_2(t), \dots, x_n(t))^T \in R^n$ ,  $\phi(\theta) \in C[-\tau, 0, R^n]$ . Functions  $f(x) = (f_1(x_1(t)), f_2(x_2(t)), \dots, f_n(x_n(t)))^T \in R^n$ ,  $g(x(t - \tau(t))) = (g_1(x_1(t - \tau(t))), g_2(x_2(t - \tau(t))), \dots, g_n(x_n(t - \tau(t))))^T \in R^n$ ,  $\rho(x(t)) = (\rho_1(x_1(t)), \rho_2(x_2(t)), \dots, \rho_n(x_n(t)))^T \in R^n$ , and time delays  $0 \leq \tau(t) \leq \tau$  for all  $i = 1, 2, \dots, n$ . The fixed impulsive moments  $t_k$  ( $k = 1, 2, \dots$ ) satisfy  $0 < t_1 < t_2 < \dots$  with  $\lim_{k \rightarrow \infty} t_k = \infty$ .  $x(t_k^+)$ , and  $x(t_k^-)$  stand for the right-hand and left-hand limits of  $x(t)$  at time  $t_k$ , respectively. We always assume  $x(t_k^-) = x(t_k)$ , for all  $k = 1, 2, \dots$ . Similarly as in [33], we assume in this paper that  $f(0) = g(0) = \rho(0) = 0 \in R^n$ .

Constant matrix  $B = \text{diag}(b_1, b_2, \dots, b_n)$  is a positive definite diagonal matrix, and both  $C = (c_{ij})_{n \times n}$  and  $D = (d_{ij})_{n \times n}$  are matrices with  $n \times n$  dimension. It is well known that the above equation admits its practical implications. For example, it can serve as a model of impulsive cellular neural networks with time-varying delays. The parameter  $c_{ij}$  denotes the connection weight of the  $j$ th neuron on the  $i$ th neuron at time  $t$ . And the parameter  $d_{ij}$  represents the connection strength of the  $j$ th neuron on the  $i$ th neuron at time  $t - \tau(t)$ . The constant  $b_i$  represents the rate with which the  $i$ th neuron will reset its potential to the resting state when disconnected from the network and external inputs.  $f_j(x_j(t))$  is the activation function of the  $j$ th neuron at time  $t$ , and  $g_j(x_j(t - \tau(t)))$  represents the activation function of the  $j$ th neuron at time  $t - \tau(t)$ .

Below, we describe the T-S fuzzy mathematical model with time delay as follows.

Fuzzy Rule  $j$ :

IF  $\omega_1(t)$  is  $\mu_{j1}$  and  $\dots \omega_s(t)$  is  $\mu_{js}$ , THEN

$$\begin{aligned} \frac{dx(t)}{dt} = & -Bx(t) + C_j f(x(t)) + D_j g(x(t - \tau(t))), \\ & t \geq 0, t \neq t_k, k = 1, 2, \dots \end{aligned} \quad (4)$$

$$x(t_k^+) - x(t_k) = \rho(x(t_k)), \quad k = 1, 2, \dots$$

$$x(\theta) = \phi(\theta), \quad \theta \in [-\tau, 0],$$

where  $\omega_k(t)$  ( $k = 1, 2, \dots, s$ ) is the premise variable,  $\mu_{jk}$  ( $j = 1, 2, \dots, r; k = 1, 2, \dots, s$ ) is the fuzzy set that is characterized

by membership function,  $r$  is the number of the IF-THEN rules, and  $s$  is the number of the premise variables. By the way of a standard fuzzy inference method, the system (4) is inferred as follows:

$$\begin{aligned} \frac{dx(t)}{dt} = & -Bx(t) + \sum_{j=1}^r h_j(\omega(t)) [C_j f(x(t)) + D_j g(x(t - \tau(t)))], \\ & t \geq 0, t \neq t_k, k = 1, 2, \dots, \\ x(t_k^+) - x(t_k) = & \rho(x(t_k)), \quad k = 1, 2, \dots, \\ x(\theta) = & \phi(\theta), \quad \theta \in [-\tau, 0], \end{aligned} \quad (5)$$

where  $\omega(t) = [\omega_1(t), \omega_2(t), \dots, \omega_s(t)]$ ,  $h_j(\omega(t)) = \omega_j / \sum_{k=1}^r \omega_k(\omega(t))$ ,  $\omega_j(\omega(t)) : R^s \rightarrow [0, 1]$  ( $j = 1, 2, \dots, r$ ) is the membership function of the system with respect to the fuzzy rule  $j$ .  $h_j$  can be regarded as the normalized weight of each IF-THEN rule, satisfying  $h_j(\omega(t)) \geq 0$  and  $\sum_{j=1}^r h_j(\omega(t)) = 1$ .

For convenience's sake, we introduce the following standard notations similarly as [10, (iii)-(X)]:

$$Q = (q_{ij})_{n \times n} > 0 \quad (< 0),$$

$$Q = (q_{ij})_{n \times n} \geq 0 \quad (\leq 0),$$

$$Q_1 \geq Q_2 \quad (Q_1 \leq Q_2),$$

$$Q_1 > Q_2 \quad (Q_1 < Q_2),$$

$$\lambda_{\min} \Phi, \text{ the identity matrix } I \text{ and the symmetric terms } *.$$

(6)

In addition, we denote  $|C| = (|c_{ij}|)_{n \times n}$  for any matrix  $C = (c_{ij})_{n \times n}$  and  $|v| = (|v_1|, |v_2|, \dots, |v_n|)^T$  for any  $v = (v_1, v_2, \dots, v_n)^T \in R^n$ . Denote the finite set  $\mathcal{N} = \{1, 2, \dots, n\}$ .

Throughout this paper, we assume

- (A1)  $f_j$  is locally Lipschitz continuous, and there exists a positive constant  $F_j > 0$  such that  $|f_j'(r)| \leq F_j$  for all  $r \in R$  at which  $f_j$  is differentiable;
- (A2)  $g_j$  is locally Lipschitz continuous, and there exists a positive constant  $G_j > 0$  such that  $|g_j'(r)| \leq G_j$  for all  $r \in R$  at which  $g_j$  is differentiable;
- (A3)  $\rho_j$  is locally Lipschitz continuous, and there exists a positive constant  $H_j > 0$  such that  $|\rho_j'(r)| \leq H_j$  for all  $r \in R$  at which  $\rho_j$  is differentiable.

**Lemma 1** (see [25]). Let  $f : R^n \rightarrow R^n$  be locally Lipschitz continuous. For any given  $x, y \in R^n$ , there exists an element  $\mathfrak{w}$  in the union  $\cup_{z \in [x, y]} \partial f(z)$  such that

$$f(y) - f(x) = \mathfrak{w}(y - x), \quad (7)$$

where  $[x, y]$  denotes the segment connecting  $x$  and  $y$ .

**Remark 2.** From Lemma 1, (A1)–(A3), and [10, equation (27)], we can similarly derive

$$\begin{aligned} |f(x) - f(y)| &\leq F|x - y|, \quad x, y \in R^n, \\ |g(x) - g(y)| &\leq G|x - y|, \quad x, y \in R^n, \\ |\rho(x) - \rho(y)| &\leq H|x - y|, \quad x, y \in R^n, \end{aligned} \quad (8)$$

where matrices  $F = \text{diag}(F_1, F_2, \dots, F_n)$ ,  $G = \text{diag}(G_1, G_2, \dots, G_n)$ , and  $H = \text{diag}(H_1, H_2, \dots, H_n)$ .

**Remark 3.** In many previous literature,  $f_i, g_i$  ( $i \in \mathcal{N}$ ) are always assumed to be globally Lipschitz continuous. However, the most common function  $f_i(r) = r^2$  is not globally Lipschitz continuous in  $R^1$ . Note that we extend the functions  $f, g$  from global Lipschitz continuous functions to locally Lipschitz continuous functions. Obviously,  $f_i(r) = r^2$  is a local Lipschitz continuous function.

Similarly as in [11, Definition 2.2], the exponential stability is defined as follows.

**Definition 4.** Dynamic equation (5) is said to be exponentially stable if, for any initial condition  $\phi(s) \in C[-\tau, 0], R^n$ , there exists a pair of positive constants  $a$  and  $b$  such that

$$\|x(t; s, \phi)\| \leq be^{-at}, \quad \forall t > 0, \quad (9)$$

where the norm  $\|x(t)\| = (\sum_{i=1}^n |x_i(t)|^2)^{1/2}$ .

### 3. Main Result

Before giving the main result of this paper, we need to define the matrix exponential function as follows.

**Definition 5.** For a diagonal constants matrix  $B = \text{diag}(b_1, b_2, \dots, b_n)$ , we denote the matrix exponential function  $e^{Bt} = \text{diag}(e^{b_1 t}, e^{b_2 t}, \dots, e^{b_n t})$  for all  $t \in R$ .

From the above definition of the matrix exponential function, we are not difficult to obtain the following lemma.

**Lemma 6.** Let  $B$  be a diagonal constants matrix, and let  $e^{Bt}$  be the matrix exponential function of  $B$ . Then, we have

$$\begin{aligned} (1) \quad (d/dt)e^{Bt} &= Be^{Bt}, \quad t \in R, \\ (2) \quad (d/dt)(e^{Bt}\eta) &= Be^{Bt}\eta, \quad t \in R, \end{aligned}$$

where  $\eta = (\eta_1, \eta_2, \dots, \eta_n)^T \in R^n$ , and each  $\eta_i \in R$  ( $i = 1, 2, \dots, n$ ) is a constant.

In addition, we need to define the rule on vectors in  $R^n$  as follows.

**Definition 7.**  $v \leq w$  if  $v_i - w_i \leq 0$  for all  $i \in \mathcal{N}$ , where  $v = (v_1, v_2, \dots, v_n)^T \in R^n$ ,  $w = (w_1, w_2, \dots, w_n)^T \in R^n$ .

Now, we present the main result of this paper as follows.

**Theorem 8.** Assume that there exists a positive constant  $\delta$  such that  $\inf_{k=1,2,\dots} (t_{k+1} - t_k) \geq \delta$ . In addition, there exists a constant  $0 < \lambda < 1$  such that

$$B^{-1} \sum_{j=1}^r (|C_j|F + |D_j|G) + \frac{1}{\delta} B^{-1}H + H < \lambda I, \quad (10)$$

where  $B^{-1}$  is the inverse matrix of  $B$ . Then, the impulsive fuzzy dynamic equation (5) is exponentially stable in the mean square.

*Proof.* To apply the fixed point theory, we firstly define a complete metric space  $\Omega$  as follows.

Let  $\Omega$  be the space consisting of functions  $q(t) : [-\tau, \infty) \rightarrow R^n$ , satisfying that

- (a)  $q(t)$  is continuous on  $t \neq t_k$  ( $k = 1, 2, \dots$ ),
- (b)  $\lim_{t \rightarrow t_k^-} q(t)$  and  $\lim_{t \rightarrow t_k^+} q(t)$  exist, and  $q(t_k^-) = q(t_k)$  for all  $k = 1, 2, \dots$ ,
- (c)  $q(t) = \phi(t)$  for  $t \in [-\tau, 0]$ ,
- (d)  $e^{\beta t} q(t) \rightarrow 0 \in R^n$  as  $t \rightarrow \infty$ , where  $\beta > 0$  is a positive constant, satisfying  $\beta < \lambda_{\min} B$ .

It is not difficult to verify that the above-mentioned space  $\Omega$  is a complete metric space if it is equipped with the following metric:

$$\text{dist}(\bar{q}, \tilde{q}) = \max_{i=1,2,\dots,n} \left( \sup_{t \geq -\tau} |\bar{q}_i(t) - \tilde{q}_i(t)| \right), \quad (11)$$

where  $\bar{q} = \bar{q}(t) = (\bar{q}_1(t), \bar{q}_2(t), \dots, \bar{q}_n(t))^T \in \Omega$ , and  $\tilde{q} = \tilde{q}(t) = (\tilde{q}_1(t), \tilde{q}_2(t), \dots, \tilde{q}_n(t))^T \in \Omega$ .

**Remark 9.** Here, we consider the above-defined metric, which is different from those of some previous related literature (see, e.g., [33]) so that the LMI-based stability criterion in this paper may be obtained expediently.

Next, we formulate and define a contraction mapping  $P : \Omega \rightarrow \Omega$ , which may be divided into three steps.

**Step 1.** Formulating the mapping.

Let  $x(t)$  be a solution of the fuzzy equation (5).

Then, for  $t \geq 0$ ,  $t \neq t_k$ , we have

$$\begin{aligned} \frac{dx(t)}{dt} (e^{Bt} x(t)) &= Be^{Bt} x(t) + e^{Bt} \frac{dx(t)}{dt} \\ &= e^{Bt} \sum_{j=1}^r h_j(\omega(t)) [C_j f(x(t)) \\ &\quad + D_j g(x(t - \tau(t)))] . \end{aligned} \quad (12)$$

Further, we get by the integral nature

$$\begin{aligned} x(t) &= e^{-Bt} \left\{ \int_0^t e^{Bs} \sum_{j=1}^r h_j(\omega(s)) [C_j f(x(s)) \right. \\ &\quad \left. + D_j g(x(s - \tau(s)))] ds \right. \\ &\quad \left. + \eta \right\}, \quad t \geq 0, \end{aligned} \quad (13)$$

where  $\eta \in R^n$  is the vector to be determined.

From (2) and (3) or  $x(0) = \phi(0)$ , it is not difficult to conclude  $\eta = \phi(0) + \sum_{0 < t_i < t} e^{Bt_i} \rho(x(t_i))$ . And, hence,

$$\begin{aligned} x(t) &= e^{-Bt} \left\{ \int_0^t e^{Bs} \sum_{j=1}^r h_j(\omega(s)) \right. \\ &\quad \times [C_j f(x(s)) + D_j g(x(s - \tau(s)))] ds \\ &\quad \left. + \phi(0) + \sum_{0 < t_i < t} e^{Bt_i} \rho(x(t_i)) \right\}, \quad t \geq 0. \end{aligned} \quad (14)$$

On the other hand, it follows from (14) that

$$\begin{aligned} x(t_k) &= e^{-Bt_k} \left\{ \int_0^{t_k} e^{Bs} \sum_{j=1}^r h_j(\omega(s)) \right. \\ &\quad \times [C_j f(x(s)) \\ &\quad \left. + D_j g(x(s - \tau(s)))] ds \right. \\ &\quad \left. + \phi(0) + \sum_{0 < t_i < t_k} e^{Bt_i} \rho(x(t_i)) \right\}, \\ x(t_{k+\varepsilon}) &= e^{-Bt_{k+\varepsilon}} \left\{ \int_0^{t_{k+\varepsilon}} e^{Bs} \sum_{j=1}^r h_j(\omega(s)) \right. \\ &\quad \times [C_j f(x(s)) \\ &\quad \left. + D_j g(x(s - \tau(s)))] ds \right. \\ &\quad \left. + \phi(0) + \sum_{0 < t_i < t_{k+\varepsilon}} e^{Bt_i} \rho(x(t_i)) \right\}. \end{aligned} \quad (15)$$

Let  $\varepsilon \rightarrow 0^+$ ; then we can derive from the two equations above that

$$\lim_{\varepsilon \rightarrow 0^+} x(t_{k+\varepsilon}) - x(t_k) = x(t_k^+) - x(t_k) = \rho(x(t_k)). \quad (16)$$

So, we may define the mapping  $P$  on the space  $\Omega$  as follows:

$$\begin{aligned} Px(t) &= e^{-Bt} \phi(0) + e^{-Bt} \int_0^t e^{Bs} \sum_{j=1}^r h_j(\omega(s)) \\ &\quad \times [C_j f(x(s)) \\ &\quad + D_j g(x(s - \tau(s)))] ds \\ &\quad + e^{-Bt} \sum_{0 < t_i < t} e^{Bt_i} \rho(x(t_i)), \end{aligned} \quad (17)$$

on  $t \geq 0$ , and  $Px(s) = \phi(s)$  on  $s \in [-\tau, 0]$ .

*Step 2.* We claim that  $Px(t) \in \Omega$  for any  $x(t) \in \Omega$ . That is,  $Px(t)$  satisfies the conditions (a)–(d) of  $\Omega$ .

Indeed, since  $Px(s) = \phi(s)$  on  $s \in [-\tau, 0]$ , the condition (c) is satisfied. It is obvious from (17) that  $Px(t)$  is continuous on  $t \neq t_k$  and  $t \geq 0$ . And then the condition (a) is satisfied.

Next, we verify the condition (b).

Indeed, for any given  $t_k$ , we can get from (17)

$$Px(t_k + \varepsilon) - Px(t_k) = \pi_1 + \pi_2 + \pi_3, \quad (18)$$

where

$$\begin{aligned} \pi_1 &= e^{-B(t_k+\varepsilon)} \phi(0) - e^{-B(t_k)} \phi(0), \\ \pi_2 &= e^{-B(t_k+\varepsilon)} \int_0^{t_k+\varepsilon} e^{Bs} \sum_{j=1}^r h_j(\omega(s)) \\ &\quad \times [C_j f(x(s)) \\ &\quad + D_j g(x(s - \tau(s)))] ds \\ &\quad - e^{-Bt_k} \int_0^{t_k} e^{Bs} \sum_{j=1}^r h_j(\omega(s)) \\ &\quad \times [C_j f(x(s)) \\ &\quad + D_j g(x(s - \tau(s)))] ds \\ \pi_3 &= e^{-B(t_k+\varepsilon)} \sum_{0 < t_i < t_k+\varepsilon} e^{Bt_i} \rho(x(t_i)) \\ &\quad - e^{-Bt_k} \sum_{0 < t_i < t_k} e^{Bt_i} \rho(x(t_i)). \end{aligned} \quad (19)$$

Obviously,  $\pi_1 \rightarrow 0$  and  $\pi_2 \rightarrow 0$  as  $\varepsilon \rightarrow 0$ . In addition, letting  $\varepsilon \rightarrow 0^+$ , we have  $\pi_3 \rightarrow 0$ . Letting  $\varepsilon \rightarrow 0^+$ , we get by (18)

$$Px(t_k^+) - Px(t_k) = \lim_{\varepsilon \rightarrow 0^+} \pi_3 = \rho(x(t_k)). \quad (20)$$

So the condition (b) is satisfied.

Finally, it is followed from (17) that

$$\begin{aligned} e^{\beta t} P x(t) &= e^{\beta t} e^{-Bt} \phi(0) \\ &+ e^{\beta t} e^{-Bt} \int_0^t e^{Bs} \sum_{j=1}^r h_j(\omega(s)) \\ &\quad \times [C_j f(x(s)) \\ &\quad + D_j g(x(s - \tau(s)))] ds \\ &+ e^{\beta t} e^{-Bt} \sum_{0 < t_i < t} e^{Bt_i} \rho(x(t_i)). \end{aligned} \quad (21)$$

Obviously,  $e^{\beta t} e^{-Bt} \phi(0) \rightarrow 0 \in R^n$  as  $t \rightarrow \infty$ .

On the other hand, it follows from  $e^{\beta t} x(t) \rightarrow 0$  that, for any given  $\varepsilon > 0$ , there exists a corresponding constant  $t_*$  such that  $|e^{\beta t} x(t)| < \varepsilon u$  for all  $t \geq t_*$ , where

$$u = (1, 1, \dots, 1)^T \in R^n. \quad (22)$$

In addition, (A3) derives

$$\begin{aligned} \left| e^{\beta t} e^{-Bt} \sum_{0 < t_i < t} e^{Bt_i} \rho(x(t_i)) \right| &\leq e^{\beta t} e^{-Bt} \sum_{0 < t_i < t} e^{Bt_i} H |x(t_i)| \\ &\leq e^{\beta t} e^{-Bt} \left( \sum_{0 < t_i \leq t_*} e^{Bt_i} H |x(t_i)| + \sum_{t_* < t_i < t} e^{Bt_i} H |x(t_i)| \right). \end{aligned} \quad (23)$$

Obviously,

$$\begin{aligned} e^{\beta t} e^{-Bt} \sum_{0 < t_i \leq t_*} e^{Bt_i} H |x(t_i)| \\ = e^{(\beta I - B)t} \sum_{0 < t_i \leq t_*} e^{Bt_i} H |x(t_i)| \rightarrow 0 \in R^n, \quad t \rightarrow \infty. \end{aligned} \quad (24)$$

Besides,

$$\begin{aligned} e^{\beta t} e^{-Bt} \sum_{t_* < t_i < t} e^{Bt_i} H |x(t_i)| \\ = e^{(\beta I - B)t} \sum_{t_* < t_i < t} e^{(B - \beta I)t_i} H (e^{\beta t_i} |x(t_i)|) \\ \leq \varepsilon e^{(\beta I - B)t} \sum_{t_* < t_i < t} e^{(B - \beta I)t_i} H u \\ = \begin{pmatrix} \varepsilon e^{(\beta - b_1)t} \sum_{t_* < t_i < t} e^{(b_1 - \beta)t_i} H_1 \\ \varepsilon e^{(\beta - b_2)t} \sum_{t_* < t_i < t} e^{(b_2 - \beta)t_i} H_2 \\ \vdots \\ \varepsilon e^{(\beta - b_n)t} \sum_{t_* < t_i < t} e^{(b_n - \beta)t_i} H_n \end{pmatrix}. \end{aligned} \quad (25)$$

To prove  $e^{\beta t} e^{-Bt} \sum_{t_* < t_i < t} e^{Bt_i} H |x(t_i)| \rightarrow 0 \in R^n$ , we only need to prove

$$\varepsilon e^{(\beta - b_1)t} \sum_{t_* < t_i < t} e^{(b_1 - \beta)t_i} H_1 \rightarrow 0, \quad t \rightarrow \infty, \quad (26)$$

for the other cases can be similarly proved.

Indeed, we may as well assume  $t_k < t \leq t_{k+1}$  and  $t_{m-1} < t_* \leq t_m$ . Then, we have

$$\begin{aligned} \varepsilon e^{(\beta - b_1)t} \sum_{t_* < t_i < t} e^{(b_1 - \beta)t_i} H_1 \\ = \frac{\varepsilon}{\delta} H_1 e^{(\beta - b_1)t} \sum_{t_m \leq t_i \leq t_k} \delta e^{(b_1 - \beta)t_i} \\ \leq \frac{\varepsilon}{\delta} H_1 e^{(\beta - b_1)t} \left( \delta e^{(b_1 - \beta)t_k} + \sum_{t_m \leq t_i \leq t_{k-1}} (t_{i+1} - t_i) e^{(b_1 - \beta)t_i} \right) \\ \leq \frac{\varepsilon}{\delta} H_1 e^{(\beta - b_1)t} \left( \delta e^{(b_1 - \beta)t_k} + \int_{t_*}^t e^{(b_1 - \beta)s} ds \right) \\ \leq \varepsilon H_1 e^{-(b_1 - \beta)(t - t_k)} + \frac{H_1}{\delta(b_1 - \beta)} \varepsilon, \end{aligned} \quad (27)$$

which implies that (26) holds. And, hence,  $e^{\beta t} e^{-Bt} \sum_{t_* < t_i < t} e^{Bt_i} H |x(t_i)| \rightarrow 0 \in R^n$ ,  $t \rightarrow \infty$ .

Below, we only need to prove

$$\begin{aligned} e^{\beta t} e^{-Bt} \int_0^t e^{Bs} \sum_{j=1}^r h_j(\omega(s)) \\ \times [C_j f(x(s)) \\ + D_j g(x(s - \tau(s)))] ds \rightarrow 0, \end{aligned} \quad (28)$$

$t \rightarrow \infty$ .

In fact, it follows from  $e^{\beta t} x(t) \rightarrow 0$  that, for any given  $\varepsilon > 0$ , there exists a corresponding constant  $t^*$  such that  $|e^{\beta t} x(t)| < \varepsilon u$  for all  $t \geq t^*$ . Then, we get by (A1)

$$\begin{aligned} \left| e^{\beta t} e^{-Bt} \int_0^t e^{Bs} \sum_{j=1}^r h_j(\omega(s)) C_j f(x(s)) ds \right| \\ \leq e^{-(B - \beta I)t} \int_0^{t^*} e^{Bs} \sum_{j=1}^r |C_j| F |x(s)| ds \\ + e^{-(B - \beta I)t} \int_{t^*}^t e^{Bs} \sum_{j=1}^r |C_j| F |x(s)| ds. \end{aligned} \quad (29)$$

On the one hand,

$$\begin{aligned}
 & e^{-(B-\beta I)t} \int_0^{t^*} e^{Bs} \sum_{j=1}^r |C_j| F|x(s)| ds \\
 & \leq e^{-(B-\beta I)t} \int_0^{t^*} e^{Bs} \sum_{j=1}^r |C_j| F \left( \sum_{i \in \mathcal{N}} \sup_{s \in [0, t^*]} |x_i(s)| \right) u ds \\
 & \leq \left( \sum_{i=1}^m w_i \right) e^{-(B-\beta I)t} \int_0^{t^*} e^{Bs} u ds \longrightarrow 0 \in R^n, \quad t \longrightarrow \infty,
 \end{aligned} \tag{30}$$

where  $w = \sum_{j=1}^r |C_j| F(\sum_{i \in \mathcal{N}} \sup_{s \in [0, t^*]} |x_i(s)|)u = (w_1, w_2, \dots, w_n)^T \in R^n$ .

On the other hand,

$$\begin{aligned}
 & e^{-(B-\beta I)t} \int_{t^*}^t e^{Bs} \sum_{j=1}^r |C_j| F|x(s)| ds \\
 & \leq \varepsilon e^{-(B-\beta I)t} \int_{t^*}^t e^{(B-\beta I)s} w^* ds \\
 & \leq \varepsilon \left( \sum_{i=1}^n w_i^* \right) \left( \frac{1}{b_1 - \beta}, \frac{1}{b_2 - \beta}, \dots, \frac{1}{b_n - \beta} \right)^T,
 \end{aligned} \tag{31}$$

where  $w^* = \sum_{j=1}^r |C_j| Fu = (w_1^*, w_2^*, \dots, w_n^*)^T \in R^n$ .

Now we can conclude from (29)–(31) that

$$\begin{aligned}
 & e^{\beta t} e^{-Bt} \int_0^t e^{Bs} \sum_{j=1}^r h_j(\omega(s)) \\
 & \quad \times [C_j f(x(s))] ds \longrightarrow 0 \in R^n, \quad t \longrightarrow \infty.
 \end{aligned} \tag{32}$$

Similarly, we can prove

$$\begin{aligned}
 & e^{\beta t} e^{-Bt} \int_0^t e^{Bs} \sum_{j=1}^r h_j(\omega(s)) \\
 & \quad \times [D_j g(x(s - \tau(s)))] ds \longrightarrow 0 \in R^n, \quad t \longrightarrow \infty.
 \end{aligned} \tag{33}$$

Indeed, we can similarly define the corresponding constant  $T_*$  for any given  $\varepsilon > 0$ , satisfying  $|e^{\beta t} x(t)| < \varepsilon u$  for all  $t \geq T_*$ . Then, we have

$$\begin{aligned}
 & \left| e^{\beta t} e^{-Bt} \int_0^t e^{Bs} \sum_{j=1}^r h_j(\omega(s)) [D_j g(x(s - \tau(s)))] ds \right| \\
 & \leq e^{\beta t} e^{-Bt} \left( \int_0^{T_*} e^{Bs} \sum_{j=1}^r |D_j| G|x(s - \tau(s))| ds \right. \\
 & \quad \left. + \int_{T_*}^t e^{Bs} \sum_{j=1}^r |D_j| G|x(s - \tau(s))| ds \right).
 \end{aligned} \tag{34}$$

Similarly, we can prove

$$\begin{aligned}
 & e^{\beta t} e^{-Bt} \int_0^{T_*} e^{Bs} \sum_{j=1}^r |D_j| G|x(s - \tau(s))| ds \\
 & \leq \left( \sum_{i=1}^m \bar{w}_i \right) e^{-(B-\beta I)t} \int_0^{T_*} e^{Bs} u ds \longrightarrow 0 \in R^n, \quad t \longrightarrow \infty,
 \end{aligned} \tag{35}$$

where  $\bar{w} = \sum_{j=1}^r |D_j| G(\sum_{i \in \mathcal{N}} \sup_{s \in [-\tau, T_*]} |x_i(s)|)u = (\bar{w}_1, \bar{w}_2, \dots, \bar{w}_n)^T \in R^n$ .

On the other hand,

$$\begin{aligned}
 & e^{\beta t} e^{-Bt} \int_{T_*}^t e^{Bs} \sum_{j=1}^r |D_j| G|x(s - \tau(s))| ds \\
 & \leq \varepsilon e^{\beta \tau} e^{-(B-\beta I)t} \int_{T_*}^t e^{(B-\beta I)s} \bar{w}^* ds \\
 & \leq \varepsilon \left( \sum_{i \in \mathcal{N}} \bar{w}_i^* \right) e^{\beta \tau} \left( \frac{1}{b_1 - \beta}, \frac{1}{b_2 - \beta}, \dots, \frac{1}{b_n - \beta} \right)^T,
 \end{aligned} \tag{36}$$

where  $\bar{w}^* = \sum_{j=1}^r |D_j| Gu = (\bar{w}_1^*, \bar{w}_2^*, \dots, \bar{w}_n^*)^T \in R^n$ .

From (34)–(36), we can conclude that (33) holds. Hence, the condition (d) is satisfied.

*Step 3.* Below, we only need to prove that  $P$  is a contraction mapping.

Indeed, for any  $x = x(t) = (x_1(t), x_2(t), \dots, x_n(t))^T$ ,  $y = y(t) = (y_1(t), y_2(t), \dots, y_n(t))^T \in \Omega$ , we estimate  $|Px(t) - Py(t)| \leq K_1 + K_2 + K_3$ , where

$$\begin{aligned}
 K_1 &= e^{-Bt} \int_0^t e^{Bs} \sum_{j=1}^r |C_j| |f(x(s)) - f(y(s))| ds \\
 &\leq e^{-Bt} \int_0^t e^{Bs} \sum_{j=1}^r |C_j| F|x(s) - y(s)| ds, \\
 K_2 &= e^{-Bt} \int_0^t e^{Bs} \sum_{j=1}^r h_j(\omega(s)) |D_j| \\
 &\quad \times |g(x(s - \tau(s))) - g(y(s - \tau(s)))| ds, \\
 K_3 &= e^{-Bt} \sum_{0 < t_i < t} e^{Bt_i} |\rho(x(t_i)) - \rho(y(t_i))|.
 \end{aligned} \tag{37}$$



From mathematical analysis and computation, we can derive

$$\begin{aligned} K_1 &= e^{-Bt} \int_0^t e^{Bs} \sum_{j=1}^r |C_j| |f(x(s)) - f(y(s))| ds \\ &\leq e^{-Bt} \int_0^t e^{Bs} \sum_{j=1}^r |C_j| F |x(s) - y(s)| ds \\ &\leq \text{dist}(x, y) e^{-Bt} \int_0^t e^{Bs} \sum_{j=1}^r |C_j| F ds \\ &\leq \left( B^{-1} \sum_{j=1}^r |C_j| F \right) \text{dist}(x, y) u. \end{aligned} \quad (38)$$

Similarly, we have

$$\begin{aligned} K_2 &= e^{-Bt} \int_0^t e^{Bs} \sum_{j=1}^r h_j(\omega(s)) |D_j| |g(x(s - \tau(s))) \\ &\quad - g(y(s - \tau(s)))| ds \\ &\leq \left( B^{-1} \sum_{j=1}^r |D_j| G \right) \text{dist}(x, y) u \\ K_3 &\leq e^{-Bt} \sum_{0 < t_i < t} e^{Bt_i} H |(x(t_i)) - (y(t_i))| \\ &\leq \frac{1}{\delta} \text{dist}(x, y) e^{-Bt} \left( \sum_{t_1 \leq t_i \leq t_{k-1}} e^{Bt_i} (t_{i+1} - t_i) Hu \right. \\ &\quad \left. + e^{Bt_k} \delta Hu \right) \\ &\leq \frac{1}{\delta} \text{dist}(x, y) e^{-Bt} \left( \int_0^t e^{Bs} Hu ds + e^{Bt} \delta Hu \right) \\ &\leq \text{dist}(x, y) \left( \frac{1}{\delta} B^{-1} H + H \right) u. \end{aligned} \quad (39)$$

Combining the above three inequalities results in

$$\begin{aligned} &|Px(t) - Py(t)| \\ &\leq \left( B^{-1} \sum_{j=1}^r (|C_j| F + |D_j| G) + \frac{1}{\delta} B^{-1} H + H \right) u \text{dist}(x, y) \\ &< \lambda u \text{dist}(x, y), \end{aligned} \quad (40)$$

and hence

$$\text{dist}(P(x), P(y)) \leq \lambda \text{dist}(x, y). \quad (41)$$

Therefore,  $P : \Omega \rightarrow \Omega$  is a contraction mapping such that there exists the fixed point  $x(t)$  of  $P$  in  $\Omega$ , which implies that  $x(t)$  is the solution for the the impulsive fuzzy dynamic equation (5), satisfying  $e^{\beta t} \|x(t)\| \rightarrow 0$  as  $t \rightarrow \infty$ . So the proof is completed.  $\square$

## 4. Numerical Example

*Example 1.* Consider the T-S fuzzy impulsive dynamic equations as follows.

Fuzzy Rule 1:

IF  $\omega_1(t)$  is  $1/e^{-2\omega_1(t)}$ , THEN

$$\begin{aligned} \frac{dx(t)}{dt} &= -Bx(t) + C_1 f(x(t)) + D_1 g(x(t - \tau(t))), \\ t &\geq 0, \quad t \neq t_k, \quad k = 1, 2, \dots \end{aligned} \quad (42)$$

$$x(t_k^+) - x(t_k) = \rho(x(t_k)), \quad k = 1, 2, \dots$$

$$x(\theta) = \phi(\theta), \quad \theta \in [-\tau, 0],$$

Fuzzy Rule 2:

IF  $\omega_2(t)$  is  $1 - 1/e^{-2\omega_1(t)}$ , THEN

$$\begin{aligned} \frac{dx(t)}{dt} &= -Bx(t) + C_2 f(x(t)) + D_2 g(x(t - \tau(t))), \\ t &\geq 0, \quad t \neq t_k, \quad k = 1, 2, \dots \end{aligned} \quad (43)$$

$$x(t_k^+) - x(t_k) = \rho(x(t_k)), \quad k = 1, 2, \dots$$

$$x(\theta) = \phi(\theta), \quad \theta \in [-\tau, 0],$$

where

$$\begin{aligned} B &= \begin{pmatrix} 2 & 0 \\ 0 & 1.9 \end{pmatrix}, \quad C_1 = \begin{pmatrix} -0.2 & 0 \\ 0 & 0.3 \end{pmatrix} = D_1, \\ C_2 &= \begin{pmatrix} 0.3 & 0 \\ 0 & -0.1 \end{pmatrix} = D_2, \\ F &= \begin{pmatrix} 0.1 & 0 \\ 0 & 0.2 \end{pmatrix} = G, \quad H = \begin{pmatrix} 0.7 & 0 \\ 0 & 0.6 \end{pmatrix}. \end{aligned} \quad (44)$$

Let  $\delta = 1.5$ . Then we can use Matlab LMI toolbox to solve the LMI condition (10) and obtain  $t_{\min} = -0.0046 < 0$  which implies it is feasible (see [10, Remark 29(3)] for details). Further, extracting the datum shows

$$\lambda = 0.9883, \quad (45)$$

which means  $0 < \lambda < 1$ . Thereby, we can conclude from Theorem 8 that this impulsive fuzzy dynamic equation is exponentially stable in the mean square.

## 5. Conclusion

By formulating a contraction mapping and the matrix exponential function, the author applies linear matrix inequality (LMI) technique to investigate and obtain the LMI-based stability criterion of a class of time-delay Takagi-Sugeno (T-S) fuzzy differential equations. It is the first time to obtain the LMI-based stability criterion derived by a fixed point theory. The LMI methods have high efficiency and other advantages in large-scale engineering calculations. And the feasibility of LMI-based stability criterion can efficiently be computed and confirmed by computer Matlab LMI toolbox. A numerical

example is presented to illustrate the effectiveness of the proposed methods. In the end of this paper, we have to point out that there are still many difficulties in obtaining the LMI-based stability criteria for some other dynamics equations, such as Cohen-Grossberg neural networks (see, e.g., [11, 19]) and other neural networks. These problems remain open and challenging.

## Acknowledgments

This work was supported by the Scientific Research Fund of Science Technology Department of Sichuan Province (2010JY0057, 2012JYZ010) and the Scientific Research Fund of Sichuan Provincial Education Department (12ZB349).

## References

- [1] L. O. Chua and L. Yang, "Cellular neural networks: theory," *IEEE Transactions on Circuits and Systems*, vol. 35, no. 10, pp. 1257–1272, 1988.
- [2] L. O. Chua and L. Yang, "Cellular neural networks: applications," *IEEE Transactions on Circuits and Systems*, vol. 35, no. 10, pp. 1273–1290, 1988.
- [3] D. Xu, X. Wang, and Z. Yang, "Further results on existence-uniqueness for stochastic functional differential equations," *Science China Mathematics*, vol. 56, no. 6, pp. 1169–1180, 2013.
- [4] D. Xu and S. Long, "Attracting and quasi-invariant sets of non-autonomous neural networks with delays," *Neurocomputing*, vol. 77, no. 1, pp. 222–228, 2012.
- [5] D. Xu and L. Xu, "New results for studying a certain class of nonlinear delay differential systems," *IEEE Transactions on Automatic Control*, vol. 55, no. 7, pp. 1641–1645, 2010.
- [6] D. Xu and X. Wang, "A new nonlinear integro-differential inequality and its application," *Applied Mathematics Letters*, vol. 22, no. 11, pp. 1721–1726, 2009.
- [7] D. Xu and Z. Yang, "Attracting and invariant sets for a class of impulsive functional differential equations," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 2, pp. 1036–1044, 2007.
- [8] D. Xu, Z. Yang, and Y. Huang, "Existence-uniqueness and continuation theorems for stochastic functional differential equations," *Journal of Differential Equations*, vol. 245, no. 6, pp. 1681–1703, 2008.
- [9] D. Xu and H. Zhao, "Invariant set and attractivity of nonlinear differential equations with delays," *Applied Mathematics Letters*, vol. 15, no. 3, pp. 321–325, 2002.
- [10] R. Rao, X. Wang, S. Zhong, and Z. Pu, "LMI approach to exponential stability and almost sure exponential stability for stochastic fuzzy Markovian-jumping Cohen-Grossberg neural networks with nonlinear  $p$ -Laplace diffusion," *Journal of Applied Mathematics*, vol. 2013, Article ID 396903, 21 pages, 2013.
- [11] X. Yang, C. Huang, D. Zhang, and Y. Long, "Dynamics of Cohen-Grossberg neural networks with mixed delays and impulses," *Abstract and Applied Analysis*, vol. 2008, Article ID 432341, 14 pages, 2008.
- [12] L. Wan and J. Sun, "Global asymptotic stability of Cohen-Grossberg neural network with continuously distributed delays," *Physics Letters A*, vol. 342, no. 4, pp. 331–340, 2005.
- [13] R. F. Rao, S. M. Zhong, and X. R. Wang, "Stochastic stability criteria with LMI conditions for Markovian jumping impulsive BAM neural networks with mode-dependent time-varying delays and nonlinear reaction-diffusion," *Communications in Nonlinear Science and Numerical Simulation*, 2013.
- [14] Q. Song and J. Zhang, "Global exponential stability of impulsive Cohen-Grossberg neural network with time-varying delays," *Nonlinear Analysis: Real World Applications*, vol. 9, no. 2, pp. 500–510, 2008.
- [15] Q. Song and J. Cao, "Impulsive effects on stability of fuzzy Cohen-Grossberg neural networks with time-varying delays," *IEEE Transactions on Systems, Man, and Cybernetics B*, vol. 37, no. 3, pp. 733–741, 2007.
- [16] Q. Song and Z. Wang, "Stability analysis of impulsive stochastic Cohen-Grossberg neural networks with mixed time delays," *Physica A*, vol. 387, no. 13, pp. 3314–3326, 2008.
- [17] C. Bai, "Stability analysis of Cohen-Grossberg BAM neural networks with delays and impulses," *Chaos, Solitons & Fractals*, vol. 35, no. 2, pp. 263–267, 2008.
- [18] B. Li and D. Y. Xu, "Exponential  $p$ -stability of stochastic recurrent neural networks with mixed delays and Markovian switching," *Neurocomputing*, vol. 103, pp. 239–246, 2013.
- [19] B. Li and D. Xu, "Existence and exponential stability of periodic solution for impulsive Cohen-Grossberg neural networks with time-varying delays," *Applied Mathematics and Computation*, vol. 219, no. 5, pp. 2506–2520, 2012.
- [20] Q. Song, Z. Wang, and J. Liang, "Analysis on passivity and passification of T-S fuzzy systems with time-varying delays," *Journal of Intelligent & Fuzzy Systems*, vol. 24, no. 1, pp. 21–30, 2013.
- [21] P. Balasubramaniam, V. Vembarasan, and R. Rakkiyappan, "Delay-dependent robust exponential state estimation of Markovian jumping fuzzy Hopfield neural networks with mixed random time-varying delays," *Communications in Nonlinear Science and Numerical Simulation*, vol. 16, no. 4, pp. 2109–2129, 2011.
- [22] S. Muralisankar, N. Gopalakrishnan, and P. Balasubramaniam, "An LMI approach for global robust dissipativity analysis of T-S fuzzy neural networks with interval time-varying delays," *Expert Systems with Applications*, vol. 39, no. 3, pp. 3345–3355, 2012.
- [23] R. F. Rao and Z. L. Pu, "Stability analysis for impulsive stochastic fuzzy  $p$ -Laplace dynamic equations under Neumann or Dirichlet boundary condition," *Boundary Value Problems*, vol. 2013, article 133, 2013.
- [24] Z. L. Pu and R. F. Rao, "Stability analysis of Takagi-Sugeno fuzzy Markovian jumping PDEs with  $p$ -Laplace under zero-boundary condition," to appear *Journal of Applied Mathematics*.
- [25] P. Balasubramaniam, V. Vembarasan, and R. Rakkiyappan, "Delay-dependent robust asymptotic state estimation of Takagi-Sugeno fuzzy Hopfield neural networks with mixed interval time-varying delays," *Expert Systems with Applications*, vol. 39, no. 1, pp. 472–481, 2012.
- [26] J. Luo, "Fixed points and stability of neutral stochastic delay differential equations," *Journal of Mathematical Analysis and Applications*, vol. 334, no. 1, pp. 431–440, 2007.
- [27] J. Luo, "Fixed points and exponential stability of mild solutions of stochastic partial differential equations with delays," *Journal of Mathematical Analysis and Applications*, vol. 342, no. 2, pp. 753–760, 2008.



- [28] J. Luo, "Stability of stochastic partial differential equations with infinite delays," *Journal of Computational and Applied Mathematics*, vol. 222, no. 2, pp. 364–371, 2008.
- [29] J. Luo and T. Taniguchi, "Fixed points and stability of stochastic neutral partial differential equations with infinite delays," *Stochastic Analysis and Applications*, vol. 27, no. 6, pp. 1163–1173, 2009.
- [30] R. Sakthivel and J. Luo, "Asymptotic stability of impulsive stochastic partial differential equations with infinite delays," *Journal of Mathematical Analysis and Applications*, vol. 356, no. 1, pp. 1–6, 2009.
- [31] R. Sakthivel and J. Luo, "Asymptotic stability of nonlinear impulsive stochastic differential equations," *Statistics & Probability Letters*, vol. 79, no. 9, pp. 1219–1223, 2009.
- [32] J. Luo, "Fixed points and exponential stability for stochastic Volterra-Levin equations," *Journal of Computational and Applied Mathematics*, vol. 234, no. 3, pp. 934–940, 2010.
- [33] X. Lai and Y. Zhang, "Fixed point and asymptotic analysis of cellular neural networks," *Journal of Applied Mathematics*, Article ID 689845, 12 pages, 2012.
- [34] T. A. Burton, *Stability by Fixed Point Theory for Functional Differential Equations*, Dover, Mineola, NY, USA, 2006.
- [35] R. F. Rao, S. M. Zhong, and X. R. Wang, "Delay-dependent exponential stability for Markovian jumping stochastic Cohen-Grossberg neural networks with p-Laplace diffusion and partially known transition rates via a differential inequality," *Advances in Difference Equations*, vol. 2013, article 183, 2013.
- [36] T. A. Burton, "Fixed points, stability, and exact linearization," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 61, no. 5, pp. 857–870, 2005.
- [37] C. Jin and J. Luo, "Fixed points and stability in neutral differential equations with variable delays," *Proceedings of the American Mathematical Society*, vol. 136, no. 3, pp. 909–918, 2008.
- [38] Y. N. Raffoul, "Stability in neutral nonlinear differential equations with functional delays using fixed-point theory," *Mathematical and Computer Modelling*, vol. 40, no. 7-8, pp. 691–700, 2004.
- [39] M. Wu, N. j. Huang, and C. W. Zhao, "Fixed points and stability in neutral stochastic differential equations with variable delays," *Fixed Point Theory and Applications*, vol. 2008, Article ID 407352, 11 pages, 2008.
- [40] B. Zhang, "Fixed points and stability in differential equations with variable delays," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 63, no. 5–7, pp. e233–e242, 2005.
- [41] Y. T. Zhang and Q. Luo, "Global exponential stability of impulsive cellular neural networks with time-varying delays via fixed point theory," *Advances in Difference Equations*, vol. 2013, article 23, 2013.
- [42] L. Ding, X. Li, and Z. Li, "Fixed points and stability in nonlinear equations with variable delays," *Fixed Point Theory and Applications*, vol. 2010, Article ID 195916, 14 pages, 2010.

