

Research Article

C^* -Algebras from Groupoids on Self-Similar Groups

Inhyeop Yi

Department of Mathematics Education, Ewha Womans University, Seoul 120-750, Republic of Korea

Correspondence should be addressed to Inhyeop Yi; yih@ewha.ac.kr

Received 10 May 2013; Accepted 2 July 2013

Academic Editor: Salvador Hernandez

Copyright © 2013 Inhyeop Yi. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We show that the Smale spaces from self-similar groups are topologically mixing and their stable algebra and stable Ruelle algebra are strongly Morita equivalent to groupoid algebras of Anantharaman-Delaroche and Deaconu. And we show that $C^*(R_\infty)$ associated to a postcritically finite hyperbolic rational function is an AT -algebra of real-rank zero with a unique trace state.

1. Introduction

Nekrashevych has developed a theory of dynamical systems and C^* -algebras for self-similar groups in [1, 2]. These groups include groups acting on rooted trees and finite automata and iterated monodromy groups of self-covering on topological spaces. From self-similar groups, Nekrashevych constructed Smale spaces of Ruelle and Putnam with their corresponding stable and unstable algebras and those of Ruelle algebras for various equivalence relations on the Smale spaces [3–7].

Main approach to C^* -algebras structures in [2] is based on Cuntz-Pimsner algebras generated by self-similar groups. However Smale spaces and their corresponding C^* -algebras have rich dynamical structures, and it is conceivable that dynamical systems associated with self-similar groups may give another way to study C^* -algebras from self-similar groups. Our intention is to elucidate self-similar groups from the perspective of dynamical systems.

This paper is concerned with groupoids and their groupoid C^* -algebras from the stable equivalence relation on the limit solenoid $(S_G, \bar{\sigma})$ of a self-similar group (G, X) . Instead of using the groupoids G_s and $G_s \rtimes \mathbb{Z}$ on the Smale space $(S_G, \bar{\sigma})$ as Putnam [3, 4] and Nekrashevych [2] did, we consider the essentially principal groupoids R_∞ and $\Gamma(J_G, \sigma)$ of Anantharaman-Delaroche [8] and Deaconu [9] on a presentation (J_G, σ) of $(S_G, \bar{\sigma})$. While G_s and $G_s \rtimes \mathbb{Z}$ are not r -discrete groupoids, R_∞ and $\Gamma(J_G, \sigma)$ are r -discrete. And R_∞ and $\Gamma(J_G, \sigma)$ are defined on (J_G, σ) so that we do not need

to entail the inverse limit structure of $(S_G, \bar{\sigma})$. Thus R_∞ and $\Gamma(J_G, \sigma)$ are more manageable than G_s and $G_s \rtimes \mathbb{Z}$ for the structures of their C^* -algebras.

In this paper, we prove that, for a self-similar group (G, X) , its limit dynamical system (J_G, σ) is topologically mixing so that $(S_G, \bar{\sigma})$ is an irreducible Smale space. And we show that R_∞ is equivalent to G_s and $\Gamma(J_G, s)$ is equivalent to $G_s \rtimes \mathbb{Z}$ in the sense of Muhly et al. [10]. Consequently, the groupoid C^* -algebras $C^*(R_\infty)$ and $C^*(\Gamma(J_G, \sigma))$ are strongly Morita equivalent to the stable algebra S and the stable Ruelle algebra R_s , respectively, of $(S_G, \bar{\sigma})$. Then we use R_∞ and $\Gamma(J_G, \sigma)$ to study structures of C^* -algebras from a self-similar group (G, X) . Finally we show that groupoid algebras of R_∞ from postcritically finite hyperbolic rational functions are AT -algebras of real-rank zero.

The outline of the paper is as follows. In Section 2, we review the notions of self-similar groups and their groupoids and show that the induced limit dynamical system and the limit solenoid of a self-similar group are topologically mixing. In Section 3, we observe that R_∞ is equivalent to G_s and $\Gamma(J_G, s)$ is equivalent to $G_s \rtimes \mathbb{Z}$. In Section 4, we give a proof that its groupoid algebra $C^*(\Gamma(J_G, \sigma))$ is simple, purely infinite, separable, stable, and nuclear and satisfies the Universal Coefficient Theorem. For R_∞ , we show that $C^*(R_\infty)$ is simple and nuclear. And, when self-similar group is defined by a postcritically finite hyperbolic rational function and its Julia set, we show that $C^*(R_\infty)$ is an AT -algebra.

2. Self-Similar Groups

We review the properties of self-similar groups. As for general references for the notions of self-similar groups, we refer to [1, 2].

Suppose that X is a finite set. We denote by X^n the set of words of length n in X with $X^0 = \{\emptyset\}$, and define $X^* = \bigcup_{n=0}^{\infty} X^n$. A *self-similar group* (G, X) consists of an X and a faithful action of a group G on X such that, for all $g \in G$ and $x \in X$, there exist unique $y \in X$ and $h \in G$ such that

$$g(xw) = yh(w) \quad \text{for every } w \in X^*. \quad (1)$$

The above equality is written formally as

$$g \cdot x = y \cdot h. \quad (2)$$

We observe that for any $g \in G$ and $v \in X^*$, there exists a unique element $h \in G$ such that $g(vw) = g(v)h(w)$ for every $w \in X^*$. The unique element h is called the *restriction* of g at v and is denoted by $g|_v$. For $u = g(v)$ and $h = g|_v$, we write

$$g \cdot v = u \cdot h. \quad (3)$$

A self-similar group (G, X) is called *recurrent* if, for all $x, y \in X$, there is a $g \in G$ such that $g \cdot x = y \cdot 1$; that is, $g(xw) = yw$ for every $w \in X^*$. We say that (G, X) is *contracting* if there is a finite subset N of G satisfying the following: for every $g \in G$, there is $n \geq 0$ such that $g|_v \in N$ for every $v \in X^*$ of length $|v| \geq n$. If the group is contracting, the smallest set N satisfying this condition is called the *nucleus* of the group.

Standing Assumption. We assume that our self-similar group (G, X) is a contracting, recurrent, and regular group and that the group G is finitely generated.

Path Spaces. For a self-similar group (G, X) , the set X^* has a natural structure of a rooted tree: the root is \emptyset , the vertices are words in X^* , and the edges are of the form v to vx , where $v \in X^*$ and $x \in X$. Then the boundary of the tree X^* is identified with the space X^ω of right-infinite paths of the form $x_1x_2 \cdots$, where $x_i \in X$. The product topology of discrete set X is given on X^ω .

We say that a self-similar group (G, X) is *regular* if, for every $g \in G$ and every $w \in X^\omega$, either $g(w) \neq w$ or there is a neighborhood of w such that every point in the neighborhood is fixed by g .

We also consider the space $X^{-\omega}$ of left-infinite paths $\cdots x_{-2}x_{-1}$ over X with the product topology. Two paths $\cdots x_{-2}x_{-1}$ and $\cdots y_{-2}y_{-1}$ in $X^{-\omega}$ are said to be *asymptotically equivalent* if there is a finite set $I \subset G$ and a sequence $g_n \in I$ such that

$$g_n(x_{-n} \cdots x_{-1}) = y_{-n} \cdots y_{-1}, \quad (4)$$

for every $n \in \mathbb{N}$. The quotient of the space $X^{-\omega}$ by the asymptotic equivalence relation is called the *limit space* of (G, X) and is denoted by J_G . Since the asymptotic equivalence relation is invariant under the shift map $\cdots x_{-2}x_{-1} \mapsto \cdots x_{-3}x_{-2}$, the shift map induces a continuous map $\sigma : J_G \rightarrow J_G$. We call the induced dynamical system (J_G, σ) the *limit dynamical system* of (G, X) (see [1, 2] for details).

Remark 1. Recurrent and finitely generated conditions imply that J_G is a compact, connected, locally connected, metrizable space of a finite dimension by Corollary 2.8.5 and Theorem 3.6.4 of [1]. And regular condition implies that σ is an $|X|$ -fold self-covering map by Proposition 6.1 of [2].

A cylinder set $Z(u)$ for each $u \in X^* = \bigcup_{n \geq 0} X^n$ is defined as follows:

$$Z(u) = \{\xi \in X^{-\omega} : \xi = \cdots x_{-n-1}x_{-n} \cdots x_{-1} \text{ such that } x_{-n} \cdots x_{-1} = u\}. \quad (5)$$

Then the collection of all such cylinder sets forms a basis for the product topology on $X^{-\omega}$. And we recall that a dynamical system (Y, f) is called *topologically mixing* if, for every pair of nonempty open sets A, B in Y , there is an $n \in \mathbb{N}$ such that $f^k(A) \cap B \neq \emptyset$ for every $k \geq n$.

Theorem 2. (J_G, σ) is a topologically mixing system.

Proof. As $X^{-\omega}$ has the product topology and J_G has the quotient topology induced from asymptotic equivalence relation, it is sufficient to show that, for arbitrary cylinder sets $Z(u)$ and $Z(v)$ of $X^{-\omega}$, there are infinite paths $\xi = \cdots x_{-2}x_{-1} \in Z(u)$ and $\eta = \cdots y_{-2}y_{-1} \in Z(v)$ such that ξ is asymptotically equivalent to η . Moreover we can assume that $u, v \in X^n$ for some $n \in \mathbb{N}$ so that $u = x_{-n} \cdots x_{-1}$ and $v = y_{-n} \cdots y_{-1}$.

We choose sufficiently large m and let $a, b \in X^{m-n}$ so that au and bv are elements of X^m . Then by recurrent condition and [1, Corollary 2.8.5], for au and bv in X^m , there is a $g \in G$ such that $g(au) = g(a)g|_a(u) = bv$. Since we chose large m , by contracting condition, $g|_a$ is an element of the nucleus of (G, X) .

We remind that the nucleus of (G, X) is a finite set and equal to

$$N = \bigcup_{g \in G} \bigcap_{n \geq 0} \{g|_v : v \in X^*, |v| \geq n\}. \quad (6)$$

So an element of the nucleus is a restriction of another element of the nucleus. Hence $g|_a \in N$ implies that there exist a letter x_{-n-1} and a $g_{-n-1} \in N$ such that $g_{-n-1}|_{x_{-n-1}} = g|_a$. Then, for $g_{-n-1}(x_{-n-1}u) = y_{-n-1}v$, we have

$$g_{-n-1}(x_{-n-1}u) = y_{-n-1}v. \quad (7)$$

So by induction there are a letter x_{-m} and a $g_{-m} \in N$ for every $m \geq n$ such that

$$g_{-m}(x_{-m} \cdots x_{-n-1}u) = y_{-m} \cdots y_{-n-1}v. \quad (8)$$

Let $\xi = \cdots x_{-2}x_{-1}$ and let $\eta = \cdots y_{-2}y_{-1}$. Then it is trivial that $\xi \in Z(u)$ and $\eta \in Z(v)$. And ξ is asymptotically equivalent to η . Therefore the limit dynamical system (J_G, σ) is topologically mixing. \square

Let $X^{\mathbb{Z}}$ be the space of bi-infinite paths $\cdots x_{-1}x_0 \cdot x_1x_2 \cdots$ over the alphabet X . The direct product topology of the discrete set X is given on $X^{\mathbb{Z}}$. We say that two paths $\cdots x_{-1}x_0 \cdot x_1x_2 \cdots$ and $\cdots y_{-1}y_0 \cdot y_1y_2 \cdots$ in $X^{\mathbb{Z}}$ are *asymptotically*

equivalent if there is a finite set $I \subset G$ and a sequence $g_n \in I$ such that

$$g_n(x_n x_{n+1} \cdots) = y_n y_{n+1} \cdots, \quad (9)$$

for every $n \in \mathbb{Z}$. The quotient of $X^{\mathbb{Z}}$ by the asymptotic equivalence relation is called the *limit solenoid* of (G, X) and is denoted by S_G . As in the case of J_G , the shift map on $X^{\mathbb{Z}}$ is transferred to an induced homeomorphism on S_G , which we will denote by $\bar{\sigma}$.

Theorem 3 (see [1, 2]). *The limit solenoid S_G is homeomorphic to the inverse limit space of (J_G, σ)*

$$J_G \xleftarrow{\sigma} J_G \xleftarrow{\sigma} \cdots = \left\{ (\xi_0, \xi_1, \xi_2, \dots) \in \prod_{n \geq 0} J_G : \sigma(\xi_{n+1}) = \xi_n \text{ for every } n \geq 0 \right\}, \quad (10)$$

and $\bar{\sigma} : S_G \rightarrow S_G$ is the induced homeomorphism defined by

$$\begin{aligned} (\xi_0, \xi_1, \xi_2, \dots) &\mapsto (\sigma(\xi_0), \sigma(\xi_1), \sigma(\xi_2), \dots) \\ &= (\sigma(\xi_0), \xi_0, \xi_1, \dots). \end{aligned} \quad (11)$$

Moreover, the limit solenoid system $(S_G, \bar{\sigma})$ is a Smale space.

Then we have the following from Theorem 2.

Corollary 4. *$(S_G, \bar{\sigma})$ is topologically mixing.*

We have a natural projection $\pi : S_G \rightarrow J_G$ induced from the map

$$\cdots x_{-1} x_0 \cdot x_1 x_2 \cdots \mapsto \cdots x_{-1} x_0, \quad (12)$$

and the relation that $\cdots x_{n-1} x_n \in X^{-\omega}$ represents $\xi_n \in J_G$. Then it is easy to check $\pi \circ \bar{\sigma} = \sigma \circ \pi$. The *stable equivalence relation* on $(S_G, \bar{\sigma})$ is defined as follows [2, Proposition 6.8]:

Definition 5. One says that two elements α and β in S_G are stably equivalent and write $\alpha \sim_s \beta$ if there is a $k \in \mathbb{Z}$ such that $\pi \bar{\sigma}^k(\alpha) = \pi \bar{\sigma}^k(\beta)$.

In other words, when α and β are represented by infinite paths $(x_n)_{n \in \mathbb{Z}}$ and $(y_n)_{n \in \mathbb{Z}}$ in $X^{\mathbb{Z}}$, $\alpha \sim_s \beta$ if and only if the corresponding left-infinite paths $\cdots x_{k-1} x_k$ and $\cdots y_{k-1} y_k$ in $X^{-\omega}$ are asymptotically equivalent for some $k \in \mathbb{Z}$.

Groupoids on (J_G, σ) and $(S_G, \bar{\sigma})$. Suppose that (G, X) is a self-similar group and $(S_G, \bar{\sigma})$ is its corresponding limit solenoid. We recall from [3] that the stable equivalence groupoid G_s on S_G and its semidirect product by \mathbb{Z} are defined to be

$$\begin{aligned} G_s &= \{(\alpha, \beta) \in S_G \times S_G : \alpha \sim_s \beta\}, \\ G_s \rtimes \mathbb{Z} &= \{(\alpha, n, \beta) \in S_G \times \mathbb{Z} \times S_G : n \in \mathbb{Z}, \\ &\quad (\bar{\sigma}^n(\alpha), \beta) \in G_s\}. \end{aligned} \quad (13)$$

Then G_s and $G_s \rtimes \mathbb{Z}$ are groupoids with the natural structure maps. The unit spaces of G_s and $G_s \rtimes \mathbb{Z}$ are identified with S_G via the maps $\alpha \in S_G \mapsto (\alpha, \alpha) \in G_s$ and $\alpha \mapsto (\alpha, 0, \alpha) \in G_s \rtimes \mathbb{Z}$, respectively.

To give topologies on these groupoids, we consider subgroupoids of G_s . For each $n \geq 0$, set

$$G_{s,n} = \{(\alpha, \beta) \in S_G \times S_G : \pi \bar{\sigma}^n(\alpha) = \pi \bar{\sigma}^n(\beta)\}. \quad (14)$$

Then $G_{s,n}$ is a subgroupoid of G_s . Note that if μ and ν in S_G are stably equivalent with $\pi \bar{\sigma}^l(\mu) = \pi \bar{\sigma}^l(\nu)$ for some negative integer l , then

$$\pi(\mu) = \sigma^{-l} \pi \bar{\sigma}^l(\mu) = \sigma^{-l} \pi \bar{\sigma}^l(\nu) = \pi(\nu) \quad (15)$$

implies that $(\mu, \nu) \in G_{s,0}$. So we obtain the stable equivalence groupoid

$$G_s = \bigcup_{n \geq 0} G_{s,n}. \quad (16)$$

Each $G_{s,n}$ is given the relative topology from $S_G \times S_G$, and G_s is given the inductive limit topology. Under this topology, it is not difficult to check that G_s is a locally compact Hausdorff principal groupoid with the natural structure maps. For $G_s \rtimes \mathbb{Z}$, we transfer the product topology of $G_s \times \mathbb{Z}$ to $G_s \rtimes \mathbb{Z}$ via the map $((\alpha, \beta), n) \mapsto (\alpha, n, \bar{\sigma}^n(\beta))$. Amenability and Haar systems on G_s and $G_s \times \mathbb{Z}$ are explained in [2–4]. We denote the groupoid C^* -algebra of G_s by S and that of $G_s \rtimes \mathbb{Z}$ by R_s and call it *stable Ruelle algebra* on $(S_G, \bar{\sigma})$.

For the limit dynamical system $(J_G \cdot \sigma)$ of a self-similar group (G, X) , we construct groupoids R_∞ and $\Gamma(J_G, \sigma)$ of Anantharaman-Delaroche [8] and Deaconu [9]. Let $R_n = \{(\xi, \eta) \in J_G \times J_G : \sigma^n(\xi) = \sigma^n(\eta)\}$ for $n \geq 0$ and define

$$R_\infty = \bigcup_{n \geq 0} R_n,$$

$$\begin{aligned} \Gamma(J_G, \sigma) &= \{(\xi, n, \eta) \in J_G \times \mathbb{Z} \times J_G : \exists k, l \geq 0, \\ &\quad n = k - l, \sigma^k(\xi) = \sigma^l(\eta)\} \end{aligned} \quad (17)$$

with the natural structure maps. The unit spaces of R_∞ and $\Gamma(J_G, \sigma)$ are identified with J_G via $\xi \mapsto (\xi, \xi)$ and $\xi \mapsto (\xi, 0, \xi)$.

We give the relative topology from $J_G \times J_G$ on R_n and the inductive limit topology on R_∞ . Then R_∞ is a second countable, locally compact, Hausdorff, r -discrete groupoid with the Haar system given by the counting measures. A topology on $\Gamma(J_G, \sigma)$ is given by basis of the form

$$\Delta(U, V, k \cdot l) = \left\{ \left(\xi, k - l, (\sigma^l|_V)^{-1} \circ \sigma^k(\xi) \right) : \xi \in U \right\}, \quad (18)$$

where U and V are open sets in J_G and $k, l \geq 0$ such that $\sigma^k|_U$ and $\sigma^l|_V$ are homeomorphisms with the same range. Then $\Gamma(J_G, \sigma)$ is a second countable, locally compact, Hausdorff, r -discrete groupoid, and the counting measure is a Haar system [9, 11]. Amenability of R_∞ and $\Gamma(J_G, \sigma)$ is explained in Proposition 2.4 of [12]. We denote the groupoid C^* -algebras of R_∞ and $\Gamma(J_G, \sigma)$ by $C^*(R_\infty)$ and $C^*(\Gamma(J_G, \sigma))$, respectively.

3. Groupoid Equivalence

We follow Kumjian and Pask [13, Section 5] to obtain equivalence of groupoids between G_s and R_∞ and between $G_s \rtimes \mathbb{Z}$ and $\Gamma(J_G, \sigma)$, respectively, in the sense of Muhly et al. [10].

We repeat Kumjian and Pask's observation [13]. Suppose that Y is a locally compact Hausdorff space and that Γ is a locally compact Hausdorff groupoid. For a continuous open surjection $\phi : Y \rightarrow \Gamma^0$, we set a topological space

$$Z = Y * \Gamma = \{(y, \gamma) : y \in Y, \gamma \in \Gamma, \phi(y) = s(\gamma)\} \quad (19)$$

with the relative topology in $Y \times \Gamma$ and a locally compact Hausdorff groupoid

$$\Gamma^\phi = \{(y_1, \gamma, y_2) : y_1, y_2 \in Y, \gamma \in \Gamma, \phi(y_1) = s(\gamma), r(\gamma) = \phi(y_2)\} \quad (20)$$

with the relative topology.

Theorem 6 (see [13, Lemma 5.1]). *Suppose that Y, Γ, ϕ, Z , and Γ^ϕ are as previous. Then Z implements an equivalence between Γ and Γ^ϕ in the sense of Muhly-Renault-Williams.*

Now we consider $\phi : S_G \rightarrow R_\infty^0$ defined by $\alpha \mapsto (\pi(\alpha), \pi(\alpha))$. Since ϕ is the composition of the projection map $\pi : S_G \rightarrow J_G$ and the identity map from J_G to R_∞^0 , ϕ is a continuous open surjection. Then we have

$$R_\infty^\phi = \{(\alpha, (\pi(\alpha), \pi(\beta)), \beta) : \alpha, \beta \in S_G, (\pi(\alpha), \pi(\beta)) \in R_\infty\}. \quad (21)$$

It is not difficult to check that $R_\infty^\phi = \cup_{n \geq 0} R_n^\phi$, where

$$R_n^\phi = \{(\alpha, (\pi(\alpha), \pi(\beta)), \beta) : \alpha, \beta \in S_G, \sigma^n(\pi(\alpha)) = \sigma^n(\pi(\beta))\}, \quad (22)$$

and that the relative topology on R_∞^ϕ is equivalent to the inductive limit topology.

Lemma 7. *Suppose that $(S_G, \bar{\sigma})$ is the limit solenoid system induced from a self-similar group (G, X) and that G_s is the stable equivalence groupoid associated with $(S_G, \bar{\sigma})$. Then $\tilde{\phi} : G_s \rightarrow R_\infty^\phi$ defined by $(\alpha, \beta) \mapsto (\alpha, (\pi(\alpha), \pi(\beta)), \beta)$ is a groupoid isomorphism.*

Proof. Remember that $G_s = \cup_{n \geq 0} G_{s,n}$ and $R_\infty^\alpha = \cup_{n \geq 0} R_n^\alpha$. From the commutative relation $\sigma\pi = \pi\bar{\sigma}$, we observe

$$\begin{aligned} (\alpha, \beta) \in G_{s,n} &\iff \pi\bar{\sigma}^n(\alpha) = \pi\bar{\sigma}^n(\beta) \iff \sigma^n\pi(\alpha) \\ &= \sigma^n\pi(\beta). \end{aligned} \quad (23)$$

Hence $\tilde{\phi}|_{G_{s,n}}$ is a well-defined bijective map between $G_{s,n}$ and R_n^ϕ .

Since topologies on $G_{s,n}$ and R_n^ϕ are relative topologies from $S_G \times S_G$, $\tilde{\phi}|_{G_{s,n}}$ is a homeomorphism. Then $\tilde{\phi}$ is a homeomorphism as the inductive limit topologies are given on G_s and R_∞^ϕ . It is routine to check that $\tilde{\phi}$ is a groupoid morphism. \square

The groupoid equivalence between R_∞ and G_s follows from Theorem 6 and Lemma 7. Strong Morita equivalence is from [10, Proposition 2.8] as both groupoids have Haar systems.

Theorem 8. *Suppose that (G, X) is a self-similar group, that R_∞ is the groupoid associated with (J_G, σ) , and that G_s is the stable equivalence groupoid associated with $(S_G, \bar{\sigma})$. Then R_∞ and G_s are equivalent in the sense of Muhly-Renault-Williams. Therefore $C^*(R_\infty)$ is strongly Morita equivalent to the stable algebra S on the limit solenoid system $(S_G, \bar{\sigma})$.*

Analogous assertions hold for $\Gamma(J_G, \sigma)$ and $G_s \rtimes \mathbb{Z}$. For $\psi : S_G \rightarrow \Gamma(J_G, \sigma)^0$ defined by $\alpha \mapsto (\pi(\alpha), 0, \pi(\alpha))$, we observe

$$\begin{aligned} \Gamma(J_G, \sigma)^\psi &= \{(\alpha, (\pi(\alpha), n, \pi(\beta)), \beta) : \alpha, \beta \in S_G, \\ &(\pi(\alpha), n, \pi(\beta)) \in \Gamma(J_G, \sigma)\}. \end{aligned} \quad (24)$$

Lemma 9. *Suppose that G_s is the stable equivalence groupoid of $(S_G, \bar{\sigma})$ and that $G_s \rtimes \mathbb{Z}$ is the semidirect product groupoid. Then $\tilde{\psi} : G_s \rtimes \mathbb{Z} \rightarrow \Gamma(J_G, \sigma)^\psi$ defined by $(\alpha, n, \beta) \mapsto (\alpha, (\pi(\alpha), n, \pi(\beta)), \beta)$ is a groupoid isomorphism.*

Proof. Recall that $(\alpha, n, \beta) \in G_s \rtimes \mathbb{Z} \iff (\bar{\sigma}^n(\alpha), \beta) \in G_s$. Then $G_s = \cup_{n \geq 0} G_{s,n}$ implies that $(\bar{\sigma}^n(\alpha), \beta) \in G_{s,l}$ for some $l \geq 0$. So from the proof of Lemma 7, we obtain that

$$\begin{aligned} (\bar{\sigma}^n(\alpha), \beta) \in G_{s,l} &\iff \sigma^{n+l}(\pi(\alpha)) \\ &= \sigma^l(\pi(\beta)) \iff (\pi(\alpha), n, \pi(\beta)) \in \Gamma(J_G, \sigma). \end{aligned} \quad (25)$$

Thus $\tilde{\psi}$ is a well-defined bijective map. As $G_s \rtimes \mathbb{Z}$ has the product topology, we notice that $\tilde{\psi}|_{G_s \rtimes \{0\}}$ is the homeomorphism $\tilde{\phi}$ defined in Lemma 7 and that $\tilde{\psi}|_{G_s \rtimes \{n\}}$ is homeomorphism onto

$$\begin{aligned} \{(\alpha, (\pi(\alpha), n, \pi(\beta)), \beta) : \alpha, \beta \in S_G, \\ \sigma^{n+l}(\pi(\alpha)) = \sigma^l(\pi(\beta))\}. \end{aligned} \quad (26)$$

It is trivial that $\tilde{\psi}$ is a groupoid morphism. \square

Theorem 10. *Suppose that (G, X) is a self-similar group. Then $\Gamma(J_G, \sigma)$ and $G_s \rtimes \mathbb{Z}$ are equivalent in the sense of Muhly-Renault-Williams. Therefore $C^*(\Gamma(J_G, \sigma))$ is strongly Morita equivalent to the stable Ruelle algebra R_s on $(S_G, \bar{\sigma})$.*

Remark 11. In [11], Chen and Hou showed similar result under an extra condition that a Smale space is the inverse limit of an expanding surjection on a compact metric space.

4. Groupoid Algebras

Suppose that (G, X) is a self-similar group. We use its corresponding R_∞ and $\Gamma(J_G, \sigma)$ to study C^* -algebraic structures of stable algebra and stable Ruelle algebra from (G, X) .

Following Renault [15], we say that a topological groupoid Γ with an open range map is *essentially principal* if Γ is locally compact and, for every closed invariant subset E of its unit

space Γ^0 , $\{u \in E : r^{-1}(u) \cap s^{-1}(u) = \{u\}\}$ is dense in E . A subset E of Γ^0 is called invariant if $r \circ s^{-1}(E) = E$. And Γ is called *minimal* if the only open invariant subsets of Γ^0 are the empty set \emptyset and Γ^0 itself. We refer [15] for details.

Proposition 12. *The groupoid $\Gamma(J_G, \sigma)$ is essentially principal.*

Proof. Let

$$A = \{\xi \in J_G : \text{for } k, l \geq 0, \sigma^k(\xi) = \sigma^l(\xi) \text{ implies } k = l\},$$

$$B = \{b \in \Gamma(J_f, f)^0 : r^{-1}(b) \cap s^{-1}(b) = \{b\}\}. \quad (27)$$

Then we observe $\xi \in A \Leftrightarrow (\xi, 0, \xi) \in B$. Hence A is dense in X implying that B is dense in $\Gamma(J_f, f)^0$ so that $\Gamma(J_G, \sigma)$ is essentially principal.

To show that A is dense in J_G , we assume A is not dense in J_G . Then we can find an open set $U \subset J_G$ such that $\bar{U} \cap A = \emptyset$ as J_G is a compact Hausdorff space. Since

$$J_G - A = \bigcup_{n=1}^{\infty} \bigcup_{k=0}^{\infty} \sigma^{-k}(\text{Per}_n), \quad (28)$$

where $\text{Per}_n = \{\xi \in J_G : \sigma^n(\xi) = \xi\}$, we have

$$\bar{U} = \bar{U} \cap (J_f - A) = \bigcup_{n=1}^{\infty} \bigcup_{k=0}^{\infty} \bar{U} \cap \sigma^{-k}(\text{Per}_n). \quad (29)$$

Then by Baire category theorem, there exist some integers $n \geq 1$ and $k \geq 0$ such that $\bar{U} \cap \sigma^{-k}(\text{Per}_n)$ has nonempty interior. But $\text{Per}_n = \{\xi \in J_G : \sigma^n(\xi) = \xi\}$ is a finite set because X is a finite set, and $\sigma^{-k}(\text{Per}_n)$ is a finite set as σ is an $|X|$ -fold covering map, a contradiction. Therefore A is dense in J_G , and $\Gamma(J_G, \sigma)$ is an essentially principal groupoid. \square

There are excellent criteria for groupoid algebras from dynamical systems to be simple and purely infinite developed by Renault [12].

Lemma 13 (see [12]). *For a topological space X and a local homeomorphism $T : X \rightarrow X$, let $\Gamma(X, T)$ be the groupoid of Anantharaman-Delaroche and Deaconu. Suppose that $\Gamma(X, T)$ is an essentially principal groupoid and $C^*(X, T)$ is its groupoid algebra.*

- (1) *Assume that for every nonempty open set $U \subset X$ and every $x \in X$, there exist $m, n \in \mathbb{N}$ such that $T^m(x) \in T^n(U)$. Then $C^*(X, T)$ is simple.*
- (2) *Assume that for every nonempty open set $U \subset X$, there exist an open set $V \subset U$ and $m, n \in \mathbb{N}$ such that $T^m(V)$ is strictly contained in $T^n(V)$. Then $C^*(X, T)$ is purely infinite.*

As $\Gamma(J_G, \sigma)$ is an essentially principal groupoid, we have an alternative proof for Theorem 6.5 of [2].

Theorem 14. *The algebra $C^*(\Gamma(J_G, \sigma))$ is simple, purely infinite, separable, stable, and nuclear and satisfies the Universal Coefficient Theorem of Rosenberg-Schochet.*

Proof. Suppose that U is an open set in J_G . Then the inverse image of U in $X^{-\omega}$, say U' , is open, and there is a cylinder set $Z(u)$ defined by some $u \in X^n$ such that $Z(u) \subset U'$. By definition of cylinder sets, we have $\sigma^n(Z(u)) = X^{-\omega} \subseteq \sigma^n(U')$, which implies that $\sigma^n(U) = J_G$ on the quotient space. Thus for every $\xi \in J_G$, $\xi \in \sigma^n(U)$ and $C^*(\Gamma(J_G, \sigma))$ is simple.

For an open set U of J_G , let V be an open subset of U such that the inverse image of V in $X^{-\omega}$ is equal to the cylinder set $Z(v)$, where $v \in X^n$ for some $n \geq 2$. Then we obtain $\sigma^n(V) = J_G$ as in the previous, and $\sigma^m(V)$ is a proper subset of $\sigma^n(V)$ for every $1 \leq m < n$. Hence $C^*(\Gamma(J_G, \sigma))$ is purely infinite.

Since $\Gamma(J_G, \sigma)$ is locally compact and second countable, $C^*(\Gamma(J_G, \sigma))$ is σ -unital, nonunital, and separable. So Zhang's dichotomy [16, Theorem 1.2] implies that $C^*(\Gamma(J_G, \sigma))$ is stable. By Proposition 2.4 of [12], nuclear is an easy consequence from amenability of $\Gamma(J_G, \sigma)$. Because $\Gamma(J_G, \sigma)$ is a locally compact amenable groupoid with Haar system, $C^*(\Gamma(J_G, \sigma))$ satisfied the Universal Coefficient Theorem by Lemma 3.5 and Proposition 10.7 of [17]. \square

Corollary 15. *$C^*(\Gamma(J_G, \sigma))$ is $*$ -isomorphic to the stable Ruelle algebra R_s .*

Proof. Because $C^*(\Gamma(J_G, \sigma))$ and R_s are stable, this is trivial from Theorem 10. \square

For $C^*(R_{\infty})$, we use the fact that $R_{\infty} = \cup R_n$ is a principal groupoid representing an AP equivalence relation [18].

Proposition 16. *The groupoid R_{∞} is minimal, and its groupoid algebra $C^*(R_{\infty})$ is simple.*

Proof. In the proof of Theorem 14, we observed that for every cylinder set $Z(u)$ of $X^{-\omega}$, there is an $n > 0$ such that $\sigma^n(Z(u)) = X^{-\omega}$. Since the inverse image of a nonempty open set U in J_G contains a cylinder set $Z(u)$, this observation induces that $\sigma^n(U) = J_G$ on the quotient space. Then R_{∞} is a minimal groupoid by [19, Proposition 2.1]. And simplicity of $C^*(R_{\infty})$ follows from [15, Proposition II.4.6] as R_{∞} is an r -discrete principal groupoid. \square

Proposition 17. *$C^*(R_{\infty})$ is the inductive limit of $C^*(R_n)$. And each $C^*(R_n)$ is strongly Morita equivalent to $C(R_n^0/R_n) = C(J_G/R_n)$.*

Proof. Note that $R_{\infty} = \cup_{n \geq 0} R_n$ is the groupoid representing an AP equivalence relation on stationary sequence $J_G \xrightarrow{\sigma} J_G \xrightarrow{\sigma} \dots$. Thus it is easy to check that Corollary 2.2 of [18] implies the inductive limit structure.

Clearly $R_n = \{(u, v) \in J_G \times J_G : \sigma^n(u) = \sigma^n(v)\}$ is the groupoid representing an equivalence relation on J_G defined by $u \sim_n v$ if and only if $\sigma^n(u) = \sigma^n(v)$. And $(s \times r)(R_n) = (\sigma^{-n} \times \sigma^{-n})(\Delta)$, where $\Delta = \{(u, u) \in J_G \times J_G\}$ implies that $(s \times r)(R_n)$ is a closed subset of $J_G \times J_G$. Thus we have strong Morita equivalence of $C^*(R_n)$ and $C(J_G/R_n)$ by [20, Proposition 2.2]. \square

Corollary 18. $C^*(R_\infty)$ is a nuclear algebra.

Proof. Since $C(J_G/R_n)$ is nuclear, $C^*(R_n)$ is also nuclear by [21, Theorem 15]. And it is a well-known fact that the class of nuclear C^* -algebras is closed under inductive limit. So $C^*(R_\infty)$ is nuclear.

Postcritically Finite Rational Maps. Suppose that $f : \mathbb{C} \rightarrow \mathbb{C}$ is a postcritically finite hyperbolic rational function of degree more than one, that is, a rational function of degree more than one such that the orbit of every critical point of f eventually belongs to a cycle containing a critical point. Then f is expanding on a neighborhood of its Julia set J_f , the group $\text{IMG}(f)$ is contracting, recurrent, regular, and finitely generated, and the limit dynamical system $\sigma : J_{\text{IMG}(f)} \rightarrow J_{\text{IMG}(f)}$ is topologically conjugate with the action of f on its Julia set J_f (see [2, Sections 2 and 6] for details). \square

We borrowed the following theorem from Theorem 3.16 and Remark 4.23 of [22].

Theorem 19. Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a postcritically finite hyperbolic rational function of degree more than one and let R_∞ be the groupoid on its limit dynamical system as in Section 2. Then $C^*(R_\infty)$ is an AT-algebra of real-rank zero with a unique trace state.

Proof. To show that $C^*(R_\infty)$ is an AT-algebra, we use the work of Gong [23, Corollary 6.7]. By Propositions 16 and 17, $C^*(R_\infty)$ is a simple algebra which is an inductive limit of an AH system with uniformly bounded dimensions of local spectra. And Nekrashevych showed that K -groups of $C^*(R_\infty)$ for postcritically finite hyperbolic rational functions are torsion free in [2, Theorem 6.6]. Hence $C^*(R_\infty)$ is an AT-algebra.

As $f : J_f \rightarrow J_f$ is an expanding local homeomorphism (see [2, Section 6.4]) and exact by Proposition 16 and [19, Proposition 2.1], $C^*(R_\infty)$ has a unique trace state by Remark 3.6 of [19]. Simplicity and uniformly bounded dimension conditions imply that $C^*(R_\infty)$ is approximately divisible in the sense of Blackadar et al. [24] as shown by Elliot et al. [14]. Therefore $C^*(R_\infty)$ has real-rank zero by Theorem 1.4 of [24]. \square

Corollary 20. $C^*(R_\infty)$ associated with postcritically finite hyperbolic rational functions of degree more than one belongs to the class of C^* -algebras covered by Elliot classification program.

Acknowledgment

The author would like to express gratitude to the referees for their kind suggestions.

References

- [1] V. Nekrashevych, *Self-Similar Groups*, vol. 117 of *Mathematical Surveys and Monographs*, American Mathematical Society, Providence, RI, USA, 2005.
- [2] V. Nekrashevych, “ C^* -algebras and self-similar groups,” *Journal für die Reine und Angewandte Mathematik*, vol. 630, pp. 59–123, 2009.
- [3] I. F. Putnam, “ C^* -algebras from Smale spaces,” *Canadian Journal of Mathematics*, vol. 48, no. 1, pp. 175–195, 1996.
- [4] I. Putnam, *Hyperbolic Systems and Generalized Cuntz-Krieger Algebras*, Lecture notes from Summer School in Operator Algebras, Odense, Denmark, 1996.
- [5] I. F. Putnam and J. Spielberg, “The structure of C^* -algebras associated with hyperbolic dynamical systems,” *Journal of Functional Analysis*, vol. 163, no. 2, pp. 279–299, 1999.
- [6] D. Ruelle, *Thermodynamic Formalism*, vol. 5, Addison-Wesley, 1978.
- [7] D. Ruelle, “Noncommutative algebras for hyperbolic diffeomorphisms,” *Inventiones Mathematicae*, vol. 93, no. 1, pp. 1–13, 1988.
- [8] C. Anantharaman-Delaroche, “Purely infinite C^* -algebras arising from dynamical systems,” *Bulletin de la Société Mathématique de France*, vol. 125, no. 2, pp. 199–225, 1997.
- [9] V. Deaconu, “Groupoids associated with endomorphisms,” *Transactions of the American Mathematical Society*, vol. 347, no. 5, pp. 1779–1786, 1995.
- [10] P. S. Muhly, J. N. Renault, and D. P. Williams, “Equivalence and isomorphism for groupoid C^* -algebras,” *Journal of Operator Theory*, vol. 17, no. 1, pp. 3–22, 1987.
- [11] X. Chen and C. Hou, “Morita equivalence of groupoid C^* -algebras arising from dynamical systems,” *Studia Mathematica*, vol. 149, no. 2, pp. 121–132, 2002.
- [12] J. Renault, “Cuntz-like algebras,” in *Operator Theoretical Methods*, pp. 371–386, Bucharest, Romania, Theta Foundation, 2000.
- [13] A. Kumjian and D. Pask, “Actions of \mathbb{Z}^k associated to higher rank graphs,” *Ergodic Theory and Dynamical Systems*, vol. 23, no. 4, pp. 1153–1172, 2003.
- [14] G. A. Elliott, G. Gong, and L. Li, “Approximate divisibility of simple inductive limit C^* -algebras,” *Contemporary Mathematics*, vol. 228, pp. 87–97, 1998.
- [15] J. Renault, *A Groupoid Approach to C^* -Algebras*, vol. 793 of *Lecture Notes in Mathematics*, Springer, Berlin, Germany, 1980.
- [16] S. Zhang, “Certain C^* -algebras with real rank zero and their corona and multiplier algebras. I,” *Pacific Journal of Mathematics*, vol. 155, no. 1, pp. 169–197, 1992.
- [17] J.-L. Tu, “La conjecture de Baum-Connes pour les feuilletages moyennables,” *K-Theory in the Mathematical Sciences*, vol. 17, no. 3, pp. 215–264, 1999.
- [18] J. Renault, “The Radon-Nikodym problem for approximately proper equivalence relations,” *Ergodic Theory and Dynamical Systems*, vol. 25, no. 5, pp. 1643–1672, 2005.
- [19] A. Kumjian and J. Renault, “KMS states on C^* -algebras associated to expansive maps,” *Proceedings of the American Mathematical Society*, vol. 134, no. 7, pp. 2067–2078, 2006.
- [20] P. S. Muhly and D. P. Williams, “Continuous trace groupoid C^* -algebras,” *Mathematica Scandinavica*, vol. 66, no. 2, pp. 231–241, 1990.
- [21] A. An Huef, I. Raeburn, and D. P. Williams, “Properties preserved under Morita equivalence of C^* -algebras,” *Proceedings of the American Mathematical Society*, vol. 135, no. 5, pp. 1495–1503, 2007.
- [22] K. Thomsen, “ C^* -algebras of homoclinic and heteroclinic structure in expansive dynamics,” *Memoirs of the American Mathematical Society*, vol. 206, no. 970, 2010.

- [23] G. Gong, "On the classification of simple inductive limit C^* -algebras—I. The reduction theorem," *Documenta Mathematica*, vol. 7, pp. 255–461, 2002.
- [24] B. Blackadar, A. Kumjian, and M. Rørdam, "Approximately central matrix units and the structure of noncommutative tori," *K-Theory*, vol. 6, no. 3, pp. 267–284, 1992.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

