

# Research Article Construction of Nodal Bubbling Solutions for the Weighted Sinh-Poisson Equation

# Yibin Zhang<sup>1</sup> and Haitao Yang<sup>2</sup>

<sup>1</sup> College of Science, Nanjing Agricultural University, Nanjing 210095, China
 <sup>2</sup> Department of Mathematics, Zhejiang University, Hangzhou 310027, China

Correspondence should be addressed to Yibin Zhang; yibin10201029@njau.edu.cn

Received 1 June 2013; Accepted 27 August 2013

Academic Editor: Marco Donatelli

Copyright © 2013 Y. Zhang and H. Yang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We consider the weighted sinh-Poisson equation  $\Delta u + 2\varepsilon^2 |x|^{2\alpha} \sinh u = 0$  in  $B_1(0)$ , u = 0 on  $\partial B_1(0)$ , where  $\varepsilon > 0$  is a small parameter,  $\alpha \in (-1, +\infty) \setminus \{0\}$ , and  $B_1(0)$  is a unit ball in  $\mathbb{R}^2$ . By a constructive way, we prove that for any positive integer *m*, there exists a nodal bubbling solution  $u_{\varepsilon}$  which concentrates at the origin and the other *m*-points  $\tilde{q}_l = (\lambda \cos (2\pi(l-1)/m), \lambda \sin (2\pi(l-1)/m))$ ,  $l = 2, \ldots, m + 1$ , such that as  $\varepsilon \to 0$ ,  $2\varepsilon^2 |x|^{2\alpha} \sinh u_{\varepsilon} \to 8\pi(1 + \alpha)\delta_0 + \sum_{l=2}^{m+1} 8\pi(-1)^{l-1}\delta_{\tilde{q}_l}$ , where  $\lambda \in (0, 1)$  and *m* is an odd integer with  $(1 + \alpha)(m + 2) - 1 > 0$ , or *m* is an even integer. The same techniques lead also to a more general result on general domains.

### 1. Introduction

We are concerned with stationary Euler equations for an incompressible, homogeneous, and inviscid fluid on a bounded, smooth planar domain  $\Omega$ , consider

where **w** is the velocity field, *p* is the pressure, and  $\nu$  is the unit outer normal vector to  $\partial\Omega$ . Let us introduce the vorticity  $\omega = \text{curl } \mathbf{w}$ . By applying the curl operator to the first equation in (1), we have

$$\mathbf{w} \cdot \nabla \boldsymbol{\omega} = \mathbf{0}, \quad \text{in } \boldsymbol{\Omega}. \tag{2}$$

On the other hand, the second equation is equivalent to rewriting the velocity field **w** as

$$\mathbf{w} = \left(\nabla \psi\right)^{\perp} := \left(-\partial_{x_2}\psi, \partial_{x_1}\psi\right). \tag{3}$$

In return, the vorticity  $\omega$  is expressed as  $\omega = -\Delta \psi$  in term of  $\psi$ , the so-called stream function. Now, the ansatz  $\omega = \omega(\psi)$  guarantees that (2) is also automatically satisfied, and then the Euler equations reduce to solving the Dirichlet elliptic problem as follows

$$-\Delta \psi = \omega (\psi), \quad \text{in } \Omega,$$
  
$$\psi = 0, \quad \text{on } \partial \Omega.$$
 (4)

Over the past decades, some vortex-type configurations for planar stationary turbulent Euler flows have aroused wide concern among the people (see [1–3]). Many functions  $\omega(\psi)$ have been chosen in the physical perspective to describe turbulent Euler flows with vorticity  $\omega$  concentrated in small "blobs". For example, on the basis of the statistical mechanics approach, Joyce and Montgomery proposed the Stuart vortex pattern  $\omega(\psi) = \varepsilon^2 e^{\psi}$  with a small positive parameter  $\varepsilon$ to describe positive vortices (see [4–8]). Meanwhile, they also proposed the Mallier-Maslowe vortex pattern  $\omega(\psi) = 2\varepsilon^2 \sinh \psi$  to describe coexisting positive and negative vortices (see [9, 10]). Recently, Tur and Yanovsky in [11] have used the singular ansatz  $\omega(\psi) = \varepsilon^2 e^{\psi} - 4\pi N \delta_q$  to describe vortex patterns of necklace type with N+1-fold symmetry in rational shear flow, where  $N \in \mathbb{N}$  and  $\delta_q$  denotes the Dirac mass at  $q \in \Omega$ . Now, we adopt another new singular ansatz in [12]  $\omega(\psi) = 2\varepsilon^2 |x|^{2\alpha} \sinh \psi$  with  $\alpha \in (-1, +\infty) \setminus \{0\}$  to study the corresponding vortices with concentrated vorticity. To do it, we hope to investigate the effect of the presence of the weight  $|x|^{2\alpha}$  on the existence of nodal bubbling solutions for the weighted sinh-Poisson equation as follows:

$$\Delta u + 2\varepsilon^2 |x|^{2\alpha} \sinh u = 0, \quad \text{in } B_1(0),$$
  
$$u = 0, \quad \text{on } \partial B_1(0),$$
  
(5)

where  $\varepsilon > 0$  is a small parameter,  $\alpha \in (-1, +\infty) \setminus \{0\}$ , and  $B_1(0)$  is a unit ball in  $\mathbb{R}^2$ .

Let us first recall the two-dimensional sinh-Poisson equation as follows:

$$\Delta u + 2\varepsilon^2 \sinh u = 0, \tag{6}$$

which relates to various dynamics of vorticity with respect to geophysical flows, rotating and stratified fluids, and fluid layers excited by electromagnetic forces (see [13–15] and the references therein) and the geometry of constant mean curvature surfaces studied by many works (see [16-20] and the references therein). Recently, the asymptotic behavior of solutions to (6) has been studied on a closed Riemann surface in [21, 22], and the authors applied the so-called "Pohozaev identity" and "symmetrization method," respectively, to show that there possibly exist two different types of blow up for a family of solutions to (6). Furthermore, Grossi and Pistoia in [23] exhibited sign-changing multiple blow-up phenomena for the Dirichlet problem (6), more precisely, if  $0 \in \Omega$  and  $\Omega$ is symmetric with respect to the origin, for any integer k if  $\varepsilon$  is small enough, there exists a family of solutions to (6), which blows up at the origin, whose positive mass is  $4\pi k(k-1)$  and negative mass is  $4\pi k(k+1)$ . This gives a complete answer to an open problem in [21]. Besides, for a similar equation, precisely the Neumann sinh-Gordon equation on a unit ball, Esposito and Wei in [24] also constructed a family of solutions with a multiple blow up at the origin. On the other hand, Bartolucci and Pistoia in [25] tried to construct blow-up solutions of (6) with Dirichlet boundary condition, and proved that for  $\varepsilon > 0$ small enough, there exist at least two pairs of solutions, which change sign exactly once, concentrate in the domain, and whose nodal lines intersect the boundary. Furthermore, Bartsch et al. in [26] obtained the existence of changing sign solutions for this equation on an arbitrary bounded domain  $\Omega$ , which have three and four alternate-sign concentration points. In particular, when  $\Omega$  has an axial symmetry they proved for each  $N \in \mathbb{N}$  there exists a nodal bubbling solution, which changes sign N-times and whose alternate-sign concentration points align on the symmetric axis of  $\Omega$ . For (6) with Neumann boundary condition, Wei et al. in [27] constructed a family of solutions concentrating positively and negatively in the domain and its boundary. As for the presence of the weight  $|x|^{2\alpha}$ , the authors in [12] showed that there exists a family of nodal bubbling solutions  $u_{s}$  to (5) only

involving  $0 < \alpha \notin \mathbb{N}$ , such that  $2\varepsilon^2 |x|^{2\alpha} \sinh u_{\varepsilon}$  not only develops many positive and negative Dirac deltas with weight  $8\pi$  and  $-8\pi$ , respectively, but also a Dirac data with weight  $8\pi(1+\alpha)$  at the origin.

We mention that an analogous blow-up analysis can be applied to multiple blow-up solutions for the Liouville equation with or without singular data as follows:

$$-\Delta u = \varepsilon^2 e^u - 4\pi \sum_{i=1}^M \alpha_i \delta_{q_i}, \quad \text{in } \Omega,$$
  
$$u = 0, \quad \text{on } \partial\Omega,$$
 (7)

where  $\Omega$  is a smooth bounded domain in  $\mathbb{R}^2$ ,  $M \ge 0$ ,  $\alpha_i > 0$ -1,  $q_i \in \Omega$ ,  $\delta_{q_i}$  defines the Dirac mass at  $q_i$ , and  $\varepsilon > 0$ is a small parameter. For the past decades, the asymptotic analysis for blow up solutions of problem (7) has been deeply studied in the vast literature (see [28-33] and the reference therein), which exhibits the quantiaztion properties of the weak limit of  $\varepsilon^2 e^u$  as  $\varepsilon \to 0^1$  if  $\varepsilon^2 \int_{\Omega} e^u$  remains uniformly bounded, and characterizes the location of concentration points as critical points of a functional in terms of Green's function. Reciprocally, an obvious problem is how to construct solutions of (7) with these properties. In [34, 35], the authors use the asymptotic analysis to construct solutions with multiple interior concentration points for (7) with  $M \ge$ 0 and 0 <  $\alpha_i \notin \mathbb{N}$ . More generally, by a constructive way, similar results related to  $-1 < \alpha_i$  can also be obtained in [36-40] under some milder notions of stability of critical points. In particular, when  $M \ge 1$  and  $\alpha_i$ s are positive numbers, D'Aprile in [37] recently established a family of solutions to (7) consisting of  $\widetilde{N} := \sum_{i=1}^{M} N_i$  blow up points in  $\Omega \setminus \{q_1, \ldots, q_M\}$  as long as  $N_i < 1 + \alpha_i$  for any *i*, provided that the weights  $\alpha_i$  avoid the integers 1, 2, ...,  $\tilde{N} - 1$ , and so the result of del Pino et al. in [39] can be extended to the case of several singular sources.

In this paper, we will continue the study of the existence of solutions to (5). We prove that there exists a family of solutions  $u_{\varepsilon}$  concentrating positively and negatively at the origin and outside the origin as long as  $\alpha \in (-1, +\infty) \setminus \{0\}$ . Concerning the sign-changing concentration at the origin and outside the origin, if we introduce the function  $f(\lambda) : (0, 1) \rightarrow \mathbb{R}$  defined as

$$f(\lambda) = -16\pi \left\{ \sum_{l,j=2,l\neq j}^{m+1} (-1)^{l+j} \times \left[ \frac{1}{2} \log \left( \lambda^4 + 1 - 2\lambda^2 \times \cos \left( 2\pi \frac{l-j}{m} \right) \right) - \log \left| 2\sin \left( \pi \frac{l-j}{m} \right) \right| \right] + \left[ (1+\alpha) (m-2A) - A^2 \right] \log \lambda + m \log \left( 1 - \lambda^2 \right) \right\},$$
(8)

with  $A = \sum_{l=2}^{m+1} (-1)^{l+1}$ , our main result for problem (5) can be stated as follows.

**Theorem 1.** For any positive integer *m*, there exists a nodal solution  $u_{\varepsilon}$  to problem (5) which concentrates at the origin and the other *m*-points  $\tilde{q}_l = (\lambda_0 \cos(2\pi(l-1)/m), \lambda_0 \sin(2\pi(l-1)/m)), l = 2, ..., m + 1$ , such that as  $\varepsilon \to 0$ ,

$$2\varepsilon^2 |x|^{2\alpha} \sinh u_{\varepsilon} \rightharpoonup 8\pi (1+\alpha) \,\delta_0 + \sum_{l=2}^{m+1} 8\pi (-1)^{l-1} \delta_{\tilde{q}_l}, \quad (9)$$

weakly in the sense of measures in  $\overline{B_1(0)}$ , where  $\lambda_0$  is an absolute minimum point of  $f(\lambda)$  in (0, 1), *m* is an odd integer with  $(1 + \alpha)(m + 2) - 1 > 0$ , or *m* is an even integer. Moreover, for any  $\delta > 0$ ,  $u_{\varepsilon}$  remains uniformly bounded on  $B_1(0) \setminus (B_{\delta}(0) \cup \bigcup_{l=2}^{m+1} B_{\delta}(\tilde{q}_l))$ , and as  $\varepsilon \to 0$ ,

$$\sup_{B_{\delta}(0)} u_{\varepsilon} \longrightarrow +\infty,$$

$$\sup_{B_{\delta}(\tilde{q}_{l})} (-1)^{l-1} u_{\varepsilon} \longrightarrow +\infty \quad \forall l = 2, \dots, m+1.$$
(10)

Theorem 1 is based on a constructive way which also works for the more generally weighted sinh-Poisson equation as follows:

$$\Delta u + 2\varepsilon^2 c(x) \prod_{i=1}^M |x - q_i|^{2\alpha_i} \sinh u = 0, \quad \text{in } \Omega,$$

$$u = 0, \quad \text{on } \partial\Omega,$$
(11)

for  $\varepsilon > 0$  small, where  $\Omega$  is a bounded smooth domain in  $\mathbb{R}^2$ , M = n + k with  $M \ge 0$ ,  $\{q_1, \ldots, q_M\}$  are different singular sources in  $\Omega$ ,  $\{\alpha_1, \ldots, \alpha_n\} \in (-1, +\infty) \setminus (\mathbb{N} \cup \{0\})$ ,  $\{\alpha_{n+1}, \ldots, \alpha_M\} \in \mathbb{N}$ , and  $c : \Omega \to \mathbb{R}$  is a continuous function such that  $c(q_i) > 0$  for any  $i = 1, \ldots, M$ .

To further state our results, we need to introduce some notations. Let G(x, y) be Green's function of  $\Delta_x$  such that for  $y \in \Omega, -\Delta_x G(x, y) = \delta_y(x)$  in  $\Omega$  and G(x, y) = 0 on  $\partial\Omega$ , and let H(x, y) be its regular part defined as  $H(x, y) = G(x, y) + (1/2\pi) \log |x-y|$ . Besides, let us denote  $S(x) = \prod_{i=1}^{M} |x-q_i|^{2\alpha_i}$ ,  $\Omega' = \{x \in \Omega : c(x) > 0\}, \Gamma = \{q_1, \dots, q_n\}, J_1 = \{1, \dots, n\}, J_2 = \{n+1, \dots, n+k\}, J_3 = \{n+k+1, \dots, n+k+m\}, \text{and } \Delta_{k+m} = \{(q_{n+1}, \dots, q_{n+k+m}) : q_i = q_j \text{ for some } i \neq j\}$ , where  $m \ge 0$  is an integer. In what follows, we fix *n* different points  $q_i, i \in J_1$ , and define

$$\varphi_{k,m}^{n}(q) = -\sum_{i \in J_{2} \cup J_{3}} d_{i} \left\{ \sum_{j \in J_{1}} a_{i}a_{j}d_{j}G\left(q_{i},q_{j}\right) + \frac{1}{2}d_{i}H\left(q_{i},q_{i}\right) + \sum_{j \in J_{2} \cup J_{3}, j \neq i} \frac{1}{2}a_{i}a_{j}d_{j}G\left(q_{i},q_{j}\right) \right\}$$
$$-\sum_{i \in \bigcup_{l=1}^{3} J_{3}} d_{i}\log c_{i}\left(q_{i}\right),$$
(12)

which is well defined on the following domain:

$$\Lambda_{k,m} = \frac{\left(\Omega' \setminus \Gamma\right)^{k+m}}{\Delta_{k+m}},\tag{13}$$

where  $q = (q_{n+1}, \dots, q_{M+m}), \alpha_i = 0$  for  $i \in J_3, c_i(x) = c(x)S(x)/|x - q_i|^{2\alpha_i}$  for  $i \in \bigcup_{l=1}^3 J_l, d_i = 8\pi(1 + \alpha_i)$  for  $i \in \bigcup_{l=1}^3 J_l$ , and  $a_i = \pm 1$  for  $i \in \bigcup_{l=1}^3 J_l$ .

Definition 2 (see [41]). We say that  $q^*$  is a  $C^0$ -stable critical point of  $\varphi_{k,m}^n$  in  $\Lambda_{k,m}$  if for any sequence of functions  $\psi_j$  such that  $\psi_j \to \varphi_{k,m}^n$  uniformly on the compact subsets of  $\Lambda_{k,m}, \psi_j$  has a critical point  $\xi_j$  such that  $\psi_j(\xi_j) \to \varphi_{k,m}^n(q^*)$ .

In particular, if  $q^*$  is a strict local maximum or minimum point of  $\varphi_{k,m}^n$ ,  $q^*$  is a  $C^0$ -stable critical point of  $\varphi_{k,m}^n$ .

**Theorem 3.** Assume that  $\{n, k, m\} \in \mathbb{N} \cup \{0\}$  and  $q^* = (q_{n+1}^*, \ldots, q_{M+m}^*)$  is a  $C^0$ -stable critical point of  $\varphi_{k,m}^n$  in  $\Lambda_{k,m}$  with  $k + m \ge 1$ . Then, for any sufficiently small  $\varepsilon > 0$ , there exists different points  $q_{\varepsilon,l} \in \Omega' \setminus \Gamma$ ,  $l \in J_2 \cup J_3$ , away from  $\partial \Omega \cup \Gamma$ , so that problem (11), for  $q_l = q_{\varepsilon,l}, l \in J_2$ , has a nodal solution  $u_{\varepsilon}$  such that as  $\varepsilon \to 0$ ,

$$2\varepsilon^{2}c(x)\prod_{i=1}^{M}|x-q_{i}|^{2\alpha_{i}}\sinh u_{\varepsilon}$$

$$-\sum_{J_{2}\cup J_{3}}a_{l}d_{l}\delta_{q_{\varepsilon,l}} \rightharpoonup \sum_{J_{1}}a_{l}d_{l}\delta_{q_{l}},$$
(14)

weakly in the sense of measures in  $\overline{\Omega}$ . Moreover, up to a subsequence, there exists  $\tilde{q} = (\tilde{q}_{n+1}, \dots, \tilde{q}_{M+m}) \in \Lambda_{k,m}$  such that

$$\varphi_{k,m}^{n}\left(\tilde{q}\right) = \varphi_{k,m}^{n}\left(q^{*}\right),$$

$$d\left(q_{\varepsilon,l},\tilde{q}_{l}\right) \longrightarrow 0, \quad \forall l \in J_{2} \cup J_{3}, \text{ as } \varepsilon \longrightarrow 0.$$
(15)

Besides,  $u_{\varepsilon}$  remains uniformly bounded on  $\overline{\Omega} \setminus (\bigcup_{i \in J_1} B_{\delta}(q_i)) \cup (\bigcup_{i \in J_2 \cup J_3} B_{\delta}(\tilde{q}_i))$  for any  $\delta > 0$ , and for any points  $q_l, l \in J_1$ , and  $\tilde{q}_l, l \in J_2 \cup J_3$ , as  $\varepsilon \to 0$ ,

$$\sup_{\overline{\Omega}\cap B_{\delta}(q_{l})} a_{l}u_{\varepsilon} \longrightarrow +\infty, \qquad \sup_{\overline{\Omega}\cap B_{\delta}(\overline{q}_{l})} a_{l}u_{\varepsilon} \longrightarrow +\infty.$$
(16)

Note that for the case n = 1 and k = 0 (or n = k = 0), Theorem 3 was partly proved in [12] (or [25]) only when c(x) = 1 and  $\alpha_l \in (0, +\infty) \setminus \mathbb{N}$ ,  $l \in J_1$ . In contrast with the results of [12, 25], this theorem provides a more complex concentration phenomenon involving the existence of changingsign solutions for problem (11) with both positive and negative bubbles near the singular sources  $q_l, l \in J_1 \cup J_2$ , and some other discrete points. Unlike the concentration set in [12] only contains singular sources  $q_l$  with  $\alpha_l \in (0, +\infty) \setminus \mathbb{N}$  and  $l \in J_1$ , and no singular source points in the domain, our concentration set also contains some singular sources  $q_l$  with  $\alpha_l \in$ (-1,0) and  $l \in J_1$ , except for singular sources  $q_l, l \in J_2$ , where concentration points and singular sources coincide at the limit. As for the latter exception, which till now is a similar but very simple concentration phenomenon it appears only in [38] for the study of the Liouville equation with a singular source of integer multiplicity.

In order to obtain multiple sign-changing blow up solutions of problem (11), we use a Lyapunov-Schmidt finitedimensional reduction scheme and convert the problem into a finite-dimensional one, for a suitable asymptotic reduced energy, related to  $\varphi_{k,m}^n$  in (12). Thus, a stable critical point of  $\varphi_{k,m}^{n}$  leads to the existence of multiple sign-changing blow up solutions to (11). However, in view of different signs of Green's functions in (12), it seems very difficult to find out a stable critical point of  $\varphi_{k,m}^n$  for a general bounded domain  $\Omega$ . A simple approach can help us to overcome this difficulty by imposing the very strong symmetry condition on the domain of the problem, namely, we use the symmetry of the unit ball  $B_1(0)$  to reduce the problem of finding solutions of (5) to that of finding an absolute minimum point of  $f(\lambda)$  defined in (8), and so we get the existence of nodal bubbling solutions for (5) in Theorem 1. On the other hand, motivated by the obtained results in [23], we believe that Theorem 1 should be valid for a general domain than a unit ball. More precisely, we suspect that, if  $0 \in \Omega$  and  $\Omega$  is symmetric with respect to the origin, it is possible to construct a family of sign-changing blow up solutions whose maxima and minima are located alternately at the origin and the vertices of a regular polygon, and so Theorem 1 will be a consequence of this general result.

It is important to point out that to prove the above results, we need to use classification solutions of the Liouville-type equation to construct approximate solutions of problem (5) (or (11)) as follows:

$$\Delta u + |x|^{2\alpha} e^u = 0, \quad \text{in } \mathbb{R}^2,$$

$$\int_{\mathbb{R}^2} |x|^{2\alpha} e^u < +\infty, \quad \alpha \in (-1, +\infty).$$
(17)

In complex notations, a complete classification of the solutions of (17) takes the following form:

$$u(z) = \log \frac{8(1+\alpha)^2 \tilde{\delta}^2}{\left(\tilde{\delta}^2 + |z^{\alpha+1} - \xi|^2\right)^2},$$
(18)

where  $\delta > 0, \xi \in \mathbb{C}$  if  $\alpha \in \mathbb{N} \cup \{0\}$  and  $\xi = 0$  if  $\alpha \in (-1, +\infty) \setminus (\mathbb{N} \cup \{0\})$  (see [33, 42–44]). Using classification solutions scaled up and projected to satisfy the Dirichlet boundary condition up to a right order, the approximate solutions can be built up as a summation of these initial approximations with some suitable signs. Thus, the nodal bubbling solutions can be constructed as a small additive perturbation of these approximations through the so-called "localized energy method," which combined the Lyapunov-Schmidt finite dimensional reduction and variational techniques. Here, we follow [12, 25], but we will overcome some of the difficulties that the nodal concentration phenomenon brings by delicate analysis.

#### 2. Construction of the Approximate Solution

In this section, we will provide a first approximation for the solutions of problem (11). Given a sufficiently small but fixed

number  $\delta > 0$ , let us first fix *n* different points  $q_i$ ,  $i \in J_1$ , and assume that points  $q = (q_{n+1}, \ldots, q_{n+k+m}) \in \Lambda_{k,m}(\delta)$ , where

$$\Lambda_{k,m}\left(\delta\right):=\left\{q\in\Lambda_{k,m}:\min_{i\in J_{2}\cup J_{3}}d\left(q_{i},\partial\left(\Omega'\setminus\Gamma\right)\right)\geq 2\delta,$$
$$\min_{i,j\in J_{2}\cup J_{3},i\neq j}d\left(q_{i},q_{j}\right)\geq 2\delta\right\}.$$
(19)

Suppose that  $\mu_i$ ,  $i \in \bigcup_{l=1}^3 J_l$ , are positive numbers to be chosen later, we define

$$\rho_{i} = \varepsilon^{(1/(\alpha_{i}+1))}, \qquad v_{i} = \mu_{i}^{(1/(\alpha_{i}+1))},$$

$$\mu_{i}(x) = \log \frac{8\mu_{i}^{2}(1+\alpha_{i})^{2}}{c_{i}(q_{i})\left[\mu_{i}^{2}\varepsilon^{2} + |x-q_{i}|^{2(1+\alpha_{i})}\right]^{2}}.$$
(20)

The ansatz is

$$U_{q}(x) = \sum_{i \in \bigcup_{l=1}^{3} J_{l}} a_{l} P u_{i}(x), \qquad (21)$$

where  $a_i = \pm 1, P : H^1(\Omega) \to H^1_0(\Omega)$  is a linear operator such that for any  $u \in H^1(\Omega)$ ,  $\Delta P u = \Delta u$  in  $\Omega$ , and P u = 0 on  $\partial \Omega$ . By harmonicity, we easily get that

$$Pu_{i}(x) = u_{i}(x) + d_{i}H(x,q_{i}) - \log \frac{8\mu_{i}^{2}(1+\alpha_{i})^{2}}{c_{i}(q_{i})} + O(\rho), \text{ in } C^{1}(\overline{\Omega}),$$
(22)

$$Pu_{i}(x) = d_{i}G(x,q_{i}) + O(\rho), \quad \text{in } C_{\text{loc}}\left(\overline{\Omega} \setminus \{q_{i}\}\right), \quad (23)$$

where  $\rho := \max\{\rho_i : i \in \bigcup_{l=1}^3 J_l\}.$ 

Consider that the scaling of solution to problem (11) is as follows:

$$v(y) = u(\varepsilon y), \quad \forall y \in \overline{\Omega}_{\varepsilon},$$
 (24)

where  $\Omega_{\varepsilon} = (1/\varepsilon)\Omega$ , then *v* satisfies

$$\Delta v + 2\varepsilon^4 c(\varepsilon y) S(\varepsilon y) \sinh v = 0, \quad \text{in } \Omega_{\varepsilon},$$

$$v = 0, \quad \text{on } \partial \Omega_{\varepsilon}.$$
(25)

We will seek solutions of problem (25) of the form  $v = V_q + \phi$ , where

$$V_q(y) = U_q(\varepsilon y) = \sum_{i \in \bigcup_{l=1}^3 J_l} a_i P u_i(\varepsilon y), \quad \forall y \in \overline{\Omega}_{\varepsilon}.$$
 (26)

In terms of  $\phi$ , problem (11) (or (25)) becomes

$$L(\phi) := \Delta \phi + W_q \phi$$
  
= - [R<sub>q</sub> + N(\phi)], in \Omega\_\varepsilon, (27)  
\phi = 0, on \frac{\partial\Omega\_\varepsilon}{\phi},

where

$$W_q = 2\varepsilon^4 c(\varepsilon y) S(\varepsilon y) \cosh V_q, \qquad (28)$$

 $R_q$  is the "error term":

$$R_q = \Delta V_q + 2\varepsilon^4 c \left(\varepsilon y\right) S\left(\varepsilon y\right) \sinh V_q, \tag{29}$$

and  $N(\phi)$  denotes the following "nonlinear term":

$$N(\phi) = 2\varepsilon^4 c(\varepsilon y) S(\varepsilon y) \times \left[\sinh\left(V_q + \phi\right) - \sinh V_q - \phi \cosh V_q\right].$$
(30)

Finally, in order to make  $R_q$  sufficiently small, we choose the parameters  $\mu_i, i \in \bigcup_{l=1}^3 J_l$ , as

$$\log \frac{8\mu_{i}^{2}(1+\alpha_{i})^{2}}{c_{i}\left(q_{i}\right)} = d_{i}H\left(q_{i},q_{i}\right) + \sum_{j\neq i}a_{i}a_{j}d_{j}G\left(q_{i},q_{j}\right), \quad (31)$$

so that from Appendix, we have

$$W_{q} = \begin{cases} \left(\frac{\varepsilon}{v_{i}\rho_{i}}\right)^{2} |z_{i}|^{2\alpha_{i}} \begin{cases} \frac{8(1+\alpha_{i})^{2} \left[1+O\left(\rho_{i} |z_{i}|\right)+O\left(\rho\right)\right]}{\left[1+|z_{i}|^{2(1+\alpha_{i})}\right]^{2}} + O\left(\varepsilon^{4}\right) \end{cases}, & \text{if } |z_{i}| \leq \delta(v_{i}\rho_{i})^{-1}, \\ O\left(\varepsilon^{4}\right), & \text{otherwise,} \end{cases}$$
(32)

$$R_{q} = \begin{cases} \left(\frac{\varepsilon}{v_{i}\rho_{i}}\right)^{2} \left|z_{i}\right|^{2\alpha_{i}} \begin{cases} \frac{8a_{i}(1+\alpha_{i})^{2}\left[O\left(\rho_{i}\left|z_{i}\right|\right)+O\left(\rho\right)\right]}{\left[1+\left|z_{i}\right|^{2(1+\alpha_{i})}\right]^{2}} + O\left(\varepsilon^{4}\right) \end{cases}, & \text{if } \left|z_{i}\right| \leq \delta\left(v_{i}\rho_{i}\right)^{-1}, \\ O\left(\varepsilon^{4}\right), & \text{otherwise,} \end{cases}$$
(33)

 $z_i := (1/v_i \rho_i)(\varepsilon y - q_i).$ 

# 3. The Linearized Problem and the Nonlinear Problem

In this section, we will first prove the bounded invertibility of the linearized operator L under suitable orthogonality conditions. Let us define

$$L_{i}(\phi) = \Delta \phi + \frac{8(1+\alpha_{i})^{2}|z|^{2\alpha_{i}}}{\left[1+|z|^{2(1+\alpha_{i})}\right]^{2}}\phi \quad \forall i \in \bigcup_{l=1}^{3} J_{l},$$

$$z_{i0} = \frac{1}{\mu_{i}} \frac{|z|^{2(1+\alpha_{i})}-1}{|z|^{2(1+\alpha_{i})}+1} \quad \forall i \in \bigcup_{l=1}^{3} J_{l},$$

$$z_{i1} = \frac{4\operatorname{Re}\left(z^{1+\alpha_{i}}\right)}{|z|^{2(1+\alpha_{i})}+1} \quad \forall i \in \bigcup_{l=2}^{3} J_{l},$$

$$z_{i2} = \frac{4\operatorname{Im}\left(z^{1+\alpha_{i}}\right)}{|z|^{2(1+\alpha_{i})}+1} \quad \forall i \in \bigcup_{l=2}^{3} J_{l}.$$
(34)

The key fact to develop a satisfactory solvability theory for the operator *L* is that *L* formally approaches  $L_i$  under dilations and translations, and any bounded solution of  $L_i(\phi) = 0$ in  $\mathbb{R}^2$  is a linear combination of  $z_{i0}$ ,  $z_{i1}$ , and  $z_{i2}$  for  $i \in J_2 \cup J_3$  (see [34, 45, 46]), or proportional to  $z_{i0}$  for  $i \in J_1$  (see [35, 36, 41]). Let us denote

$$Z_{i0}(y) := z_{i0}(z_i), \quad \forall i \in \bigcup_{l=1}^{3} J_l,$$

$$Z_{ij}(y) := z_{ij}(z_i), \quad \forall i \in \bigcup_{l=2}^{3} J_l, \ j = 1, 2,$$

$$\theta_i(y) := \arg(z_i), \quad \forall i \in J_2,$$

$$\chi_i(y) := \chi(|z_i|), \quad \forall i \in \bigcup_{l=1}^{3} J_l,$$
(35)

where  $\chi(r)$  is a smooth, nonincreasing cut-off function such that for a large but fixed number  $R_0 > 0$ ,  $\chi(r) = 1$  if  $r \le R_0$ , and  $\chi(r) = 0$  if  $r \ge R_0 + 1$ .

Additionally, we consider the following Banach space:

$$\mathscr{C}_* = \left\{ h : \|h\|_* < +\infty \right\},\tag{36}$$

with the norm

$$\|h\|_{*} = \sup_{y \in \overline{\Omega}_{\varepsilon}} \left( \|h(y)\| \times \left( \varepsilon^{2} + \sum_{\alpha_{i} < 0, i \in J_{1}} \left( \frac{\varepsilon}{\nu_{i} \rho_{i}} \right)^{2} \right)$$

$$\times \frac{\left|z_{i}\right|^{2\alpha_{i}}}{\left(1+\left|z_{i}\right|\right)^{4+2\alpha+2\alpha_{i}}} + \sum_{\alpha_{i}\geq0, i\in J_{1}\cup J_{2}\cup J_{3}} \left(\frac{\varepsilon}{\nu_{i}\rho_{i}}\right)^{2} \times \frac{1}{\left(1+\left|z_{i}\right|\right)^{4+2\alpha}}\right)^{-1}\right),$$

$$(37)$$

where  $\alpha \in (-1, \alpha_0)$  and  $\alpha_0 := \min\{\alpha_i : i \in \bigcup_{i=1}^3 J_i\}.$ 

Given that  $h \in \mathcal{C}_*$  and  $q \in \Lambda_{k,m}(\delta)$ , we consider the linear problem of finding a function  $\phi$  and scalars  $c_{ij}$ ,  $i \in J_2 \cup J_3$ , j = 1, 2, such that

$$L(\phi) = h + \sum_{j=1}^{2} \sum_{i=n+1}^{n+k+m} c_{ij} \chi_i \cos^2\left(\frac{1}{2}\alpha_i \theta_i\right) Z_{ij}, \quad \text{in } \Omega_{\varepsilon},$$
  
$$\phi = 0, \quad \text{on } \partial \Omega_{\varepsilon},$$
  
$$\int_{\Omega_{\varepsilon}} \chi_i \cos^2\left(\frac{1}{2}\alpha_i \theta_i\right) Z_{ij} \phi dy = 0, \quad \forall i \in J_2 \cup J_3, \ j = 1, 2.$$
(38)

Our main interest in this problem is its bounded solvability for any  $h \in \mathcal{C}_*$ , uniform in small  $\varepsilon$  and points  $q \in \Lambda_{k,m}(\delta)$ , as the following result states.

**Proposition 4.** There exist positive numbers  $\varepsilon_0$  and C such that for any  $h \in \mathcal{C}_*$ , there is a unique solution  $\phi \in L^{\infty}(\Omega_{\varepsilon})$ , scalars  $c_{ij} \in \mathbb{R}, i \in J_2 \cup J_3, j = 1, 2$ , to problem (38) for all  $\varepsilon < \varepsilon_0$  and points  $q \in \Lambda_{k,m}(\delta)$ , which satisfies

$$\|\phi\|_{\infty} \le C\left(\log\frac{1}{\varepsilon}\right) \|h\|_{*}.$$
(39)

Moreover, the map  $q' \mapsto \phi$  is  $C^1$ , precisely for  $l \in J_2 \cup J_3$ , s = 1, 2,

$$\left\|\partial_{q_{l,s}'}\phi\right\|_{\infty} \le C \frac{\varepsilon}{\nu_l \rho_l} \left(\log \frac{1}{\varepsilon}\right)^2 \|h\|_*, \tag{40}$$

where  $q' := (1/\varepsilon)q = ((1/\varepsilon)q_{n+1}, \dots, (1/\varepsilon)q_{n+k+m}).$ 

We begin by stating a priori estimate for solutions of (38) satisfying orthogonality conditions with respect to  $Z_{i0}$ ,  $i \in J_1$ , and  $Z_{ij}$ ,  $i \in J_2 \cup J_3$ , j = 0, 1, 2.

**Lemma 5.** There exist positive numbers  $\varepsilon_0$  and C such that for any points  $q \in \Lambda_{k,m}(\delta)$ ,  $h \in \mathcal{C}_*$  and any solution  $\phi$  to the following equation:

$$L(\phi) = h, \quad in \ \Omega_{\varepsilon},$$
  

$$\phi = 0, \quad on \ \partial\Omega_{\varepsilon},$$
(41)

with the orthogonality conditions

$$\int_{\Omega_{\varepsilon}} \chi_i Z_{i0} \phi dy = 0, \quad \forall i \in J_1,$$

$$\int_{\Omega_{\varepsilon}} \chi_i \cos^2\left(\frac{1}{2}\alpha_i \theta_i\right) Z_{ij} \phi dy = 0,$$
  
$$\forall i \in J_2 \cup J_3, \ j = 0, 1, 2,$$
(42)

one has

$$\left\|\phi\right\|_{\infty} \le C \|h\|_*,\tag{43}$$

for all  $\varepsilon < \varepsilon_0$ .

*Proof.* We have the following steps.

*Step 1.* We first construct a suitable barrier. To realize it, we claim that for  $\varepsilon > 0$  small enough, there exist  $R_d > 0$  and

$$\psi: \overline{\Omega_{\varepsilon} \setminus \bigcup_{i=1}^{n+k+m} B_{i,R_d}} \longmapsto \mathbb{R}$$
(44)

positive and uniformly bounded so that

$$L(\psi) \le -\sum_{i=1}^{n+k+m} \left(\frac{\varepsilon}{\nu_i \rho_i}\right)^2 \frac{1}{\left|z_i\right|^{2\alpha+4}} - \varepsilon^2, \qquad (45)$$

where  $-1 < \alpha < \alpha_0$  and  $B_{i,R_d} = \{y \in \Omega_{\varepsilon} : |z_i| < R_d\}$ . Take

$$g_{0}(z) = \frac{|z|^{2(1+\alpha)} - 1}{|z|^{2(1+\alpha)} + 1}, \qquad R_{d} = \frac{1}{d} 3^{(1/2(1+\alpha))},$$

$$g_{1i}(y) = g_{0}(d|z_{i}|), \quad \text{where } |z_{i}| \ge R_{d}.$$
(46)

Thus, we have  $1/2 \le g_{1i}(y) \le 1$ , and

$$\Delta g_{1i}(y) = -\left(\frac{\varepsilon}{v_i \rho_i}\right)^2 d^2 \frac{8(1+\alpha)^2 |dz_i|^{2\alpha}}{\left[1+|dz_i|^{2(1+\alpha)}\right]^2} \times g_0(d|z_i|)$$

$$< -\left(\frac{\varepsilon}{v_i \rho_i}\right)^2 \frac{d^{-2(1+\alpha)}(1+\alpha)^2}{|z_i|^{2\alpha+4}}.$$
(47)

By (32),

$$W_q \leq C_1 \sum_{i=1}^{n+k+m} \left(\frac{\varepsilon}{v_i \rho_i}\right)^2 \frac{1}{|z_i|^{2\alpha_i+4}},$$

$$\overline{\forall y \in \Omega_{\varepsilon} \setminus \bigcup_{i=1}^{n+k+m} B_{i,R_d}}.$$
(48)

So, if *d* is taken small and fixed, by (47)-(48), it follows that for  $|z_i| \ge R_d$ ,

$$L(g_{1i})(y) < -\sum_{i=1}^{n+k+m} \left(\frac{\varepsilon}{\nu_i \rho_i}\right)^2 \frac{1}{\left|z_i\right|^{2\alpha+4}} < 0.$$
(49)

Let  $\tilde{g}_2$  be a positive, bounded solution of  $-\Delta \tilde{g}_2 = 1$  in  $\Omega$ and  $\tilde{g}_2 = 0$  on  $\partial \Omega$ . Set  $g_2(y) = \tilde{g}_2(\varepsilon y)$ . Then,  $g_2$  is a positive, uniformly bounded function in  $\Omega_{\varepsilon}$  such that

$$-\Delta g_2(y) = \varepsilon^2, \quad \text{in } \Omega_{\varepsilon},$$
  
$$g_2 = 0, \quad \text{on } \partial \Omega_{\varepsilon}.$$
 (50)

Consider

$$\psi(y) = C_2 \sum_{i=1}^{n+k+m} g_{1i}(y) + g_2(y),$$

$$(51)$$

$$\forall y \in \Omega_{\varepsilon} \setminus \bigcup_{i=1}^{n+k+m} B_{i,R_d},$$

where  $C_2 > 0$  is a sufficiently large constant. Obviously,  $\psi$  is a positive and uniformly bounded function. Moreover, in view of (48)–(50), it is easy to check that  $\psi$  satisfies the estimate (45), and the claim follows.

Step 2. Consider the following "inner norm":

$$\left\|\phi\right\|_{l} = \sup_{\bigcup_{i=1}^{n+k+m}\overline{B_{i,R_{d}}}} \left|\phi\right|.$$
(52)

Let us claim that there exists a constant  $C_3 > 0$ , such that

$$\|\phi\|_{\infty} \le C_3 \left(\|\phi\|_l + \|h\|_*\right).$$
(53)

Let us take

q

where  $C_4 > 0$  is chosen larger if necessary. Then, for  $y \in \Omega_{\varepsilon} \setminus \bigcup_{i=1}^{n+k+m} B_{i,R_d}$ ,

$$L\left(\tilde{\phi} \pm \phi\right)(y) \leq -C_{4}\left(\left\|\phi\right\|_{l} + \left\|h\right\|_{*}\right)$$

$$\times \left[\sum_{i=1}^{n+k+m} \left(\frac{\varepsilon}{v_{i}\rho_{i}}\right)^{2} \frac{1}{\left|z_{i}\right|^{2\alpha+4}} + \varepsilon^{2}\right] \quad (55)$$

$$\pm h\left(y\right)$$

$$\leq -\left|h\left(y\right)\right| \pm h\left(y\right) \leq 0,$$

for  $y \in \partial \Omega_{\varepsilon}$ ,

$$\left(\widetilde{\phi} \pm \phi\right)(y) = \widetilde{\phi}(y) > 0, \tag{56}$$

and for  $y \in \bigcup_{i=1}^{n+k+m} \partial B_{i,R_d}$ ,

$$\left(\widetilde{\phi} \pm \phi\right)(y) > \left\|\phi\right\|_{l} \pm \phi(y) \ge 0.$$
(57)

From the maximum principle, it follows that  $-\phi(y) \leq \widetilde{\phi}(y) \leq \phi(y)$  on  $\overline{\Omega_{\varepsilon} \setminus \bigcup_{i=1}^{n+k+m} B_{i,R_d}}$ , which provides the estimate (53).

7

Step 3. We prove the priori estimate (43) by contradiction. Let us assume the existence of a sequence  $\varepsilon_j \rightarrow 0$ , and points  $q^j = (q_{n+1}^j, \dots, q_{n+k+m}^j) \in \Lambda_{k,m}(\delta)$ , functions  $h_j$  with  $||h_j||_* \rightarrow 0$ , solutions  $\phi_j$  with  $||\phi_j||_{\infty} = 1$ , such that (41)-(5) hold. From the estimate (53),  $||\phi_j||_l \ge \kappa$  for some  $\kappa > 0$ , namely,  $\sup_{B_{i,R_d}} |\phi_j| \ge \kappa$  for some *i*. To simplify the notation, let us denote  $\varepsilon = \varepsilon_j$  and  $q_i = q_i^j$ . Set  $\hat{\phi}_j(z) = \phi_j((v_i\rho_i z + q_i)/\varepsilon)$  and  $\hat{h}_j(z) = h_j((v_i\rho_i z + q_i)/\varepsilon)$ . By (32),  $\hat{\phi}_j$  satisfies

$$\Delta \widehat{\phi}_{j} + \frac{8(1+\alpha_{i})^{2}|z|^{2\alpha_{i}}}{\left[1+|z|^{2(1+\alpha_{i})}\right]^{2}} \left[1+O\left(\rho_{i}|z|\right)\right]$$

$$+ O\left(\rho\right) \widehat{\phi}_{j} = \left(\frac{v_{i}\rho_{i}}{\varepsilon}\right)^{2} \widehat{h}_{j},$$
(58)

for  $z \in B_{R_d}(0)$ . Obviously, for  $\beta \in [1, -(1/\alpha)]$ ,  $(v_i\rho_i/\varepsilon)^2 \hat{h}_j \rightarrow 0$  in  $L^{\beta}(B_{R_d}(0))$ . Since  $8(1 + \alpha_i)^2 |z|^{2\alpha_i}/[1 + |z|^{2(1+\alpha_i)}]^2$  is bounded in  $L^{\beta}(B_{R_d}(0))$  and  $\|\hat{\phi}_j\|_{\infty} = 1$ , the elliptic regularity theory implies that  $\hat{\phi}_j$  converges uniformly over compact sets near the origin to a bounded nontrivial solution of  $L_i(\hat{\phi}) = 0$  in  $\mathbb{R}^2$ . Then,  $\hat{\phi}$  is proportional to  $z_{i0}$  for  $i \in J_1$  (see [35, 36, 41]), or a linear combination of  $z_{i0}, z_{i1}$ , and  $z_{i2}$  for  $i \in J_2 \cup J_3$  (see [34, 45, 46]). However, our assumed conditions (5) on i and  $\phi_j$  pass to the limit and yield  $\int \chi z_{i0} \hat{\phi} dz = 0$  for  $i \in J_1$ , or  $\int \chi \cos^2((1/2)\alpha_i \theta) z_{il} \hat{\phi} dz = 0$  for  $i \in J_2 \cup J_3$ , l = 0, 1, 2, which implies that  $\hat{\phi} \equiv 0$ . This is absurd because  $\hat{\phi}$  is nontrivial.  $\Box$ 

We will give next a priori estimate for the solutions of (41) satisfying orthogonality conditions with respect to  $Z_{ij}$ ,  $i \in J_2 \cup J_3$ , j = 1, 2.

**Lemma 6.** There exist positive numbers  $\varepsilon_0$  and C such that for any points  $q \in \Lambda_{k,m}(\delta)$ ,  $h \in \mathcal{C}_*$  and any solution  $\phi$  to (41) with the following orthogonality conditions:

$$\int_{\Omega_{\varepsilon}} \chi_i \cos^2\left(\frac{1}{2}\alpha_i \theta_i\right) \ Z_{ij} \phi dy = 0, \quad \forall i \in J_2 \cup J_3, \ j = 1, 2,$$
(59)

one has

$$\|\phi\|_{\infty} \le C\left(\log\frac{1}{\varepsilon}\right) \|h\|_{*},\tag{60}$$

for all  $\varepsilon < \varepsilon_0$ .

*Proof.* Consider the radial solution  $\psi_i$ ,  $i \in \bigcup_{l=1}^3 J_l$ , for the following equation:

$$\Delta \psi_i (r) = 0, \quad \text{in } R < r < \delta (v_i \rho_i)^{-1},$$
  

$$\psi_i (r) = 1, \quad \text{on } r = R,$$
  

$$\psi_i (r) = 0, \quad \text{on } r = \delta (v_i \rho_i)^{-1},$$
(61)

where  $R > R_0 + 1$  is a large number. Then,  $\psi_i(r)$  is explicitly given by

$$\psi_i(r) = \frac{\log\left[\delta(v_i\rho_i)^{-1}\right] - \log r}{\log\left[\delta(v_i\rho_i)^{-1}\right] - \log R}.$$
(62)

Let  $\eta_1$  and  $\eta_2$  be radial smooth cut-off functions on  $\mathbb{R}^2$  so that

$$0 \leq \eta_{1} \leq 1; \quad |\nabla \eta_{1}| \leq C \quad \text{in } \mathbb{R}^{2};$$
  

$$\eta_{1} \equiv 1 \quad \text{in } B_{R}(0);$$
  

$$\eta_{1} \equiv 0 \quad \text{in } \mathbb{R}^{2} \setminus B_{R+1}(0);$$
  

$$0 \leq \eta_{2} \leq 1; \quad |\nabla \eta_{2}| \leq C \quad \text{in } \mathbb{R}^{2};$$
  

$$\eta_{2} \equiv 1 \quad \text{in } B_{\delta/4}(0);$$
  

$$\eta_{2} \equiv 0 \quad \text{in } \mathbb{R}^{2} \setminus B_{\delta}(0).$$
  
(63)

Set

$$\widetilde{\psi}_{i}(y) = \psi_{i}(z_{i}), 
\eta_{1i}(y) = \eta_{1}(z_{i}), 
\eta_{2i}(y) = \eta_{2}(v_{i}\rho_{i}z_{i}).$$
(64)

$$\widetilde{\phi} = \phi + \sum_{i=1}^{n+k+m} d_i \widetilde{Z}_{i0}, \tag{65}$$

where

$$\widetilde{Z}_{i0}(y) = \eta_{1i} Z_{i0} + (1 - \eta_{1i}) \eta_{2i} \widetilde{\psi}_i Z_{i0}, \qquad (66)$$

and  $d_i$  is chosen such that for  $i \in J_1$ ,

$$d_i \int_{\Omega_{\varepsilon}} \chi_i |Z_{i0}|^2 + \int_{\Omega_{\varepsilon}} \chi_i Z_{i0} \phi = 0, \qquad (67)$$

and for  $i \in J_2 \cup J_3$ ,

$$d_{i} \int_{\Omega_{\varepsilon}} \chi_{i} \cos^{2} \left(\frac{1}{2}\alpha_{i}\theta_{i}\right) Z_{i0}^{2}$$

$$+ \int_{\Omega_{\varepsilon}} \chi_{i} \cos^{2} \left(\frac{1}{2}\alpha_{i}\theta_{i}\right) Z_{i0}\phi = 0.$$
(68)

Thus,

$$L\left(\tilde{\phi}\right) = h + \sum_{i=1}^{n+k+m} d_i L \widetilde{Z}_{i0}, \quad \text{in } \Omega_{\varepsilon},$$
  
$$\tilde{\phi} = 0, \quad \text{on } \partial\Omega_{\varepsilon},$$
(69)

and  $\tilde{\phi}$  satisfies the orthogonality conditions (5). By (43),

$$\left\|\widetilde{\phi}\right\|_{\infty} \le C \left\{ \left\|h\right\|_{*} + \sum_{i=1}^{n+k+m} \left|d_{i}\right| \cdot \left\|L\widetilde{Z}_{i0}\right\|_{*} \right\}.$$
(70)

Multiplying (69) by  $\widetilde{Z}_{i0}, i \in \bigcup_{l=1}^{3} J_l$ , and integrating by parts, it follows that

$$\left\langle L\left(\widetilde{Z}_{i0}\right),\widetilde{\phi}\right\rangle = \left\langle \widetilde{Z}_{i0},h\right\rangle + d_i\left\langle L\left(\widetilde{Z}_{i0}\right),\widetilde{Z}_{i0}\right\rangle,$$
 (71)

where  $\langle f, g \rangle = \int_{\Omega_{\varepsilon}} fg$ . Then, for  $i \in \bigcup_{l=1}^{3} J_l$ , by (70) and (71),  $\left| d_i \left\langle L\left(\widetilde{Z}_{i0}\right), \widetilde{Z}_{i0} \right\rangle \right| \leq C \|h\|_* \left\{ 1 + \left\| L\left(\widetilde{Z}_{i0}\right) \right\|_* \right\}$  $+ C \left\| L\left(\widetilde{Z}_{i0}\right) \right\|_* \sum_{l=1}^{n+k+m} |d_l| \cdot \left\| L\left(\widetilde{Z}_{l0}\right) \right\|_*.$ (72)

Let us claim that for  $i \in \bigcup_{l=1}^{3} J_l$ , there exists some constant C > 0 independent of  $\varepsilon$ , such that

$$\left\|L\left(\widetilde{Z}_{i0}\right)\right\|_{*} = O\left(\frac{1}{\left|\log\varepsilon\right|}\right),\tag{73}$$

$$\left\langle L\left(\widetilde{Z}_{i0}\right), \widetilde{Z}_{i0}\right\rangle < -\frac{C}{\left|\log\varepsilon\right|} \left\{ 1 + O\left(\frac{1}{\left|\log\varepsilon\right|}\right) \right\}.$$
 (74)

Once these estimates (73)-(74) are proven, it easily follows that

$$\left|d_{i}\right| \leq C\left(\log\frac{1}{\varepsilon}\right) \|h\|_{*}.$$
(75)

This, together with (65), (70), and (73), easily gives the estimate (60) of  $\phi$ .

Proofs of (73) and (74). Let us first define that

$$W_{i0} = \frac{8(1+\alpha_i)^2 |z_i|^{2\alpha_i}}{\left[1+|z_i|^{2(1+\alpha_i)}\right]^2},$$
(76)

$$\tilde{z}_{i0}(z_i) = \eta_1 z_{i0} + (1 - \eta_1) \eta_2(v_i \rho_i z_i) \psi_i z_{i0}.$$

For  $r_i := |z_i| \le R$ , by (32),

$$L\left(\widetilde{Z}_{i0}\right) = \left(\frac{\varepsilon}{\nu_i \rho_i}\right)^2 W_{i0} z_{i0} \left\{O\left(\rho_i \left|z_i\right|\right) + O\left(\rho\right)\right\},$$

$$I_1 = \int_{r_i \le R} \widetilde{Z}_{i0} \cdot L\left(\widetilde{Z}_{i0}\right) \qquad (77)$$

$$= \int_{|z| \le R} W_{i0} z_{i0}^2 \left[O\left(\rho_i \left|z\right|\right) + O\left(\rho\right)\right] = O\left(\rho\right).$$

For  $R < r_i \le R + 1$ ,  $\widetilde{Z}_{i0}(y) = \eta_1 z_{i0} + (1 - \eta_1) \psi_i z_{i0}$ ,  $1 - \psi_i = O(1/|\log \varepsilon|)$ , and  $|\nabla \psi_i| = O(1/|\log \varepsilon|)$ . Then,

$$L\left(\widetilde{Z}_{i0}\right) = \left(\frac{\varepsilon}{v_i\rho_i}\right)^2 \left\{2\left(1-\psi_i\right)\nabla\eta_1\nabla z_{i0} + 2\left(1-\eta_1\right)\nabla\psi_i\nabla z_{i0} - 2z_{i0}\nabla\eta_1\nabla\psi_i + \left(1-\psi_i\right)z_{i0}\Delta\eta_1 + W_{i0}\widetilde{z}_{i0}\left[O\left(\rho_i\left|z_i\right|\right) + O\left(\rho\right)\right]\right\}(z_i)$$
$$= \left(\frac{\varepsilon}{v_i\rho_i}\right)^2 O\left(\frac{1}{\left|\log\varepsilon\right|}\right).$$
(78)

Furthermore,

$$I_{2} = \int_{R \leq r_{i} \leq R+1} \widetilde{Z}_{i0} \cdot L\left(\widetilde{Z}_{i0}\right)$$

$$= \int_{R \leq |z| \leq R+1} \widetilde{z}_{i0} \left\{ \left(1 - \psi_{i}\right) z_{i0} \Delta \eta_{1} \right.$$

$$+ 2 \left(1 - \psi_{i}\right) \nabla \eta_{1} \nabla z_{i0} - 2 z_{i0} \nabla \eta_{1} \nabla \psi_{i}$$

$$+ 2 \left(1 - \eta_{1}\right) \nabla \psi_{i} \nabla z_{i0} \right\} dz + O\left(\rho\right).$$

$$(79)$$

Integrating by parts the first term of  $I_2$ , it follows that

$$I_{2} = \int_{R \le |z| \le R+1} 2(1 - \eta_{1}) \tilde{z}_{i0} \nabla \psi_{i} \nabla z_{i0} - z_{i0}^{2} \nabla \psi_{i} \nabla \eta_{1} - z_{i0}^{2} (1 - \psi_{i})^{2} |\nabla \eta_{1}|^{2} + O(\rho).$$
(80)

Integrating by parts the first term again, it also follows that

$$I_{2} = \int_{|z|=R+1} \psi_{i'} (|z|) \psi_{i} z_{i0}^{2}$$
  
- 
$$\int_{R \le |z| \le R+1} z_{i0}^{2} |(1 - \eta_{1}) \nabla \psi_{i}$$
(81)  
+ 
$$(1 - \psi_{i}) \nabla \eta_{1} |^{2} + O(\rho).$$

Then,

$$I_{2} = \int_{|z|=R+1} \psi_{i'}(|z|) \psi_{i} z_{i0}^{2} + O\left(\frac{1}{|\log \varepsilon|^{2}}\right).$$
(82)

For  $R + 1 < r_i \le \delta(4\nu_i\rho_i)^{-1}$ ,  $\widetilde{Z}_{ij}(y) = \psi_i z_{ij}(z_i)$ . Then,

$$L\left(\widetilde{Z}_{i0}\right) = \left(\frac{\varepsilon}{\nu_i \rho_i}\right)^2 \left\{ 2\nabla \psi_i \nabla z_{i0} + \psi_i \Delta z_{i0} + W_{i0} \psi_i z_{i0} \left[1 + O\left(\rho_i \left|z_i\right|\right) + O\left(\rho\right)\right] \right\} (83)$$

$$+ O\left(\rho\right) \right\} (z_i) .$$

Note that  $|\nabla z_{i0}| \leq C/|z_i|^{3+2\alpha_i}$ ,  $|\nabla \psi_i| \leq C/(|z_i|\log(1/\varepsilon))$ , and  $|W_{i0}| \leq C/|z_i|^{4+2\alpha_i}$ . Then,

$$L\left(\widetilde{Z}_{i0}\right) = \left(\frac{\varepsilon}{\nu_i \rho_i}\right)^2 \frac{1}{\left|z_i\right|^{4+2\alpha_i}} \times \left\{O\left(\frac{1}{\left|\log\varepsilon\right|}\right) + O\left(\rho_i\left|z_i\right|\right) + O\left(\rho\right)\right\}.$$
(84)

Besides,

$$I_{3} = \int_{R+1 \leq r_{i} \leq \delta(4v_{i}\rho_{i})^{-1}} \widetilde{Z}_{i0} \cdot L\left(\widetilde{Z}_{i0}\right)$$

$$= \int_{R+1 \leq |z| \leq \delta(4v_{i}\rho_{i})^{-1}} \psi_{i}z_{i0}$$

$$\times \left\{ 2\nabla\psi_{i}\nabla z_{i0} + W_{i0}\psi_{i}z_{i0} + \sum_{i=1}^{N} \left[O\left(\rho_{i}|z|\right) + O\left(\rho\right)\right]\right\}$$

$$(85)$$

$$= \left( \int_{|z|=\delta(4v_i\rho_i)^{-1}} - \int_{|z|=R+1} \right) \psi'_i(|z|) \\ \times \psi_i z_{i0}^2 - \int_{R+1 \le |z| \le \delta(4v_i\rho_i)^{-1}} z_{i0}^2 |\nabla \psi_i|^2 + O(\rho) \,.$$

Some simple computations show that

$$\int_{|z|=\delta(4\nu_{i}\rho_{i})^{-1}} \psi_{i}'(|z|) \psi_{i}z_{i0}^{2} = O\left(\frac{1}{|\log\varepsilon|^{2}}\right), \quad (86)$$

$$\int_{R+1 \le |z| \le \delta(4\nu_{i}\rho_{i})^{-1}} z_{i0}^{2} |\nabla\psi_{i}|^{2}$$

$$\ge \frac{\pi}{2} \frac{\log(\delta/4\nu_{i}\rho_{i}) - \log(R+1)}{(\log(\delta/\nu_{i}\rho_{i}) - \log R)^{2}}.$$
(87)

Then, there exists some constant C>0 independent of  $\varepsilon$  and R such that

$$I_{3} < -\int_{|z|=R+1} \psi_{i}'(|z|) \psi_{i} z_{i0}^{2} - \frac{C}{|\log \varepsilon|} \left\{ 1 + O\left(\frac{1}{|\log \varepsilon|}\right) \right\}.$$

$$(88)$$

For  $\delta(4v_i\rho_i)^{-1} < r_i \le \delta(v_i\rho_i)^{-1}$ ,  $\overline{Z}_{i0}(y) = \eta_2(v_i\rho_i z_i)\psi_i z_{i0}$ . Then,

$$L\left(\widetilde{Z}_{i0}\right) = \left(\frac{\varepsilon}{v_i\rho_i}\right)^2 \left\{\psi_i z_{i0} \Delta \eta_2 \left(v_i\rho_i z_i\right) + 2\nabla \eta_2 \left(v_i\rho_i z_i\right) \nabla \psi_i z_{i0} + 2\psi_i \nabla \eta_2 \left(v_i\rho_i z_i\right) \nabla z_{i0} + 2\eta_2 \left(v_i\rho_i z_i\right) \nabla \psi_i \nabla z_{i0} + W_{i0} \eta_2 \left(v_i\rho_i z_i\right) \psi_i z_{i0} \times \left[O\left(\rho_i \left|z_i\right|\right) + O\left(\rho\right)\right]\right\}.$$
(89)

Note that  $\psi_i = O(1/|\log \varepsilon|)$ ,  $|\nabla \psi_i| = O(\rho_i/|\log \varepsilon|)$ , and  $|\nabla z_{i0}| = O(\rho_i^{3+2\alpha_i})$ . Then,

$$L\left(\widetilde{Z}_{i0}\right) = \left(\frac{\varepsilon}{v_i \rho_i}\right)^2 O\left(\frac{\rho_i^2}{\left|\log \varepsilon\right|}\right).$$
(90)

Furthermore,

$$I_4 = \int_{\delta(4\nu_i\rho_i)^{-1} < r_i \le \delta(\nu_i\rho_i)^{-1}} \widetilde{Z}_{i0} \cdot L\left(\widetilde{Z}_{i0}\right) = O\left(\frac{1}{\left|\log\varepsilon\right|^2}\right).$$
(91)

As a result, the estimates (73)-(74) can be easily derived from (77)–(91).  $\hfill \Box$ 

*Proof of Proposition 4.* We first establish the a priori estimate (39). Testing (38) against  $\eta_{2i}Z_{ij}$ ,  $i \in J_2 \cup J_3$ , j = 1, 2, and integrating by parts, it follows that

$$\left|c_{ij}\right| \leq C\left(\frac{\varepsilon}{\nu_{i}\rho_{i}}\right)^{2} \left(\left|\left\langle\phi, L\left(\eta_{2i}Z_{ij}\right)\right\rangle\right| + \left|\left\langle h, \eta_{2i}Z_{ij}\right\rangle\right|\right).$$
(92)

Observe that

$$L\left(\eta_{2i}Z_{ij}\right) = \left(\frac{\varepsilon}{\nu_i\rho_i}\right)^2 \left\{ O\left(\frac{\rho_i}{\left(1+|z_i|\right)^{2+\alpha_i}}\right) + O\left(\frac{\rho_i^2}{\left(1+|z_i|\right)^{1+\alpha_i}}\right) \right\} + \eta_2\left(\nu_i\rho_iz_i\right)z_{ij}\left(W_q - W_{i0}\right).$$

$$(93)$$

Then,

$$\left\langle \phi, L\left(\eta_{2i}Z_{ij}\right) \right\rangle = O\left(\rho \|\phi\|_{\infty}\right).$$
 (94)

Note that  $\langle h, \eta_{2i} Z_{ij} \rangle = O(||h||_*)$ . This, together with (92)-(94), implies that

$$\left|c_{ij}\right| \le C \left(\frac{\varepsilon}{\nu_i \rho_i}\right)^2 \left(\|h\|_* + \rho \|\phi\|_{\infty}\right).$$
(95)

By (60),

$$\begin{aligned} \|\phi\|_{\infty} &\leq C\left(\log\frac{1}{\varepsilon}\right) \left\{ \|h\|_{*} \\ &+ \sum_{j=1}^{2} \sum_{i=n+1}^{n+k+m} \left(\frac{\nu_{i}\rho_{i}}{\varepsilon}\right)^{2} |c_{ij}| \right\}. \end{aligned}$$
(96)

Combining this with (95), we find that the a priori estimate (39) holds. Furthermore,

$$\left|c_{ij}\right| \le C \left(\frac{\varepsilon}{\nu_i \rho_i}\right)^2 \|h\|_*,\tag{97}$$

which implies that there exists a unique trivial solution to problem (38) with  $h \equiv 0$ .

Next, we prove the solvability of problem (38). Consider the following Hilbert space:

$$\mathbb{H} = \left\{ \phi \in H_0^1 \left( \Omega_{\varepsilon} \right) : \\ \phi \text{ satisfies the orthogonality conditions (59)} \right\},$$
(98)

endowed with the usual norm  $\|\psi\|_{H_0^1} = \int_{\Omega_{\epsilon}} |\nabla \psi|^2$ . Problem (38) is equivalent to that of finding  $\phi \in \mathbb{H}$  such that

$$\int_{\Omega_{\varepsilon}} \nabla \phi \nabla \psi - W_q \phi \psi + h \psi = 0, \quad \forall \psi \in \mathbb{H}.$$
(99)

By Fredholm's alternative, this is equivalent to the uniqueness of solutions to this problem, which is guaranteed by (39). As a consequence, there exists a unique solution  $\phi = T(h)$ , scalars  $c_{ij}, i \in J_2 \cup J_3, j = 1, 2$ , for problem (38) with  $h \in \mathscr{C}_*$ , where  $T : \mathscr{C}_* \mapsto L^{\infty}(\Omega_{\varepsilon})$  is a continuous linear map satisfying  $||T(h)||_{\infty} \leq C(\log(1/\varepsilon))||h||_*$ .

Finally, we give the a priori estimate (40). Let us Differentiate (38) with respect to the parameters  $q'_{l,s}$ ,  $l \in J_2 \cup J_3$ , s = 1, 2. Formally,  $Z = \partial_{q'_s} \phi$  should satisfy

$$L(Z) = -\phi \ \partial_{q'_{i,s}} W_q$$

$$+ \sum_{j=1}^{2} \sum_{i=n+1}^{n+k+m} \left\{ c_{ij} \partial_{q'_{i,s}} \right\}$$

$$\times \left[ \chi_i \cos^2 \left( \frac{1}{2} \alpha_i \theta_i \right) Z_{ij} \right]$$

$$+ d_{ij} \chi_i \cos^2 \left( \frac{1}{2} \alpha_i \theta_i \right) Z_{ij} \right], \qquad (100)$$

where (still formally)  $d_{ij} = \partial_{q'_{l,s}} c_{ij}$ . The orthogonality conditions now become

$$\int_{\Omega_{\varepsilon}} \chi_{i} \cos^{2}\left(\frac{1}{2}\alpha_{i}\theta_{i}\right) Z_{ij}Z$$
  
=  $-\int_{\Omega_{\varepsilon}} \phi \partial_{q'_{i,s}} \left[\chi_{i} \cos^{2}\left(\frac{1}{2}\alpha_{i}\theta_{i}\right) Z_{ij}\right],$  (101)  
 $\forall i \in J_{2} \cup J_{3}, \ j = 1, 2.$ 

We consider the constants  $b_{ij}$ ,  $i \in J_2 \cup J_3$ , j = 1, 2, defined as

$$b_{ij} \int_{\Omega_{\varepsilon}} \chi_i \cos^2\left(\frac{1}{2}\alpha_i \theta_i\right) |Z_{ij}|^2$$
  
=  $-\int_{\Omega_{\varepsilon}} \phi \partial_{q'_{l,s}} \left[\chi_i \cos^2\left(\frac{1}{2}\alpha_i \theta_i\right) Z_{ij}\right].$  (102)

By (31), it follows that  $|\partial_{q'_{l,s}} \log \mu_i| = O(\varepsilon)$  uniformly holds on  $\Lambda_{k,m}(\delta)$ , which implies that for  $i \in J_2 \cup J_3$ , j = 1, 2,

$$\left| b_{ij} \right| \leq \begin{cases} C \frac{\varepsilon}{v_l \rho_l} \| \phi \|_{\infty}, & \text{if } i = l, \\ C \varepsilon \| \phi \|_{\infty}, & \text{if } i \neq l. \end{cases}$$
 (103)

Define

$$\widetilde{Z} = Z - \sum_{j=1}^{2} \sum_{i=n+1}^{n+k+m} b_{ij} \eta_{2i} Z_{ij},$$

$$f = \sum_{j=1}^{2} \sum_{i=n+1}^{n+k+m} \left\{ c_{ij} \partial_{q'_{i,s}} \right\}$$

$$\times \left[ \chi_i \cos^2 \left( \frac{1}{2} \alpha_i \theta_i \right) Z_{ij} \right]$$

$$- b_{ij} L \left( \eta_{2i} Z_{ij} \right)$$

$$- \phi \partial_{q'_{i,s}} W_q.$$
(104)

We then have

$$L\left(\widetilde{Z}\right) = f + \sum_{j=1}^{2} \sum_{i=n+1}^{n+k+m} d_{ij} \chi_i \cos^2\left(\frac{1}{2}\alpha_i \theta_i\right), \quad \text{in } \Omega_{\varepsilon},$$
  
$$\phi = 0, \quad \text{on } \partial\Omega_{\varepsilon}, \tag{105}$$

$$\int_{\Omega_{\varepsilon}} \chi_i \cos^2\left(\frac{1}{2}\alpha_i \theta_i\right) Z_{ij} \widetilde{Z} = 0, \quad \forall i \in J_2 \cup J_3, \ j = 1, 2.$$

Thus,  $Z = \partial_{q'_{1s}} \phi$  can be uniquely expressed as

$$Z = T(f) + \sum_{j=1}^{2} \sum_{i=n+1}^{n+k+m} b_{ij} \eta_{2i} Z_{ij}.$$
 (106)

Furthermore, elliptic regularity theory implies that  $\phi = T(h)$  is differentiable with respect to  $q'_{l,s}$ ,  $l \in J_2 \cup J_3$ , s = 1, 2. Note that  $\|L(\eta_{2i}Z_{ij})\|_* = O(\rho)$ ,  $\|\partial_{q'_{l,s}}W_q\|_* = O(\rho_l^{\alpha_l})$ ,  $\|\partial_{q'_{l,s}}[\chi_l \cos^2((1/2)\alpha_l\theta_l)Z_{lj}]\|_* = O(\rho_l^{-\alpha_l})$ , and  $\|\partial_{q'_{l,s}}[\chi_i \cos^2((1/2)\alpha_i\theta_l)Z_{ij}]\|_* = O(\rho_l^{1-\alpha_i})$  for  $i \neq l$ . This, together with (39) and (97)–(106), implies that

$$\begin{split} \|Z\|_{\infty} &\leq C \left(\log \frac{1}{\varepsilon}\right) \|f\|_{*} + C \sum_{j=1}^{2} \sum_{i=n+1}^{n+k+m} \left|b_{ij}\right| \\ &\leq C \left(\log \frac{1}{\varepsilon}\right) \\ &\times \sum_{j=1}^{2} \sum_{i=n+1}^{n+k+m} \left\{ \left(\frac{\varepsilon}{\nu_{i}\rho_{i}}\right)^{2} \|h\|_{*} \\ &\times \left\|\partial_{q_{l,s}'} \left[\chi_{i} \cos^{2}\left(\frac{1}{2}\alpha_{i}\theta_{i}\right) Z_{ij}\right]\right\|_{*} \right. (107) \\ &+ \rho \left|b_{ij}\right| \right\} \\ &+ C \left(\log \frac{1}{\varepsilon}\right) \|\phi\|_{\infty} \left\|\partial_{q_{l,s}'} W_{q}\right\|_{*} \\ &+ C \frac{\varepsilon}{\nu_{l}\rho_{l}} \|\phi\|_{\infty} \\ &\leq C \left. \frac{\varepsilon}{\nu_{l}\rho_{l}} \left(\log \frac{1}{\varepsilon}\right)^{2} \|h\|_{*}, \end{split}$$

which implies that the estimate (40) holds.

Now, we solve the auxiliary nonlinear problem: for  $q \in \Lambda_{k,m}(\delta)$ , we find a function  $\phi$  and scalars  $c_{ij}$ ,  $i \in J_2 \cup J_3$ , j = 1, 2, such that

$$\begin{split} L(\phi) &= -\left[R_q + N(\phi)\right] \\ &+ \sum_{j=1}^{2} \sum_{i=n+1}^{n+k+m} c_{ij} \chi_i \cos^2\left(\frac{1}{2} \alpha_i \theta_i\right) Z_{ij}, \\ &\quad \text{in } \Omega_{\varepsilon}, \end{split}$$

$$\phi = 0$$
, on  $\partial \Omega_{\varepsilon}$ ,

$$\int_{\Omega_{\varepsilon}} \chi_i \cos^2\left(\frac{1}{2}\alpha_i \theta_i\right) Z_{ij} \phi dy = 0,$$
  
$$\forall i \in J_2 \cup J_3, \ j = 1, 2.$$
(108)

The following result can be proved through arguments similar to these of [39].

**Proposition 7.** There exist positive numbers  $\varepsilon_0$  and C such that for any  $\varepsilon < \varepsilon_0$  and points  $q \in \Lambda_{k,m}(\delta)$ , there is a unique solution  $\phi \in L^{\infty}(\Omega_{\varepsilon})$ , scalars  $c_{ij} \in \mathbb{R}$ ,  $i \in J_2 \cup J_3$ , j = 1, 2, of problem (3), which satisfies

$$\|\phi\|_{\infty} \le C\rho \log \frac{1}{\varepsilon}.$$
 (109)

Moreover, the map  $q' \mapsto \phi$  is  $C^1$ , precisely for  $l \in J_2 \cup J_3$ , s = 1, 2,

$$\left\| \partial_{q_{l,s}'} \phi \right\|_{\infty} \le C \rho \frac{\varepsilon}{v_l \rho_l} \left( \log \frac{1}{\varepsilon} \right)^2, \tag{110}$$

where  $\rho = \max\{\rho_i : i \in \bigcup_{l=1}^3 J_l\}.$ 

### 4. Variational Reduction

In what follows, we only need to find a solution of problem (27) (or (25)) with  $k + m \ge 1$ , and hence to problem (3) if points  $q \in \Lambda_{k,m}(\delta)$  satisfy

$$c_{ij}(q') = 0, \quad \forall i \in J_2 \cup J_3, \ j = 1, 2.$$
 (111)

To realize it, we consider the energy functional  $J_{\varepsilon}$  associated with problem (25), namely,

$$J_{\varepsilon}(v) = \frac{1}{2} \int_{\Omega_{\varepsilon}} |\nabla v|^2 - 2\varepsilon^4 \int_{\Omega_{\varepsilon}} c(\varepsilon y) S(\varepsilon y) \cosh v,$$
  
$$\forall u \in H_0^1(\Omega_{\varepsilon}).$$
 (112)

We define

$$F_{\varepsilon}(q) = J_{\varepsilon}(V_q + \phi), \qquad (113)$$

where  $V_q$  is defined in (26) and  $\phi$  is the unique solution of problem (3). Critical points of  $F_{\varepsilon}$  correspond to solutions of (111) for small  $\varepsilon$ , as the following result states.

**Lemma 8.** For any points  $q \in \Lambda_{k,m}(\delta)$ , the functional  $F_{\varepsilon}(q)$  is of class  $C^1$ . Moreover, for  $\varepsilon > 0$  small enough, if  $D_q F_{\varepsilon}(q) = 0$ , then points  $q \in \Lambda_{k,m}(\delta)$  satisfy (112).

*Proof.* Observe that for  $l \in J_2 \cup J_3$ , s = 1, 2,

$$\partial_{q_{l,s}} F_{\varepsilon} (q) = \varepsilon^{-1} D J_{\varepsilon} (V_q + \phi) \times \left[ \partial_{q'_{l,s}} (V_q + \phi) \right]$$
(114)  
$$= \varepsilon^{-1} \sum_{j=1}^{2} \sum_{i=n+1}^{n+k+m} c_{ij} K_{il} (j, s) ,$$

where

$$K_{il}(j,s) = \int_{\Omega_{\varepsilon}} \chi_i \cos^2\left(\frac{1}{2}\alpha_i \theta_i\right) Z_{ij}\left(\partial_{q'_{l,s}} V_q + \partial_{q'_{l,s}}\phi\right) dy.$$
(115)

Then, if  $D_q F_{\varepsilon}(q) = 0$ , we have

$$\sum_{j=1}^{2} \sum_{i=n+1}^{n+k+m} c_{ij} K_{il}(j,s) = 0, \quad \forall l \in J_2 \cup J_3, \ s = 1, 2.$$
(116)

Set

$$F_{l} = \int_{0}^{R_{0}+1} \frac{16(1+\alpha_{l})r^{3(1+\alpha_{l})}}{\left[1+r^{2(1+\alpha_{l})}\right]^{2}}\chi(r)dr,$$

$$\forall l \in J_{2} \cup J_{3}.$$
(117)

A simple computation shows that

$$\partial_{q_{l,s}^{\prime}} V_{q} = \frac{\varepsilon}{\nu_{l} \rho_{l}} \frac{4 \left(1 + \alpha_{l}\right) a_{l} \left|z_{l}\right|^{2\alpha_{l}} z_{l,s}}{1 + \left|z_{l}\right|^{2\left(1 + \alpha_{l}\right)}} + O\left(\varepsilon\right),$$

$$\forall l \in J_{2} \cup J_{3}, \ s = 1, 2.$$
(118)

This, together with the estimate (110) of  $\partial_{q'_{ls}}\phi$ , implies that

$$K_{il}(j,s) = \begin{cases} \left(\frac{v_i \rho_i}{\varepsilon}\right)^2 O\left(\rho \frac{\varepsilon}{v_l \rho_l} |\log \varepsilon|^2\right), & \forall i \neq l \text{ or } j \neq s, \\ \frac{v_l \rho_l}{\varepsilon} \left[A_l + O\left(\rho |\log \varepsilon|^2\right)\right], & \forall i = l \text{ and } j = s, \end{cases}$$
(119)

where

$$A_{l} = \begin{cases} \pi a_{l} F_{l}, & \forall l \in J_{3}, \\ \frac{1}{4} \pi a_{l} F_{l}, & \forall l \in J_{2}. \end{cases}$$
(120)

Set

$$\tilde{c}_{ij} = \left(\frac{\nu_i \rho_i}{\varepsilon}\right)^2 c_{ij}, \quad \forall i \in J_2 \cup J_3, \ j = 1, 2.$$
(121)

Thus, (116) can be rewritten as: for any  $l \in J_2 \cup J_3$ , s = 1, 2,

$$\widetilde{c}_{ls}a_{l}F_{l} = \sum_{j=1}^{2}\sum_{i=n+1}^{n+k+m} \widetilde{c}_{ij}a_{i}O\left(\rho \left|\log \varepsilon\right|^{2}\right),$$

$$\forall l \in J_{2} \cup J_{3}, \ s = 1, 2.$$
(122)

This is a diagonal dominant system and we thus get  $c_{ij} = 0$  for all *i*, *j*.

Next, we give a precise asymptotic expansion of  $F_{\varepsilon}(q)$  defined in (113).

Lemma 9. The following precise asymptotic expansion holds

$$F_{\varepsilon}(q) = -\sum_{i \in J_{1}} d_{i} \left\{ \frac{1}{2} d_{i} H(q_{i}, q_{i}) + \sum_{j \in J_{1}, i \neq j} \frac{1}{2} a_{i} a_{j} d_{j} G(q_{j}, q_{i}) \right\}$$
(123)  
+ 
$$\sum_{i \in \bigcup_{l=1}^{3} J_{l}} d_{i} \left\{ \log 8(1 + \alpha_{i})^{2} - 2(1 + \log \varepsilon) \right\}$$
+ 
$$q_{k,m}^{n}(q) + O(\rho),$$

uniformly for any points  $q \in \Lambda_{k,m}(\delta)$ , where  $\varphi_{k,m}^n(q)$  is defined in (12).

*Proof.* Let us first give a priori estimate of  $\theta_{\varepsilon}(q)$ , where  $\theta_{\varepsilon}(q) = F_{\varepsilon}(q) - J_{\varepsilon}(V_q)$ . Using  $DJ_{\varepsilon}(V_q + \phi)[\phi] = 0$ , a Taylor expansion and an integration by parts, it follows that

$$\begin{aligned} \theta_{\varepsilon}(q) &= -\int_{0}^{1} D^{2} J_{\varepsilon} \left( V_{q} + s\phi \right) \left[\phi\right]^{2} s ds \\ &= -\int_{0}^{1} \left( \int_{\Omega_{\varepsilon}} \left[ N\left(\phi\right) + R_{q} \right] \phi \right. \\ &+ 2\varepsilon^{4} c\left(\varepsilon y\right) S\left(\varepsilon y\right) \\ &\times \left[ \cosh V_{q} - \cosh \right. \\ &\left. \times \left( V_{q} + s\phi \right) \right] \phi^{2} \right) s ds. \end{aligned}$$
(124)

Note that  $||N(\phi)||_{*} = O(||\phi||_{\infty}^{2}), ||R_{q}||_{*} = O(\rho),$ and  $||2\varepsilon^{4}c(\varepsilon y)S(\varepsilon y)[\cosh V_{q} - \cosh(V_{q} + s\phi)]||_{*} = O(||\phi||_{\infty}).$ These, together with (109), imply that  $\theta_{\varepsilon}(q) = O(\rho^{2}|\log \varepsilon|),$ and so  $F_{\varepsilon}(q) = J_{\varepsilon}(V_{q}) + O(\rho^{2}|\log \varepsilon|).$ 

Next, we only need to give an asymptotic expansion of  $J_{\varepsilon}(V_q)$ . Observe that

$$J_{\varepsilon}\left(V_{q}\right) = -\frac{1}{2} \sum_{i,j \in \bigcup_{l=1}^{3} J_{l}} a_{i}a_{j} \int_{\Omega} Pu_{i}\Delta u_{j}dx - \int_{\Omega_{\varepsilon}} W_{q}dy.$$
(125)

By (22),

$$-\int_{\Omega} Pu_{i}\Delta u_{j} = \int_{\Omega} \varepsilon^{2} c_{j}(q_{j}) \left| x - q_{j} \right|^{2\alpha_{j}} e^{u_{j}}$$

$$\times \left\{ u_{i} + d_{i}H(x, q_{i}) - \log \frac{8\mu_{i}^{2}(1 + \alpha_{i})^{2}}{c_{i}(q_{i})} + O(\rho) \right\}$$

$$= I_{ij} + J_{ij},$$
(126)

where

$$\begin{split} I_{ij} &= \int_{\Omega_{\nu_{j}\rho_{j}}} \frac{8(1+\alpha_{j})^{2} |z_{j}|^{2\alpha_{j}}}{\left[1+|z_{j}|^{2(1+\alpha_{j})}\right]^{2}} \\ &\times \log \frac{1}{\left[\mu_{i}^{2}\varepsilon^{2}+|\nu_{j}\rho_{j}z_{j}+q_{j}-q_{i}|^{2(1+\alpha_{i})}\right]^{2}} dz_{j}, \end{split}$$
(127)  
$$J_{ij} &= \int_{\Omega_{\nu_{j}\rho_{j}}} \frac{8(1+\alpha_{j})^{2} |z_{j}|^{2\alpha_{j}}}{\left[1+|z_{j}|^{2(1+\alpha_{j})}\right]^{2}} \\ &\times \left\{d_{i}H\left(q_{j},q_{i}\right)+O\left(\rho_{j}|z_{j}|\right)+O\left(\rho\right)\right\} dz_{j}. \end{split}$$

Note that

$$\int_{\Omega_{\nu_{j}\rho_{j}}} \frac{8(1+\alpha_{j})^{2}|z|^{2\alpha_{j}}}{\left[1+|z|^{2(1+\alpha_{j})}\right]^{2}} dz = d_{j} + O\left(\varepsilon^{2}\right),$$
(128)

$$\int_{\Omega_{\nu_{j}\rho_{j}}} \frac{8\left(1+\alpha_{j}\right)^{2}|z|^{2\alpha_{j}}}{\left[1+|z|^{2(1+\alpha_{j})}\right]^{2}}O\left(\rho_{j}|z|\right)dz = O\left(\rho_{j}\right) + O\left(\varepsilon^{2}\right),$$
(129)

and for  $i \neq j$ ,

$$\log \left[ \mu_i^2 \varepsilon^2 + \left| v_j \rho_j z_j + q_j - q_i \right|^{2(1+\alpha_i)} \right]$$
  
= 2 (1 + \alpha\_i) log |q\_j - q\_i| (130)  
+ O (\rho\_j |z\_j|) + O (\varepsilon^2).

Then, for any  $i \neq j$ , by (128)–(130),

$$I_{ij} = -4\left(1 + \alpha_i\right) d_j \log \left|q_j - q_i\right| + O\left(\rho\right).$$
(131)

By (128),

$$I_{jj} = -2 \int_{\Omega_{\nu_{j}\rho_{j}}} \frac{8(1+\alpha_{j})^{2}|z|^{2\alpha_{j}}}{[1+|z|^{2(1+\alpha_{j})}]^{2}} \times \log \left[1+|z|^{2(1+\alpha_{j})}\right] dz -4 \left[d_{j}+O\left(\varepsilon^{2}\right)\right] \log \left(\mu_{j}\varepsilon\right).$$
(132)

Making the complex changes of variables  $\omega = z^{1+\alpha_j}$  with  $\omega = |\omega|e^{i\tilde{\theta}}$ , we get

$$I_{jj} = -2 \int_{0}^{2\pi(1+\alpha_{j})} d\tilde{\theta} \int_{0}^{(\mu_{j}\varepsilon)^{-1}} \frac{8 |\omega|}{\left[1+|\omega|^{2}\right]^{2}} \times \log\left[1+|\omega|^{2}\right] d |\omega| \qquad (133)$$
$$-4d_{j} \log\left(\mu_{j}\varepsilon\right) + O\left(\rho\right)$$
$$= -4d_{j} \log\left(\mu_{j}\varepsilon\right) - 2d_{j} + O\left(\rho\right).$$

On the other hand, by (128)-(129), we have

$$J_{ij} = d_i d_j H\left(q_i, q_j\right) + O\left(\rho\right). \tag{134}$$

Thus, by (131)–(134),

$$-\int_{\Omega} Pu_{i}\Delta u_{j}$$

$$=\begin{cases} d_{i}d_{j}G(q_{i},q_{j}) + O(\rho), & \forall i \neq j, \\ d_{j}\{d_{j}H(q_{j},q_{j}) - 4\log(\mu_{j}\varepsilon) - 2\} + O(\rho), & \forall i = j. \end{cases}$$
(135)

Besides, by (32),

$$\int_{\Omega_{\varepsilon}} W_q = \sum_{\bigcup_{l=1}^{3} J_l} d_l + O(\rho).$$
(136)

Using the choice for  $\mu_i$ 's by (31), together with (125), (135), and (136), it follows that

$$J_{\varepsilon}\left(V_{q}\right) = \sum_{i \in \bigcup_{l=1}^{3} J_{l}} d_{i} \left\{ \log \frac{8(1+\alpha_{i})^{2}}{c_{i}\left(q_{i}\right)} - 2\left(1+\log\varepsilon\right) - \frac{1}{2}d_{i}H\left(q_{i},q_{i}\right) - \frac{1}{2}\sum_{j \neq i, j \in \bigcup_{l=1}^{3} J_{l}} a_{i}a_{j}d_{j}G\left(q_{j},q_{i}\right) \right\} + O\left(\rho\right).$$

$$(137)$$

This, together with (12), easily gives the asymptotic expansion (123) of  $F_{\varepsilon}(q)$ .

# 5. Proofs of Theorems

*Proof of Theorem 3.* According to Lemma 8, we only need to find a critical point of the function, consider

$$\widetilde{F}_{\varepsilon}(q) = F_{\varepsilon}(q)$$

$$-\sum_{i \in \bigcup_{l=1}^{3} J_{l}} d_{i} \left\{ \log 8(1+\alpha_{i})^{2} - 2(1+\log \varepsilon) \right\}$$

$$+\sum_{i \in J_{1}} d_{i} \left\{ \frac{1}{2} d_{i} H\left(q_{i},q_{i}\right) + \sum_{j \in J_{1}, i \neq j} \frac{1}{2} a_{i} a_{j} d_{j} G\left(q_{j},q_{i}\right) \right\}.$$
(138)

By (123),  $\tilde{F}_{\varepsilon} = \varphi_{k,m}^n + O(\rho) \rightarrow \varphi_{k,m}^n$  uniformly holds on  $\Lambda_{k,m}(\delta)$ . By Definition 2, there exists a critical point  $q_{\varepsilon}$  of  $F_{\varepsilon}$  such that  $\varphi_{k,m}^n(q_{\varepsilon}) \rightarrow \varphi_{k,m}^n(q^*)$ . Moreover, up to a subsequence, there exists points  $\tilde{q} = (\tilde{q}_{n+1}, \dots, \tilde{q}_{M+m}) \in \Lambda_{k,m}$  such that  $q_{\varepsilon} \rightarrow \tilde{q} \in \Lambda_{k,m}$  as  $\varepsilon \rightarrow 0$ , and  $\varphi_{k,m}^n(q^*) = \varphi_{k,m}^n(\tilde{q})$ . Thus,  $v_{\varepsilon} = V_{q_{\varepsilon}} + \phi(q_{\varepsilon})$  is a family of solutions of problem (25) (or (27)). Set  $u_{\varepsilon}(x) = V_{q_{\varepsilon}}((1/\varepsilon)x) + \phi(q_{\varepsilon})((1/\varepsilon)x)$  for any  $x \in \Omega$ . As a result,  $u_{\varepsilon}$  is a family of solutions of problem (11) with the qualitative properties predicted by the theorem, as it can be easily shown.

*Proof of Theorem 1.* Set  $q_1 = (0, 0)$ , c(x) = 1, and  $a_l = (-1)^{l-1}$  for l = 1, ..., m + 1. If  $\alpha \in \mathbb{N}$ ,

$$\begin{split} \varphi_{1,m}^{0}\left(q\right) &= -32\pi^{2} \\ &\times \left\{ \frac{\alpha}{2\pi} \sum_{l=2}^{m+1} \log \left|q_{l}\right| + (1+\alpha)^{2} H\left(0,0\right) \\ &+ \sum_{l=2}^{m+1} H\left(q_{l},q_{l}\right) + 2\left(1+\alpha\right) \\ &\times \sum_{l=2}^{m+1} (-1)^{l+1} G\left(0,q_{l}\right) \\ &+ \sum_{l,j=2,l\neq j}^{m+1} (-1)^{l+j} G\left(q_{l},q_{j}\right) \right\}, \end{split}$$
(139)

and if  $\alpha \in (-1, +\infty) \setminus (\mathbb{N} \cup \{0\})$ ,

$$\begin{split} \varphi_{0,m}^{1}\left(q\right) &= -32\pi^{2} \\ &\times \left\{ \frac{\alpha}{2\pi} \sum_{l=2}^{m+1} \log \left|q_{l}\right| + 2\left(1 + \alpha\right) \\ &\times \sum_{l=2}^{m+1} (-1)^{l+1} G\left(0, q_{l}\right) + \sum_{l=2}^{m+1} H\left(q_{l}, q_{l}\right) \right. \end{split}$$
(140)
$$&+ \left. \sum_{l,j=2, \, l \neq j}^{m+1} (-1)^{l+j} G\left(q_{l}, q_{j}\right) \right\} . \end{split}$$

We will seek a nodal solution for problem (5) with the concentration points 0 and  $\tilde{q}_l = (\lambda \cos(2\pi(l-1)/m))$ ,  $\lambda \sin(2\pi(l-1)/m)$ ), l = 2, ..., m + 1. Note that

$$G(x, y) = -\frac{\log |x - y|}{2\pi} + \frac{\log (|x|^2 |y|^2 + 1 - 2 \langle x, y \rangle)}{4\pi}, \quad (141)$$
$$H(x, x) = \frac{\log (1 - |x|^2)}{2\pi}.$$

By Theorem 3, we can reduce the problem of finding solutions of (5) to that of finding a  $C^0$ -stable critical point of the function  $f(\lambda) : (0,1) \rightarrow \mathbb{R}$  defined in (8). Obviously,  $\lim_{\lambda \to 0^+} f(\lambda) = \lim_{\lambda \to 1^-} f(\lambda) = +\infty$ , which implies that  $f(\lambda)$  has an absolute minimum point  $\lambda_0$  in (0, 1). Then,  $\lambda_0$  is a  $C^0$ -stable critical point of  $f(\lambda)$ .

# Appendix

*Proofs of* (32) *and* (33). For  $|z_i| \le \delta(v_i \rho_i)^{-1}$ , by (22)-(23),

$$\begin{split} V_{q} &= a_{i}Pu_{i}\left(\varepsilon y\right) + \sum_{j\neq i}a_{j}Pu_{j}\left(\varepsilon y\right) \\ &= a_{i}\left[u_{i}\left(\varepsilon y\right) + d_{i}H\left(\varepsilon y, q_{i}\right)\right. \\ &\left. -\log\frac{8\mu_{i}^{2}\left(1+\alpha_{i}\right)^{2}}{c_{i}\left(q_{i}\right)} + O\left(\rho\right)\right] \\ &+ \sum_{j\neq i}a_{j}\left[d_{j}G\left(\varepsilon y, q_{j}\right) + O\left(\rho\right)\right] \\ &= a_{i}\left\{u_{i}\left(\varepsilon y\right) + d_{i}H\left(q_{i}, q_{i}\right)\right. \\ &\left. -\log\frac{8\mu_{i}^{2}\left(1+\alpha_{i}\right)^{2}}{c_{i}\left(q_{i}\right)} \\ &+ \sum_{j\neq i}a_{i}a_{j}d_{j}G\left(q_{i}, q_{j}\right) \\ &+ O\left(\left|\varepsilon y - q_{i}\right|\right) + O\left(\rho\right)\right\}, \end{split}$$

which, together with the definition (31) of  $\mu_i$ , implies that

$$V_{q} = a_{i} \left\{ u_{i}\left(\varepsilon y\right) + O\left(\left|\varepsilon y - q_{i}\right|\right) + O\left(\rho\right) \right\}.$$
(A.2)

Furthermore, for  $|z_i| \leq \delta(v_i \rho_i)^{-1}$ , a direct computation shows that

$$\begin{split} W_{q} &= \varepsilon^{4} c\left(\varepsilon y\right) S\left(\varepsilon y\right) \left[ e^{V_{q}} + e^{-V_{q}} \right] \\ &= \varepsilon^{4} c_{i}\left(\varepsilon y\right) \left| \varepsilon y - q_{i} \right|^{2\alpha_{i}} \left[ e^{u_{i}(\varepsilon y)} + e^{-u_{i}(\varepsilon y)} \right] \\ &\cdot \left[ 1 + O\left( \left| \varepsilon y - q_{i} \right| \right) + O\left(\rho\right) \right] \\ &= \varepsilon^{4} c_{i}\left(q_{i}\right) \left| \varepsilon y - q_{i} \right|^{2\alpha_{i}} \\ &\times \left[ \frac{8\mu_{i}^{2}(1 + \alpha_{i})^{2}}{c_{i}\left(q_{i}\right) \left[ \mu_{i}^{2}\varepsilon^{2} + \left| x - q_{i} \right|^{2(1 + \alpha_{i})} \right]^{2}} + O\left(1\right) \right] \\ &\cdot \left[ 1 + O\left( \left| \varepsilon y - q_{i} \right| \right) + O\left(\rho\right) \right] \\ &= \left( \frac{\varepsilon}{v_{i}\rho_{i}} \right)^{2} \left| z_{i} \right|^{2\alpha_{i}} \\ &\times \left\{ \frac{8(1 + \alpha_{i})^{2} \left[ 1 + O\left(\rho_{i} \left| z_{i} \right| \right) + O\left(\rho\right) \right]}{\left[ 1 + \left| z_{i} \right|^{2(1 + \alpha_{i})} \right]^{2}} + O\left(\varepsilon^{4}\right) \right\}. \end{split}$$
(A.3)

Similarly, for 
$$|z_i| \leq \delta(v_i \rho_i)^{-1}$$
,  
 $2\varepsilon^4 c(\varepsilon y) S(\varepsilon y) \sinh V_q$   
 $= \left(\frac{\varepsilon}{v_i \rho_i}\right)^2 |z_i|^{2\alpha_i}$   
 $\times \left\{ \frac{8a_i(1+\alpha_i)^2 [1+O(\rho_i |z_i|)+O(\rho)]}{[1+|z_i|^{2(1+\alpha_i)}]^2} + O(\varepsilon^4) \right\}.$ 
(A.4)

On the other hand, if  $|z_i| \ge \delta(v_i \rho_i)^{-1}$ , for any  $i \in \bigcup_{l=1}^3 J_l$ , it is easy to check that

$$W_q = O(\varepsilon^4), \qquad 2\varepsilon^4 c(\varepsilon y) S(\varepsilon y) \sinh V_q = O(\varepsilon^4).$$
 (A.5)

This, together with (A.3), implies (32). Next, by our definitions,

$$\begin{aligned} \Delta V_{q} &= \sum_{i \in \bigcup_{l=1}^{3} J_{l}} \varepsilon^{2} a_{i} \Delta u_{i} \left( \varepsilon y \right) \\ &= \sum_{i \in \bigcup_{l=1}^{3} J_{l}} \varepsilon^{4} a_{i} c_{i} \left( q_{i} \right) \left| \varepsilon y - q_{i} \right|^{2\alpha_{i}} e^{u_{i} \left( \varepsilon y \right)} \\ &= \sum_{i \in \bigcup_{l=1}^{3} J_{l}} \varepsilon^{4} a_{i} c_{i} \left( q_{i} \right) \left| \varepsilon y - q_{i} \right|^{2\alpha_{i}} \\ &\times \frac{8 \mu_{i}^{2} (1 + \alpha_{i})^{2}}{c_{i} \left( q_{i} \right) \left[ \mu_{i}^{2} \varepsilon^{2} + \left| \varepsilon y - q_{i} \right|^{2(1 + \alpha_{i})} \right]^{2}} \end{aligned}$$
(A.6)  
$$&= \sum_{i \in \bigcup_{l=1}^{3} J_{l}} \left( \frac{\varepsilon}{v_{i} \rho_{i}} \right)^{2} \left| z_{i} \right|^{2\alpha_{i}} \\ &\times \frac{8 a_{i} (1 + \alpha_{i})^{2}}{\left[ 1 + \left| z_{i} \right|^{2(1 + \alpha_{i})} \right]^{2}}. \end{aligned}$$

So, if  $|z_i| \ge \delta(v_i \rho_i)^{-1}$ , for all *i*,

$$\Delta V_q = O\left(\varepsilon^4\right),\tag{A.7}$$

while if  $|z_i| \leq \delta(v_i \rho_i)^{-1}$ ,

$$\Delta V_{q} = \left(\frac{\varepsilon}{\nu_{i}\rho_{i}}\right)^{2} |z_{i}|^{2\alpha_{i}} \times \frac{8a_{i}(1+\alpha_{i})^{2}}{\left[1+|z_{i}|^{2(1+\alpha_{i})}\right]^{2}} + O\left(\varepsilon^{4}\right).$$
(A.8)

As a result, combining (A.4)-(A.5) with (A.7)-(A.8), we get (33).  $\hfill \Box$ 

### **Conflict of Interests**

The authors declare that there is no conflict of interests regarding the publication of this paper.

#### Acknowledgments

We would like to thank the anonymous referees for their helpful comments which improved this paper. This research is supported by the National Natural Science Foundation of China (Grant no. 11171214) and the foundation of Nanjing Agricultural University (Grant no. LXYQ201300106).

# References

- A. J. Chorin, "Vorticity and turbulence," in *Applied Mathemati*cal Sciences, vol. 103, Springer, New York, NY, USA, 1994.
- [2] A. J. Chorin and J. E. Marsden, "A mathematical introduction to fluid mechanics," in *Texts in Applied Mathematics*, vol. 4, Springer, New York, NY, USA, 2nd edition, 1990.
- [3] C. Marchioro and M. Pulvirenti, *Mathematical Theory of Incompressible Nonviscous Fluids*, vol. 96, Springer, New York, NY, USA, 1994.
- [4] E. Caglioti, P.-L. Lions, C. Marchioro, and M. Pulvirenti, "A special class of stationary flows for two-dimensional Euler equations: a statistical mechanics description," *Communications in Mathematical Physics*, vol. 143, no. 3, pp. 501–525, 1992.
- [5] E. Caglioti, P.-L. Lions, C. Marchioro, and M. Pulvirenti, "A special class of stationary flows for two-dimensional Euler equations: a statistical mechanics description. II," *Communications in Mathematical Physics*, vol. 174, no. 2, pp. 229–260, 1995.
- [6] G. L. Eyink and H. Spohn, "Negative-temperature states and large-scale, long-lived vortices in two-dimensional turbulence," *Journal of Statistical Physics*, vol. 70, no. 3-4, pp. 833–886, 1993.
- [7] M. K.-H. Kiessling, "Statistical mechanics of classical particles with logarithmic interactions," *Communications on Pure and Applied Mathematics*, vol. 46, no. 1, pp. 27–56, 1993.
- [8] J. T. Stuart, "On finite amplitude oscillations in laminar mixing layers," *Journal of Fluid Mechanics*, vol. 29, pp. 417–440, 1967.
- [9] D. Gurarie and K. W. Chow, "Vortex arrays for sinh-Poisson equation of two-dimensional fluids: equilibria and stability," *Physics of Fluids*, vol. 16, no. 9, pp. 3296–3305, 2004.
- [10] R. Mallier and S. A. Maslowe, "A row of counter-rotating vortices," *Physics of Fluids A*, vol. 5, no. 4, pp. 1074–1075, 1993.
- [11] A. Tur and V. Yanovsky, "Point vortices with a rational necklace: new exact stationary solutions of the two-dimensional Euler equation," *Physics of Fluids*, vol. 16, no. 8, pp. 2877–2885, 2004.
- [12] Y. Wang and L. Wei, "Nodal bubbling solutions to a weighted sinh-Poisson equation," *Advances in Differential Equations*, vol. 13, no. 9-10, pp. 881–906, 2008.
- [13] G. R. Flierl, "Isolated eddy models in geophysics," Annual Review of Fluid Mechanics, vol. 19, pp. 493–530, 1987.
- [14] E. J. Hopfinger and G. J. F. van Heijst, "Vortices in rotating fluids," *Annual Review of Fluid Mechanics*, vol. 25, pp. 241–289, 1993.
- [15] D. Marteau, O. Cardoso, and P. Tabeling, "Equilibrium states of twodimensional turbulence: an experimental study," *Physical Review E*, vol. 51, pp. 5124–5127, 1995.
- [16] H. Brezis and J.-M. Coron, "Multiple solutions of H-systems and Rellich's conjecture," *Communications on Pure and Applied Mathematics*, vol. 37, no. 2, pp. 149–187, 1984.

- [17] K. Steffen, "On the nonuniqueness of surfaces with constant mean curvature spanning a given contour," *Archive for Rational Mechanics and Analysis*, vol. 94, no. 2, pp. 101–122, 1986.
- [18] J. Spruck, "The elliptic sinh Gordon equation and the construction of toroidal soap bubbles," in *Calculus of Variations and Partial Differential Equations*, vol. 1340 of *Lecture Notes in Mathematics*, pp. 275–301, Springer, Berlin, Germany, 1988.
- [19] M. Struwe, "Nonuniqueness in the Plateau problem for surfaces of constant mean curvature," *Archive for Rational Mechanics and Analysis*, vol. 93, no. 2, pp. 135–157, 1986.
- [20] H. C. Wente, "Counterexample to a conjecture of *H. Hopf*," *Pacific Journal of Mathematics*, vol. 121, no. 1, pp. 193–243, 1986.
- [21] J. Jost, G. Wang, D. Ye, and C. Zhou, "The blow up analysis of solutions of the elliptic sinh-Gordon equation," *Calculus of Variations and Partial Differential Equations*, vol. 31, no. 2, pp. 263–276, 2008.
- [22] H. Ohtsuka and T. Suzuki, "Mean field equation for the equilibrium turbulence and a related functional inequality," *Advances in Differential Equations*, vol. 11, no. 3, pp. 281–304, 2006.
- [23] M. Grossi and A. Pistoia, "Multiple blow-up phenomena for the sinh-Poisson equation," *Archive for Rational Mechanics and Analysis*, vol. 209, no. 1, pp. 287–320, 2013.
- [24] P. Esposito and J. Wei, "Non-simple blow-up solutions for the Neumann two-dimensional sinh-Gordon equation," *Calculus of Variations and Partial Differential Equations*, vol. 34, no. 3, pp. 341–375, 2009.
- [25] D. Bartolucci and A. Pistoia, "Existence and qualitative properties of concentrating solutions for the sinh-Poisson equation," *IMA Journal of Applied Mathematics*, vol. 72, no. 6, pp. 706–729, 2007.
- [26] T. Bartsch, A. Pistoia, and T. Weth, "N-vortex equilibria for ideal fluids in bounded planar domains and new nodal solutions of the sinh-Poisson and the Lane-Emden-Fowler equations," *Communications in Mathematical Physics*, vol. 297, no. 3, pp. 653–686, 2010.
- [27] J. Wei, L. Wei, and F. Zhou, "Mixed interior and boundary nodal bubbling solutions for a sinh-Poisson equation," *Pacific Journal* of *Mathematics*, vol. 250, no. 1, pp. 225–256, 2011.
- [28] H. Brezis and F. Merle, "Uniform estimates and blow-up behavior for solutions of  $-\Delta u = V(x)e^u$  in two dimensions," *Communications in Partial Differential Equations*, vol. 16, no. 8-9, pp. 1223–1253, 1991.
- [29] C.-C. Chen and C.-S. Lin, "Mean field equations of Liouville type with singular data: sharper estimates," *Discrete and Continuous Dynamical Systems A*, vol. 28, no. 3, pp. 1237–1272, 2010.
- [30] Y. Y. Li and I. Shafrir, "Blow-up analysis for solutions of −△u = Ve<sup>u</sup> in dimension two," *Indiana University Mathematics Journal*, vol. 43, no. 4, pp. 1255–1270, 1994.
- [31] L. Ma and J. C. Wei, "Convergence for a Liouville equation," *Commentarii Mathematici Helvetici*, vol. 76, no. 3, pp. 506–514, 2001.
- [32] K. Nagasaki and T. Suzuki, "Asymptotic analysis for two-dimensional elliptic eigenvalue problems with exponentially dominated nonlinearities," *Asymptotic Analysis*, vol. 3, no. 2, pp. 173– 188, 1990.
- [33] G. Tarantello, "Analytical, geometrical and topological aspects of a class of mean field equations on surfaces," *Discrete and Continuous Dynamical Systems A*, vol. 28, no. 3, pp. 931–973, 2010.
- [34] S. Baraket and F. Pacard, "Construction of singular limits for a semilinear elliptic equation in dimension 2," *Calculus of Variations and Partial Differential Equations*, vol. 6, no. 1, pp. 1–38, 1998.

- [35] P. Esposito, "Blowup solutions for a Liouville equation with singular data," *SIAM Journal on Mathematical Analysis*, vol. 36, no. 4, pp. 1310–1345, 2005.
- [36] Y. Chang and H. Yang, "Multiple blowing-up and concentrating solutions for Liouville-type equations with singular sources under mixed boundary conditions," *Boundary Value Problems*, vol. 2012, article 33, 2012.
- [37] T. D. 'Aprile, "Multiple blow-up solutions for the Liouville equation with singular data," *Communications in Partial Differential Equations*, vol. 38, no. 8, pp. 1409–1436, 2013.
- [38] M. del Pino, P. Esposito, and M. Musso, "Two-dimensional Euler flows with concentrated vorticities," *Transactions of the American Mathematical Society*, vol. 362, no. 12, pp. 6381–6395, 2010.
- [39] M. del Pino, M. Kowalczyk, and M. Musso, "Singular limits in Liouville-type equations," *Calculus of Variations and Partial Differential Equations*, vol. 24, no. 1, pp. 47–81, 2005.
- [40] P. Esposito, M. Grossi, and A. Pistoia, "On the existence of blowing-up solutions for a mean field equation," *Annales de l'Institut Henri Poincaré. Analyse Non Linéaire*, vol. 22, no. 2, pp. 227–257, 2005.
- [41] P. Esposito, A. Pistoia, and J. Wei, "Concentrating solutions for the Hénon equation in R<sup>2</sup>," *Journal d'Analyse Mathématique*, vol. 100, pp. 249–280, 2006.
- [42] W. X. Chen and C. Li, "Classification of solutions of some nonlinear elliptic equations," *Duke Mathematical Journal*, vol. 63, no. 3, pp. 615–622, 1991.
- [43] W. X. Chen and C. Li, "Qualitative properties of solutions to some nonlinear elliptic equations in ℝ<sup>2</sup>," *Duke Mathematical Journal*, vol. 71, no. 2, pp. 427–439, 1993.
- [44] J. Prajapat and G. Tarantello, "On a class of elliptic problems in  $\mathbb{R}^2$ : symmetry and uniqueness results," *Proceedings of the Royal Society of Edinburgh A*, vol. 131, no. 4, pp. 967–985, 2001.
- [45] C.-C. Chen and C.-S. Lin, "Sharp estimates for solutions of multi-bubbles in compact Riemann surfaces," *Communications* on *Pure and Applied Mathematics*, vol. 55, no. 6, pp. 728–771, 2002.
- [46] M. del Pino, P. Esposito, and M. Musso, "Nondegeneracy of entire solutions of a singular Liouvillle equation," *Proceedings of the American Mathematical Society*, vol. 140, no. 2, pp. 581–588, 2012.



The Scientific World Journal





**Decision Sciences** 







Journal of Probability and Statistics



Hindawi Submit your manuscripts at http://www.hindawi.com



(0,1),

International Journal of Differential Equations





International Journal of Combinatorics





Mathematical Problems in Engineering



Abstract and Applied Analysis



Discrete Dynamics in Nature and Society







Function Spaces



International Journal of Stochastic Analysis



Journal of Optimization