

Research Article

A Remark on the Regularity Criterion for the 3D Boussinesq Equations Involving the Pressure Gradient

Zujin Zhang

School of Mathematics and Computer Science, Gannan Normal University, Ganzhou 341000, China

Correspondence should be addressed to Zujin Zhang; zhangzujin361@163.com

Received 10 November 2013; Accepted 28 December 2013; Published 21 January 2014

Academic Editor: Giovanni P. Galdi

Copyright © 2014 Zujin Zhang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We consider the three-dimensional Boussinesq equations and obtain a regularity criterion involving the pressure gradient in the Morrey-Companato space $M_{p,q}$. This extends and improves the result of Gala (Gala 2013) for the Navier-Stokes equations.

1. Introduction

This paper concerns itself with the following three-dimensional (3D) Boussinesq equations:

$$\mathbf{u}_{t} + (\mathbf{u} \cdot \nabla) \,\mathbf{u} - \Delta \mathbf{u} + \nabla \pi = \theta \mathbf{e}_{3}, \quad \text{in } \mathbb{R}^{3} \times (0, T) \,,$$

$$\theta_{t} + (\mathbf{u} \cdot \nabla) \,\theta - \Delta \theta = 0, \quad \text{in } \mathbb{R}^{3} \times (0, T) \,,$$

$$\nabla \cdot \mathbf{u} = 0, \quad \text{in } \mathbb{R}^{3} \times (0, T) \,,$$

$$\mathbf{u} (0) = \mathbf{u}_{0}, \quad \theta (0) = \theta_{0}, \quad \text{on } \mathbb{R}^{3},$$

(1)

where T > 0 is given time, $\mathbf{u} = (u_1(x, t), u_2(x, t), u_3(x, t))$ is the fluid velocity, $\pi = \pi(x, t)$ is a scalar pressure, and $\theta = \theta(x, t)$ is the temperature, while \mathbf{u}_0 and θ_0 are the prescribed initial velocity field and temperature, respectively.

When $\theta = 0$, (1) reduces to the incompressible Navier-Stokes equations. The regularity of its weak solutions and the existence of global strong solutions are challenging open problems; see [1–3]. Starting with [4, 5], there have been a lot of literature devoted to finding sufficient conditions to ensure the smoothness of the solutions; see [6–15] and the references cited therein. Since the convective terms are similar in the Navier-Stokes equations and Boussinesq equations, the authors also consider the regularity conditions for (1); see [16–20] and so forth.

In [6], Gala uses intricate decomposition technique to obtain the following regularity criterion for the Navier-Stokes equations:

$$\nabla \pi \in L^{2/(3-r)}\left(0,T;\dot{X}_r\right) \quad \text{with } 0 \le r \le 1.$$

Here, \dot{X}_r is the point-wise multiplier space from \dot{H}^r to L^2 , which is strictly larger than $L^{3/r}(\mathbb{R}^3)$ (see [6, Lemma 1.2]).

In this paper, we will extend and improve the regularity condition (2) to the Boussineq equations (1).

Before stating the precise result, let us recall the weak formulation of (1).

Definition 1. Let $\mathbf{u}_0 \in L^2(\mathbb{R}^3)$, $\theta_0 \in L^1 \cap L^\infty(\mathbb{R}^3)$. A measurable pair (\mathbf{u}, θ) is said to be a weak solution of (1) in (0, T), provided that

- (1) $(\mathbf{u}, \theta) \in L^{\infty}(0, T; L^{2}(\mathbb{R}^{3})) \cap L^{2}(0, T; H^{1}(\mathbb{R}^{3})), \theta \in L^{\infty}$ $(0, T; L^{1} \cap L^{\infty}(\mathbb{R}^{3}));$
- (2) $(1)_{1,2,3}$ are satisfied in the sense of distributions;
- (3) the energy inequality

$$\begin{aligned} \|(\mathbf{u},\theta)\|_{L^{2}}^{2} + 2 \int_{0}^{t} \|\nabla(\mathbf{u},\theta)\|_{L^{2}}^{2} ds \\ \leq \|(\mathbf{u}_{0},\theta_{0})\|_{L^{2}}^{2} \\ + 2 \int_{0}^{t} \int_{\mathbb{R}^{3}} \theta u_{3} dx ds, \end{aligned}$$
(3)

for all $0 \le t \le T$.

Now, our main result reads the following.

Theorem 2. Let $\mathbf{u}_0 \in L^2(\mathbb{R}^3)$ with $\nabla \cdot \mathbf{u}_0 = 0$ in the sense of distributions, $\theta_0 \in L^1 \cap L^\infty(\mathbb{R}^3)$. Supposing that (\mathbf{u}, θ) is a weak solution of (1) in [0, T), and the pressure gradient $\nabla \pi$ satisfies

$$\nabla \pi \in L^{2/(3-r)}\left(0, T; \dot{M}_{2,3/r}\right) \quad \text{with } 0 < r \le 1, \qquad (4)$$

then the solution $(\mathbf{u}, \theta) \in C^{\infty}((0, T) \times \mathbb{R}^3)$.

Here, $\dot{M}_{p,q}$ is the Morrey-Campanato space, which will be introduced in Section 2. And Section 3 is devoted to the proof of Theorem 2.

Remark 3. Noticing that $X_r \,\subset M_{2,3/r}$ for 0 < r < 1 (see (10)), we indeed improve the result of [6] for the Navier-Stokes equations.

2. Preliminaries

In this section, we will introduce the definition of Morrey-Campanato space $\dot{M}_{p,q}$ and recall its fundamental properties. The space plays an important role in studying the regularity of solutions to partial differential equations (see [21–23], e.g.).

Definition 4. For $1 , the Morrey-Campanato space <math>\dot{M}_{p,q}$ is defined as

$$\dot{M}_{p,q} = \left\{ f \in L^{p}_{\text{loc}} \left(\mathbb{R}^{3} \right); \| f \|_{\dot{M}_{p,q}} \right.$$
$$= \sup_{x \in \mathbb{R}^{3}, R > 0} \frac{1}{R^{(3/p) - (3/q)}} \left(\int_{B(x,R)} \left| f(y) \right|^{p} \, \mathrm{d}y \right)^{1/p} (5)$$
$$< +\infty \right\},$$

where $B(x, R) \in \mathbb{R}^3$ is the ball with center x and radius R.

One sees readily that $\dot{M}_{p,q}$ is a Banach space under the norm $\|\cdot\|_{\dot{M}_{p,q}}$ and contains the classical Lebesgue space as a subspace:

$$L^q = \dot{M}_{q,q} \subset \dot{M}_{p,q}.$$
 (6)

Moreover, the following scaling property holds:

$$\|f(\lambda \cdot)\|_{\dot{M}_{p,q}} = \frac{1}{\lambda^{3/q}} \|f\|_{\dot{M}_{p,q}}, \quad \text{for } \lambda > 0.$$
 (7)

Due to the following characterization in [24].

Lemma 5. For $0 \le r < 3/2$, the space \dot{Z}_r is defined as the space of all functions $f \in L^2_{loc}(\mathbb{R}^3)$ such that

$$\|f\|_{\dot{Z}_{r}} = \sup_{\|g\|_{\dot{B}_{2,1}^{r}} \le 1} \|fg\|_{L^{2}} < +\infty.$$
(8)

Then $f \in \dot{M}_{2,3/r}$ if and only if $f \in \dot{Z}_r$ with equivalent norm.

And with the fact that

$$L^{2} \cap \dot{H}^{r} \subset \dot{B}^{r}_{2,1} \subset \dot{H}^{r} \quad \text{for } 0 < r < 1,$$
(9)

we have

$$\dot{X}_r \in \dot{M}_{2,3/r}.\tag{10}$$

Here $\dot{B}_{2,1}^r$ is the Besov space, which is intermediate between L^2 and \dot{H}^1 (see [25]):

$$\|f\|_{\dot{B}^{r}_{2,1}} \le C \|f\|_{L^{2}}^{1-r} \|\nabla f\|_{L^{2}}^{r}, \quad \text{for } 0 < r < 1.$$
(11)

3. Proof of Theorem 2

In this section, we will prove Theorem 2.

Due to the Serrin type regularity criterion

$$\mathbf{u} \in L^{p}\left(0, T; L^{q}\left(\mathbb{R}^{3}\right)\right) \quad \text{with } \frac{2}{p} + \frac{3}{q} = 1, \ 3 < q \le +\infty$$

$$(12)$$

in [19], we need only to prove

$$\mathbf{u} \in L^{\infty}\left(0, T; L^{4}\left(\mathbb{R}^{4}\right)\right) \subset L^{8}\left(0, T; L^{4}\left(\mathbb{R}^{3}\right)\right).$$
(13)

We just do a priori estimates, with the justification being from passage to limits for the Galerkin approximated solutions.

Taking the inner product of $(1)_2$ with 2θ in $L^2(\mathbb{R}^3)$, we find

$$\frac{d}{dt}\|\theta\|_{L^2}^2 + 2\|\nabla\theta\|_{L^2}^2 = 0.$$
 (14)

Thus,

$$\|\theta\|_{L^2} \le \|\theta_0\|_{L^2}.$$
 (15)

One can also take the inner product of $(1)_2$ with $p\theta^{p-1}$ $(1 \le p < \infty)$ in $L^2(\mathbb{R}^3)$ to derive the estimate of θ in L^p -norm and invoke the maximum principle to bound the L^∞ -norm of θ , as stated in Definition 1.

Taking the divergence of $(1)_1$, we get

$$-\Delta \pi = \sum_{i,j=1}^{3} \partial_i \partial_j \left(u_i u_j \right) - \partial_3 \theta.$$
 (16)

Consequently,

$$\|\nabla \pi\|_{L^{2}} \leq C \|\|\mathbf{u}\| \cdot \|\nabla \mathbf{u}\|\|_{L^{2}} + \|\theta\|_{L^{2}} \leq C \left(\|\|\mathbf{u}\| \cdot \|\nabla \mathbf{u}\|\|_{L^{2}} + 1 \right).$$
(17)

Taking the inner product of $(1)_1$ with $4|\mathbf{u}|^2\mathbf{u}$ in $L^2(\mathbb{R}^3)$, we get

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}t} \left\| \left| \mathbf{u} \right|^{2} \right\|_{L^{2}}^{2} + 4 \int_{\mathbb{R}^{3}} \left| \mathbf{u} \right|^{2} \left| \nabla \mathbf{u} \right|^{2} \mathrm{d}x + 2 \int_{\mathbb{R}^{3}} \left| \nabla \left| \mathbf{u} \right|^{2} \right| \mathrm{d}x \\ &\leq 4 \int_{\mathbb{R}^{3}} \left| \nabla \pi \right| \cdot \left| \mathbf{u} \right|^{3} \mathrm{d}x + 4 \int_{\mathbb{R}^{3}} \left| \theta \right| \cdot \left| \mathbf{u} \right|^{3} \mathrm{d}x \\ &\equiv I_{1} + I_{2}. \end{aligned}$$
(18)

For I_1 , we estimate as

The term I_2 can be dominated as

$$\begin{split} I_{2} &\leq 4 \|\theta\|_{L^{2}} \left\| |\mathbf{u}|^{3} \right\|_{L^{2}} \\ &\leq C \left\| |\mathbf{u}|^{2} \right\|_{L^{3}}^{3/2} \quad (by (15)) \\ &\leq C \left\| |\mathbf{u}|^{2} \right\|_{L^{2}}^{3/4} \left\| \nabla |\mathbf{u}|^{2} \right\|_{L^{2}}^{3/4} \quad (by \text{ interpolation inequality}) \\ &\leq C \left\| |\mathbf{u}|^{2} \right\|_{L^{2}}^{6/5} + \frac{1}{2} \left\| \nabla |\mathbf{u}|^{2} \right\|_{L^{2}}^{2} \\ &\leq C + C \left\| |\mathbf{u}|^{2} \right\|_{L^{2}}^{2} + \frac{1}{2} \left\| \nabla |\mathbf{u}|^{2} \right\|_{L^{2}}^{2}. \end{split}$$

$$(20)$$

Plugging (19) and (20) into (18), we deduce that

$$\frac{d}{dt} \left\| \left| \mathbf{u} \right|^{2} \right\|_{L^{2}}^{2} + \int_{\mathbb{R}^{3}} \left| \mathbf{u} \right|^{2} \left| \nabla \mathbf{u} \right|^{2} dx + \int_{\mathbb{R}^{3}} \left| \nabla \left| \mathbf{u} \right|^{2} \right| dx
\leq C + C \left(\left\| \nabla \pi \right\|_{\dot{M}_{2,3/r}}^{2/(3-r)} + 1 \right) \left\| \left| \mathbf{u} \right|^{2} \right\|_{L^{2}}^{2}.$$
(21)

Applying Gronwall inequality, we see that

$$\|\mathbf{u}(t)\|_{L^{4}}^{4} \leq \left(\|\mathbf{u}_{0}\|_{L^{4}}^{4} + CT\right) \\ \cdot \exp\left\{C\int_{0}^{T}\left(\|\nabla\pi\|_{\dot{M}_{2,3/r}}^{2/(3-r)} + 1\right)\mathrm{d}s\right\},$$
(22)

for every $t \in [0, T]$. Recalling (13), we complete the proof of Theorem 2.

Conflict of Interests

The author declares that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

Zujin Zhang was partially supported by the Youth Natural Science Foundation of Jiangxi Province (20132BAB211007),

the Science Foundation of Jiangxi Provincial Department of Education (GJJ13658, GJJ13659), and the National Natural Science Foundation of China (11326138, 11361004, and 11326238).

References

- E. Hopf, "Über die anfangswertaufgabe für die hydrodynamischen Grundgleichungen," *Mathematische Nachrichten*, vol. 4, pp. 213–231, 1951.
- [2] P. G. Lemarié-Rieusset, Recent Developments in the Navier-Stokes Problem, Chapman & Hall, Boca Raton, Fla, USA, 2002.
- [3] J. Leray, "Sur le mouvement d'un liquide visqueux emplissant l'espace," Acta Mathematica, vol. 63, no. 1, pp. 193–248, 1934.
- [4] J. Serrin, "On the interior regularity of weak solutions of the Navier-Stokes equations," *Archive for Rational Mechanics and Analysis*, vol. 9, no. 1, pp. 187–195, 1962.
- [5] J. Serrin, "The initial value problem for the Navier-Stokes equations," in *Nonlinear Problems*, R. E. Langer, Ed., University of Wisconsin Press, Madison, Wis, USA, 1963.
- [6] S. Gala, "Remarks on regularity criterion for weak solutions to the Navier-Stokes equations in terms of the gradient of the pressure," *Applicable Analysis*, vol. 92, no. 1, pp. 96–103, 2013.
- [7] X. He and S. Gala, "Regularity criterion for weak solutions to the Navier-Stokes equations in terms of the pressure in the class $L^2(0, T; \dot{B}_{\infty,\infty}^{-1}(\mathbb{R}^3)$," *Nonlinear Analysis: Real World Applications*, vol. 12, no. 6, pp. 3602–3607, 2011.
- [8] J. Neustupa, A. Novotný, and P. Penel, "An interior regularity of a weak solution to the Navier-Stokes equations in dependence on one component of velocity, topics in mathematical uid mechanics," *Quaderni di Matematica*, vol. 10, pp. 163–183, 2002.
- [9] Z. Zhang, "A serrin-type regularity criterion for the Navier-Stokes equations via one velocity component," *Communications* on Pure and Applied Analysis, vol. 12, no. 1, pp. 117–124, 2013.
- [10] Z. Zhang, Z.-A. Yao, P. Li, C. Guo, and M. Lu, "Two new regularity criteria for the 3D Navier-Stokes equations via two entries of the velocity gradient tensor," *Acta Applicandae Mathematicae*, vol. 123, no. 1, pp. 43–52, 2013.
- [11] Z. J. Zhang, D. X. Zhong, and L. Hu, "A new regularity criterion for the 3D Navier-Stokes equations via two entries of the velocity gradient tensor," *Acta Applicandae Mathematicae*, 2013.

- [12] Z. Zhang, Z.-A. Yao, M. Lu, and L. Ni, "Some serrin-type regularity criteria for weak solutions to the Navier-Stokes equations," *Journal of Mathematical Physics*, vol. 52, no. 5, Article ID 053103, 2011.
- [13] Y. Zhou, "A new regularity criterion for weak solutions to the Navier-Stokes equations," *Journal des Mathematiques Pures et Appliquees*, vol. 84, no. 11, pp. 1496–1514, 2005.
- [14] Y. Zhou and M. Pokorný, "On a regularity criterion for the Navier-Stokes equations involving gradient of one velocity component," *Journal of Mathematical Physics*, vol. 50, no. 12, Article ID 123514, 2009.
- [15] Y. Zhou and M. Pokorný, "On the regularity of the solutions of the Navier-Stokes equations via one velocity component," *Nonlinearity*, vol. 23, no. 5, pp. 1097–1107, 2010.
- [16] J. S. Fan and Y. Zhou, "A note on regularity criterion for the 3D Boussinesq system with partial viscosity," *Applied Mathematics Letters*, vol. 22, no. 5, pp. 802–805, 2009.
- [17] N. Ishimura and H. Morimoto, "Remarks on the blow-up criterion for the 3-D Boussinesq equations," *Mathematical Models and Methods in Applied Sciences*, vol. 9, no. 9, pp. 1323– 1332, 1999.
- [18] H. Qiu, Y. Du, and Z. Yao, "Blow-up criteria for 3D Boussinesq equations in the multiplier space," *Communications in Nonlinear Science and Numerical Simulation*, vol. 16, no. 4, pp. 1820– 1824, 2011.
- [19] H. Qiu, Y. Du, and Z. Yao, "Serrin-type blow-up criteria for 3D Boussinesq equations," *Applicable Analysis*, vol. 89, no. 10, pp. 1603–1613, 2010.
- [20] Z. J. Zhang, "A logarithmically improved regularity criterion for the 3D Boussinesq equations via the pressure," Acta Applicandae Mathematicae, 2013.
- [21] T. Kato, "Strong solutions of the Navier-Stokes equation in Morrey spaces," *Boletim da Sociedade Brasileira de Matemática*, vol. 22, no. 2, pp. 127–155, 1992.
- [22] M. E. Taylor, "Analysis on Morrey spaces and applications to Navier-Stokes and other evolution equations," *Communications in Partial Differential Equations*, vol. 17, no. 9-10, pp. 1407–1456, 1992.
- [23] Z. Zhang, "Remarks on the regularity criteria for generalized MHD equations," *Journal of Mathematical Analysis and Applications*, vol. 375, no. 2, pp. 799–802, 2011.
- [24] P. G. Lemarié-Rieusset, Recent Developments in the Navier-Stokes Problem, Chapman & Hall, London, UK, 2002.
- [25] Y. Zhou and S. Gala, "A new regularity criterion for weak solutions to the viscous MHD equations in terms of the vorticity field," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 72, no. 9-10, pp. 3643–3648, 2010.



The Scientific World Journal





Decision Sciences







Journal of Probability and Statistics



Hindawi Submit your manuscripts at http://www.hindawi.com



(0,1),

International Journal of Differential Equations





International Journal of Combinatorics





Mathematical Problems in Engineering



Abstract and Applied Analysis



Discrete Dynamics in Nature and Society







Function Spaces



International Journal of Stochastic Analysis



Journal of Optimization