

Research Article

Differential Subordination Results for Analytic Functions in the Upper Half-Plane

Huo Tang,^{1,2} M. K. Aouf,³ Guan-Tie Deng,² and Shu-Hai Li¹

¹ School of Mathematics and Statistics, Chifeng University, Chifeng, Inner Mongolia 024000, China

² School of Mathematical Sciences, Beijing Normal University, Beijing 100875, China

³ Department of Mathematics, Faculty of Science, Mansoura University, Mansoura 35516, Egypt

Correspondence should be addressed to Huo Tang; thth2009@tom.com

Received 11 November 2013; Accepted 19 January 2014; Published 25 February 2014

Academic Editor: V. Ravichandran

Copyright © 2014 Huo Tang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

There are many articles in the literature dealing with differential subordination problems for analytic functions in the unit disk, and only a few articles deal with the above problems in the upper half-plane. In this paper, we aim to derive several differential subordination results for analytic functions in the upper half-plane by investigating certain suitable classes of admissible functions. Some useful consequences of our main results are also pointed out.

1. Introduction

Let Δ denote the upper half-plane; that is,

$$\Delta = \{z \in \mathbb{C} : \operatorname{Im}(z) > 0\}, \quad (1)$$

and let $\mathcal{H}[\Delta]$ denote the class of functions f which are analytic in Δ and which satisfy the so-called hydrodynamic normalization (see [1–3]):

$$\lim_{\Delta \ni z \rightarrow \infty} [f(z) - z] = 0. \quad (2)$$

Also let $\mathcal{S}[\Delta]$ denote the class of all functions in $\mathcal{H}[\Delta]$ which are univalent in Δ . Various basic properties concerning functions belonging to the class $\mathcal{S}[\Delta]$ were developed in a series of articles (see, for details, [4–6]).

A function $f \in \mathcal{H}[\Delta]$, with $f(z) \neq 0$, is said to be starlike in Δ if and only if

$$\operatorname{Im} \left\{ \frac{f'(z)}{f(z)} \right\} < 0. \quad (3)$$

We denote by $\mathcal{S}^*[\Delta]$ the subclass of $\mathcal{H}[\Delta]$ which consists of functions which are starlike in Δ .

A function $f \in \mathcal{H}[\Delta]$, with $f(z) \neq z$, is said to be convex in Δ if and only if

$$\operatorname{Im} \left\{ \frac{f''(z)}{f'(z)} \right\} > 0. \quad (4)$$

Also, we denote by $\mathcal{K}[\Delta]$ the subclass of $\mathcal{H}[\Delta]$ which consists of functions which are convex in Δ . The classes $\mathcal{S}^*[\Delta]$ and $\mathcal{K}[\Delta]$ were introduced by Stankiewicz [3].

We first need to recall the notion of subordination in the upper half-plane.

Let f and g be members of $\mathcal{H}[\Delta]$. The function f is subordinate to g , written as $f < g$ or $f(z) < g(z)$, if there exists a function $\varphi \in \mathcal{H}[\Delta]$ with $\varphi[\Delta] \subset \Delta$ such that $f(z) = g(\varphi(z))$. Furthermore, if the function g is univalent in Δ , then we have the following equivalence (cf. [7]):

$$f(z) < g(z) \quad (z \in \Delta) \iff f(\Delta) \subset g(\Delta). \quad (5)$$

Using methods similar to those used in the unit disk, Răducanu and Pascu [7] have extended the theory of differential subordinations to the upper half-plane. In the following, we will list some definitions and theorems, which are required to prove our main results.

Definition 1 (see [8, Definition 8.3i, p.403]). Denote by $\mathcal{Q}(\Delta)$ the set of functions $q \in \mathcal{H}[\Delta]$ that are analytic and injective on $\bar{\Delta} \setminus E(q)$, where

$$E(q) = \left\{ \xi \in \partial\Delta : \lim_{z \rightarrow \xi} q(z) = \infty \right\}, \quad (6)$$

and are such that $q'(\xi) \neq 0$ for $\xi \in \partial\Delta \setminus E(q)$.

Definition 2 (see [7]). Let Ω be a set in \mathbb{C} and $q \in \mathcal{Q}(\Delta)$. The class of admissible functions $\Psi_{\Delta}[\Omega, q]$ consists of those functions $\psi : \mathbb{C}^3 \times \Delta \rightarrow \mathbb{C}$ that satisfy the following admissibility condition:

$$\psi(r, s, t; z) \notin \Omega, \quad (7)$$

whenever

$$r = q(\xi), \quad s = kq'(\xi), \quad \operatorname{Im} \left\{ \frac{t}{q'(\xi)} \right\} \geq k^2 \operatorname{Im} \left\{ \frac{q''(\xi)}{q'(\xi)} \right\}, \quad (8)$$

where $z \in \Delta$, $\xi \in \partial\Delta \setminus E(q)$, and $k \geq 0$.

If $\psi : \mathbb{C}^2 \times \Delta \rightarrow \mathbb{C}$, then the admissibility condition reduces to

$$\psi(q(\xi), kq'(\xi); z) \notin \Omega, \quad (9)$$

where $z \in \Delta$, $\xi \in \partial\Delta \setminus E(q)$, and $k \geq 0$.

Theorem 3 (see [7]). Let $\psi \in \Psi_{\Delta}[\Omega, q]$ and $p \in \mathcal{H}[\Delta]$. If

$$\psi(p(z), p'(z), p''(z); z) \in \Omega, \quad (10)$$

for $z \in \Delta$, then

$$p(z) < q(z). \quad (11)$$

In the present paper, by making use of the differential subordination results in the upper half-plane of Răducanu and Pascu [7] (which is a generalization of results in the unit disk obtained by Miller and Mocanu [8]), we determine certain appropriate classes of admissible functions and investigate some differential subordination properties of analytic functions in the upper half-plane. It should be remarked in passing that, in recent years, several authors obtained many interesting results associated with differential subordination and superordination in the unit disk; the interested reader may refer to, for example, [9–18].

2. The Main Subordination Results

We first define the following class of admissible functions that are required in proving our first result.

Definition 4. Let Ω be a set in \mathbb{C} and $q \in \mathcal{Q}(\Delta) \cap \mathcal{H}[\Delta]$. The class of admissible functions $\Phi_{\Delta}[\Omega, q]$ consists of those functions $\phi : \mathbb{C}^3 \times \Delta \rightarrow \mathbb{C}$ that satisfy the following admissibility condition:

$$\phi(u, v, w; z) \notin \Omega, \quad (12)$$

whenever

$$\begin{aligned} u &= q(\xi), \quad v = \frac{kq'(\xi)}{q'(\xi)} \quad (q(\xi) \neq 0), \\ \operatorname{Im} \left\{ \frac{u(wv + v^2)}{q'(\xi)} \right\} \\ &\geq k^2 \operatorname{Im} \left\{ \frac{q''(\xi)}{q'(\xi)} \right\} \quad (z \in \Delta; \xi \in \partial\Delta \setminus E(q); k \geq 0). \end{aligned} \quad (13)$$

Theorem 5. Let $\phi \in \Phi_{\Delta}[\Omega, q]$. If $f \in \mathcal{H}[\Delta]$ satisfies

$$\begin{aligned} &\left\{ \phi \left(\frac{f'(z)}{f(z)}, \frac{f''(z)}{f'(z)} - \frac{f'(z)}{f(z)}, \right. \right. \\ &\quad \left. \frac{f(z) [f'''(z) f'(z) - (f''(z))^2]}{f'(z) [f(z) f''(z) - (f'(z))^2]} \right. \\ &\quad \left. \left. - \frac{f'(z)}{f(z)}; z \right) : z \in \Delta \right\} \subset \Omega, \end{aligned} \quad (14)$$

then

$$\frac{f'(z)}{f(z)} < q(z) \quad (z \in \Delta). \quad (15)$$

Proof. Define the function $p(z)$ in Δ by

$$p(z) = \frac{f'(z)}{f(z)}. \quad (16)$$

A simple calculation yields

$$\frac{f''(z)}{f'(z)} - \frac{f'(z)}{f(z)} = \frac{p'(z)}{p(z)}. \quad (17)$$

Further computations show that

$$\begin{aligned} &\frac{f(z) [f'''(z) f'(z) - (f''(z))^2]}{f'(z) [f(z) f''(z) - (f'(z))^2]} - \frac{f'(z)}{f(z)} \\ &= \frac{p''(z)}{p'(z)} - \frac{p'(z)}{p(z)}. \end{aligned} \quad (18)$$

We now define the transformation from \mathbb{C}^3 to \mathbb{C} by

$$u(r, s, t) = r, \quad v(r, s, t) = \frac{s}{r}, \quad w(r, s, t) = \frac{rt - s^2}{rs}. \quad (19)$$

Let

$$\psi(r, s, t; z) = \phi(u, v, w; z) = \phi \left(r, \frac{s}{r}, \frac{rt - s^2}{rs}; z \right). \quad (20)$$

Using (16)–(18), and from (20), we obtain

$$\begin{aligned} & \psi(p(z), p'(z), p''(z); z) \\ &= \phi\left(\frac{f'(z)}{f(z)}, \frac{f''(z)}{f'(z)} - \frac{f'(z)}{f(z)}, \right. \\ & \quad \left. \frac{f(z) \left[f'''(z) f'(z) - (f''(z))^2 \right]}{f'(z) \left[f(z) f''(z) - (f'(z))^2 \right]} \right. \\ & \quad \left. - \frac{f'(z)}{f(z)}; z\right). \end{aligned} \quad (21)$$

Hence, (14) becomes

$$\psi(p(z), p'(z), p''(z); z) \in \Omega. \quad (22)$$

From (19), we easily get

$$t = u(wv + v^2). \quad (23)$$

Thus, the admissibility condition for $\phi \in \Phi_\Delta[\Omega, q]$ in Definition 4 is equivalent to the admissibility condition for ψ as given in Definition 2. Therefore $\psi \in \Psi_\Delta[\Omega, q]$, and by Theorem 3, we have $p(z) \prec q(z)$, or, equivalently, $f'(z)/f(z) \prec q(z)$, which evidently completes the proof of Theorem 5. \square

If $\Omega \neq \mathbb{C}$ is a simply connected domain, then $\Omega = h(\Delta)$ for some conformal mapping $h(z)$ of Δ onto Ω . In this case, the class $\Phi_\Delta[h(\Delta), q]$ is written as $\Phi_\Delta[h, q]$. The following result is an immediate consequence of Theorem 5.

Theorem 6. Let $\phi \in \Phi_\Delta[h, q]$. If $f \in \mathcal{H}[\Delta]$ satisfies

$$\begin{aligned} & \phi\left(\frac{f'(z)}{f(z)}, \frac{f''(z)}{f'(z)} - \frac{f'(z)}{f(z)}, \right. \\ & \quad \left. \frac{f(z) \left[f'''(z) f'(z) - (f''(z))^2 \right]}{f'(z) \left[f(z) f''(z) - (f'(z))^2 \right]} - \frac{f'(z)}{f(z)}; z\right) \\ & \prec h(z), \end{aligned} \quad (24)$$

then

$$\frac{f'(z)}{f(z)} \prec q(z) \quad (z \in \Delta). \quad (25)$$

Our next result is an extension of Theorem 5 to the case where the behavior of $q(z)$ on $\partial\Delta$ is not known.

Theorem 7. Let h and q be univalent in Δ with $q \in \mathcal{Q}(\Delta)$, and set $q_\rho(z) = q(\rho z)$ and $h_\rho(z) = h(\rho z)$. Let $\phi : \mathbb{C}^3 \times \Delta \rightarrow \mathbb{C}$ satisfy one of the following conditions:

- (1) $\phi \in \Phi_\Delta[h, q_\rho]$, for some $\rho \in (0, 1)$, or
- (2) there exists $\rho_0 \in (0, 1)$ such that $\phi \in \Phi_\Delta[h_\rho, q_\rho]$ for all $\rho \in (\rho_0, 1)$.

If $f \in \mathcal{H}[\Delta]$ satisfies (24), then

$$\frac{f'(z)}{f(z)} \prec q(z) \quad (z \in \Delta). \quad (26)$$

Proof. The proof of Theorem 7 is similar to that of [8, Theorem 2.3d, p.30] and so we choose to omit it. \square

The next theorem yields the best dominant of the differential subordination (24).

Theorem 8. Let h be univalent in Δ and $\phi : \mathbb{C}^3 \times \Delta \rightarrow \mathbb{C}$. Suppose that the following differential equation:

$$\phi\left(q(z), \frac{q'(z)}{q(z)}, \frac{q''(z)}{q'(z)} - \frac{q'(z)}{q(z)}; z\right) = h(z), \quad (27)$$

has a solution $q(z)$ and satisfies one of the following conditions:

- (1) $q \in \mathcal{Q}(\Delta)$ and $\phi \in \Phi_\Delta[h, q]$,
- (2) q is univalent in Δ and $\phi \in \Phi_\Delta[h, q_\rho]$, for some $\rho \in (0, 1)$, or
- (3) q is univalent in Δ and there exists $\rho_0 \in (0, 1)$ such that $\phi \in \Phi_\Delta[h_\rho, q_\rho]$ for all $\rho \in (\rho_0, 1)$.

If $f \in \mathcal{H}[\Delta]$ satisfies (24), then

$$\frac{f'(z)}{f(z)} \prec q(z), \quad (28)$$

and q is the best dominant.

Proof. Following the same arguments as in [8, Theorem 2.3e, p.31], we deduce that q is a dominant from Theorems 6 and 7. Since q satisfies (27), it is also a solution of (24) and therefore q will be dominated by all dominants. Hence, q is the best dominant. \square

In the particular case $q(z) = z$, and in view of Definition 4, the class of admissible functions $\Phi_\Delta[\Omega, q]$, denoted by $\Phi_\Delta[\Omega, z]$, is described below.

Definition 9. Let Ω be a set in \mathbb{C} . The class of admissible functions $\Phi_\Delta[\Omega, z]$ consists of those functions $\phi : \mathbb{C}^3 \times \Delta \rightarrow \mathbb{C}$ such that

$$\phi\left(\xi, \frac{k}{\xi}, \frac{L\xi - k^2}{k\xi}; z\right) \notin \Omega, \quad (29)$$

whenever $z \in \Delta$, $\text{Im}(L) = 0$, $\xi \in \mathbb{R} \setminus \{0\}$, and $k > 0$.

Corollary 10. Let $\phi \in \Phi_\Delta[\Omega, z]$. If $f \in \mathcal{H}[\Delta]$ satisfies

$$\begin{aligned} & \phi\left(\frac{f'(z)}{f(z)}, \frac{f''(z)}{f'(z)} - \frac{f'(z)}{f(z)}, \right. \\ & \quad \left. \frac{f(z) \left[f'''(z) f'(z) - (f''(z))^2 \right]}{f'(z) \left[f(z) f''(z) - (f'(z))^2 \right]} - \frac{f'(z)}{f(z)}; z\right) \\ & \in \Omega, \end{aligned} \quad (30)$$

then

$$\frac{f'(z)}{f(z)} < z \quad (z \in \Delta). \quad (31)$$

For the special case $\Omega = q(\Delta) = \{\omega : \text{Im}(\omega) > 0\}$, the class $\Phi_{\Delta}[\Omega, z]$ is simply denoted by $\Phi_{\Delta}[\Delta, z]$. Corollary 10 can now be written in the following form.

Corollary 11. Let $\phi \in \Phi_{\Delta}[\Delta, z]$. If $f \in \mathcal{H}[\Delta]$ satisfies

$$\begin{aligned} & \text{Im} \left\{ \phi \left(\frac{f'(z)}{f(z)}, \frac{f''(z)}{f'(z)} - \frac{f'(z)}{f(z)}, \right. \right. \\ & \quad \left. \left. \frac{f(z) [f'''(z) f'(z) - (f''(z))^2]}{f'(z) [f(z) f''(z) - (f'(z))^2]} - \frac{f'(z)}{f(z)}; z \right) \right\} \\ & > 0, \end{aligned} \quad (32)$$

then

$$\text{Im} \left\{ \frac{f'(z)}{f(z)} \right\} > 0 \quad (z \in \Delta). \quad (33)$$

Example 12. Let the functions $A, B : \Delta \rightarrow \mathbb{C}$ be analytic in Δ and satisfy $\text{Im } A(z) \leq 0$ and $\text{Im } B(z) \leq 0$. Then, the functions

$$\begin{aligned} \phi_1(u, v, w; z) &= u + v + A(z), \\ \phi_2(u, v, w; z) &= \alpha(u + v) + (1 - \alpha)u + B(z) \quad (0 \leq \alpha \leq 1) \end{aligned} \quad (34)$$

satisfy the admissibility condition (29) and hence Corollary 10 yields

$$\begin{aligned} & \text{Im} \left\{ \frac{f''(z)}{f'(z)} + A(z) \right\} > 0 \implies \text{Im} \left\{ \frac{f'(z)}{f(z)} \right\} > 0, \\ & \text{Im} \left\{ \alpha \frac{f''(z)}{f'(z)} + (1 - \alpha) \frac{f'(z)}{f(z)} + B(z) \right\} \\ & > 0 \implies \text{Im} \left\{ \frac{f'(z)}{f(z)} \right\} > 0. \end{aligned} \quad (35)$$

Next, we introduce the following class of admissible functions.

Definition 13. Let Ω be a set in \mathbb{C} and $q \in \mathcal{Q}(\Delta) \cap \mathcal{H}[\Delta]$. The class of admissible functions $\Phi_{\Delta,1}[\Omega, q]$ consists of those functions $\phi : \mathbb{C}^2 \times \Delta \rightarrow \mathbb{C}$ that satisfy the admissibility condition

$$\phi \left(q(\xi), q(\xi) + \frac{kq'(\xi)}{q(\xi)}; z \right) \notin \Omega, \quad (36)$$

where $z \in \Delta$, $\xi \in \partial\Delta \setminus E(q)$, and $k \geq 0$.

Theorem 14. Let $\phi \in \Phi_{\Delta,1}[\Omega, q]$. If $f \in \mathcal{H}[\Delta]$ satisfies

$$\left\{ \phi \left(\frac{f'(z)}{f(z)}, \frac{f''(z)}{f'(z)}; z \right) : z \in \Delta \right\} \subset \Omega, \quad (37)$$

then

$$\frac{f'(z)}{f(z)} < q(z) \quad (z \in \Delta). \quad (38)$$

Proof. Define the function $p(z)$ in Δ by

$$p(z) = \frac{f'(z)}{f(z)}. \quad (39)$$

A simple calculation yields

$$\frac{f''(z)}{f'(z)} = p(z) + \frac{p'(z)}{p(z)}. \quad (40)$$

Define the transformation from \mathbb{C}^2 to \mathbb{C} by

$$u(r, s) = r, \quad v(r, s) = r + \frac{s}{r}. \quad (41)$$

Let

$$\psi(r, s; z) = \phi(u, v; z) = \phi \left(r, r + \frac{s}{r}; z \right). \quad (42)$$

The proof will make use of Theorem 3. Using (39) and (40), and from (42), we get

$$\psi(p(z), p'(z); z) = \phi \left(\frac{f'(z)}{f(z)}, \frac{f''(z)}{f'(z)}; z \right). \quad (43)$$

Hence, (37) becomes

$$\psi(p(z), p'(z); z) \in \Omega. \quad (44)$$

From (42), we see that the admissibility condition for $\phi \in \Phi_{\Delta,1}[\Omega, q]$ in Definition 13 is equivalent to the admissibility condition for ψ as given in Definition 2. Hence $\psi \in \Psi_{\Delta}[\Omega, q]$, and by Theorem 3, we have $p(z) < q(z)$ or $f'(z)/f(z) < q(z)$. \square

We will denote the class $\Phi_{\Delta,1}[h(\Delta), q]$ by $\Phi_{\Delta,1}[h, q]$, where h is the conformal mapping of Δ onto $\Omega \neq \mathbb{C}$.

Theorem 15. Let $\phi \in \Phi_{\Delta,1}[h, q]$. If $f \in \mathcal{H}[\Delta]$ satisfies

$$\phi \left(\frac{f'(z)}{f(z)}, \frac{f''(z)}{f'(z)}; z \right) < h(z), \quad (45)$$

then

$$\frac{f'(z)}{f(z)} < q(z). \quad (46)$$

We extend Theorem 15 to the case in which the behavior of $q(z)$ on $\partial\Delta$ is not known.

Theorem 16. Let $\Omega \subset \mathbb{C}$ and q be univalent in Δ with $q \in \mathcal{Q}(\Delta)$. Let $\phi \in \Phi_{\Delta,1}[h, q_\rho]$, for some $\rho \in (0, 1)$, where $q_\rho(z) = q(\rho z)$. If $f \in \mathcal{H}[\Delta]$ satisfies (37), then (46) holds.

As a special case, when $q(z) = z$, we get the following corollary.

Corollary 17. Let Ω be a set in \mathbb{C} and let $\phi : \mathbb{C}^2 \times \Delta \rightarrow \mathbb{C}$ satisfy

$$\phi\left(\xi, \xi + \frac{k}{\xi}; z\right) \notin \Omega, \quad (47)$$

whenever $z \in \Delta$, $\xi \in \mathbb{R} \setminus \{0\}$, and $k \geq 0$. If $f \in \mathcal{H}[\Delta]$ satisfies

$$\phi\left(\frac{f'(z)}{f(z)}, \frac{f''(z)}{f'(z)}; z\right) \in \Omega, \quad (48)$$

then

$$\operatorname{Im} \left\{ \frac{f'(z)}{f(z)} \right\} > 0 \quad (z \in \Delta). \quad (49)$$

In the special case $\Omega = q(\Delta) = \{\omega : \operatorname{Im}(\omega) > 0\}$, Corollary 17 reduces to the following corollary.

Corollary 18. Let $\phi : \mathbb{C}^2 \times \Delta \rightarrow \mathbb{C}$ satisfy

$$\operatorname{Im} \left\{ \phi\left(\xi, \xi + \frac{k}{\xi}; z\right) \right\} \leq 0, \quad (50)$$

whenever $z \in \Delta$, $\xi \in \mathbb{R} \setminus \{0\}$, and $k \geq 0$. If $f \in \mathcal{H}[\Delta]$ satisfies

$$\operatorname{Im} \left\{ \phi\left(\frac{f'(z)}{f(z)}, \frac{f''(z)}{f'(z)}; z\right) \right\} > 0, \quad (51)$$

then

$$\operatorname{Im} \left\{ \frac{f'(z)}{f(z)} \right\} > 0 \quad (z \in \Delta). \quad (52)$$

Example 19. Let the function $C : \Delta \rightarrow \mathbb{C}$ be analytic in Δ and satisfy $\operatorname{Im} C(z) \leq 0$. Then, the function

$$\phi(u, v; z) = uv + C(z) \quad (53)$$

satisfies the admissibility condition (47) and hence Corollary 18 becomes

$$\operatorname{Im} \left\{ \frac{f''(z)}{f'(z)} + C(z) \right\} > 0 \implies \operatorname{Im} \left\{ \frac{f'(z)}{f(z)} \right\} > 0. \quad (54)$$

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The present investigation was partly supported by the Natural Science Foundation of China under Grant 11271045, the Higher School Doctoral Foundation of China under Grant 20100003110004, the Natural Science Foundation of Inner Mongolia of China under Grant 2010MS0117, and the Higher School Foundation of Inner Mongolia of China under Grant NJZY13298. The authors would like to thank Professor V. Ravichandran for his valuable suggestions and the referees for their careful reading and helpful comments to improve their paper.

References

- [1] I. A. Aleksander and V. V. Sobolev, "Extremal problems for some classes of univalent functions in the half plane," *Ukrainskii Matematicheskii Zhurnal*, vol. 22, pp. 291–307, 1970 (Russian).
- [2] V. G. Moskvina, T. N. Selakova, and V. V. Sobolev, "Extremal properties of some classes of conformal self-mapping of the half plane with fixed coefficients," *Sibirskii Matematicheskii Zhurnal*, vol. 21, no. 2, pp. 139–154, 1980 (Russian).
- [3] J. Stankiewicz, "Geometric properties of functions regular in a half-plane," in *Current Topics in Analytic Function Theory*, pp. 349–362, World Scientific, Singapore, 1992.
- [4] G. Dimkov, J. Stankiewicz, and Z. Stankiewicz, "On a class of starlike functions defined in a half-plane," *Annales Polonici Mathematici*, vol. 55, pp. 81–86, 1991.
- [5] J. Stankiewicz and Z. Stankiewicz, "On the classes of functions regular in a half-plane. I," *Bulletin of the Polish Academy of Sciences*, vol. 39, no. 1-2, pp. 49–56, 1991.
- [6] J. Stankiewicz and Z. Stankiewicz, "On the classes of functions regular in a half-plane. II," *Folia Scientiarum Universitatis Technicae Resoviensis*, vol. 60, no. 9, pp. 111–123, 1989.
- [7] D. Răducanu and N. N. Pascu, "Differential subordinations for holomorphic functions in the upper half-plane," *Mathematica*, vol. 36, no. 59, pp. 215–217, 1994.
- [8] S. S. Miller and P. T. Mocanu, *Differential Subordinations*, vol. 225 of *Monographs and Textbooks in Pure and Applied Mathematics*, Marcel Dekker, New York, NY, USA, 2000.
- [9] R. M. Ali, V. Ravichandran, and N. Seenivasagan, "Subordination and superordination on Schwarzian derivatives," *Journal of Inequalities and Applications*, vol. 2008, Article ID 712328, 18 pages, 2008.
- [10] R. M. Ali, V. Ravichandran, and N. Seenivasagan, "Subordination and superordination of the Liu-Srivastava linear operator on meromorphic functions," *Bulletin of the Malaysian Mathematical Sciences Society*, vol. 31, no. 2, pp. 193–207, 2008.
- [11] R. M. Ali, V. Ravichandran, and N. Seenivasagan, "Differential subordination and superordination of analytic functions defined by the multiplier transformation," *Mathematical Inequalities & Applications*, vol. 12, no. 1, pp. 123–139, 2009.
- [12] R. M. Ali, V. Ravichandran, and N. Seenivasagan, "Differential subordination and superordination of analytic functions defined by the Dziok-Srivastava linear operator," *Journal of the Franklin Institute*, vol. 347, no. 9, pp. 1762–1781, 2010.
- [13] R. M. Ali, V. Ravichandran, and N. Seenivasagan, "On subordination and superordination of the multiplier transformation for meromorphic functions," *Bulletin of the Malaysian Mathematical Sciences Society*, vol. 33, no. 2, pp. 311–324, 2010.

- [14] M. K. Aouf and T. M. Seoudy, "Subordination and superordination of a certain integral operator on meromorphic functions," *Computers & Mathematics with Applications*, vol. 59, no. 12, pp. 3669–3678, 2010.
- [15] N. E. Cho, O. S. Kwon, S. Owa, and H. M. Srivastava, "A class of integral operators preserving subordination and superordination for meromorphic functions," *Applied Mathematics and Computation*, vol. 193, no. 2, pp. 463–474, 2007.
- [16] N. E. Cho and H. M. Srivastava, "A class of nonlinear integral operators preserving subordination and superordination," *Integral Transforms and Special Functions*, vol. 18, no. 1-2, pp. 95–107, 2007.
- [17] T. N. Shanmugam, S. Sivasubramanian, and H. Srivastava, "Differential sandwich theorems for certain subclasses of analytic functions involving multiplier transformations," *Integral Transforms and Special Functions*, vol. 17, no. 12, pp. 889–899, 2006.
- [18] H. M. Srivastava, D.-G. Yang, and N.-E. Xu, "Subordinations for multivalent analytic functions associated with the Dziok-Srivastava operator," *Integral Transforms and Special Functions*, vol. 20, no. 7-8, pp. 581–606, 2009.

