

Research Article

Polynomial and Rational Approximations and the Link between Schröder's Processes of the First and Second Kind

François Dubeau

Département de Mathématiques, Faculté des Sciences, Université de Sherbrooke, 2500 boulevard de l'Université, Sherbrooke, QC, Canada J1K 2R1

Correspondence should be addressed to François Dubeau; francois.dubeau@usherbrooke.ca

Received 28 January 2014; Revised 18 March 2014; Accepted 24 April 2014; Published 7 May 2014

Academic Editor: Elena Berdysheva

Copyright © 2014 François Dubeau. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We show that Schröder's processes of the first kind and of the second kind to obtain a simple root of a nonlinear equation are related by polynomial and rational approximations.

1. Introduction

In [1, 2], Schröder proposed two fixed point processes to find a simple root α of a nonlinear equation $f(x) = 0$. These two processes have been reconsidered in Kalantari et al. [3] and Kalantari [4] from a modern point of view. Iteration functions (IF) of arbitrary order $p \geq 2$ associated with the two processes will be noted $E_p(x)$ for Schröder's process of the first kind and $S_p(x)$ for Schröder's process of the second kind.

Order 2 processes are

$$E_2(x) = S_2(x) = N_f(x) = x - \frac{f(x)}{f'(x)}, \quad (1)$$

where $N_f(x)$ is the Newton's IF, and order $p > 2$ processes can be expressed as

$$E_p(x) = x - \frac{f(x)}{f^{(1)}(x)} \Lambda_{p-2}(\xi) \Big|_{\xi=f(x)/f^{(1)}(x)}, \quad (2)$$

$$S_p(x) = x - \frac{f(x)}{f^{(1)}(x)} \left[\frac{\Gamma_{p-3}(\xi)}{\Gamma_{p-2}(\xi)} \right] \Big|_{\xi=f(x)/f^{(1)}(x)},$$

where $\Lambda_q(\xi)$ and $\Gamma_q(\xi)$ are polynomials of degree $q \geq 0$, such that $\Lambda_q(0) = 1 = \Gamma_q(0)$.

A question raised and discussed in [3–6] is to find and explain a possible link between the two processes. The main result of this paper is to show that $E_p(x)$ is a polynomial

approximation of $S_p(x)$, and $S_p(x)$ is a rational approximation of $E_p(x)$. More precisely, we explain the relation between the polynomials $\Lambda_{p-2}(\xi)$, $\Gamma_{p-3}(\xi)$, and $\Gamma_{p-2}(\xi)$ in (2). This paper completes the work done previously in [3–6].

Other links or comparisons could be established between these two families, for example, between their basins of attraction, their asymptotic constants, and their complexities. First results in these directions appeared in [7, 8], for example, but are not the object of the present paper.

In the next section we present notations and definitions used in this paper. Sections 3 and 4 present the two processes of Schröder and their corresponding polynomials $\Lambda_q(\xi)$ and $\Gamma_q(\xi)$. We prove the main result in the last section.

2. Preliminaries

Throughout the paper we consider real valued functions which are regular enough to be differentiated sufficiently many times. The l th derivative will be noted $f^{(l)}(x)$ for $l = 1, 2, \dots$. We will use the notation $g(x) = O(f(x))$ (and $g(x) = o(f(x))$) for two functions $f(x)$ and $g(x)$ defined around $x = \alpha$, when the following limit exists (and is finite):

$$\lim_{x \rightarrow \alpha} \frac{g(x)}{f(x)} = c \quad (3)$$

for $c \neq 0$ (for $c = 0$, resp.).

Let α be a fixed point of an IF $\Phi(x)$, and let the sequence $\{x_{k+1} = \Phi(x_k)\}_{k=0}^{+\infty}$ converge to α . Let p be a positive integer such that the following limit exists (and is finite):

$$\frac{x_{k+1} - \alpha}{(x_k - \alpha)^p} \xrightarrow{k \rightarrow +\infty} K_p(\alpha; \Phi). \tag{4}$$

We say that the convergence of the sequence to α is of (integer) order p if and only if $K_p(\alpha; \Phi) \neq 0$. We also say that $\Phi(x)$ is of order p .

Let α be a simple root of $f(x)$, which means that $f(\alpha) = 0$ and $f^{(1)}(\alpha) \neq 0$; then $g(x) = O((x - \alpha)^p)$ is equivalent to $g(x) = O(f^p(x))$ or $g(x) = O((f/f^{(1)})^p(x))$. Moreover if α is a fixed point of an IF $\Phi_p(x)$ of order p , then we can write [9]

$$\begin{aligned} \Phi_p(x) &= \alpha + O((x - \alpha)^p) = \alpha + O(f^p(x)) \\ &= \alpha + O\left(\left(\frac{f}{f^{(1)}}\right)^p(x)\right). \end{aligned} \tag{5}$$

3. Schröder’s Process of the First Kind

3.1. The Process. Schröder’s process of the first kind, proposed to increase the order of convergence of a fixed-point method [1–3, 9, 10], can be obtained by considering Taylor’s expansion of the inverse $g(y)$ of $f(x)$ around $y = 0$ [9]. It is also associated with Chebyshev and Euler [11–13]. The IF $E_p(x)$ of order p is defined by the series

$$E_p(x) = x + \sum_{l=1}^{p-1} c_l(x) f^l(x) = \sum_{l=0}^{p-1} c_l(x) f^l(x), \tag{6}$$

where $c_0(x) = x$ and

$$\begin{aligned} c_l(x) &= -\frac{1}{l} \left(\frac{1}{f^{(1)}(x)} \frac{d}{dx} \right) c_{l-1}(x) \\ &= \frac{(-1)^l}{l!} \left(\frac{1}{f^{(1)}(x)} \frac{d}{dx} \right)^l c_0(x) \end{aligned} \tag{7}$$

for $l = 1, 2, \dots$

Relation (7) implies that $c_l(x)$ is a rational function of the form

$$\begin{aligned} c_l(x) &= \left(\text{polynomial of degree } l - 1 \text{ w.r.t.} \right. \\ &\quad \left. f^{(1)}(x), f^{(2)}(x), \dots, f^{(l)}(x) \right) \\ &\quad \times \left([f^{(1)}(x)]^{2l-1} \right)^{-1}. \end{aligned} \tag{8}$$

Then we can write

$$E_p(x) = x - \frac{f(x)}{f^{(1)}(x)} \Lambda_{p-2}(\xi) \Big|_{\xi=f(x)/f^{(1)}(x)}, \tag{9}$$

where

$$\Lambda_{p-2}(\xi) = \sum_{l=0}^{p-2} \lambda_l(x) \xi^l, \tag{10}$$

$$\lambda_l(x) = \begin{cases} 1 & \text{for } l = 0, \\ -\frac{[f^{(1)}(x)]^l}{l+1} \frac{d}{dx} \left(\frac{\lambda_{l-1}(x)}{[f^{(1)}(x)]^l} \right) & \text{for } l \geq 1. \end{cases} \tag{11}$$

Consequently

$$\begin{aligned} \lambda_l(x) &= \left(\left\{ \text{polynomial of degree } l \text{ w.r.t.} \right. \right. \\ &\quad \left. \left. f^{(1)}(x), f^{(2)}(x), \dots, f^{(l+1)}(x) \right\} \right. \\ &\quad \left. \times \left([f^{(1)}(x)]^l \right)^{-1} \right). \end{aligned} \tag{12}$$

3.2. Examples. The first 4 IFs are given here. For

(a) $p = 2$:

$$E_2(x) = x - \frac{f(x)}{f^{(1)}(x)}, \tag{13}$$

which corresponds to Newton’s IF of order 2;

(b) $p = 3$:

$$E_3(x) = x - \frac{f(x)}{f^{(1)}(x)} - \frac{1}{2!} \frac{f^{(2)}(x)}{f^{(1)}(x)} \left[\frac{f(x)}{f^{(1)}(x)} \right]^2, \tag{14}$$

which corresponds to Chebyshev’s IF order 3 [11];

(c) $p = 4$:

$$\begin{aligned} E_4(x) &= x - \frac{f(x)}{f^{(1)}(x)} - \frac{1}{2!} \frac{f^{(2)}(x)}{f^{(1)}(x)} \left[\frac{f(x)}{f^{(1)}(x)} \right]^2 \\ &\quad - \frac{1}{3!} \left[\frac{3[f^{(2)}(x)]^2 - f^{(1)}(x) f^{(3)}(x)}{[f^{(1)}(x)]^2} \right] \\ &\quad \times \left[\frac{f(x)}{f^{(1)}(x)} \right]^3; \end{aligned} \tag{15}$$

(d) $p = 5$:

$$\begin{aligned}
 E_5(x) = & x - \frac{1}{f^{(1)}(x)} f(x) - \frac{1}{2!} \frac{f^{(2)}(x)}{f^{(1)}(x)} \left[\frac{f(x)}{f^{(1)}(x)} \right]^2 \\
 & - \frac{1}{3!} \left[\frac{3[f^{(2)}(x)]^2 - f^{(1)}(x) f^{(3)}(x)}{[f^{(1)}(x)]^2} \right] \\
 & \times \left[\frac{f(x)}{f^{(1)}(x)} \right]^3 \\
 & - \frac{1}{4!} \left[\frac{[f^{(1)}(x)]^2 f^{(4)}(x)}{[f^{(1)}(x)]^3} + \frac{15[f^{(2)}(x)]^3}{[f^{(1)}(x)]^3} \right. \\
 & \quad \left. - \frac{10 f^{(1)}(x) f^{(2)}(x) f^{(3)}(x)}{[f^{(1)}(x)]^3} \right] \\
 & \times \left[\frac{f(x)}{f^{(1)}(x)} \right]^4.
 \end{aligned} \tag{16}$$

4. Schröder's Process of the Second Kind

4.1. *The Process.* Different equivalent formulations exist for Schröder's process of the second kind [4, 6, 13, 14]. One such form is based on a determinantal identity. Let $\Delta_0(x) = 1$ and for $p \geq 1$

$$\Delta_p(x) = \begin{vmatrix} f^{(1)}(x) & \frac{f^{(2)}(x)}{2!} & \dots & \dots & \dots & \frac{f^{(p)}(x)}{p!} \\ f(x) & f^{(1)}(x) & \frac{f^{(2)}(x)}{2!} & \dots & \dots & \frac{f^{(p-1)}(x)}{(p-1)!} \\ 0 & f(x) & f^{(1)}(x) & \frac{f^{(2)}(x)}{2!} & \dots & \frac{f^{(p-2)}(x)}{(p-2)!} \\ \vdots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & f(x) & f^{(1)}(x) & \frac{f^{(2)}(x)}{2!} \\ 0 & \dots & \dots & 0 & f(x) & \frac{f^{(1)}(x)}{1!} \end{vmatrix}. \tag{17}$$

Expanding this determinant along the first line, we obtain

$$\Delta_p(x) = \sum_{j=1}^p (-1)^{j+1} \frac{f^{(j)}(x)}{j!} f^{j-1}(x) \Delta_{p-j}(x) \tag{18}$$

for $p \geq 1$. Using $R_p(x) = \Delta_p(x)/f^{p+1}(x)$, Schröder's process of the second kind [1, 2] of order p is defined by

$$S_p(x) = x - \frac{R_{p-2}(x)}{R_{p-1}(x)} = x - f(x) \frac{\Delta_{p-2}(x)}{\Delta_{p-1}(x)}. \tag{19}$$

We prove by mathematical induction that

$$\Delta_p(x) = \frac{(-1)^p f^{p+1}(x)}{p!} \left(\frac{1}{f(x)} \right)^{(p)}, \tag{20}$$

$$f(x) \Delta_{p-1}^{(1)}(x) = p f^{(1)}(x) \Delta_{p-1}(x) - p \Delta_p(x).$$

Using

$$\left(\frac{1}{f(x)} \right)^{(p)} = \sum_{l=2}^{p+1} d_{p,l}(x) \left(\frac{1}{f(x)} \right)^l, \tag{21}$$

then we have $d_{1,2}(x) = -f^{(1)}(x)$, and we can obtain recursively $d_{p,l}(x)$ for $p \geq 2$ by the formulae

$$d_{p,l}(x) = \begin{cases} d_{p-1,l}^{(1)}(x) & \text{for } l = 2, \\ d_{p-1,l}^{(1)}(x) - (l-1) d_{p-1,l-1}(x) f^{(1)}(x) & \text{for } l = 3, \dots, p, \\ -p d_{p-1,p}(x) f^{(1)}(x) & \text{for } l = p+1. \end{cases} \tag{22}$$

Consequently

$$d_{p,l}(x) = \begin{cases} -f^{(p)}(x), & \text{for } l = 2, \\ \{\text{polynomial of degree } l-1 \text{ w.r.t. } f^{(1)}(x)\}, & \text{for } l = 3, \dots, p, \\ f^{(2)}(x), \dots, f^{(p+2-l)}(x) & \text{for } l = 3, \dots, p, \\ (-1)^p p! [f^{(1)}(x)]^p & \text{for } l = p+1. \end{cases} \tag{23}$$

It follows that $\Delta_p(x)$ is a polynomial of degree $p-1$ with respect to $f(x)$ and the term not depending on $f(x)$ is $[f^{(1)}(x)]^p$. Hence

$$\Delta_p(x) = [f^{(1)}(x)]^p \Gamma_{p-1}(\xi) \Big|_{\xi=f(x)/f^{(1)}(x)}, \tag{24}$$

where $\Gamma_{p-1}(\xi)$ is a polynomial of degree $p-1$ such that $\Gamma_{p-1}(0) = 1$. Let us set

$$\Gamma_{p-1}(\xi) = \sum_{l=0}^{p-1} \gamma_{p-1,l}(x) \xi^l, \tag{25}$$

where the coefficients $\gamma_{p-1,l}(x)$ are rational functions of the form

$$\gamma_{p-1,l}(x) = (-1)^p \frac{d_{p,p+1-l}(x)}{p! [f^{(1)}(x)]^{p-l}} \tag{26}$$

$$\begin{aligned}
 &= \left(\{\text{polynomial of degree } p-l \text{ w.r.t. } f^{(1)}(x), f^{(2)}(x), \dots, f^{(l+1)}(x)\} \right. \\
 & \quad \left. \times \left([f^{(1)}(x)]^{p-l} \right)^{-1} \right). \tag{27}
 \end{aligned}$$

Then we obtain

$$S_p(x) = x - \frac{f(x)}{f^{(1)}(x)} \left[\frac{\Gamma_{p-3}(\xi)}{\Gamma_{p-2}(\xi)} \right] \Big|_{\xi=f(x)/f^{(1)}(x)}. \tag{28}$$

4.2. *Examples.* The first 4 IFs are presented here. For

(a) $p = 2$:

$$S_2(x) = x - \frac{f(x)}{f^{(1)}(x)}, \tag{29}$$

which corresponds to Newton's IF of order 2;

(b) $p = 3$:

$$S_3(x) = x - \frac{f(x)}{f^{(1)}(x)} \times \left[\frac{1}{1 - (1/2) \frac{f^{(2)}(x)}{f^{(1)}(x)} \frac{f(x)}{f^{(1)}(x)}} \right], \tag{30}$$

which is Halley's IF of order 3 [15];

(c) $p = 4$:

$$S_4(x) = x - \frac{f(x)}{f^{(1)}(x)} \times \left[\left(1 - \frac{1}{2} \frac{f^{(2)}(x)}{f^{(1)}(x)} \frac{f(x)}{f^{(1)}(x)} \right) \times \left(1 - \frac{f^{(2)}(x)}{f^{(1)}(x)} \frac{f(x)}{f^{(1)}(x)} + \frac{1}{3!} \frac{f^{(3)}(x)}{f^{(1)}(x)} \left[\frac{f(x)}{f^{(1)}(x)} \right]^2 \right)^{-1} \right]; \tag{31}$$

(d) $p = 5$:

$$S_5(x) = x - \frac{f(x)}{f^{(1)}(x)} \times \left[\left(1 - \frac{f^{(2)}(x)}{f^{(1)}(x)} \frac{f(x)}{f^{(1)}(x)} + \frac{1}{3!} \frac{f^{(3)}(x)}{f^{(1)}(x)} \left[\frac{f(x)}{f^{(1)}(x)} \right]^2 \right) \times \left(1 - \frac{3}{2} \frac{f^{(2)}(x)}{f^{(1)}(x)} \frac{f(x)}{f^{(1)}(x)} + \left[\frac{1}{3} \frac{f^{(3)}(x)}{f^{(1)}(x)} + \frac{1}{4} \left[\frac{f^{(2)}(x)}{f^{(1)}(x)} \right]^2 \right) \left[\frac{f(x)}{f^{(1)}(x)} \right]^2 - \frac{1}{4!} \frac{f^{(4)}(x)}{f^{(1)}(x)} \left[\frac{f(x)}{f^{(1)}(x)} \right]^3 \right)^{-1} \right]. \tag{32}$$

5. Proof of the Main Result

Since both processes $E_p(x)$ and $S_p(x)$ are of order p , following [9], the next result holds.

Lemma 1. *Let α be a simple root of $f(x)$; then*

$$E_p(x) - S_p(x) = O(f^p(x)) = O\left(\left(\frac{f(x)}{f^{(1)}(x)}\right)^p\right). \tag{33}$$

From this lemma

$$\frac{f(x)}{f^{(1)}(x)} \left[\Lambda_{p-2}(\xi) - \frac{\Gamma_{p-3}(\xi)}{\Gamma_{p-2}(\xi)} \right] \Big|_{\xi=f(x)/f^{(1)}(x)} = O\left(\left(\frac{f(x)}{f^{(1)}(x)}\right)^p\right), \tag{34}$$

and so

$$\left[\Lambda_{p-2}(\xi) - \frac{\Gamma_{p-3}(\xi)}{\Gamma_{p-2}(\xi)} \right] \Big|_{\xi=f(x)/f^{(1)}(x)} = O\left(\left(\frac{f(x)}{f^{(1)}(x)}\right)^{p-1}\right). \tag{35}$$

The next step is to consider the following basic result about polynomial and rational approximations.

Lemma 2. *Let us consider the expression*

$$\Lambda_p(\xi) = \frac{\Gamma_{p-1}(\xi)}{\Gamma_p(\xi)} + O(\xi^{p+1}), \tag{36}$$

where $\Lambda_p(\xi)$ and $\Gamma_p(\xi)$ are polynomials of degree p such that $\Lambda_p(0) = 1 = \Gamma_p(0)$.

- (a) *If $\Gamma_{p-1}(\xi)$ and $\Gamma_p(\xi)$ are given, there exists one and only one polynomial $\Lambda_p(\xi)$ such that (36) holds.*
- (b) *If $\Gamma_{p-1}(\xi)$ and $\Lambda_p(\xi)$ are given, there exists one and only one polynomial $\Gamma_p(\xi)$ such that (36) holds.*

Proof. This result is based on the following identity:

$$\frac{\Gamma_{p-1}(\xi)}{\Gamma_p(\xi)} = \frac{\Gamma_{p-1}(\xi)}{1 - (1 - \Gamma_p(\xi))} = \Gamma_{p-1}(\xi) \sum_{l=0}^{+\infty} (1 - \Gamma_p(\xi))^l = \Gamma_{p-1}(\xi) \sum_{l=0}^p (1 - \Gamma_p(\xi))^l + O(\xi^{p+1}), \tag{37}$$

and we would like to have

$$\Lambda_p(\xi) = \Gamma_{p-1}(\xi) \sum_{l=0}^p (1 - \Gamma_p(\xi))^l + O(\xi^{p+1}). \tag{38}$$

We know that $\lambda_0 = \gamma_{p-1,0} = \gamma_{p,0} = 1$. Moreover the coefficient of ξ^l on the left-hand side is λ_l and on the right-hand side is

$\gamma_{p,l}$

$$+ \text{an expression in terms of } \begin{cases} \gamma_{p-1,j} & \text{for } j = 0, \dots, l, \\ \gamma_{p,j} & \text{for } j = 0, \dots, l-1 \end{cases} \tag{39}$$

for $l = 1, \dots, p$. This expression shows that if the γ_p 's and the γ_{p-1} 's are given, we can obtain λ_p 's, and conversely if the λ_p 's and the γ_{p-1} 's are given, we can obtain the γ_p 's. \square

In view of these two lemmas we obtain the main result of this paper.

Theorem 3. $E_p(x)$ and $S_p(x)$ are related as follows.

- (a) For $S_p(x)$ given by (28), one can obtain the form (9) of $E_p(x)$ by expanding the denominator in (28), multiplying, and truncating to keep powers of $f(x)/f^{(1)}(x)$ up to $p - 1$.
- (b) Since $S_2(x) = E_2(x) = N_f(x)$, one can obtain recursively $S_p(x)$ given by (28) from $E_p(x)$ given by (9).

Proof. (a) Indeed, if $S_p(x)$ is given, which means we know $\Gamma_{p-3}(\xi)$ and $\Gamma_{p-2}(\xi)$, we can write

$$\begin{aligned}
 S_p(x) &= x - \frac{f(x)}{f^{(1)}(x)} \left[\frac{\Gamma_{p-3}(\xi)}{\Gamma_{p-2}(\xi)} \right] \Bigg|_{\xi=f(x)/f^{(1)}(x)} \\
 &= x - \frac{f(x)}{f^{(1)}(x)} \left[\frac{\Gamma_{p-3}(\xi)}{1 - (1 - \Gamma_{p-2}(\xi))} \right] \Bigg|_{\xi=f(x)/f^{(1)}(x)} \\
 &= x - \frac{f(x)}{f^{(1)}(x)} \left[\Gamma_{p-3}(\xi) \sum_{l=0}^{p-2} (1 - \Gamma_{p-2}(\xi))^l + O(\xi^{p-1}) \right] \Bigg|_{\xi=f(x)/f^{(1)}(x)} \\
 &= x - \frac{f(x)}{f^{(1)}(x)} \Lambda_{p-2}(\xi) \Big|_{\xi=f(x)/f^{(1)}(x)} + O(f^p(x)) \\
 &= E_p(x) + O(f^p(x)),
 \end{aligned} \tag{40}$$

which follows from part (a) of Lemma 1.

(b) As already observed, $E_2(x) = S_2(x) = N_f(x)$. If, for $p > 2$, we have $S_{p-1}(x)$, and we know $\Gamma_{p-4}(\xi)$ and $\Gamma_{p-3}(\xi)$, and $E_p(x)$, and we know also $\Lambda_{p-2}(\xi)$, then we can determine $\Gamma_{p-2}(\xi)$ from part (b) of Lemma 1. Consequently we obtain $S_p(x)$. \square

The computation of the polynomials $\Lambda_p(\xi)$ and $\Gamma_p(\xi)$, and their coefficients λ_p 's and the γ_p 's, can be done explicitly using (11) for the λ_p 's and (22) and (26) for the γ_p 's. The verification of the link between the two Schröder's processes has already been done using symbolic computation up to order 20 [5, 6].

Conflict of Interests

The author declares that there is no conflict of interests regarding the publication of this paper.

Acknowledgment

This work has been financially supported by an individual discovery grant from NSERC (Natural Sciences and Engineering Research Council of Canada).

References

- [1] E. Schröder, "Ueber unendlich viele Algorithmen zur Auflösung der Gleichungen," *Mathematische Annalen*, vol. 2, no. 2, pp. 317–365, 1870.
- [2] E. Schröder, "On infinitely many algorithms for solving equations," Tech. Rep. TR-92-121, TR-2990, Institute for advanced Computer Studies, University of Maryland, 1992, Translated by G. W. Stewart.
- [3] B. Kalantari, I. Kalantari, and R. Zaare-Nahandi, "A basic family of iteration functions for polynomial root finding and its characterizations," *Journal of Computational and Applied Mathematics*, vol. 80, no. 2, pp. 209–226, 1997.
- [4] B. Kalantari, *Polynomial Root-Finding and Polynomiography*, World Scientific, Singapore, 2009.
- [5] L. Petković and M. Petković, "The link between Schröder's iteration methods of the first and second kind," *Novi Sad Journal of Mathematics*, vol. 38, no. 3, pp. 55–63, 2008.
- [6] M. S. Petković, L. D. Petković, and D. Herceg, "On Schröder's families of root-finding methods," *Journal of Computational and Applied Mathematics*, vol. 233, no. 8, pp. 1755–1762, 2010.
- [7] B. Kalantari, "Polynomial root-finding methods whose basins of attraction approximate Voronoi diagram," *Discrete & Computational Geometry*, vol. 46, no. 1, pp. 187–203, 2011.
- [8] F. Dubeau, "On comparisons of Chebyshev-Halley iteration functions based on their asymptotic constants," *International Journal of Pure and Applied Mathematics*, vol. 58, pp. 965–981, 2013.
- [9] J. F. Traub, *Iterative Methods for the Solution of Equations*, Prentice Hall, Englewood Cliffs, NJ, USA, 1964.
- [10] A. S. Householder, *Principles of Numerical Analysis*, McGraw-Hill, Columbus, NY, USA, 1953.
- [11] E. Bodewig, "On types of convergence and on the behavior of approximations in the neighborhood of a multiple root of an equation," *Quarterly of Applied Mathematics*, vol. 7, pp. 325–333, 1949.
- [12] M. Shub and S. Smale, "Computational complexity. On the geometry of polynomials and a theory of cost. I," *Annales Scientifiques de l'École Normale Supérieure*, vol. 18, no. 1, pp. 107–142, 1985.
- [13] M. Petković and D. Herceg, "On rediscovered iteration methods for solving equations," *Journal of Computational and Applied Mathematics*, vol. 107, no. 2, pp. 275–284, 1999.
- [14] B. Kalantari and J. Gerlach, "Newton's method and generation of a determinantal family of iteration functions," *Journal of Computational and Applied Mathematics*, vol. 116, no. 1, pp. 195–200, 2000.
- [15] W. Gander, "On Halley's iteration method," *The American Mathematical Monthly*, vol. 92, no. 2, pp. 131–134, 1985.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

